

# A NOTE ON COMPLEX $\mathcal{P}_1$ SPACES

BY

OTTE HUSTAD

## ABSTRACT

For a family of closed balls in a normed space we define the concept of weak intersection property, and we show that a complex Banach space is a  $\mathcal{P}_1$  space if and only if every family of closed balls with the weak intersection property has a non-empty intersection.

According to established terminology, a Banach space  $A$  is a  $\mathcal{P}_1$  space if for every pair of Banach spaces  $X$  and  $Y$  with  $Y \subset X$  and for every bounded linear operator  $T: Y \rightarrow A$  there exists a norm preserving extension  $\tilde{T}: X \rightarrow A$ . Nachbin shows in [3] that if  $A$  is a real Banach space, then  $A$  is a  $\mathcal{P}_1$  space if and only if every family of closed balls in  $A$  with the binary intersection property (that is, any two balls of the family have a common element) has a non-empty intersection. In a recent paper [1] we have shown that in the case of complex Banach spaces the following intersection property corresponds in several ways to the binary intersection property. A family  $\{B(a_j, r_j)\}_{j \in J}$  of closed balls in a normed space  $A$  has the *weak intersection property* if for any linear functional  $\phi \in A^*$  with norm  $\leq 1$  the family  $\{B(\phi(a_j), r_j)\}_{j \in J}$  of balls in  $\mathbb{C}$  (or in  $\mathbb{R}$ ) has a non-empty intersection. It is easy to see [1, Cor. 1.10] that in the case of real normed spaces the weak intersection property is equivalent with the binary intersection property. Hence the real version of the following theorem is exactly the theorem of Nachbin [3] mentioned above.

**THEOREM 1.** *Let  $A$  be a complex Banach space. Then  $A$  is a  $\mathcal{P}_1$  space if and only if any family of closed balls in  $A$  with the weak intersection property has a non-empty intersection.*

---

Received May 23, 1973

The proof of this theorem will be given after we have proved the following three lemmata. In what follows

$$H^3 = \left\{ (z_j) \in \mathbb{C}^3 : \sum_{j=1}^3 z_j = 0 \right\}.$$

LEMMA 2. *A family  $\mathcal{F}$  of closed balls in a complex normed space  $A$  has the weak intersection property if and only if for any subfamily of three balls  $\{B(a_j, r_j)\}_{j=1}^3 \subset \mathcal{F}$  it is true that*

$$(1) \quad \left\| \sum_{j=1}^3 z_j a_j \right\| \leq \sum_{j=1}^3 |z_j| r_j, \quad (z_j) \in H^3.$$

PROOF. This follows easily from [1, Cor. 1.3, 1.4] together with the Helly theorem on intersection of compact convex sets in  $\mathbb{C}$ .

LEMMA 3. *A family  $\mathcal{F} = \{B(a_j, r_j)\}_{j \in J}$  of closed balls in a normed space  $A$  has the weak intersection property if and only if there exists a Banach space  $X \supset A$  and an  $x \in X$  such that*

$$(2) \quad \|x - a_j\| \leq r_j, \quad j \in J.$$

PROOF. Assume that  $X \supset A$  is a normed space and that  $x \in X$  satisfies (2). Let  $\phi \in A^*$  with  $\|\phi\| \leq 1$  be given. Choose  $\tilde{\phi} \in X^*$  as a norm preserving extension of  $\phi$ . It then follows from (2) that

$$|\tilde{\phi}(x) - \phi(a_j)| = |\tilde{\phi}(x - a_j)| \leq \|x - a_j\| \leq r_j, \quad j \in J.$$

Hence  $\mathcal{F}$  has the weak intersection property. Assume conversely that  $\mathcal{F}$  has this property. It is well known that there exists a set  $\Gamma$  such that  $A$  can be considered as a subspace of  $l_\infty(\Gamma)$ . Let  $\gamma \in \Gamma$ . Since the evaluation map  $x \mapsto x(\gamma)$  is a linear functional on  $l_\infty(\Gamma)$  with norm  $\leq 1$ , it follows by hypothesis that there exists a complex number  $x_\gamma$  such that

$$(3) \quad |x_\gamma - a_j(\gamma)| \leq r_j, \quad j \in J.$$

Define  $x \in l_\infty(\Gamma)$  by  $x(\gamma) = x_\gamma$ ,  $\gamma \in \Gamma$ . We then obtain from (3) that

$$\|x - a_j\| = \sup_{\gamma \in \Gamma} \{|x_\gamma - a_j(\gamma)|\} \leq r_j, \quad j \in J.$$

LEMMA 4. *Let  $X$ ,  $Y$  and  $A$  be normed spaces with  $Y \subset X$ . Let  $T: Y \rightarrow A$  be a bounded linear operator and let  $x \in X \setminus Y$ . Then the family  $\{B(Ty, \|T\| \cdot \|x - y\|) : y \in Y\}$  has the weak intersection property.*

PROOF. Let  $y_1, y_2, y_3 \in Y$  and let  $(z_j) \in H^3$ . Then

$$\left\| \sum_{j=1}^3 z_j T y_j \right\| \leq \|T\| \cdot \left\| \sum_{j=1}^3 z_j (y_j - x) \right\| \leq \sum_{j=1}^3 |z_j| \|T\| \cdot \|x - y_j\|.$$

The desired conclusion now follows from Lemma 2.

PROOF OF THEOREM 1. Assume that  $A$  is a  $\mathcal{P}_1$  space. Let  $\mathcal{F} = \{B(a_j, r_j)\}_{j \in J}$  be a family of balls in  $A$  with the weak intersection property. Choose, according to Lemma 3, the Banach space  $X \supset A$  such that (2) is fulfilled. Since  $A$  is a  $\mathcal{P}_1$  space, there exists a projection  $P$  from  $X$  onto  $A$  such that  $\|P\| = 1$ . It follows that  $Px \in A$  and that

$$\|Px - a_j\| = \|P(x - a_j)\| \leq \|x - a_j\| \leq r_j, \quad j \in J.$$

Hence  $\mathcal{F}$  has a non-empty intersection. Assume conversely that any family of balls in  $A$  with the weak intersection property has a non-empty intersection. Let  $X$  and  $Y$  be Banach spaces with  $Y \subset X$  and let  $T: Y \rightarrow A$  be a bounded linear operator. In order to show that  $T$  admits a norm-preserving extension  $\tilde{T}: X \rightarrow A$ , it is sufficient, by a standard application of the Zorn's lemma, to assume that  $\dim X/Y = 1$ . Let  $x \in X \setminus Y$ . By a basic lemma of Nachbin (see [2, Lem. 5.2]), the operator  $T$  admits a norm-preserving extension  $\tilde{T}: X \rightarrow A$  if and only if

$$(4) \quad \bigcap_{y \in Y} B(Ty, \|T\| \cdot \|x - y\|) \neq \emptyset.$$

But it follows from Lemma 4 and from our assumption that (4) is fulfilled.

#### REFERENCES

1. O. Hustad, *Intersection properties of balls in complex Banach spaces whose duals are  $L_1$  spaces*, Preprint Series, Inst. Math., University of Oslo, No. 9, (1973).
2. J. Lindenstrauss, *Extension of compact operators*, Mem. Amer. Math. Soc. **48** (1964).
3. L. Nachbin, *A theorem of the Hahn-Banach type for linear transformations*, Trans. Amer. Math. Soc. **68** (1950), 28–46.

UNIVERSITY OF OSLO  
OSLO, NORWAY