

comparative data show that the hydrogen atoms which are split off in the initial reaction step from the  $\text{CH}_3$  group and from the phenyl ring have different reactivities. Basically, the hydrogen atoms from phenyl groups enter into addition reactions, forming radicals of cyclohexadienyl-type; whereas those from the  $\text{CH}_3$  group—due to their greater kinetic energy content—are primarily consumed in the secondary reactions of atomic hydrogen split-off from toluene molecules.

#### Kinetics of Heterogeneous Catalytic Reactions in Chromatographic Pulsed Flow Units at Conditions of Ideal Linear Chromatography

By G. A. GAZEYEV, V. YU. FELEENOVSKII,  
AND M. I. YANOVSKII

*Institute of Chemical Physics of the  
Academy of Sciences of USSR*

The authors analyze from a theoretical standpoint the kinetics of heterogeneous catalytic reactions of different orders. The reactions are assumed to be carried out in pulsed flow units at the conditions of ideal linear chromatography. The equations derived to evaluate the rate constants and activation energies of the first order reactions are used in processing the experimental data on dehydrogenation of cyclohexane to benzene over a number of catalysts. The effect of various types of pulsed feed on conversion is also shown.

#### Kinetics of Liquid Phase Hydrochlorination of Acetylene in Presence of Cuprous Chloride

By R. M. FLEED, N. F. ALEKSEIYEVA,  
T. G. KHMELEVSKAYA, AND  
N. A. GAIDAI

*M. V. Lomonosov Institute of Fine  
Chemical Technology in Moscow*

The kinetics of hydrochlorination of acetylene in liquid phase at  $40^\circ\text{--}98^\circ$  was studied, using  $\text{CuCl--MeCl--HCl--H}_2\text{O}$  and  $\text{CuCl--HCl--H}_2\text{O}$  as the contact solvents. The reaction rate constants bear linear relationship to activity of the catalyzing ions. The conditions to select contact solvents of optimum composition are described.

#### Chemiluminescence in Slow Chemical Reactions: Investigation of Kinetics of Accumulation of Hydroperoxide in Catalytic Oxidation of Ethyl Benzene by a Chemiluminescence Technique

By I. V. ZAKHAROV AND  
V. YA. SHLIAPEENTOKH

*Moscow Physico-Technical Institute,  
Institute of Chemical Physics of the  
Academy of Sciences of USSR*

This study of the chemiluminescence-control-

ling factors during initial stages of oxidation of ethyl benzene in a catalytic solution of cobalt acetate in glacial acetic acid shows that the change in chemiluminescence intensity with reaction time is directly tied to the kinetics of hydroperoxide accumulation. A chemiluminescence technique is proposed to determine the hydroperoxide decomposition rate constants in the course of the oxidation process. The rates of oxidation and concentrations of hydroperoxide at steady state conditions were also determined by use of this technique.

#### Mechanism and Kinetic Principles of Dimerization of 1,1-Diphenylethylene in Aqueous Sulfuric Acid

By S. G. ENTELEES AND K. S. KAZANSKII

*Institute of Chemical Physics of the  
Academy of Sciences of USSR*

A spectrographic study of kinetics of dimerization of 1,1-diphenylethylene in aqueous (64–68%) sulfuric acid was carried out at temperatures of  $5^\circ$ ,  $20^\circ$  and  $40^\circ$ . In terms of decreasing carbonium ion concentration, the rate of the second order reaction decreases with increasing concentration of the acid. At the conditions studied, the reaction temperature has practically no effect upon the rate.

To explain observed regularities of the process, production of the dimer is assumed to occur by two parallel routes: (1) interaction of 1,1-diphenylethylene with carbonium ions and (2) interaction of carbonium ions with diphenylmethylcarbinol, which is produced from the olefin in presence of aqueous sulfuric acid. The values of activation energies and reaction rate constants for the two processes are listed.

The conclusions presented satisfactorily agree with the earlier-published (1) principles of ionization of 1,1-diphenylethylene.

#### Liquid Phase Dehydrogenation of Isopropyl Alcohol

By S. L. KEEPERMAN, N. V. NEEKOLAYEVA  
AND I. P. DAVIDOVA

*N. D. Zelenskii Institute of Organic  
Chemistry of the Academy of Sciences  
of USSR*

A study of the mechanism and kinetics of liquid phase dehydrogenation of isopropyl alcohol was carried out over a skeletal nickel catalyst. The reaction which in the initial stage is of zero order, approaches a near-zero order at the subsequent steady state conditions—although at a much slower rate.

The data on reaction kinetics, catalytic ac-