

ently, belongs to the chain-radical class of reactions.

### High-Dosage Radiolysis of Neutral Solutions of Nitrate Salts

By A. K. PEEKAYEV, P. YA. GLAZONOV,  
AND A. A. YAKOOBOVECH

*Institute of Physical Chemistry of the  
Academy of Sciences of USSR*

Radiolysis of neutral aqueous solutions of  $\text{NaNO}_3$  was investigated at various solution concentrations and x-ray dosage rates. The  $\text{NO}_2^-$  ion yields were determined by irradiation of solutions containing nitrite and hydrogen peroxide. With dilute- and medium-strength solutions, increasing the x-ray dosage rates also increases the values of  $G(\text{NO}_2^-)$  and  $G(\text{H}_2\text{O}_2)$ . For 1M solutions of  $\text{NaNO}_3$ , the values of  $G(\text{NO}_2^-)$  are independent of the dosage rates. A mechanism is proposed to explain the radiolytic reduction of the nitrate ions. The distinguishing feature of the postulated mechanism is competition between the reverse oxidation of the  $\text{NO}_2^-$  ions by the OH radicals and the recombination of the radicals.

In irradiation of relatively dilute  $\text{NaNO}_3$  solutions by powerful pulsed beams, recombination of the hydrogen atoms, originating from various tracks, plays an important role.

### Determination of Absolute Rate Constants of Free Radical Reactions: Addition of Trichloro-Bromo-Methane to Cyclohexane and to Heptene-1

By H. S. BAGDASAR'YAN AND  
A. F. REVZEEN

*L. Ya. Karpov Physico-Chemical Institute*

A new method is proposed to investigate the non-stationary kinetics of chain reactions. This method involves measuring the non-adiabatic temperature rise following complete or partial cessation of the photochemical initiation. Using this method, the following rate constants (as  $1/\text{mol} \times \text{sec}$ ) were obtained:  $\text{CCl}_3^{\cdot} + \text{cyclohexane}$ , 773 (at 25°);  $\text{CCl}_3^{\cdot} + \text{Heptene-1}$ ,  $2.30 \times 10^9 \exp(-7000/RT)$ ;  $\text{C}_7\text{H}_{14}\text{CCl}_3^{\cdot} + \text{CCl}_3\text{Br}$ ,  $2.48 \times 10^9 \exp(-8500/RT)$ ;  $\text{CCl}_3^{\cdot} + \text{CCl}_3^{\cdot}$ ,  $1 \times 10^8$ ;  $\text{C}_7\text{H}_{14}\text{CCl}_3^{\cdot} + \text{C}_7\text{H}_{14}\text{CCl}_3^{\cdot}$ ,  $1.0 \times 10^6$ . A comparison with published data shows that the constants of addition of  $\text{CH}_3^{\cdot}$  and  $\text{CCl}_3^{\cdot}$  radicals to cyclohexene, heptene, vinyl acetate, anthracene, and styrene increase in the order mentioned. The absolute values of the constants for these radicals differ but little.

### Synthetic Zeolites as Ion Exchangers: Study of Kinetics of Ionic Exchange

By G. M. PANCHENKOV AND  
A. M. TOLMACHOV

*Department of Chemistry of  
M. V. Lomonosov State Uni-  
versity in Moscow*

The results of a study of kinetics of ionic exchange processes are summarized for various systems of synthetic zeolites and solutions of  $\text{LiCl}$ ,  $\text{NH}_4\text{Cl}$ ,  $\text{NH}_4\text{NO}_3$ ,  $\text{CaCl}_2$ ,  $\text{MgCl}_2$ , and  $\text{PC}(\text{NO}_3)_2$ . For the solution concentrations of 0.1 to 1.0N, external diffusion processes control the ion exchange rates. Agglomeration of the original particles into larger aggregates does not substantially decrease the exchange rates.

### Effect of Reaction Temperatures on the Rates of Addition of Atomic Hydrogen to Some Solid Unsaturated Hydrocarbons

By A. N. PONAMAR'YOV

*Institute of Chemical Physics of the  
Academy of Sciences of USSR*

A method is described to measure the rates of consumption of atomic hydrogen (being produced in the gaseous phase) by certain solid unsaturated hydrocarbons. For the adsorbed-hydrogen layers at 63° to 160°K, the adsorption rates are temperature-dependent. Effective activation energies were determined in hydrogenation of a number of unsaturated hydrocarbons. For some of these compounds, the relatively minor differences in the heterogeneous hydrogenation rates are accompanied by significant differences in the effective activation energy values.

### Effect of Additives and Conditions of Preparation of Zinc Oxide on Its Performance in Isotopic Oxygen Exchange Reactions

By V. J. GORCORAKY, L. A. KASATKEENA,  
AND V. YU. LEVEEN

*V. I. Mend'yevyev Chemo-Tech-  
nological Institute in Moscow*

A study was made of the effects of Li, Ga, and In additives in ZnO catalyst and of the catalyst preparation conditions upon the kinetics of isotopic oxygen exchange. Increasing calcination temperature of pure ZnO from 850° to 1200°, increases the exchange rate to some extent. Li added to ZnO in amounts of 0.5–0.75 atom % at 850° increases the rate of isotopic oxygen exchange with the ZnO catalyst so that it exceeds