

Possible Mechanisms of Some Homogeneous Catalytic Oxidation-Reduction Reactions

K. B. YATZEEMEERSKY

*Institute of General and Inorganic Chemistry
of the Academy of Sciences of USSR*

The action mechanisms of different catalysts in oxidation-reduction reactions are examined, including the formation of a complex to effect transfer of electrical charge. In complexes of this type, an electron is transferred from a filled d -orbital of a catalyst to an anti-bonding orbital of the reactant ($d \rightarrow p\delta^*$). This step is followed by rupture of the δ -bond of the reacting molecule.

An electron can also be transferred to a d -orbital of a catalyst either from the π -orbital of the reactant (the $\pi \rightarrow d^*$ mechanism) or, else, from one of the non-bonding orbitals of the reactant (the $\delta \rightarrow d^*$ mechanism).

Realization of any one of the three mechanisms, above, requires appropriate orientation of the atomic orbitals of the catalyst and of the reactant. The substances which can facilitate these orientations, are also the ones which accelerate the reactions examined in this study.

The Essential Changes and Refinements in Classification of Reactions of Thermal Decomposition of Solids

V. V. BOLDIR'YEV

*Institute of Combustion and Chemical Kinetics
of Siberian Division of the Academy of Sciences of USSR*

(No abstract.)

Radiolysis of Water Vapor Over Semiconductor Oxides, ZnO and V₂O₅

N. S. BOOBIR'YEVA, P. I. DOLEEN,
A. I. KONONOVITCH, N. D. ROZENBL'YUM

Institution—Not given

In radiolysis of pure water vapor, the yields of molecular products can be significantly increased in presence of an electronic semiconductor, such as zinc oxide. It is believed that in the case of zinc oxide, the sensibilization reaction occurs via (a) capture of the excited electrons of the semiconductor by the adsorbed molecules of water, and (b) photodecomposition of the water molecules due to thermoluminescence.

Hydrogenation of Carbon Monoxide and Carbon Dioxide Over a Nickel Catalyst

V. M. VLAS'YENKO, G. E. YUZ'YEFOVITCH,
M. T. ROOSOV

*L. V. Peesariyevsky
Institute of Physical Chemistry of
the Ukrainian SSR*

The results of hydrogenation of the oxides of carbon in small concentrations show that (a) CO₂ (in the reaction system) has no effect on the conversion of CO; (b) hydrogenation of CO₂ is retarded by presence of CO; (c) initiation of CO₂ hydrogenation is observed in the course of transitory hydrogenation of CO on the catalyst surface; (d) the CO₂ hydrogenation rate increases with increasing effect of the CO diffusing from the surface, on the rate of CO methanation; (e) During hydrogenation of CO in the outer zone of free diffusion, the rate of CO₂ hydrogenation should approximate its rate of hydrogenation in the absence of CO.

The Problem of Modification of Ziegler Catalysts

G. A. RAZOOVA'YEV, R. P. CHERNOVSKAYA,
K. A. MEENSKER

Institution—Not shown

The effect of modifiers on activity of α -TiCl₃ + (C₂H₅)₃Al catalyst (Ti:Al ratio, 1:5 to 1:10) was studied in polymerization of propylene with aromatics and their derivatives benzene, toluene, ethyl benzene, chlorobenzene, naphthalene, tetrahydronaphthalene) at a temperature of 45°. The modifiers used with the α -TiCl₃ + (C₂H₅)₃Al catalyst were nucleophilic (C₂H₅)₃N and electrophilic (*iso*-C₄H₉)₃Al and (C₂H₅)₃Al. The modifying effect of the (C₂H₅)₃H · (C₂H₅)₃Al complex on α -TiCl₃, alone, was also determined.

The results show that the polymerization rate varies with the concentration of each modifier used. In general, the action mechanism of the additives on catalysts of Ziegler-Natta type can be explained in terms of the concepts of heterogeneous catalysis.

Nature of Oxidation of Individual *n*-Butenes and of Their Mixtures Over Copper Catalysts

N. N. POPOVA, F. A. MEEL'MAN

Institute for Coal- and Petro-Chemical Synthesis

Oxidation of mixtures of *n*-butenes over copper catalysts was investigated in a continuous-flow unit. The gaseous- and liquid chromatographic analyses show that oxidation of butene-1, butene-2, and of mixtures of isomeric *n*-butenes proceeds