

Corrigendum to “Synthesis and characterization of TM-MCM-48 (TM = Mn, V, Cr) and their catalytic activity in the oxidation of styrene” [J. Catal. 233 (1) (2005) 60–67]

S. Gómez ^a, L.J. Garces ^a, J. Villegas ^a, R. Ghosh ^b, O. Giraldo ^d, Steven L. Suib ^{a,b,c,*}

^a *Institute of Materials Science, University of Connecticut, Storrs, CT 06269-3060, USA*

^b *Department of Chemistry, University of Connecticut, Storrs, CT 06269-3060, USA*

^c *Department of Chemical Engineering, University of Connecticut, Storrs, CT 06269-3060, USA*

^d *Departamento de Física y Química, Universidad Nacional de Colombia, Manizales, Colombia*

Available online 21 June 2005

A column containing pore volume was inadvertently omitted from Table 1 during editing. For the reader's convenience, the corrected Table 1 appears here.

Table 1
N₂ sorption results for TM-MCM-48 materials

Catalyst	Si/TM ^a	BET S.A. ^b (m ² /g)	Average P.D. ^c (Å)	Pore volume ^d (cm ³ /g)
MCM-48	n.a. ^e	1056	29	0.88
Mn-MCM-48	7	930	34	0.86
V-MCM-48	13	1007	37	0.88
Cr-MCM-48	19	828	37	0.77

^a Atomic ratios calculated by EDS.

^b BET surface area.

^c BJH average pore diameter.

^d Single-point pore volume at $P/P_0 = 0.99$.

^e n.a., no applicable.

In the first paragraph, line 4, of Section 3.3 should have read:

For Mn-MCM-48, desorption peaks are observed around 190, 630, and 730 °C and a shoulder at 450 °C.

DOI of original article: 10.1016/j.jcat.2005.04.015.

* Corresponding author.

E-mail address: steven.suib@uconn.edu (S.L. Suib).