

Organic Synthesis with Anion Exchange Resins: Synthesis of α -Imino Nitriles from Nitrones[†]

Dilip Konwar,* Birendra Nath Goswami and Naleen Borthakur


Synthetic Organic Chemistry Division, Regional Research Laboratory, Jorhat-785 006, India

J. Chem. Research (S),
1999, 242–243[†]

Nitrones react with amberlite IRA 400 [CN[−]] in acetonitrile to produce α -iminonitriles in good yield.

α -Iminonitriles are important intermediates for the synthesis of α -amino acid derivatives and heterocyclic compounds.^{1,2} These compounds are prepared by reacting acetals with isocyanates in presence of an AlClEt₂ catalyst,³ reacting dithioacetals with amines,⁴ refluxing trimethyl silyl cyanide (TMSCN) with bis-anils,² oxidising conjugated imines with chromic anhydrides in the presence of TMSCN,⁵ reacting α -chloroimines with hydrogen cyanide⁶ and by reacting cyanogen bromide with imines.⁷


In recent years anion exchange resins have been used in many organic transformations. In continuation of our work on anion exchange resin,⁸ we now report here the reaction of amberlite IRA 400 [CN[−]] **2** with nitrones **1** to produce α -imino nitriles. The reaction was carried out by stirring a mixture of nitrones **1a–h** and **2** in acetonitrile under reflux. The products **3a–h** were obtained by simple filtration, evaporation of the solvent under reduced pressure and crystallization from appropriate solvents (Scheme 1). The results are summarised in Table 1.



Scheme 1

Regarding the mechanism of the reaction, it is proposed that it proceeds *via* formation of a cyanohydroxylamine intermediate **A** by taking a proton from the trace amount of water present in the solvent which immediately loses a water molecule in presence of the HO[−] +N[−]–P resin formed *in situ* (Scheme 2). In conclusion, we propose that it is a new method for the preparation of α -imino nitriles using amberlite IRA 400 [CN[−]] as cyanating agent. The reaction has the following advantages over the existing methods: (1) The polymer supported reagent is non-toxic and can

be stored for longer periods. (2) It does not require multi-step synthesis, anhydrous conditions, and the product can be obtained simply by filtering out the resin and evaporating off the solvent.



Scheme 2

Experimental

The m.p.s. were measured in a Buchi apparatus and are uncorrected. IR spectra were recorded on a Perkin-Elmer 237B spectrophotometer; ¹H NMR spectra on a Varian T-60 spectrometer with TMS as internal standard; and mass spectra on an AEIMS-30 spectrometer. The anion exchange resin amberlite IRA 400 was purchased from Aldrich chemicals as a chloride salt (16–60 mesh). The conversion of chloride into cyanide was accomplished by following the literature procedure.⁹ The nitrones were prepared following a standard method.¹⁰

Preparation of α -Imino Nitriles.—In a typical experiment a mixture of α -phenyl-*N*-phenylnitrone (**1a**) (1.97 g, 0.01 mol), acetonitrile (50 ml) and amberlite IRA 400 [CN[−]] (10 ml, 0.01 equiv)⁹ was stirred magnetically for 3 h in a 100 ml flask at reflux temperature. The reaction was monitored by tlc. The resin was filtered and the solvent distilled under reduced pressure. The residue was crystallized from ethanol to obtain **3a**, 1.89 g, 90%, m.p. 71 °C (lit. 72 °C⁴). Compounds **3b–h** were prepared in similar fashion (Table 1).

We acknowledge Dr J. C. S. Katak, scientist, and Dr J. S. Sandhu, Acting Director, regional research laboratory, Jorhat for their encouragement and helpful suggestions.

Table 1 Synthesis of α -iminonitriles using amberlite IRA 400 [CN[−]]^a

Entry	Product (3)	R ¹	R ²	Yield(%)	Time/h	m.p./°C found (lit.)
1	a	Ph	Ph	90	3	71 (72 ⁴)
2	b	<i>p</i> -Cl-C ₆ H ₄	Me	80	3.5	50 (52 ⁴)
3	c	<i>p</i> -Me-C ₆ H ₄	C ₆ H ₁₁	75	4	67 (67 ⁴)
4	d	<i>o</i> -Cl-C ₆ H ₄	<i>p</i> -NO ₂ -C ₆ H ₄	78	3.6	117 (116 ⁴)
5	e	<i>p</i> -NO ₂ -C ₆ H ₄	<i>m</i> -NO ₂ -C ₆ H ₄	82	4	168 (166 ⁴)
6	f	PhCH=CH	Ph	76	6	72 (73 ⁵)
7	g	PhCH=CH	<i>p</i> -Br-C ₆ H ₄	77	5	141 (142 ⁵)
8	h	PhCH=CH	<i>p</i> -OMe-C ₆ H ₄	81	5.5	122 (124 ⁵)

^aAll the compounds gave satisfactory IR, NMR, mass spectral and elemental analyses.

* To receive any correspondence.

[†] This is a **Short Paper** as defined in the Instructions for Authors, Section 5.0 [see *J. Chem. Research (S)*, 1999, Issue 1]; there is therefore no corresponding material in *J. Chem. Research (M)*.

Received, 17th September 1998; Accepted, 2nd December 1998
Paper E/8/07253A

References

- 1 R. M. Williams *Synthesis of Optically Active α -Amino Acids*, Pergamon Press, Oxford, 1989, p. 208.
- 2 M. Takahashi, H. Miyahara and N. Yoshida, *Heterocycles*, 1988, **27**, 155.
- 3 H. Pellissier and H. Gill, *Tetrahedron Lett.*, 1989, **30**, 171.
- 4 F. Pochat, *Tetrahedron Lett.*, 1981, **22**, 955.
- 5 D. Karmakar, D. Prajapati and J. S. Sandhu, *J. Chem. Res. (S)*, 1996, 538.
- 6 J. Manfred and L. Kurt, *German Offen*, 1973, 2,221,771 (*Chem. Abstr.*, 1974, **80**, 36883f).
- 7 M. Rai, K. Kewel and A. Singh, *Indian J. Chem.*, 1976, **14Bs**, 376.
- 8 D. Konwar, D. K. Dutta and B. N. Goswami, *J. Chem. Res. (S)*, 1998, 342 and references cited therein.
- 9 M. Gordon, M. L. Depamphilis and C. E. Griffin, *J. Org. Chem.*, 1963, **28**, 698.
- 10 A. Roe and J. A. Montgomery, *J. Am. Chem. Soc.*, 1953, **75**, 910.