

Bibliography Section

Liquid Column Chromatography

1. REVIEWS AND BOOKS

- 2094 Antia, F.D. and Horvath, C.: Operational modes of chromatographic separation processes. *Ber. Bunsen-Ges. Phys. Chem.*, 93 (1989) 961-968; *C.A.*, 111 (1989) 211208x - a review with 46 refs.
- 2095 Brinkman, U.A.T.: (GC, LC, TLC: which one is best?) *Lab. 2000*, 2 (1988) 40-46; *C.A.*, 111 (1989) 224539f - a review with no refs.
- 2096 Buydens, L.M.C., van Leeuwen, J.A., Mulholland, M., Vandeginste, B.G.M. and Kateman, G.: An expert system for the validation of high-performance liquid chromatographic methods. *TrAC*, 9 (1990) 58-62.
- 2097 Caiken, I.M., Fassina, G. and Caliceti, P.: High performance affinity chromatography. *Chem. Anal.*, 98 (1989) 317-335; *C.A.*, 111 (1989) 211180g - a review with 66 refs.
- 2098 Citterio, A., Fancelli, D., Grossi, G. and Brambilla, S.: (New developments in HPLC technology). *Lab. 2000*, 1 (1987) 38-42; *C.A.*, 111 (1989) 247191a - a review with no refs.
- 2099 Collins, C.H., Radler de Aquino Neto, F. and Pereira da Silva, J.R.: (Terminology in chromatography. Part I. Vocabulary for chromatography). *Quim. Nova*, 11 (1988) 443-464; *C.A.*, 112 (1990) 29870a - a review with 4 refs.
- 2100 Fetzner, J.C.: Gas- and liquid-chromatographic techniques. *Chem. Anal.*, 101 (1989) 59-109; *C.A.*, 111 (1989) 247192b - a review with 229 refs.
- 2101 Graciani, C.E., Janer del Valle, M.L., Albi, V.T. and Lanzon, R.A.: Bibliographical summary on high performance liquid chromatography of lipids: I. 1975-1979. *Grasas Aceites (Seville)*, 39 (1988) 373-386; *C.A.*, 111 (1989) 211203s - a bibliography with 218 refs.
- 2102 Krull, I.S., Colgan, S.T. and Selavka, C.M.: A theoretical approach to derivatizations for HPLC. *Chem. Anal.*, 98 (1989) 393-413; *C.A.*, 111 (1989) 224537d - a review with 42 refs.
- 2103 Meriluoto, J. and Bjorklund, H.: Regenerating internal surface reversed-phase columns: two case histories. *LC-GC*, 7 (1989) 738-740; *C.A.*, 111 (1989) 211199v - a review with 5 refs.
- 2104 Mueller, W.: Aqueous two-phase polymer systems for liquid/liquid partition-chromatography of biopolymers. *Ber. Bunsen-Ges. Phys. Chem.*, 93 (1989) 956-961; *C.A.*, 111 (1989) 211207w - a review with 9 refs.
- 2105 Ramsay, J.D.F.: Oxide gels: applications in chromatography and investigations of structure by neutron scattering. *Pure Appl. Chem.*, 61 (1989) 1963-1986; *C.A.*, 112 (1990) 26096s - a review with 21 refs.
- 2106 Sagliano, N., Jr., Raglione, T.V. and Hartwick, R.A.: Multidimensional techniques in chromatography. *Chem. Anal.*, 98 (1989) 643-668; *C.A.*, 111 (1989) 224538e - a review with 32 refs.
- 2107 Simpson, R.C.: High speed liquid chromatography. *Chem. Anal.*, 98 (1989) 375-391; *C.A.*, 111 (1989) 224172f - a review with 25 refs.

- 2108 Sing, K.S.W. (Editor): *International Symposium on Surface Chemistry, Adsorption and Chromatography (Dedicated to the Memory of A.V. Kiselev)*, held in Moscow, USSR, 13-17 November 1988, *Pure Appl. Chem.*, Vol. 61, Blackwell, Oxford, 1989, 235 p.; *C.A.*, 112 (1990) 26255t.
- 2109 Sofer, G.K. and Nystroem, L.E.: *Process Chromatography. A Practical Guide*. Academic, San Francisco, 1989, 160 p.; *C.A.*, 111 (1989) 236013t.
- 2110 Timerbaev, A.R. and Petrukhin, O.M.: *(Liquid Adsorption Chromatography of Chelates)*. Nauka, Moscow, 1989, 284 p.; *C.A.*, 112 (1990) 30001z.
- 2111 Van Leeuwen, J.A., Buydens, L.M.C., Vandeginste, B.G.M. and Kateman, G.: Expert systems in chemical analysis. *TrAC*, 9 (1990) 49-54.
- 2112 Williams, S.S.: Application of expert systems programming to high-performance liquid chromatography method development. *TrAC*, 9 (1990) 63-65.
- 2113 Witkiewicz, Z., Mazurek, M. and Szulc, J.: Chromatographic analysis of chemical warfare agents. *J. Chromatogr.*, 503 (1990) 292-357 - a review with 413 refs.
- 2114 Yang, F.J.: Microbore column chromatography: a unified approach to chromatography. *Chromatography Sci.*, 45 (1989) 1-36; *C.A.*, 111 (1989) 224535b - a review with 83 refs.
- 2115 Yau, W.W., Kirkland, J.J. and Bly, D.D.: Size-exclusion liquid chromatography. *Chem. Anal.*, 98 (1989) 277-316; *C.A.*, 111 (1989) 224536c - a review with 34 refs.

For additional information see:

C.A., 112 (1990) 63100j.

See also 2121, 2164, 2170, 2173, 2178, 2179, 2184, 2186, 2206, 2222, 2231, 2234, 2236, 2243, 2245, 2248, 2249, 2254, 2255, 2257, 2259, 2260, 2265, 2271, 2272, 2273, 2295, 2319, 2325, 2408, 2413, 2417, 2420, 2427, 2474, 2539, 2561, 2567, 2574, 2579, 2584, 2589, 2595, 2596, 2601, 2613, 2801, 2808, 2811, 2826, 2836, 2920, 2931, 2932, 2933, 2934, 2935, 2937, 2938, 2943, 2944, 3069, 3078, 3079, 3082, 3090, 3093, 3097.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 2116 Ahuja, S.: Optimization of selectivity, detectability and analysis time in high-performance liquid chromatography. *J. Chromatogr.*, 499 (1990) 489-498.
- 2117 Bartha, A., Vigh, G. and Varga-Fuchony, Z.: Basis of the rational selection of the hydrophobicity and concentration of the ion-pairing reagent in reversed-phase ion-pair high-performance liquid chromatography. *J. Chromatogr.*, 499 (1990) 423-434.
- 2118 Cross, R.F.: Retention time deconvolution in high performance liquid chromatography. *LC-GC*, 7 (1989) 418-423; *C.A.*, 111 (1989) 224579u.
- 2119 Crow, J.A. and Foley, J.P.: Optimization of separations in supercritical fluid chromatography using a modified simplex algorithm and short capillary columns. *Anal. Chem.*, 62 (1990) 378-386.
- 2120 Dolan, J.W.: Column flushing demystified. *LC-GC*, 7 (1989) 556-560; *C.A.*, 112 (1990) 69131s.
- 2121 Dolan, J.W.: Peak tailing. *LC-GC*, 7 (1989) 476-482; *C.A.*, 112 (1990) 47981s - a review with 10 refs.
- 2122 Dolan, J.W.: LC troubleshooting: readers' questions. *LC-GC*, 7 (1989) 384-387; *C.A.*, 112 (1990) 48023t.

- 2123 Hayashi, Y., Yoshioka, S. and Takeda, Y.: Shannon information retrieved from overlapped chromatographic peaks with Kalman filter. *Anal. Sci.*, 5 (1989) 329-343; *C.A.*, 112 (1990) 47684x.
- 2124 Kaibara, A., Nakagawa, T., Hohda, C., Hirata, N. and Tanaka, H.: Molecular posture of solute adsorbed to stationary alkyl ligand in reversed-phase liquid chromatography. *J. Chromatogr. Sci.*, 27 (1989) 716-720.
- 2125 Korenaga, T., Shen, F., Yoshida, H. and Takahashi, T.: Dispersion behavior of solutes in an ideal laminar flow with small-bore glass tubes. *Bull. Chem. Soc. Jpn.*, 62 (1989) 1492-1500; *C.A.*, 112 (1990) 63429y.
- 2126 MacLeod, S.K., Fagan, D.T. and Scholl, J.P.: Comments on solute-solvent interactions in quantitative high-performance liquid chromatography. *J. Chromatogr.*, 502 (1990) 236-240.
- 2127 Mueller, E. and Jork, H.: (Hydrophobic chromatography on a gel filtration medium. Effect of salts on the separation of low-molecular-weight compounds). *GIT-Suppl.*, (1989) 97-101; *C.A.*, 111 (1989) 247218q.
- 2128 Strasters, J.K., Billiet, H.A.H., de Galan, L. and Vandeginste, B.G.M.: Strategy for peak tracking in liquid chromatography on the basis of a multivariate analysis of spectral data. *J. Chromatogr.*, 499 (1990) 499-522.
- 2129 Wilson, W.H., McNair, H.M., Maa, Y.F. and Hyver, K.J.: The effect of the Knox-Parcher ratio in microcolumn LC. *J. High Resolut. Chromatogr.*, 13 (1990) 18-21.

For additional information see:

C.A., 111 (1989) 193726h.

See also 2099, 2108, 2109, 2114, 2115, 2165, 2199, 2200, 2208, 2212, 2240, 2281, 2282, 2997.

2b. Thermodynamics and theoretical relationships

- 2130 Atamna, I.Z., Muschik, G.M. and Issaq, H.J.: Effect of alcohol chain length, concentration and polarity on separations in high-performance liquid chromatography using bonded cyclodextrin columns. *J. Chromatogr.*, 499 (1990) 477-488.
- 2131 Benedek, P. and Janko, T.: Investigation and improvement of the Horvath-Lin equation of state. *J. Chromatogr.*, 499 (1990) 463-467.
- 2132 Berek, D. and Macko, T.: Pressure effects in high performance liquid chromatography. *Pure Appl. Chem.*, 61 (1989) 2041-2046; *C.A.*, 112 (1990) 12473a.
- 2133 Cheong, W.J. and Carr, P.W.: Study of partition models in reversed-phase liquid chromatography based on measured mobile phase solute activity coefficients. *J. Chromatogr.*, 499 (1990) 373-393.
- 2134 Czok, M. and Guiochon, G.: The physical sense of simulation models of liquid chromatography: propagation through a grid or solution of the mass balance equation? *Anal. Chem.*, 62 (1990) 189-200.
- 2135 Davidova, H. and Schneider, P.: Elimination of dead-volume contributions to moment of chromatographic peaks. *Collect. Czech. Chem. Commun.*, 54 (1989) 2974-2984; *C.A.*, 112 (1990) 38888q.
- 2136 Dore, E.V. and Guiochon, G.: Normalised measure of overlap between non-gaussian chromatographic peaks. *Anal. Chem.*, 62 (1990) 174-181.
- 2137 Engelhardt, H. and Jungheim, M.: Comparison and characterization of reversed phases. *Chromatographia*, 29 (1990) 59-68.
- 2138 Huss, V., Chevalier, J.L. and Siouffi, A.M.: Diffusion coefficients in water-acetonitrile mixtures. *J. Chromatogr.*, 500 (1990) 241-255.
- 2139 Jandera, P. and Kubat, J.: Possibilities of determination and prediction of solute capacity factors in reversed-phase systems with pure water as the mobile phase. *J. Chromatogr.*, 500 (1990) 281-299.

- 2140 Kawasaki, T.: A fundamental structure of the general theory of overload quasi-static linear gradient chromatography. *Sep. Sci Technol.*, 24 (1989) 1109-1158; *C.A.*, 112 (1990) 26229n.
- 2141 Landau, V.V., Zabokritskii, M.P. and Sterkhov, N.V.: (Capacity for separation in chromatography). *Zh. Fiz. Khim.*, 63 (1989) 2978-2983; *C.A.*, 112 (1990) 63544g.
- 2142 Lin, B., Ma, Z., Golshan-Shirazi, S. and Guiochon, G.: Theoretical analysis of non-linear preparative liquid chromatography. *J. Chromatogr.*, 500 (1990) 185-213.
- 2143 Michels, J.J. and Dorsey, J.G.: Estimation of the reversed-phase liquid chromatographic lipophilicity parameter $\log k_w$ using ET-30 solvatochromism. *J. Chromatogr.*, 499 (1990) 435-451.
- 2144 Naish-Chamberlain, P.J. and Lynch, R.J.: Evaluation of a complete HPLC solvent optimization system involving piece-wise quadratic modelling. *Chromatographia*, 29 (1990) 79-89.
- 2145 Peichang, L. and Hongxin, H.: An intelligent search method for HPLC optimization. *J. Chromatogr. Sci.*, 27 (1989) 690-697.
- 2146 Sobolev, D.M., Sharikov, Yu.V. and Selekhnova, G.B.: (Use of regression models for the calculation of chromatograms with overlapping peaks). *Zh. Anal. Khim.*, 44 (1989) 1266-1273; *C.A.*, 111 (1989) 247274e.
- 2147 Sreenivasan, K.: Negentropy and retention in reversed phase liquid chromatography. *Chromatographia*, 29 (1990) 90-92.

see also 2117, 2232, 2233, 2563, 2583, 2783, 2921.

2d. Measurement of physico-chemical and related values

- 2148 Golshan-Shirazi, S. and Guiochon, G.: General solution of the ideal model of chromatography for a single-compound band. *J. Phys. Chem.*, 94 (1990) 495-500; *C.A.*, 112 (1990) 43359x.
- 2149 Johnson, D.K., Chum, H.L. and Hyatt, J.A.: Molecular weight distribution studies using lignin model compounds. *ACS Symp. Ser.*, 397 (1989) 109-123; *C.A.*, 111 (1989) 216164q.
- 2150 Kazakevich, Yu.V. and Vorob'eva, L.D.: (Effect of the composition of the mobile phase and the amount of the modifier in the eluent on the retention of model compounds). *Zh. Fiz. Khim.*, 63 (1989) 420-425; *C.A.*, 111 (1989) 224564k.
- 2151 Kazakevich, Yu.V., Larionov, O.G. and El'tekov, Yu.A.: Physico-chemical aspects of liquid chromatography. *Pure Appl. Chem.*, 61 (1989) 2037-2040; *C.A.*, 112 (1990) 12472z.
- 2152 Li, H., Chen, D., He, J. and Huang, G.: Measurement of the Mark-Houwink constants and the unperturbed molecular parameters of polyepichlorohydrin by gel permeation chromatography and low-angle laser light scattering. *Eur. Polym. J.*, 25 (1989) 1065-1067; *C.A.*, 112 (1990) 37146j.
- 2153 McCoy, B.J.: Adsorption chromatography of a heterogeneous mixture. *Chem. Eng. Sci.*, 44 (1989) 993-996; *C.A.*, 112 (1990) 15775m.
- 2154 Muchnik, V.L. and Shaevich, A.B.: (Ways of improving chromatographic identification based on the use of reference retention data and reference materials). *Zh. Anal. Khim.*, 44 (1989) 707-711; *C.A.*, 111 (1989) 224276t.
- 2155 Omorodin, S.N.E. and Hamielec, A.E.: Newly proposed relationship between peak dispersion coefficient and width of the experimental chromatogram based on theories of proposed instrumental spreading shape function in size exclusion chromatography. *Angew. Macromol. Chem.*, 172 (1989) 1-14; *C.A.*, 112 (1990) 57166u.
- 2156 Vit, I., Fahnrich, J. and Popl, M.: Determination of the dead volume columns in reversed phase liquid chromatography. *Collect. Czech. Chem. Commun.*, 54 (1989) 953-966; *C.A.*, 112 (1990) 63428x.

- 2157 Yamamoto, F.M., Rokushika, S. and Hatano, H.: Comparison of thermodynamic retention behaviors on various C₁₈ columns different in their hydrophobicity. *J. Chromatogr. Sci.*, 27 (1989) 704-709.

For additional information see:

C.A., 112 (1990) 23273m, 26228m, 36991u, 63426v.

See also 2283, 2502, 2815, 2917, 2918, 2920, 2922, 2923, 2928, 2930, 3091.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 2158 Biemann, K. and Gagel, J.J.: Method and apparatus for continuous collection of chromatograph effluent. *U.S. Pat.* US 4,843,243 (Cl. 250-341; G01N30/02), 27 Jun. 1989; Appl. 851,445, 14 Apr. 1986; 11 p.; C.A., 111 (1989) 224621c.
- 2159 Conlon, R.D., Schmid, C.E. and Perrone, P.A.: An LC pump family. *Am. Lab. (Fairfield)*, 21 (1989) 50-57; C.A., 112 (1990) 47996a.
- 2160 Duffin, K.L., Flurer, R.A., Busch, K.L., Sexton, L.W. and Dorsett, J.L.: Automated sample transport system for chromatography/secondary ion mass spectrometry. *Rev. Sci. Instrum.*, 60 (1989) 1071-1074; C.A., 112 (1990) 47710c.
- 2161 Goodley, P.C.: On-axis electron-acceleration electrode for liquid chromatograph/mass spectrometer. *U.S. Pat.* US 4,851,700 (Cl.250-288; D01D59/44), 25 Jul. 1989, Appl. 194,591, 16 May 1988; 10 p.; C.A., 111 (1989) 224629k.
- 2162 Joshua, H. and Kath, G.S.: Automated sequential actuation of two autoinjectors. *LC-GC*, 7 (1989) 272-275; C.A., 111 (1989) 224554g.
- 2163 Krejci, M. and Kahle, V.: Method and device for chromatographic analysis. *Czech. Pat.* CS 257,657 (Cl. G01N30/16), 09 Jan. 1989, Appl. 85/8,421, 22 Nov. 1985; 7 p.; C.A., 111 (1989) 247140h.
- 2164 Majors, R.E.: Highlights of column developments at the 13th Symposium on Column Liquid Chromatography. *LC-GC*, 7 (1989) 812-820; C.A., 111 (1989) 219858e - a review with 1 ref.
- 2165 Mason, A.Z.: In ICP-MS study of saline-induced corrosion of stainless steel HPLC components: is there a need to go to non-metallic systems? *Chromatogram*, 10 (1989) 7-10; C.A., 112 (1990) 9074r.
- 2166 Myers, P. and Roberts, J.C.: Tubing assembly for chromatographic separation columns. *PCT Int. Pat.* Appl. WO 89 07,759 (Cl. G01N30/60), 24 Aug. 1989, GB Appl. 88/3,993, 20 Feb. 1988; 11 p.; C.A., 111 (1989) 224623d.
- 2167 Podolak, M. and Zezulka, A.: (Filling reactor for microcolumn high-efficiency liquid chromatographs). *Czech. Pat.* CS 260,896 (Cl. G01N30/02), 15 May 1989, Appl. 86/7,965, 04 Nov. 1986; 3 p.; C.A., 111 (1989) 247280d.
- 2168 Pozhidaev, V.M., Korolev, A.A. and Berezkin, V.G.: (Device for sample feed into a capillary column for the analysis of polycyclic aromatic hydrocarbons). *Zavod. Lab.*, 55 (1989) 26-28; C.A., 111 (1989) 247224p.
- 2169 Rericha, L., Klir, L., Hajkova, I. and Ehrenberger, J.: (Filter for liquids, especially for chromatographic analysis of biologically active materials). *Czech. Pat.* CS 258,771 (Cl. B01D15/08), 14 Apr. 1989, Appl. 86/3,392, 08 May 1986; 3 p.; C.A., 111 (1989) 216400p.

For additional information see:

C.A., 111 (1989) 224477j, 224483h, 224504r, 224507u, 228564h, 247177a, 247225q, 247228t, 247229u, 247286k;
112 (1990) 15748e, 15755e, 15762e, 30010b, 46778u, 47944g.

See also 2189, 2279.

3b. Detectors and detection reagents

- 2170 Austin-Harrison, D.S. and Johnson, D.C.: Pulsed amperometric detection based on direct and indirect anodic reactions: a review. *Electroanalysis*, 1 (1989) 189-197; *C.A.*, 112 (1990) 47976u - a review with 26 refs.
- 2171 Baeyens, W.R.G., Ling, B.L., Brinkman, U.A.T. and Schulman, S.G.: Principles and applications of luminescence spectrometry coupled to liquid chromatographic techniques. *J. Biolumin. Chemilumin.*, 4 (1989) 484-499; *C.A.*, 111 (1989) 228149b.
- 2172 Berthod, A., Glick, M. and Winefordner, J.D.: Sensitive, indirect photometric detector for high-performance liquid chromatography using a light-emitting diode. *J. Chromatogr.*, 502 (1990) 305-315.
- 2173 Birs, J.W. (Editor): *Chemiluminescence and Photochemical Reaction Detection in Chromatography*. VCH, New York, 1989, 291 p.; *C.A.*, 112 (1990) 48081k.
- 2174 Brotherton, T.J., Phannerstill, P.E., Creed, J.T., Heitkemper, D.T., Caruso, J.A. and Pratsinis, S.E.: Evaluation of three low-volume interfaces for organic solvent introduction to the inductively coupled plasma-applications to flow injection. *J. Anal. At. Spectrom.*, 4 (1989) 341-345; *C.A.*, 112 (1990) 68770n.
- 2175 Chervet, J.P., Ursem, M., Salzmann, J.P. and Vannort, R.W.: Performance comparison of flow cells for UV detection in capillary liquid chromatography. *LC-GC*, 7 (1989) 514-518; *C.A.*, 112 (1990) 48006q.
- 2176 Cobb, W.T. and McGown, L.B.: On-line fluorescence lifetime detector for chromatographic peak resolution. *Anal. Chem.*, 62 (1990) 186-189.
- 2177 Evans, C.E. and McGuffin, V.L.: Evaluation of refractive index artifacts in liquid chromatographic absorbance detection. Extension to non-ideal injection and gradient elution. *J. Chromatogr.*, 503 (1990) 127-154.
- 2178 Gao, C.X. and Krull, I.S.: Solid-phase reaction detection in HPLC. *Biochromatography*, 4 (1989) 222-233; *C.A.*, 112 (1990) 3671m - a review with 55 refs.
- 2179 Gin, G.: (Analysis of refractory metals). *Fenxi Shiyanshi*, 7 (1988) 38-53; *C.A.*, 111 (1989) 224167h - a review with 308 refs.
- 2180 Gooijer, C., van Zoonen, P., Velthorst, N.H. and Frei, R.W.: On the mechanism of peroxyoxalate chemiluminescence. Quenched chemiluminescence as a detection method in HPLC. *J. Biolumin. Chemilumin.*, 4 (1989) 479-483; *C.A.*, 111 (1989) 228148a.
- 2181 Hanaoka, N.: Post-column adjustment of conditions for peroxyoxalate chemiluminescence detection for high-performance liquid chromatography. *J. Chromatogr.*, 503 (1990) 155-165.
- 2182 Hernandez, L., Marquina, R., Escalona, J. and Guzman, N.: Detection and quantification of capillary electrophoresis zones by fluorescence microscopy. *J. Chromatogr.*, 502 (1990) 247-255.
- 2183 Hippe, H., Schirmer, A. and von Laffert, S.: Applications of online biosensors in HPLC. *CLS, Chem. Labor. Betr.*, 40 (1989) 470-474; *C.A.*, 112 (1990) 3741j.
- 2184 Imai, K., Nishitani, T., Akimoto, H. and Tsukamoto, Y.: Chemiluminescence in HPLC. *J. Biolumin. Chemilumin.*, 4 (1989) 500-504; *C.A.*, 111 (1989) 228150v - a review with 36 refs.
- 2185 Kientz, C.E., Verweij, A., de Jong, G.J. and Brinkman, U.A.T.: The potential of on-line flame photometric detection in microcolumn liquid chromatography. *J. High Resolut. Chromatogr.*, 12 (1989) 793-796.
- 2186 Lunte, C.E.: Voltametric detection for liquid chromatography. *LC-GC*, 7 (1989) 492-498; *C.A.*, 112 (1990) 47628g - a review with 17 refs.

- 2187 Marr, J.G.D.: Computer-aided detection systems for HPLC: development, assessment and application of digital techniques for peak purity, validation in HPLC utilizing photodiode array detection. Avail. *Univ. Microfilms Int.*, Order No BRDX85077, 1988, 369 p.; C.A., 111 (1989) 239616c.
- 2188 Okawa, Y., Tsuzuki, H., Yoshida, S. and Watanabe, T.: Glucose sensor carrying monomolecular layer of glucose oxidase covalently bound to tin(IV) oxide electrode. *Anal. Sci.*, 5 (1989) 507-512; C.A., 111 (1989) 211234c.
- 2189 Paradis, R.C.: Detector cell for liquid chromatographs. *Eur. Pat. Appl. EP 335,268* (Cl. G01N21/05), 04 Oct. 1989, US Appl. 176,524, 01 Apr. 1988; 7 p.; C.A., 111 (1989) 224630d.
- 2190 Ramstad, T.: Pulse-enhanced electrosorptive detection of ions at silver in ion chromatography. *Anal. Lett.*, 22 (1989) 2123-2143.
- 2191 Rhoads, R.: High-sensitivity chromatography detector. *U.S. Pat. US 4,838,688* (Cl.356-72; G01N21/05), 13 Jun. 1989, US Appl. 109,456, 16 Oct. 1987; 6 p.; C.A., 111 (1989) 224622c.
- 2192 Ruban, V.F. and Belen'kii, B.G.: (High-sensitivity amperometric microdetector for capillary liquid chromatography). *Zh. Anal. Khim.*, 44 (1989) 725-727; C.A., 111 (1989) 224556j.
- 2193 Ruban, V.F.: Characteristics of an amperometric detector for capillary HPLC. *J. High Resolut. Chromatogr.*, 13 (1990) 112-115.
- 2194 Sheehan, T.L., Adaskaveg, M. and Excoffier, J.L.: Powerful implementation of spectral processing for photodiode array detection. *Am. Lab. (Fairfield)*, 21 (1989) 66-74; C.A., 112 (1990) 47653m.
- 2195 Shintani, H. and Dasgupta, P.H.: Ion chromatography using pH-detection. *Fresenius' Z. Anal. Chem.*, 336 (1990) 38.
- 2196 Stolowitz, M.L.: Solid-phase oxime reagent for liquid chromatography. *U.S. Pat. US 4,812,532* (Cl. 525-420; C08G69/48), 14 Mar. 1989, Appl. 79,258, 29 Jul. 1987; 5 p.; C.A., 111 (1989) 247284h.

For additional information see:

C.A., 111 (1989) 224271n, 224591s;
112 (1990) 30009h, 30052s, 68723z, 68729f, 68762m.

See also 2327, 2328, 2346, 2378, 2402, 2415, 2479, 2494, 2507, 2511, 2533, 2789, 2876, 2901, 3069, 3145.

3c. Sorbents, carriers, column and layer performance, packing procedures

- 2197 Barcza, S.: Chiral stationary phases and columns for chromatographic resolution. *U.S. Pat. US 4,824,950* (Cl. 546-14; C07F7/02), 25 Apr. 1989, US Appl. 776,251, 16 Sep. 1985; 9 p.; C.A., 111 (1989) 219271h.
- 2198 Boschetti, E.: Polyacrylamide derivatives to the service of bioseparations. *J. Biochem. Biophys. Methods*, 19 (1989) 21-36; C.A., 111 (1989) 228466c.
- 2199 Braggio, R.: (When little is better (precolumn HPLC)). *Lab. 2000*, 1 (1987) 61-62; C.A., 111 (1989) 247212h.
- 2200 Buszewski, B., Suprynowicz, Z., Staszuk, P., Albert, K., Pfeleiderer, B. and Bayer, E.: Effect of coverage density on the retention mechanism in reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 499 (1990) 305-316.
- 2201 Connor, M.P., Wang, J., Kubiak, W. and Smyth, M.R.: Tissue- and microbe-based electrochemical detectors for liquid chromatography. *Anal. Chim. Acta*, 229 (1990) 139-143.
- 2202 Crane, L.J. and Henry, M.P.: Design requirements for silica-based-chromatographic matrixes. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (1989) 59-69; C.A., 111 (1989) 170325n.

- 2203 Eisenbeiss, F.: (The effect of support materials on chromatographic properties.) *Ber. Bunsen-Ges. Phys. Chem.*, 93 (1989) 1019-1026; *C.A.*, 111 (1989) 211285v.
- 2204 Hirt, R. and Panning, H.: (Practical comparison of regular bore and narrowbore HPLC). *LaborPraxis*, 13 (1989) 68-75; *C.A.*, 111 (1989) 247221k.
- 2205 Huber, J.F.K., Pawlowska, M. and Markl, P.: Selection of the solid support - a crucial step for the realization of solvent generated liquid-liquid chromatography. *J. Chromatogr.*, 500 (1990) 257-280.
- 2206 Ivanov, A.E., Saburov, V.V. and Zubov, V.P.: (Modification of mineral carriers by oligomers and polymers - path for the synthesis of sorbents for chromatography of biopolymers.) *Zh. Vses. Khim. O-va. im. D.I. Mendeleeva*, 34 (1989) 368-377; *C.A.*, 111 (1989) 190552n - a review with 58 refs.
- 2207 Jinno, K.: CP-MAS ^{13}C nuclear magnetic resonance spectra for identification of functionality of octadecylsilica bonded phases. *J. Chromatogr. Sci.*, 27 (1989) 729-734.
- 2208 Kaliszan, R., Osmialowski, K., Bassler, B.J. and Hartwick, R.A.: Mechanism of retention in high-performance liquid chromatography on porous graphitic carbon as revealed by principal component analysis of structural descriptors of solutes. *J. Chromatogr.*, 499 (1990) 333-344.
- 2209 Kaufmann, A.: Crown ether-alkali salt complexes as modifiers for normal phase HPLC. *Chromatographia*, 29 (1990) 76-78.
- 2210 Khaledi, M.G., Strasters, J.K., Rodgers, A.H. and Breyer, E.D.: Simultaneous enhancement of separation selectivity and solvent strength in reversed-phase liquid chromatography using micelles in hydro-organic solvents. *Anal. Chem.*, 62 (1990) 130-136.
- 2211 Kimata, K., Iwaguchi, K., Onishi, S., Jinno, K., Eksteen, R., Hosoya, K., Araki, M. and Tanaka, N.: Chromatographic characterization of silica C_{18} packing materials. Correlation between a preparation method and retention behavior of stationary phase. *J. Chromatogr. Sci.*, 27 (1989) 721-728.
- 2212 Majors, R.E.: A comparative study of European and U.S. HPLC column trends. *LC-GC*, 7 (1989) 468-475; *C.A.*, 112 (1990) 47679z.
- 2213 Mazza, J.A., Outumuro, P., Moroux, Y. and Boschetti, E.: Polymer design in dye chromatography: from the definition of monomers to the evaluation of polymeric support. In: Vijayalakshmi, M.A. and Bertrand, O. (Editors), *Protein-Dye Interact.*, [Proc. Int. Conf.], 1st 1988, Elsevier, Barking, 1989, pp. 126-136; *C.A.*, 111 (1989) 211270m.
- 2214 Menger, F.M., Tsuno, T. and Hammond, G.S.: Crosslinked polystyrene incorporating water pools. *J. Am. Chem. Soc.*, 112 (1990) 1263-1264; *C.A.*, 112 (1990) 56837b.
- 2215 Pedigo, S. and Bowers, L.D.: Comparison of retention behavior on polymeric resins and an alkyl-bonded silica phase in reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 499 (1990) 279-290.
- 2216 Pfannkoch, E.A., Switzer, B.S. and Kopaciewicz, W.: Aluminium ion mediated stabilization of silica-based anion-exchange packing to caustic regenerants. *J. Chromatogr.*, 503 (1990) 385-401.
- 2217 Rigney, M.P., Funkenbusch, E.F. and Carr, P.W.: Physical and chemical characterization of microporous zirconia. *J. Chromatogr.*, 499 (1990) 291-304.
- 2218 Roberts, D.W., Ruane, R.J. and Wilson, I.D.: The use of graphitized carbon black in solid phase extraction: comparison with C_{18} bonded silica gel. *J. Pharm. Biomed. Anal.*, 7 (1989) 1077-1086.
- 2219 Sarkar, B. and Basu, S.: Studies on zirconium tungstate ion exchanger. *Indian J. Chem., Sect. A*, 28A (1989) 346-348; *C.A.*, 112 (1990) 47727p.
- 2220 Sewell, R.F. and Brechley, P.E.C.: Coupling of mono- and polyamino ligands to solid phases by nucleophilic attack on coupling groups that may be spontaneously hydrolysed. *J. Chromatogr.*, 525 (1990) 246-248.

- 2221 Simonzadeh, N. and Schilt, A.A.: Chelation properties of silica-bound 1,10-phenanthroline. *J. Coord. Chem.*, 20 (1989) 117-120; *C.A.*, 112 (1990) 12429r.
- 2222 Smolkova-Keulemansova, E.: Cyclodextrins in chromatography. In: Duchene, D. (Editor), *Cyclodextrins Their Ind. Uses*, Ed. Sante, Paris, 1987, pp. 259-295; *C.A.*, 111 (1989) 224540z - a review with 88 refs.
- 2223 Stillian, J.R. and Pohl, C.A.: New latex-bonded pellicular anion exchangers with multi-phase selectivity for high-performance chromatographic separations. *J. Chromatogr.*, 499 (1990) 249-266.
- 2224 Suttmar, J.: (Purification of acetonitrile for high-performance liquid chromatography). *Chem. Listy*, 83 (1989) 435-436; *C.A.*, 111 (1989) 224561g.
- 2225 Szabo, G., Csato, E., Keresztes, P. and Pallos, J.P.: Preparation and retention characteristics of phenylmethylpolysiloxane phase for reversed-phase liquid chromatography. *J. Liq. Chromatogr.*, 12 (1989) 2839-2853.
- 2226 Tanaka, M., Inada, H., Kawaguchi, Y. and Shono, T.: 2,5-Di-O-methylated cyclodextrin-bonded stationary phases for liquid chromatography. *Anal. Sci.*, 5 (1989) 211-213; *C.A.*, 112 (1990) 48018v.
- 2227 Tani, K. and Suzuki, Y.: Comparison of the adsorption isotherms for monomeric and polymeric C₁₈ bonded phases. *J. Chromatogr. Sci.*, 27 (1989) 698-703.
- 2228 Van Berkel-Geldof, O., Kraak, J.C. and Poppe, H.: Preparation of silicone-coated 5-25- μ m I.D. fused-silica capillary columns for open-tubular liquid chromatography. *J. Chromatogr.*, 499 (1990) 345-359.
- 2229 Wang, J., Chen, S.P. and Lin, M.S.: Use of different electropolymerization conditions for controlling the size-exclusion selectivity at polyaniline, polypyrrole and polyphenol films. *J. Electroanal. Chem. Interfacial. Electrochem.*, 273 (1989) 231-242; *C.A.*, 112 (1990) 65303b.

For additional information see:

C.A., 111 (1989) 190658b, 190993g, 211478k, 224491j, 228235b, 228557h,
228561e, 233207y, 235430q, 235431r;
112 (1990) 11258x, 15756f, 23134s, 39075j, 39251p, 48084p,
48086r, 57111x, 57433d, 57816f, 57817g, 57927t,
58338p, 59102u, 59151j, 61543g, 61575u, 69037r,
69215x.

See also 2103, 2105, 2129, 2157, 2275, 2288, 2299, 2312, 2576, 2783,
2920, 2921, 2935, 3119, 3121, 3125.

3d. Quantitative analysis

- 2230 Vigdergauz, M.S. and Krauze, I.M.: (Quantitative interpretation of chromatograms using the product of peak height and retention time). *Zh. Anal. Khim.*, 44 (1989) 1080-1084; *C.A.*, 111 (1989) 247270a.

For additional information see:

C.A., 112 (1990) 15770f.

See also 3102.

3e. Preparative scale chromatography

- 2231 Anonymous.: (High-resolution liquid chromatography on a preparative scale.) *Tec. Lab.*, 12 (1989) 18-21; *C.A.*, 111 (1989) 193744n.
- 2232 Jacobson, J.M. and Frenz, J.: Determination of competitive adsorption isotherms for modeling large-scale separations in liquid chromatography. *J. Chromatogr.*, 499 (1990) 5-19.

- 2233 Katti, A.M. and Guiochon, G.: Quantitative comparison between the experimental band profiles of binary mixtures in overloaded elution chromatography and their profiles predicted by the semi-ideal model. *J. Chromatogr.*, 499 (1990) 21-35.
- 2234 Kowalczyk, J.S.: (The purity of compounds isolated by preparative liquid chromatography.) *GIT-Suppl.*, No. 3 (1989) 7-12; *C.A.*, 111 (1989) 173171v - a review with 23 refs.
- 2235 Sada, E., Kato, S., Yamanaka, K. and Sohma, Y.: Effects of antigen molecular weight on immunoadsorption equilibrium in immunoaffinity chromatography. *J. Ferment. Bioeng.*, 68 (1989) 37-39; *C.A.*, 111 (1989) 170486r.
- 2236 Speckenbach, F.: (Preparative medium pressure chromatography.) *GIT-Suppl.*, No. 3 (1989) 69-78; *C.A.*, 111 (1989) 173170u - a review with 22 refs.
- 2237 Werner, A.: Chromatographische Enantiomerentrennung an Cellulosetriacetat in Grammatstab. *Kontakte (Darmstadt)*, No. 3 (1989) 50-56.

See also 2142, 2258, 2274, 2481, 2587, 2605, 2637, 2778.

3f. Programmed temperature, pressure, vapors, gradients

- 2238 Cole, L.A. and Dorsey, J.G.: Reduction of the reequilibration time following gradient elution reversed-phase liquid chromatography. *Anal. Chem.*, 62 (1990) 16-21.

See also 2913.

3g. High performance procedures

See 2204, 3124.

4. SPECIAL TECHNIQUES

4a. Automation and computerization

- 2239 Nasyrov, M.S. and Rasskazova, R.B.: (Algorithm for determining the boundary points of individual chromatogram peaks). *Zavod. Lab.*, 55 (1989) 18-22; *C.A.*, 112 (1990) 15647w.

For additional information see:

C.A., 112 (1990) 15764g, 15767k, 60079f.

See also 2096, 2111, 2112, 2145, 2162, 2187, 2247, 2309, 2428, 2455, 2564, 2883, 3120.

4b. Combination of various chromatographic techniques

- 2240 Niyredy, S., Dallenbach-Toelke, K., Zogg, G.C. and Sticher, O.: Strategies of mobile phase transfer from thin-layer to medium pressure liquid chromatography with silica as the stationary phase. *J. Chromatogr.*, 499 (1990) 453-462.
- 2241 Schmarr, H.-G., Mosandl, A. and Grob, K.: Coupled LC-GC: Evaporation rates for partially concurrent eluent evaporation using an early solvent vapor exit. *J. High Resolut. Chromatogr.*, 12 (1989) 721-726.
- 2242 Vreuls, J.J., Cuppen, W.J.G.M., Dolecka, E., Maris, F.A., de Jong, G.J. and Brinkman, U.A.T.: Polar solvents for the desorption of a liquid chromatographic precolumn coupled on-line with a capillary gas chromatograph. *J. High Resolut. Chromatogr.*, 12 (1989) 807-812.

See also 2455, 2564, 2678.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 2243 Basset, F.: (New developments in mass spectrometry and combined techniques). *Inf. Chim.*, 302 (1989) 151-156; *C.A.*, 111 (1989) 224156d - a review with no refs.
- 2244 Fraser, D.J.J., Norton, K.L. and Griffiths, P.R.: HPLC/FT-IR measurements by transmission, reflection-absorption, and diffuse reflection microscopy. *Pract. Spectrosc.*, 6 (1988) 197-210; *C.A.*, 111 (1989) 247223n.
- 2245 Goates, S.R. and Sin, C.H.: Supersonic jet spectroscopy coupled to chromatography for very high resolution chemical analysis. *Appl. Spectrosc. Rev.*, 25 (1989) 82-126; *C.A.*, 112 (1990) 47977v - a review with 103 refs.
- 2246 Hofstraet, J.W., Gooijer, C., Brinkman, U.A.T. and Velthorst, N.H.: Fluorescence line-narrowing spectroscopy: a new and highly selective detection technique for thin-layer and liquid chromatography. In: Dallas, F.A.A. (Editor), *Recent Adv. Thin-Layer Chromatogr. (Proc. Chromatogr. Soc. Int. Symp.) 1987*, Plenum, New York, 1988, pp. 29-36; *C.A.*, 112 (1990) 69149d.
- 2247 Jinno, K. and Fujimoto, C.: Solvent elimination approach for microcolumn liquid chromatography-infrared spectroscopy: buffer-memory technique. *Prog. HPLC*, 4 (1989) 273-296; *C.A.*, 112 (1990) 48028y.
- 2248 Jinno, K.: Practice and application of microcolumn LC with inductively coupled plasma atomic emission spectrometric detection. *Chromatogr. Sci.*, 45 (1989) 175-210; *C.A.*, 111 (1989) 224165f.
- 2249 Oxford, J. and Lant, M.S.: LCMS - its role in drug metabolism. *Chem. Br.*, 25 (1989) 889-898; *C.A.*, 111 (1989) 208525t - a review with 22 refs.
- 2250 Paryjczak, T., Kocemba, I. and Karski, S.: (Study of nickel oxide reduction in a chromatographic system with simultaneous electric conductivity measurements). *Przem. Chem.*, 68 (1989) 259-261; *C.A.*, 111 (1989) 203886b.
- 2251 Robins, R.H. and Crow, F.W.: Factor affecting high-mass sensitivity and ion-current stability in a thermospray source. *Rapid Commun. Mass Spectrom.*, 2 (1988) 30-34; *C.A.*, 112 (1990) 30036q.
- 2252 Roboz, J., Holland, J.F., McDowell, M.A. and Hillmer, M.J.: Accurate mass measurement in continuous flow fast atom bombardment quadrupole mass spectrometry. *Rapid Commun. Mass Spectrom.*, 2 (1988) 64-66; *C.A.*, 112 (1990) 47991v.
- 2253 Ross, M.S.F., Leisten, O. and Bar-Shad, Y.: Real-time validation of analytical results in HPLC using a rapid scan UV-VIS detector. *Am. Lab. (Fairfield)*, 21 (1989) 64-73; *C.A.*, 112 (1990) 69091d.
- 2254 Salzer, R. and Meyer, V.: (Combination of chromatography and IR spectroscopy). *Teubner-Texte Phys.*, 20 (1988) 39-53; *C.A.*, 112 (1990) 47983u - a review with 42 refs.

For additional information see:

C.A., 111 (1989) 224257n;
112 (1990) 30093f, 69082b, 69212u.

See also 2160, 2161, 2278, 2333, 2515, 2830, 2999, 3082, 3101.

4d. Affinity chromatography

- 2255 Birkenmeier, G. and Kopperschlaeger, G.: (New affinity chromatography techniques for down-stream processing.) *Wiss. Z. - Karl-Marx-Univ. Leipzig, Math.-Naturwiss. Reihe*, 38 (1989) 349-356; *C.A.*, 111 (1989) 172348w - a review with 19 refs.
- 2256 Gordon, J., Thompson, R.E. and Kapsalis, A.A.: Method for the study of steady-state and non-steady state kinetics of immobilized enzymes. *Eur. Pat. Appl.* EP 315,058 (Cl. C12Q1/00), 10 May 1989, US Appl. 117,278, 05 Nov. 1987; 12 pp.; *C.A.*, 111 (1989) 227989p.

- 2257 Liapis, A.I.: Theoretical aspects of affinity chromatography. *J. Biotechnol.*, 11 (1989) 143-160; *C.A.*, 111 (1989) 190542j - a review with 58 refs.
- 2258 Narayanan, S.R., Knoch, S., Jr. and Crane, L.J.: Preparative affinity chromatography of proteins. Influence of the physical properties of the base matrix. *J. Chromatogr.*, 503 (1990) 93-102.
- 2259 Ochoa, J.L.: Applications of the lectin-glycoprotein biospecific interaction on immobilized systems. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (1989) 293-301; *C.A.*, 111 (1989) 170257s - a review with 22 refs.
- 2260 Sulkowski, E.: The saga of IMAC and MIT. *BioEssays*, 10 (1989) 170-175; *C.A.*, 111 (1989) 170256r - a review with 40 refs.
- 2261 Turkova, J., Petkov, L., Sajdok, J., Kas, J. and Benes, M.J.: Carbohydrates as a tool for oriented immobilization of antigens and antibodies. *J. Chromatogr.*, 500 (1990) 585-593.
- 2262 Varghese, G. and Diwan, A.M.: Electrophoretic desorption of concanavalin A from an affinity matrix: a rapid method for the preparation of carbohydrate free concanavalin A. *Biotechnol. Tech.*, 3 (1989) 233-234; *C.A.*, 111 (1989) 190889c.
- 2263 Velander, W.H., Orthner, C.L., Tharakan, J.P., Maduraware, R.D., Ralston, A.H., Strickland, D.K. and Drohan, W.N.: Process implications for metal-dependent immunoaffinity interactions. *Biotechnol. Prog.*, 5 (1989) 119-125; *C.A.*, 112 (1990) 62413b.
- 2264 Wilchek, M. and Bayer, E.A.: A universal affinity column using avidin-biotin technology. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (1989) 83-90; *C.A.*, 111 (1989) 170326p.

For additional information see:

C.A., 111 (1989) 190970x, 191016c, 191023c;
112 (1990) 47980r.

See also 2323, 2358, 2421, 2425, 2519, 2548, 2569, 2580, 2590, 2593, 2613, 2614, 2617, 2621, 2628, 2637, 2667, 2670, 2675, 2680, 2682, 2712, 2722, 2723, 2724, 2752, 3095.

4f. Trace analysis and pre separation techniques

- 2265 Carel, A.B. and Thompson, J.C.: Sample pretreatment for ion chromatography. *Lab. Pract.*, 38 (1989) 77-79; *C.A.*, 112 (1990) 47614z - a review with 4 refs.
- 2266 Fowles, I.A.: Solute focusing in LC by on-line dilution. *J. High Resolut. Chromatogr.*, 13 (1990) 138-141.
- 2267 Grazhulene, S.S., Popandopulo, Yu.I., Chapligina, N.I. and Telegin, G.F.: (Extraction-chromatographic separation of trace impurities from zinc and its high-purity chalcogenides). *Zh. Anal. Khim.*, 44 (1989) 444-449; *C.A.*, 111 (1989) 224264n.

For additional information see:

C.A., 111 (1989) 246965u, 246987c;
112 (1990) 15728y, 47824t, 68799d.

See also 2218, 2937, 3100, 3115, 3116, 3122, 3128, 3131, 3132, 3133.

4g. Separation of enantiomers

- 2268 Allenmark, S. and Andersson, S.: Optical resolution of some biologically active compounds by chiral liquid chromatography on BSA-silica (Resolvosil) columns. *Chirality*, 1 (1989) 154-160; *C.A.*, 111 (1989) 239629j.

- 2269 Belov, Yu.P., Sidorov, A.P. and Martynov, I.V.: (Ligand exchange chromatography of amino acid enantiomers on achiral phases). *Dokl. Akad. Nauk SSSR*, 308 (1989) 632-634; *C.A.*, 112 (1990) 69207w.
- 2270 Cabrera, K. and Schwinn, G.: Cyclodextrine als chirale Selektoren für die Enantiomerentrennung in der HPLC (Teil 1): Dynamische Belegung von RP-Phasen. *Kontakte (Darmstadt)*, No. 3 (1989) 3-8.
- 2271 Kinkel, J.N., Reichert, K. and Knoell, P.: Preparative liquid chromatographic separation of enantiomers. Microcrystalline cellulose triacetate and chiral polyacrylamide/silica compounds materials. *GIT-Suppl.*, No. 3 (1989) 104-112; *C.A.*, 111 (1989) 213941e - a review with 40 refs.
- 2272 Lough, W.J. (Editor): *Chiral Liquid Chromatography*. Blackie, Glasgow, 1989, 276 p.; *C.A.*, 112 (1990) 42751g.
- 2273 Meyer, V.R.: Chromatographic separation of enantiomers: the phenomenon of chirality. *Pharm. Unserer Zeit*, 18 (1989) 140-145; *C.A.*, 112 (1990) 3668r - a review with 12 refs.
- 2274 Perry, J.A. and Rateike, J.D.: Product purity as a function of elution order. *J. Chromatogr.*, 503 (1990) 403-409.
- 2275 Selligren, B.: Molecular imprinting by noncovalent interactions: tailor-made chiral stationary phases of high selectivity and sample load capacity. *Chirality*, 1 (1989) 63-68; *C.A.*, 112 (1990) 69106n.
- 2276 Tambute, A., Canceill, J. and Collet, A.: Optical resolution of C3 cyclotrivenatrylenes and D3 cryptophanes by liquid chromatography on chiral stationary phase Chiralpak-OT(+). *Bull. Chem. Soc. Jpn.*, 62 (1989) 1390-1392; *C.A.*, 111 (1989) 232256b.
- 2277 Wiesner, R. and Kuehn, H.: (Enantiomer separation of hydroxy polyenoic fatty acids on PIRKLE-type chiral phases). *GIT-Suppl.*, (1989) 32-38; *C.A.*, 112 (1990) 48045b.

For additional information see:

C.A., 112 (1990) 25776b, 47988z.

See also 2182, 2197, 2237, 2301, 2321, 2368, 2395, 2454, 2485, 2486, 2933, 2942, 2951, 2954, 2960, 2967, 2993, 3005, 3010, 3054, 3066.

4h. Other special techniques

- 2278 Aerts, M.M.L., Beek, W.M.J. and Brinkman, U.A.T.: On-line combination of dialysis and column-switching liquid chromatography as a fully automated sample preparation technique for biological samples. Determination of nitrofurans residues in edible products. *J. Chromatogr.*, 500 (1990) 453-468.
- 2279 Fish, B.B. and Carr, R.W.: An experimental study of the countercurrent moving-bed chromatographic reactor. *Chem. Eng. Sci.*, 44 (1989) 1773-1783; *C.A.*, 111 (1989) 216575z.
- 2280 Idei, M., Pajor, A. and Menyhart, J.: Bioseparation: application of isolated organs in the preparative scale isolation of biologically active compounds. *J. Lig. Chromatogr.*, 12 (1989) 2769-2791.
- 2281 Okada, T.: Ion-exclusion chromatography with mixed micellar mobile phases. *Anal. Chim. Acta*, 230 (1990) 9-15.
- 2282 Vigh, G., Irgens, L.H. and Farkas, G.: Displacement chromatography on packed capillary columns. *J. Chromatogr.*, 502 (1990) 11-19.

See also 2205, 3135.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

- 2283 Kuningas, K., Rang, S., Kailas, T., Korol, A. and Dubovik, M.: (Interlaboratory reproducibility of the retention characteristics of *n*-alkenes and *n*-alkynes on capillary columns). *Eesti NSV Tead. Akad. Toim., Keem.*, 38 (1989) 159-166; *C.A.*, 112 (1990) 48046c.

5b. Cyclic hydrocarbons

- 2284 Beltra, F.J. and Beltran de Heredia, J.: (Application of high resolution liquid chromatography to the analysis of cumene oxidation products). *Afinidad*, 46 (1989) 185-187; *C.A.*, 112 (1990) 48062e.
- 2285 Brodfuehrer, J.I., Chapman, D.E., Wilke, T.J. and Powis, G.: Comparative studies of the *in vitro* metabolism and covalent binding of ¹⁴C-benzene by liver slices and microsomal fraction of mouse, rat, and human. *Drug Metab. Disp.*, 18 (1990) 20-27.
- 2286 Li, S.F.Y., Lee, H.K. and Ong, C.P.: Retention behavior of PAH in reversed-phase HPLC. *J. Liq. Chromatogr.*, 12 (1989) 3251-3264.
- 2287 Shatz, V.D. and Sahartova, O.V.: HPLC of benzene derivatives on Silasorb 600 dynamically modified by cetrinide. *J. Liq. Chromatogr.*, 12 (1989) 2801-2819.
- 2288 Tanaka, N., Hosoya, K., Tachibana, Y., Araki, M., Tanaka, K. and Kaji, A.: Selectivity of 2-(1-pyrenyl)ethylsilylated silica gel in the isomer separation of cyclohexane derivatives. *J. Chromatogr. Sci.*, 27 (1989) 735-740.

For additional information see:

C.A., 111 (1989) 231208a;
112 (1990) 39582x.

See also 2168, 2176, 2209, 2276, 2807.

5c. Halogen derivatives

For additional information see:

C.A., 111 (1989) 235589y.

See also 2883.

5d. Complex hydrocarbon mixtures

- 2289 Aleksandrov, I.V., Povalyaev, A.N., Rogova, L.P. and Shcheblanova, T.N.: (Molecular weight distribution and properties of extracts of gas coals). *Khim. Tverd. Topl. (Moscow)*, (1989) 21-24; *C.A.*, 112 (1990) 9656g.
- 2290 Gey, M. and Riis, V.: (Group type analysis of complex hydrocarbon mixtures by HPLC. Part 2. Petroleum fractions of different boiling ranges, provenience and processing). *Chem. Tech. (Leipzig)*, 41 (1989) 435-437; *C.A.*, 112 (1990) 9590f.
- 2291 Kithinji, J.P., Raynor, M.W., Egia, B., Davies, I.L., Bartle, K.D. and Clifford, A.A.: Analysis of coal tar polycyclic aromatic hydrocarbon LC-fractions by capillary SFC on a liquid crystalline stationary phase. *J. High Resolut. Chromatogr.*, 13 (1990) 27-33.

See also 2798.

6. ALCOHOLS

- 2292 Gupta, R.N.: Determination of trichloroethanol, the active metabolite of chloral hydrate, in plasma by liquid chromatography. *J. Chromatogr.*, 500 (1990) 655-659.

For additional information see:

C.A., 111 (1989) 247244v.

See also 2274, 2288.

7. PHENOLS

- 2293 Fischer, I.U. and Dengler, H.J.: Sensitive high-performance liquid chromatographic assay for the determination of eugenol in body fluids. *J. Chromatogr.*, 525 (1990) 369-377.
- 2294 Lanin, S.N. and Nikitin, Yu.S.: Normal-phase high-performance liquid chromatographic determination of phenol. *Talanta*, 36 (1989) 573-579; C.A., 112 (1990) 69114p.
- 2295 Lauterbach, M., Jancke, H., Jaeger, G., Zimmermann, V. and Radig, W.: (Advances in the field of analysis of phenol mixtures). *Freiberg. Forschungsh. A*, 781 (1989) 44-58; C.A., 111 (1989) 247190z - a review with 16 refs.
- 2296 Scalbert, A., Duval, L., Peng, S., Monties, B. and du Penhoat, C.: Polyphenols of *Quercus robur* L. II. Preparative isolation by low-pressure and high-pressure liquid chromatography of heartwood ellagitannins. *J. Chromatogr.*, 502 (1990) 107-119.
- 2297 Scott, D.O., Bell, M.A. and Lunte, C.E.: Microdialysis-perfusion sampling for the investigation of phenol metabolism. *J. Pharm. Biomed. Anal.*, 7 (1989) 1249-1259.
- 2298 Vincze, A. and Huszar, M.L.: Determination of 2-naphthol and its derivatives in the presence of each other by HPLC. *J. Liq. Chromatogr.*, 12 (1989) 2887-2892.
- 2299 Yamaguchi, J. and Hanai, T.: Selectivity related to carbon loading and endcapping of octadecyl bonded silica gels in the reversed-phase liquid chromatography of phenolic compounds. *J. Chromatogr. Sci.*, 27 (1989) 710-715.
- 2300 Zul'figarov, O.S., Yurchenko, V.V. and Pilipenko, A.T.: (Reaction of diazo cations with azophenols). *Ukr. Khim. Zh.*, 55 (1989) 265-267; C.A., 112 (1990) 48055e.

For additional information see:

C.A., 111 (1989) 230795c.

See also 2137, 2150, 2282, 2807.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 2301 Krause, M. and Galensa, R.: Improved chiral stationary phase based on cellulose triacetate supported on non-macroporous silica gel diol for the high-performance liquid chromatographic separation of racemic flavanones and diastereomeric flavanone glycosides. *J. Chromatogr.*, 502 (1990) 287-296.

- 2302 Steinhilber, B. and Engelhardt, U.H.: Theaflavins in black tea. Comparison of the flavonoid and HPLC analysis. Preliminary results. *Z. Lebensm.-Unters. Forsch.*, 188 (1989) 509-511; *C.A.*, 111 (1989) 193137s.
- 2303 Wang, G., Kuan, S.S., Francis, O.J., Ware, G.M. and Carman, A.S.: A simplified HPLC method for the determination of phytoestrogens in soybean and its processed products. *J. Agric. Food Chem.*, 38 (1990) 185-190.

See also 2687, 3073, 3076.

8b. Aflatoxins and other mycotoxins

- 2304 Dorner, J.W. and Cole, R.J.: Comparison of two ELISA screening tests with liquid chromatography for determination of aflatoxins in raw peanuts. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 962-964.
- 2305 Nagashima, Y., Noguchi, T. and Hashimoto, K.: Detection of paralytic shellfish poisons by HPLC. *Bioact. Mol.*, 10 (Mycotoxins, Phycotoxins '88) (1989) 311-318; *C.A.*, 112 (1990) 6212y.
- 2306 Sydenham, E.W., Gelderblom, W.C.A., Thiel, P.G. and Marasas, W.F.O.: Evidence for the natural occurrence of fumonisin B₁, a mycotoxin produced by *Fusarium moniliforme*, in corn. *J. Agric. Food Chem.*, 38 (1990) 285-290.
- 2307 Tanaka, T. and Ueno, Y.: Worldwide natural occurrence of *Fusarium* mycotoxins, nivalenol, deoxynivalenol and zearalenone. *Bioact. Mol.*, 10 (Mycotoxins, Phycotoxins '88) (1989) 51-56; *C.A.*, 111 (1989) 230805f.
- 2308 Tresel, R., Hommel, E. and Helak, B.: (Determination of ochratoxin A in barley, malt, and beer by high pressure liquid chromatography with fluorescence detection.) *Monatsschr. Brauwiss.*, 42 (1989) 331-335; *C.A.*, 111 (1989) 172370x.
- 2309 Tuinstra, L.G.M.T., Kienhuis, P.G.M., Traag, W.A., Aerts, M.M.L. and Beek, W.M.J.: Fully automated column-switching HPLC determination of aflatoxin M₁ in milk using dialysis as sample preparation. *J. High Resolut. Chromatogr.*, 12 (1989) 709-713.
- 2310 Tutelyan, V.A., Eller, K.I. and Sobolev, V.S.: A survey using normal-phase high-performance liquid chromatography of aflatoxins in domestic and imported foods in the USSR. *Food Addit. Contam.*, 6 (1989) 459-465; *C.A.*, 112 (1990) 6256r.
- 2311 Ware, G.M., Francis, O.J., Kuan, S.S. and Carman, A.S.: Determination of zearalenol and zearalenone using electrochemical detection. *Anal. Lett.*, 22 (1989) 2335-2352.

For additional information see:

C.A., 112 (1990) 6193t.

8c. Other compounds with heterocyclic oxygen (including tannins)

- 2312 Kimata, K., Hosoya, K., Tanaka, N., Araki, T., Barnhart, E.R., Patterson, D.G., Jr. and Terabe, S.: A newly developed stationary phase, nitrated phenylethyl silica gel, and HPLC separation of 1,2,4,6-/1,2,4,9-TCDD isomer pair. *J. High Resolut. Chromatogr.*, 13 (1990) 137-138.
- 2313 Rizzolo, A., Nani, R. and Picariello, M.: (HPLC determination of HMF, chlorogenic acid and caffeic acid in clear fruit juices). *Ind. Bevande*, 18 (1989) 183-186; *C.A.*, 112 (1990) 6203w.

For additional information see:

C.A., 112 (1990) 42553u.

See also 2432, 3075.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 2314 Benassi, C.A., Semenzato, A., Zaccaria, F. and Bettero, A.: High-performance liquid chromatographic determination of free formaldehyde in cosmetics preserved with Diwiciil 200. *J. Chromatogr.*, 502 (1990) 193-200.
- 2315 Bittencourt de Andrade, J., Bravo, Ruiess van Fossen: Analysis of aldehydes in alcoholic drinks: a high-performance liquid chromatographic experiment for graduate courses. *Quim. Nova*, 12 (1989) 196-198; *C.A.*, 111 (1989) 231385f.
- 2316 Katayama, M., Mukai, Y. and Taniguchi, H.: High-performance liquid chromatographic determination of cinnamaldehyde. *Analyst (London)*, 115 (1990) 9-11.
- 2317 Matsushita, S., Terao, J., Yamada, K. and Shibata, S.S.: Specific detection of lipid hydroperoxides using HPLC-EC methods. *Basic Life Sci.*, 49 (Oxygen Radicals Biol. Med.) (1988) 163-174; *C.A.*, 112 (1990) 3761r.
- 2318 Miksik, I., Struzinsky, R., Macek, K. and Deyl, Z.: Liquid column chromatography of oxo compounds as their 2,4-dinitrophenylhydrazones with special reference to glucose-glycine Maillard reaction products. *J. Chromatogr.*, 500 (1990) 607-613.
- 2319 Miyazawa, T.: (Picomole level microanalysis of biological lipid hydroperoxides development and application of a chemiluminescence-high performance liquid chromatography). *Yukagaku*, 38 (1989) 800-808; *C.A.*, 112 (1990) 3699b - a review with 22 refs.
- 2320 Schäfer-Lüderßen, U. and Mauss, M.: Determination of free formaldehyde in nonwovens in the presence of glyoxalic acid and aldehydes. *Chromatographia*, 29 (1990) 21-23.

For additional information see:

C.A., 111 (1989) 219171a;
112 (1990) 25764w.

See also 2149, 2298, 2415, 2629.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 2321 Armstrong, D.W. and Jin, H.L.: Liquid chromatographic separation of anomeric forms of saccharides with cyclodextrin bonded phases. *Chirality*, 1 (1989) 27-37; *C.A.*, 111 (1989) 233385e.
- 2322 Basa, L.J. and Spellman, M.W.: Analysis of glycoprotein-derived oligosaccharides by high-pH anion-exchange chromatography. *J. Chromatogr.*, 499 (1990) 205-220.
- 2323 Bierhuizen, M.F.A., Hansson, M., Odin, F., Debray, H., Oebrick, B. and van Dijk, W.: Structural assesment of the N-linked oligosaccharides of cell-CAM 105 by lectin-agarose affinity chromatography. *Glycoconjugate J.*, 6 (1989) 195-208; *C.A.*, 111 (1989) 189872k.
- 2324 Biffen, M. and Hanke, D.E.: Reduction in the level of intracellular myo-inositol in cultured soybean (*Glycine max.*) cells inhibits cell division. *Biochem. J.*, 265 (1990) 809-814.
- 2325 Churms, S.C.: Recent developments in the chromatographic analysis of carbohydrates. *J. Chromatogr.*, 500 (1990) 555-583 - a review with 228 refs.
- 2326 Kuptsevich, Yu.E., Larionov, O.G. and Pronin, A.Ya.: (Chromatographic separation of glucose and fructose on calcium-type cation exchangers). *Zh. Fiz. Khim.*, 63 (1989) 2719-2721; *C.A.*, 112 (1990) 15780j.

- 2327 LaCourse, W.R., Mead, D.A., Jr. and Johnson, D.C.: Anion-exchange separation of carbohydrates with pulsed amperometric detection using a pH-selective reference electrode. *Anal. Chem.*, 62 (1990) 220-224.
- 2328 Martens, D.A. and Frankenberger, W.T., Jr.: Determination of saccharides by high performance anion-exchange chromatography with pulsed amperometric detection. *Chromatographia*, 29 (1990) 7-12.
- 2329 Murata, K., Yokoyama, Y. and Yoshida, K.: Heparitinases facilitate separation of disaccharides of heparan sulfate isomers in human arteries using high-performance liquid chromatography. *Biochem. Int.*, 19 (1989) 155-162; *C.A.*, 111 (1989) 170302c.
- 2330 Muto, N., Nakamura, T. and Yamamoto, I.: Enzymatic formation of a nonreducing L-ascorbic acid α -glucoside: purification and properties of α -glucosidases catalyzing site-specific transglucosylation from rat small intestine. *J. Biochem. (Tokyo)*, 107 (1990) 222-227.
- 2331 Oku, H., Hase, S. and Ikenaka, T.: Separation of oligomannose-type sugar chains having one to five mannose residues by high-performance liquid chromatography as their pyridylamino derivatives. *Anal. Biochem.*, 185 (1990) 331-334.
- 2332 Prabhu, S.V. and Baldwin, R.P.: High-pH ion-exchange separation and electrochemical detection of alditols, carbohydrates and acidic sugars. *J. Chromatogr.*, 503 (1990) 227-235.
- 2333 Simpson, R.C., Fenselau, C.C., Hardy, M.R., Townsend, R.R., Lee, Y.C. and Cotter, R.J.: Adaptation of a thermospray liquid chromatography/mass spectrometry interface for use with alkaline anion exchange liquid chromatography of carbohydrates. *Anal. Chem.*, 62 (1990) 248-252.
- 2334 Thomas, R.L., Abbas, S.A., Piskorz, C.F. and Matta, K.L.: Separation by liquid chromatography (under elevated pressure) of benzyl and nitrophenyl glycosides of oligosaccharides. *Carbohydr. Res.*, 189 (1989) 13-19; *C.A.*, 112 (1990) 7806g.
- 2335 van Pelt, J., Hokke, C.H., Dorland, L., Duran, M., Kamerling, J.P. and Vliegthart, J.F.G.: Accumulation of mannosyl-beta(1 $\rightarrow$ 4)-N-acetylglucosamine in fibroblasts and leukocytes of patients with a deficiency of beta-mannosidase. *Clin. Chim. Acta*, 187 (1990) 55-60.
- 2336 Wang, W.T. and Zopf, D.: Liquid ion-exchange chromatography under pressure of milk oligosaccharides using a pulsed amperometric detector. *Carbohydr. Res.*, 189 (1989) 1-11; *C.A.*, 112 (1990) 7805f.
- 2337 Yokoyama, K. and Ballou, C.E.: Synthesis of α 1 $\rightarrow$ 6-mannooligosaccharides *Mycobacterium smegmatis*. Function of β -mannosylphosphoryldecaprenol as the mannosyl donor. *J. Biol. Chem.*, 264 (1989) 21621-21628.
- 2338 Zaton, A., Quindos, G. and Ponton, J.: Determination of monosaccharides by high-performance liquid chromatography in systemic candidosis. *J. Chromatogr.*, 525 (1990) 169-175.

For additional information see:

C.A., 111 (1989) 174553h, 190615k, 190996k, 228556g.

See also 2313, 2348, 2361, 2738.

10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 2339 Brun, P., de Galateo, A., Camporese, A., Cortivo, R. and Abatangelo, G.: Analysis of hyaluronic acid in synovial fluid by reversed-phase liquid chromatography. *J. Chromatogr.*, 526 (1990) 530-534.
- 2340 Burg, M. and Cole, G.J.: Characterization of cell-associated proteoglycans synthesized by embryonic neural retinal cells. *Arch. Biochem. Biophys.*, 276 (1990) 396-404.

- 2341 Hotchkiss, A.T., Jr. and Hicks, K.B.: Analysis of oligogalacturonic acids with 50 or fewer residues by high-performance anion-exchange chromatography and pulsed amperometric detection. *Anal. Biochem.*, 184 (1990) 200-206.
- 2342 Hronowski, L.J.J. and Anastassiades, T.P.: Non-specific interaction of proteoglycans with surfaces and matrices. *Biochem. Biophys. Res. Commun.*, 167 (1990) 81-88.
- 2343 Kiho, T., Ito, M., Yoshida, I., Nagai, K., Hara, C. and Ukai, S.: Polysaccharides in fungi. XXIV. A(1→3)- β -D-glucan from the alkaline extract of the insect-body portion of *Chan huà* (fungus: *Cordyceps cicadae*). *Chem. Pharm. Bull.*, 37 (1989) 2770-2772.
- 2344 Kolset, S.O., Sakurai, K., Ivhed, I., Qvervatn, A. and Suzuki, S.: The effect of β -D-xylosides on the proliferation and proteoglycan biosynthesis of monoblastic U-937 cells. *Biochem. J.*, 265 (1990) 637-645.
- 2345 Ohtani, K., Hatano, S., Mizutani, K., Kasai, R. and Tanaka, O.: Reticuloendothelial system-activating polysaccharides from rhizomes of *Panax japonicus*. I. Tochibanan-A and -B. *Chem. Pharm. Bull.*, 37 (1989) 2587-2591.
- 2346 Takeuchi, T., Murayama, M. and Ishii, D.: Indirect detection of cyclodextrins in micro-HPLC. *J. High Resolut. Chromatogr.*, 13 (1990) 69-70.
- 2347 Takeuchi, Y., Matsumoto, T., Ogata, E. and Shishiba, Y.: Isolation and characterization of proteoglycans synthesized by mouse osteoblastic cells in culture during the mineralization process. *Biochem. J.*, 266 (1990) 15-24.
- 2348 Turnbull, J.E. and Gallagher, J.T.: Molecular organization of heparan sulphate from human skin fibroblasts. *Biochem. J.*, 265 (1990) 715-724.

For additional information see:

C.A., 112 (1990) 38519v, 42692p.

See also 2329, 2352.

10c. Glycoproteins and their components

- 2349 Bom, V.J.J. and Bertina, R.M.: The contributions of Ca^{2+} , phospholipids and tissue-factor apoprotein to the activation of human blood-coagulation factor X by activated factor VII. *Biochem. J.*, 265 (1990) 327-336.
- 2350 Imai, N., Kawamura, A., Higuchi, M., Oh-eda, M., Orita, T., Kawaguchi, T. and Ochi, N.: Physicochemical and biological comparison of recombinant human erythropoietin with human urinary erythropoietin. *J. Biochem. (Tokyo)*, 107 (1990) 352-359.
- 2351 Johe, K.K., Smith, A.J., Vengelen-Tyler, V. and Blumenfeld, O.O.: Amino acid sequence of an α - δ -glycophorin hybrid. A structure reciprocal to St^a δ - α -glycophorin hybrid. *J. Biol. Chem.*, 264 (1989) 17486-17493.
- 2352 Karamanos, N.K., Wikström, B., Antonopoulos, C.A. and Hjerpe, A.: Determination of N-acetyl- and N-glycolylneuraminic acids in glycoconjugates by reversed-phase high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr.*, 503 (1990) 421-429.
- 2353 Kumazaki, T., Hoshiba, N., Yokosawa, H. and Ishii, S.-i.: Primary structure of ascidian trypsin inhibitors in the hemolymph of a solitary ascidian, *Halocynthia roretzi*. *J. Biochem. (Tokyo)*, 107 (1990) 409-413.
- 2354 Malm, J., Cohen, E., Dackowski, W., Dahlbäck, B. and Wydro, R.: Expression of completely γ -carboxylated and β -hydroxylated recombinant human vitamin-K-dependent protein S with full biological activity. *Eur. J. Biochem.*, 187 (1990) 737-743.
- 2355 Nakano, M., Hatanaka, Y., Kobayashi, N., Noguchi, S., Ishikawa, S. and Tobita, T.: Further fractionation of the glycoprotein families of porcine zona pellucida by anion-exchange HPLC and some characterization of the separated fractions. *J. Biochem. (Tokyo)*, 107 (1990) 144-150.

- 2356 Olechno, J.D., Carter, S.R., Edwards, W.T., Gillen, D.G., Townsend, R.R., Lee, Y.C. and Hardy, M.R.: Chromatographic analysis of glycoprotein derived carbohydrates; pellicular ion exchange resins and pulsed electrochemical detection. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 364-376; *C.A.*, 111 (1989) 228250c.
- 2357 Rice, K.G., Rao, N.B.N. and Lee, Y.C.: Large-scale preparation and characterization of N-linked glycopeptides from bovine fetuin. *Anal. Biochem.*, 184 (1990) 249-258.
- 2358 Sawhney, S.Y. and Bhide, S.V.: Immobilized mucin: an affinity matrix for the isolation of winged bean acidic and basic lectins. *J. Chromatogr.*, 503 (1990) 272-276.
- 2359 Shimamura, M., Ohara, T., Imazawa, M. and Miyamoto, K.: Purification of myelin-associated glycoprotein from calf brain using high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 535-539.
- 2360 Wang, T., Chen, T. and Barofsky, D.F.: Reversed phase high performance liquid chromatography of glycopeptides. *Biomed. Chromatogr.*, 3 (1989) 241-245.
- 2361 Zamze, S.E., Wooten, E.W., Ashford, D.A., Ferguson, M.A.J., Dwek, R.A. and Rademacher, T.W.: Characterisation of the asparagine-linked oligosaccharides from *Trypanosoma brucei* type-I variant surface glycoproteins. *Eur. J. Biochem.*, 187 (1990) 657-663.
- 2362 Zelikman, I., Aakerblom, L., Hjerten, S. and Morein, B.: Free gp70 from FeLV: enrichment from cell culture fluid by ferric oxide-agarose chromatography. *Biotechnol. Appl. Biochem.*, 11 (1989) 209-216; *C.A.*, 111 (1989) 190613h.
- 2363 Zhu, B.C.-R. and Laine, R.A.: Purification of acetyllactosamine-specific tomato lectin by erythroglycan-Sepharose affinity chromatography. *Prepar. Biochem.*, 19 (1989) 341-350.
- 2364 Zhu, B.C.-R. and Laine, R.A.: Novel hyperglycosylated weak gelatin-binding fibronectin from human fetal placenta. Fractionation of a high poly(N-acetyllactosamine) fragment by tomato lectin affinity chromatography. *Eur. J. Biochem.*, 188 (1990) 67-71.

See also 2658, 2728.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 2365 Allenmark, S., Chelminska-Bertilsson, M. and Thompson, R.A.: N-(9-Acridinyl)-bromoacetamide - a powerful reagent for phase-transfer-catalyzed fluorescence labeling of carboxylic acids for liquid chromatography. *Anal. Biochem.*, 185 (1990) 279-285.
- 2366 Baer, A.N., Costello, P.B. and Green, F.A.: Free and esterified 13(R,S)-hydroxyoctadecadienoic acids: principal oxygenase products in psoriatic skin scales. *J. Lipid Res.*, 31 (1990) 125-130.
- 2367 Blanchard, J.R.T. and Tasker, R.A.R.: High-performance liquid chromatographic assay for domoic acid in serum of different species. *J. Chromatogr.*, 526 (1990) 546-549.
- 2368 Bojarski, J.: Chromatography of enantiomers of 2-arylpropionic acids. *J. Liq. Chromatogr.*, 12 (1989) 2685-2706.
- 2369 Christopoulou, C.N. and Perkins, E.G.: Chromatographic studies on fatty acid dimers: gas-liquid chromatography, high performance liquid chromatography and thin-layer chromatography. *J. Am. Oil Chem. Soc.*, 66 (1989) 1353-1359; *C.A.*, 112 (1990) 8964u.

- 2370 Gey, M., Thiersch, A., Rietzschel, A., Nattermueller, W., Stottmeister, U. and Nagel, B.: Characterization of biotechnological processes and products using high-performance liquid chromatography (HPLC). V. Analysis of organic acids in 2-oxo-D-gluconic acid synthesis. *Acta Biotechnol.*, 9 (1989) 491-496; *C.A.*, 112 (1990) 5982n.
- 2371 Hallstroem, A., Carlsson, A., Hillered, L. and Ungerstedt, U.: Simultaneous determination of lactate, pyruvate, and ascorbate in microdialysis samples from rat brain, blood, fat, and muscle using high-performance liquid chromatography. *J. Pharmacol. Methods*, 22 (1989) 113-124; *C.A.*, 111 (1989) 228279u.
- 2372 Ishida, J., Yamaguchi, M., Nakahara, T. and Nakamura, M.: 4,5-Diaminophthalhydrazide as a highly sensitive chemiluminescence reagent for α -keto acids in liquid chromatography. *Anal. Chim. Acta*, 231 (1990) 1-6.
- 2373 Itoh, H.: Determination of trichloroacetate in human serum and urine by ion chromatography. *Analyst (London)*, 114 (1989) 1637-1640.
- 2374 Kargas, G., Rudy, T., Spennetta, T., Takayama, K., Querishi, N. and Shrago, E.: Separation and quantitation of long-chain free fatty acids in human serum by high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 331-340.
- 2375 Kaule, G. and Günzler, V.: Assay for 2-oxoglutarate decarboxylating enzymes based on the determination of [1- 14 C]succinate: application to prolyl 4-hydroxylase. *Anal. Biochem.*, 184 (1990) 291-297.
- 2376 Kondo, K., Higuchi, Y., Sakuda, S., Nihira, T. and Yamada, Y.: New virginiae butanolides from *Streptomyces virginiae*. *J. Antibiot.*, 42 (1989) 1873-1876.
- 2377 Kralovsky, J., Placek, K. and Kalhousova, M.: (High-performance liquid chromatography of benzoic acid derivatives). *Chem Prum.*, 39 (1989) 356-359; *C.A.*, 111 (1989) 224589x.
- 2378 Lee, Y.M., Nakamura, H. and Nakajima, T.: Monodansylcadaverine as a versatile reagent for fluorogenic precolumn derivatization of carboxylic acids. *Anal. Sci.*, 5 (1989) 209-210; *C.A.*, 112 (1990) 48017u.
- 2379 Leung, C.P. and Leung, W.K.H.: Determination of undecenoic acid in dermatological preparations by liquid chromatography with direct UV detection. *Anal. Chim. Acta*, 229 (1990) 255-259.
- 2380 Marcé, R.M., Calull, M., Manchobas, M., Borrull, F. and Rius, F.X.: An optimized direct method for the determination of carboxylic acids in beverages by HPLC. *Chromatographia*, 29 (1990) 54-58.
- 2381 Marini, D. and Balestrieri, F.: Liquid chromatographic determination of free fatty acids and triglycerides in butter. *Riv. Soc. Ital. Sci. Aliment.*, 17 (1989) 477-480; *C.A.*, 111 (1989) 193132m.
- 2382 Miwa, H., Yamamoto, M. and Asano, T.: High-performance liquid chromatographic analyses of hydroxymonocarboxylic acids and dicarboxylic acids in urine as their 2-nitrophenylhydrazides. *Anal. Biochem.*, 185 (1990) 17-23.
- 2383 Quilliam, M.A., Sim, P.G., McCulloch, A.W. and McInnes, A.G.: High-performance liquid chromatography of domoic acid, a marine neurotoxin, with application to shellfish and plankton. *Int. J. Environ. Anal. Chem.*, 36 (1989) 139-154; *C.A.*, 111 (1989) 210158u.
- 2384 Ramakrishna, M.B.V., Mital, B.K., Gupta, K.C. and Sand, N.K.: Determination of phenolic acids in different soybean varieties by reversed-phase high-performance liquid chromatography. *J. Food Sci. Technol.*, 26 (1989) 154-155; *C.A.*, 111 (1989) 172544g.
- 2385 Runge, G. and Steinhart, H.: (Analysis of tocopherols and fatty acids in fish). *Fett Wiss. Technol.*, 91 (1989) 281-287; *C.A.*, 111 (1989) 230759u.
- 2386 Sok, D.-E. and Kim, M.R.: Enzymatic formation of 9,16-dihydro(peroxy)octadecatrienoic acid isomers from α -linolenic acid. *Arch. Biochem. Biophys.*, 277 (1990) 86-93.
- 2387 Van der Horst, F.A.L., Post, M.H., Holthuis, J.J.M. and Brinkman, U.A.T.: Automated high-performance liquid chromatographic determination of plasma free fatty acids using on-line derivatization with 9-bromomethylacridine based on micellar phase-transfer catalysis. *J. Chromatogr.*, 500 (1990) 443-452.

- 2388 Wainwright, S., Falck, J.R., Yadagiri, P. and Powell, W.S.: Stereochemistry of leukotriene B₄ metabolites formed by the reductase pathway in porcine polymorphonuclear leukocytes: inversion of stereochemistry of the 12-hydroxyl group. *Biochemistry*, 29 (1990) 1180-1185.
- 2389 Wiesend, B.: (The determination of the fatty acid composition of fish oil, fat of marmot, linseed oil and jojoba oil by a new high performance liquid chromatographic method). *PZ Wiss.*, 2 (1989) 47-53; *C.A.*, 112 (1990) 22672d.
- 2390 Wolf, J.H. and Korf, J.: Improved automated precolumn derivatization reaction of fatty acids with bromomethylmethoxycoumarin as label. *J. Chromatogr.*, 502 (1990) 423-430.
- 2391 Yamaguchi, M., Takehiro, O., Ishida, J. and Nakamura, M.: High-performance liquid chromatography of arachidonic acid metabolites and its application to the determination of leukotriene B₄ in stimulated leukocytes. *Chem. Pharm. Bull.*, 37 (1989) 2846-2848.

For additional information see:

C.A., 111 (1989) 170336s, 211293w, 230802c, 232710v;
112 (1990) 3760q, 48029z.

See also 2147, 2181, 2209, 2281, 2288, 2313, 2320, 2412, 2418, 2497, 3081.

11b. Prostaglandins

- 2392 Brockman, M.J., Eiroa, A.M. and Marcus, A.J.: Albumin redirects platelet eicosanoid metabolism toward 12(S)-hydroxyeicosatetraenoic acid. *J. Lipid Res.*, 30 (1989) 1925-1932.
- 2393 Dimov, V. and Green, K.: Metabolism of prostaglandin F_{2α} in sheep. *Drug Metab. Disp.*, 18 (1990) 107-114.
- 2394 Nicoll-Griffith, D., Zamboni, R., Rasmussen, J.B., Ethier, D., Charleson, S. and Tagari, P.: BIO-Fully Automated Sample Treatment high-performance liquid chromatography and radioimmunoassay for leukotriene E₄ in human urine from asthmatics. *J. Chromatogr.*, 526 (1990) 341-354.
- 2395 Oliw, E.H.: Resolution of enantiomers of 19-hydroxyeicosatetraenoate and 18-hydroxyeicosatetraenoate by chiral phase high-performance liquid chromatography of naphthoyl ester derivatives. *J. Chromatogr.*, 526 (1990) 525-529.
- 2396 Siegenthaler, C.: (HPLC method for the analysis of prostaglandin E₂ in Tylose gel). *Pharm. Acta Helv.*, 64 (1989) 345-347; *C.A.*, 112 (1990) 62746n.
- 2397 Sugiyama, S., Kawabe, Y., Kuroiwa, M., Goto, H., Tsukamoto, Y. and Ozawa, T.: Effects of the synthesized prostaglandin E₂-analogue araprostil on gastric mucosal lesions in rats. *Arzneim.-Forsch.*, 39 (1989) 1571-1573.
- 2398 Tagari, P., Ethier, D., Ford-Hutchinson, A. and Cirino, M.: Application of reversed phase high performance liquid chromatography to the analysis of sulphidopeptide leukotrienes in pig bile. *Biomed. Chromatogr.*, 4 (1990) 39-42.

For additional information see:

C.A., 111 (1989) 209242k.

See also 2391.

11c. Lipids and their constituents

- 2399 Asmis, R. and Jörg, A.: Calcium-ionophore-induced formation of platelet-activating factor and leukotrienes by horse eosinophils: a comparative study. *Eur. J. Biochem.*, 187 (1990) 475-480.

- 2400 Balesdent, D., Kapseu, C. and Kayem, G.J.: (Characterization of triglycerides of cottonseed oil by high performance liquid chromatography (HPLC) in one step. Laboratory note). *Rev. Fr. Corps Gras*, 36 (1989) 27-33; *C.A.*, 112 (1990) 6185s.
- 2401 Banerji, B., Subbaiah, P.V., Gregg, R.E. and Bagdade, J.D.: Molecular species of phosphatidylcholine in betalipoproteinemia: effect of lecithin:cholesterol acyltransferase and lysolecithin acyltransferase. *J. Lipid Res.*, 30 (1989) 1907-1916.
- 2402 Becart, J., Chevalier, C. and Biesse, J.P.: Quantitative analysis of phospholipids by HPLC with a light scattering evaporating detector - application to raw materials for cosmetic use. *J. High Resolut. Chromatogr.*, 13 (1990) 126-129.
- 2403 Benfenati, E., Macconi, D., Noris, M., Icardi, G., Bettazzoli, L., de Bellis, G., Gavinelli, M., Rotondo, S. and Remuzzi, G.: Development of a mass spectrometric method to quantitate platelet activating factor in mouse urine. *J. Lipid Res.*, 30 (1989) 1977-1981.
- 2404 Bonafede, D.M., Macala, L.J., Constantine-Paton, M. and Yu, R.K.: Isolation and characterization of ganglioside 9-O-acetyl-GD₃ from bovine buttermilk. *Lipids*, 24 (1989) 680-684; *C.A.*, 111 (1989) 228253f.
- 2405 Boyle, L.E., Sklar, L.A. and Traynor-Kaplan, A.E.: An HPLC procedure for separating polyphosphoinositides on hydroxylapatite. *J. Lipid Res.*, 31 (1990) 157-159.
- 2406 Cohen, D.E., Angelico, M. and Carey, M.C.: Structural alternations in lecithin-cholesterol vesicles following interactions with monomeric and micellar bile salts: physical-chemical basis for subselection of biliary lecithin species and aggregative states of biliary lipids during bile formation. *J. Lipid Res.*, 31 (1990) 55-70.
- 2407 Eick, S., Brummel, M. and Spener, F.: (Analysis and *in vitro* reconstitution of triglycerides of the Siberian marmot oil from *Marmota bobac*). *Fett. Wiss. Technol.*, 91 (1989) 231-233; *C.A.*, 111 (1989) 156315c.
- 2408 Gertz, C.: (The analysis and evaluation of vegetable fats and oils.) *GIT-Suppl.*, No. 2 (1989) 5-15; *C.A.*, 111 (1989) 193114g - a review with 49 refs.
- 2409 Heinze, T., Kynast, G., Dudenhausen, J.W., Schmitz, C. and Saling, E.: Quantitative determination of phospholipids in amniotic fluid by HPLC. *GIT Fachz. Lab.*, 33 (1989) 553-555; *C.A.*, 111 (1989) 170367c.
- 2410 Hullin, F., Kim, H.-Y. and Salem, N., Jr.: Analysis of aminophospholipid molecular species by high performance liquid chromatography. *J. Lipid Res.*, 30 (1989) 1963-1975.
- 2411 Janero, D.R. and Burghardt, C.: Solid-phase extraction on silica cartridges as an aid to platelet-activating factor enrichment and analysis. *J. Chromatogr.*, 526 (1990) 11-24.
- 2412 Ljusberg-Wahren, H. and Johansson, L.: Saponification of emulsified triglyceride oil in presence of rosin and fatty acid soaps. *Fat Sci. Technol.*, 91 (1989) 456-458.
- 2413 McCluer, R.H., Ullman, M.D. and Jungalwala, F.B.: High-performance liquid chromatography of membrane lipids: glycosphingolipids and phospholipids. *Methods Enzymol.*, 172 (Biomembranes, Pt. S), (1989) 538-575; *C.A.*, 111 (1989) 190626q.
- 2414 Miyazawa, T., Saeki, R. and Inaba, H.: Detection of chemiluminescence in lipid peroxidation of biological systems and its application to HPLC. *J. Biolumin. Chemilumin.*, 4 (1989) 475-478; *C.A.*, 111 (1989) 228341h.
- 2415 Miyazawa, T., Yasuda, K., Fujimoto, K. and Kaneda, T.: Chemiluminescence-high performance liquid chromatography of phosphatidylcholine hydroperoxide in human blood plasma. *Basic Life Sci.*, 49 (Oxygen Radicals Biol. Med.) (1988) 191-194; *C.A.*, 112 (1990) 3762s.

- 2416 Morgan, R.O. and Catt, K.J.: HPLC methodology with flow detection of radio-labeled neutral and phospholipids, polyphosphoinositides, phosphoinositol isomers and arachidonic acid metabolites from pituitary gonadotrophs. *Prog. HPLC*, 3 (1988) 27-55; *C.A.*, 111 (1989) 170315j.
- 2417 Phelkin, V.P. and Vereshchagin, A.G.: (Determination by liquid chromatography of types of membrane lipids and their derivatives). *Khim. Prir. Soedin.*, (1989) 453-477; *C.A.*, 112 (1990) 3684t - a review with 159 refs.
- 2418 Robinson, B.S., Johnson, D.W. and Poulos, A.: Unique molecular species of phosphatidylcholine containing very-long-chain (C₂₄-C₃₈) polyenoic fatty acids in rat brain. *Biochem. J.*, 265 (1990) 763-767.
- 2419 Saeed, T., Ali, S.G., Rahman, H.A.A. and Sawaya, W.N.: Detection of pork and lard as adulterants in processed meat: liquid chromatographic analysis of derivatized triglycerides. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 921-925.
- 2420 Shukla, V.K.S.: Application of modern analytical techniques for evaluation of lipid structures. *J. Dispersion Sci. Technol.*, 10 (1988) 581-616; *C.A.*, 111 (1989) 228183h - a review with 48 refs.
- 2421 Tomono, Y., Abe, K. and Watanabe, K.: High-performance liquid affinity chromatography and *in situ* fluorescent labeling on thin-layer chromatography of glycosphingolipids. *Anal. Biochem.*, 184 (1990) 360-368.

For additional information see:

- C.A.*, 111 (1989) 170354w, 170608g, 211291u, 228257k;
112 (1990) 12426n, 38674s.

See also 2277, 2317, 2319, 2473.

11d. Lipoproteins and their constituents

- 2422 Caines, P.S., Thibert, R.J., Draisey, T.F., Thomas, F., Foreback, C.C. and Chu, J.W.: Determination of nonenzymically glycated apolipoprotein A1. *Clin. Biochem.*, 22 (1989) 285-287; *C.A.*, 111 (1989) 190916j.
- 2423 Cocuzzi, E., Piacentini, M., Beninati, S. and Chung, S.I.: Post-translational modification of apolipoprotein B by transglutaminases. *Biochem. J.*, 265 (1990) 707-713.
- 2424 Junker, L.H. and Davis, R.A.: Receptor-mediated uptake of low density lipoprotein stimulates bile acid synthesis by cultured rat hepatocytes. *J. Lipid Res.*, 30 (1989) 1933-1941.
- 2425 Kortlandt, W., Benschop, C., Erkelens, D.W. and Thijssen, J.H.H.: A simple method for the measurement of total and glycated apolipoprotein B and its relevance to apolipoprotein-B metabolism in diabetes mellitus. *Clin. Chim. Acta*, 186 (1989) 109-118.
- 2426 Van Gent, T. and van Tol, A.: Automated gel permeation chromatography of plasma lipoproteins by preparative fast protein liquid chromatography. *J. Chromatogr.*, 525 (1990) 433-441.

See also 2631.

13. STEROIDS

- 2427 Robards, K. and Towers, P.: Chromatography as a reference technique for the determination of clinically important steroids. *Biomed. Chromatogr.*, 4 (1990) 1-19 - a review with 255 refs.

- 2428 Wei, J.-Q., Wei, J.-L. and Zhou, X.-T.: Optimization of an isocratic reversed phase liquid chromatographic system for the separation of fourteen steroids using factorial design and computer simulation. *Biomed. Chromatogr.*, 4 (1990) 34-38.

13a. Pregnane and androstane derivatives

- 2429 Aoyama, T., Korzekwa, K., Nagata, K., Adesnik, M., Reiss, A., Lapenson, D.P., Gillette, J., Gelboin, H.V., Waxman, D.J. and Gonzalez, F.J.: Sequence requirements for cytochrome P-450IIB1 catalytic activity. Alteration of the stereospecificity and regioselectivity of steroid hydroxylation by a simultaneous change of two hydrophobic amino acid residues to phenylalanine. *J. Biol. Chem.*, 264 (1989) 21327-21333.
- 2430 Karinova, I.N. and Dvorkin, V.I.: (Chromatographic determination of 7-dehydrocholesterol in human blood serum). *Anal. Khim.*, 44 (1989) 1902-1905; *C.A.*, 112 (1990) 3796f.
- 2431 Khatsenko, O.G., Gulyaeva, L.F., Mishin, V.M. and Lyakhovich, V.V.: (Estimation of the catalytic activity of multiple forms of cytochrome P-450 in Wistar rat liver microsomes by androstenedione metabolism). *Biokhimiya (Moscow)*, 55 (1990) 308-314.
- 2432 Lampert, B.L. and Stewart, J.T.: Determination of anabolic steroids and zeranol in human serum by isocratic reversed phase HPLC on silica. *J. Liq. Chromatogr.*, 12 (1989) 3231-3249.
- 2433 Oka, K., Hirano, T., Shimodaira, H., Homma, M., Sakurai, E., Tamaki, T. and Kozaki, M.: Suppression of endogenous cortisol for evaluating pharmacodynamics of prednisolone in early allograft rejection in renal transplantation. *Clin. Chem. (Winston-Salem)*, 36 (1990) 481-486.
- 2434 Staneva, L., Taraganska, G. and Simeonova, K.: (Reversed-phase HPLC of methyltestosterone tablets). *Izv. Durzh. Inst. Kontrol Lek. Sredstva*, 22 (1989) 12-15; *C.A.*, 112 (1990) 42703t.
- 2435 Underwood, R.H., Bradwin, G.R., Moore, T.J. and Williams, G.H.: Semi-automated high-performance liquid chromatographic method for the simultaneous assay of plasma cortisol and 11-deoxycortisol in the metyrapone test. *J. Chromatogr.*, 526 (1990) 180-185.

For additional information see:

C.A., 111 (1989) 167504q, 209252p, 225388t.

13b. Estrogens

- 2436 Brind, J.L., Chervinsky, K., Vogleman, J.H. and Orentreich, N.: Radioimmunoassay of estrone sulfate in the serum of normal men after a chromatographic procedure that eliminates dehydroepiandrosterone sulfate interference. *Steroids*, 54 (1989) 21-35; *C.A.*, 111 (1989) 187684b.

13c. Sterols

- 2437 Araki, N., Horiuchi, S., Rahim, A.T.M.A., Takata, K. and Morino, Y.: Microquantification of cholesterol and cholesteryl esters in rat peritoneal macrophages by reversed-phase high-performance liquid chromatography. *Anal. Biochem.*, 185 (1990) 339-345.
- 2438 Batta, A.K., Salen, G. and Shefer, S.: Chromatographic separation of putative precursors of cholestanol. *Steroids*, 52 (1988) 109-122; *C.A.*, 111 (1989) 195230r.
- 2439 Shah, Y., Khanna, S. and Gighe, V.S.: High performance liquid chromatographic method for clobetasol propionate in dosage forms. *Indian Drugs*, 27 (1989) 132-133; *C.A.*, 112 (1990) 42705v.

- 2440 Sonoda, Y., Sekigawa, Y. and Sato, Y.: Metabolism of 32-oxo-24,25-dihydrola-nosterols by partially purified cytochrome P-450_{14DM} from rat liver microsomes. *Chem. Pharm. Bull.*, 37 (1989) 2762-2765.
- 2441 Vercaemst, R., Union, A., Rosseneu, M., De Craene, I., de Backer, G. and Kornitzer, M.: Quantitation of plasma free cholesterol and cholesteryl esters by high performance liquid chromatography. Study of a normal population. *Atherosclerosis (Shannon)*, 78 (1989) 245-250; *C.A.*, 111 (1989) 228254g.
- 2442 Yan, J. and Bittman, H.: A facile synthesis of [¹⁴C]epicholesterol. *J. Lipid Res.*, 31 (1990) 160-162.

For additional information see:

- C.A.*, 111 (1989) 190622k;
112 (1990) 22747g.

13d. Bile acids and alcohols

- 2443 Axelsson, M., Mörk, B., Aly, A., Walldius, G. and Sjövall, J.: Concentrations of cholestenic acids in plasma from patients with reduced intestinal reabsorption of bile acids. *J. Lipid Res.*, 30 (1989) 1883-1887.
- 2444 Axelsson, M., Mörk, B., Aly, A., Wisén, O. and Sjövall, J.: Concentrations of cholestenic acids in plasma from patients with liver disease. *J. Lipid Res.*, 30 (1989) 1877-1882.
- 2445 Dengler, T., Kellner, S. and Fuerst, G.: Quantitative determination of residual sodium cholate in tri(*n*-butyl) phosphate/sodium cholate treated factor VIII preparations. *Vox Sang.*, 57 (1989) 161-163; *C.A.*, 112 (1990) 25755u.
- 2446 Eckers, C., Hashins, N.J. and Large, T.: On-line negative ion liquid chromatography/mass spectrometry for the analysis of bile acids using low- and high-resolution mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 702-706; *C.A.*, 111 (1989) 190940n.
- 2447 Hiraoka, T., Kohda, T., Kosaka, D., Yamauchi, T., Kihira, K., Kuramoto, T., Hoshita, T. and Kajiyama, G.: Identification of bile alcohols in rat bile. *J. Lipid Res.*, 30 (1989) 1889-1893.
- 2448 Kuipers, F., Radomska, A., Zimniak, P., Little, J.M., Havinga, R., Vonk, R.J. and Lester, R.: Defective biliary secretion of bile acid 3-O-glucuronides in rats with hereditary conjugated hyperbilirubinaemia. *J. Lipid Res.*, 30 (1989) 1835-1845.
- 2449 Li, B., Rui, L., Chen, Y. and Sheng, H.: (High performance liquid chromatographic quantitation of conjugated bile acids in human body fluids with immobilized enzyme detection system.) *Sepu*, 7 (1989) 165-167; *C.A.*, 111 (1989) 190648y.
- 2450 Wahlen, E., Eggestad, B., Strandvik, B. and Sjöval, J.: Ketonic bile acids in urine of infants during the neonatal period. *J. Lipid Res.*, 30 (1989) 1847-1857.

For additional information see:

- C.A.*, 111 (1989) 170606e;
112 (1990) 11978g, 42697u.

See also 2406.

14. STEROID GLYCOSIDES AND SAPONINS

- 2451 Embree, L. and McErlane, K.M.: Electrochemical detection of the 3,5-dinitrobenzoyl derivative of digoxin by high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 439-446.

- 2452 Rauwald, H.W. and Janssen, B.: (TLC analysis of *Ruscus aculeatus* L. and HPLC determination of deglucoruscin and deglucoruscoid in crude drug and pharmaceutical formulations). *PZ Wiss.*, 1 (1988) 61-68; *C.A.*, 112 (1990) 25742n.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 2453 Barrow, K.D. and Chuck, J.-A.: Determination of hopanoid levels in bacteria using high-performance liquid chromatography. *Anal. Biochem.*, 184 (1990) 395-399.
- 2454 Italia, A., Schiavi, M. and Ventura, P.: Direct liquid chromatographic separation of enantiometric and diastereometric terpenic alcohols as β -cyclodextrin inclusion complexes. *J. Chromatogr.*, 503 (1990) 266-271.

See also 2412, 2819.

15b. Essential oils

- 2455 Munari, F., Dugo, G. and Cotroneo, A.: Automated on-line HPLC-HRGC with gradient elution and multiple GC transfer applied to the characterization of citrus essential oils. *J. High Resolut. Chromatogr.*, 13 (1990) 56-61.

For additional information see:

C.A., 112 (1990) 42601h.

See also 2293.

16. NITRO AND NITROSO COMPOUNDS

- 2456 Chen, G.L., Hu, M.K., Chan, L.Y., Wang, H.J. and Chan, C.H.: A stability indicating method for isoamyl nitrite by high-performance liquid chromatography/diode array detector. *Chung-hua Yao Hsueh Tsa Chih*, 41 (1989) 331-337; *C.A.*, 112 (1990) 42721x.
- 2457 Jenkins, T.F., Walsh, M.E., Schumacher, P.W., Miyares, P.H., Bauer, C.F. and Grant, C.L.: Liquid chromatographic method for determination of extractable nitroaromatic and nitramine residues in soil. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 890-899.
- 2458 Rickert, D.E. and Held, S.D.: Metabolism of chloronitrobenzenes by isolated rat hepatocytes. *Drug Metab. Disp.*, 18 (1990) 5-9.

For additional information see:

C.A., 111 (1989) 216764k.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 2459 Gutierrez Bouzan, M.del C. and Crespi Rossell, M.: (Determination of aromatic amines by high-resolution liquid chromatography). *Bol. INTXTAR Inst. Invest. Text. Coop. Ind.*, 95 (1989) 75-82; *C.A.*, 111 (1989) 215960j.

- 2460 Lancaster, F.E., and Lawrence, J.F.: Determination of total non-sulfonated aromatic amines in the food color amaranth by dithionite reduction followed by derivatization and high-performance liquid chromatography. *Food Addit. Contam.*, 6 (1989) 415-423; *C.A.*, 111 (1989) 213436u.
- 2461 Lee, H.: Variability obtained with different batches of acetonitrile in the HPLC of polyamines. *Chromatogram*, 10 (1989) 4-6; *C.A.*, 112 (1990) 3767x.

See also 2113, 2137, 2457.

17b. Catecholamines and their metabolites

- 2462 Aravagiri, M., Marder, S.R. and van Putten, T.: Determination of plasma homovanillic acid by column liquid chromatography with electrochemical detection. *J. Chromatogr.*, 525 (1990) 419-425.
- 2463 Candito, M., Krstulovic, A.M., Sbirazzuoli, V. and Chambon, P.: Proposal for the standardization of the calibration method for the assay of plasma catecholamines. *J. Chromatogr.*, 526 (1990) 194-202.
- 2464 Gupta, R.N.: Improved sample preparation in determination of urinary metanephines by liquid chromatography with electrochemical detection. *Clin. Chem. (Winston-Salem)*, 36 (1990) 538-540.
- 2465 Matsumoto, M., Togashi, H., Yoshioka, M., Hirokami, M., Morii, K. and Saito, H.: Simultaneous high-performance liquid chromatographic determination of norepinephrine, serotonin, acetylcholine and their metabolites in the cerebrospinal fluid of anaesthetized normotensive rats. *J. Chromatogr.*, 526 (1990) 1-10.
- 2466 Wetherell, J.R., Fosbraey, P. and French, M.C.: A comparison of the distribution of neurotransmitters in brain regions of the rat and guinea pig using a chemiluminescent method and HPLC with electrochemical detection. *J. Neurochem.*, 53 (1989) 1519-1526; *C.A.*, 111 (1989) 225395t.

For additional information see:

C.A., 111 (1989) 210167w, 211280q.

17c. Urea and guanidine derivatives

- 2467 Boppana, V.K., Rhodes, G.R. and Brooks, D.P.: Determination of methylguanidine in plasma and urine by high-performance liquid chromatography with fluorescence detection following postcolumn derivatization. *Anal. Biochem.*, 184 (1990) 213-218.
- 2468 Chao, W.-F., Kai, M. and Ohkura, Y.: High-performance liquid chromatographic determination of a cysteine protease inhibitor and its ethyl ester in mouse serum and muscle by precolumn fluorescence derivatization. *J. Chromatogr.*, 526 (1990) 77-86.
- 2469 Ding, W., Veening, H. and van Effen, R.M.: Determination of guanidino compounds by anion-exchange chromatography and amperometric detection. *J. Chromatogr.*, 526 (1990) 355-366.

17d. Other amine derivatives and amides (excluding peptides)

- 2470 Bauer, R. and Remiger, P.: *Echinacea*. Part V. TLC and HPLC analysis of alkanamides in *Echinacea* drugs. *Planta Med.*, 55 (1989) 367-371; *C.A.*, 112 (1990) 11975d.
- 2471 Dowle, C.J., Malyan, A.P. and Matheson, A.M.: Separation of the *ortho*, *meta* and *para* isomers of aminophenol by high-performance liquid chromatography. *Analyst (London)*, 115 (1990) 105-107.

- 2472 Minkler, P.E., Ingalls, S.T. and Hoppel, C.L.: High-performance liquid chromatographic separation of acylcarnitines following derivatization with 4'-bromophenacyl trifluoromethanesulfonate. *Anal. Biochem.*, 185 (1990) 29-35.
- 2473 Tyrefors, N. and Carlsson, A.: Improvements in the separation and detection of acetylcholine and choline using liquid chromatography and electrochemical detection. *J. Chromatogr.*, 502 (1990) 337-349.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 2474 Algermissen, B., Neudel, M. and Riedel, E.: (Determination of amino acids by fluorescence HPLC). *GIT Fachz. Lab.*, 33 (1989) 783-790; *C.A.*, 111 (1989) 247207k - a review with 28 refs.
- 2475 Amiss, T.J., Tyczkowska, K.L. and Aucoin, D.P.: Analysis of taurine in feline plasma and whole blood by liquid chromatography with fluorimetric detection and confirmation by thermospray mass spectrometry. *J. Chromatogr.*, 526 (1990) 375-382.
- 2476 Andersson, A., Brattstroem, L., Isaksson, A., Israelsson, B. and Hultberg, B.: Determination of homocysteine in plasma by ion-exchange chromatography. *Scand. J. Clin. Lab. Invest.*, 49 (1989) 445-449; *C.A.*, 111 (1989) 228271k.
- 2477 Atherton, D.: Successful PTC amino acid analysis at the picomole level. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 273-283; *C.A.*, 111 (1989) 228247g.
- 2478 Beale, S.C., Hsieh, Y.-Z., Wiesler, D. and Novotny, M.: Application of 3-(2-furoyl)quinoline-2-carbaldehyde as a fluorogenic reagent for the analysis of primary amines by liquid chromatography with laser-induced fluorescence detection. *J. Chromatogr.*, 499 (1990) 579-587.
- 2479 Begg, G.S. and Simpson, R.J.: An improved injection system for on-line liquid chromatographs: application to high sensitivity phenylthiohydantoin amino acid analysis. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 79-88; *C.A.*, 111 (1989) 228187n.
- 2480 Bizzarri, M., Catizone, A., Pompei, M., Chiappini, L., Curini, L. and Lagana, A.: Determination of urinary tryptophan and its metabolites along the nicotinic acid pathway by high performance liquid chromatography with ultraviolet detection. *Biomed. Chromatogr.*, 4 (1990) 24-27.
- 2481 Cardinali, F., Ziggotti, A. and Viscomi, G.C.: Scaling-up procedure from the range of milligrams to grams for the purification of amino acid derivatives in displacement chromatography. *J. Chromatogr.*, 499 (1990) 37-45.
- 2482 Castellanos, L.: (Separation of PTH-amino acids on short RP-columns.) *Interferon Biotechnol.*, 6 (1989) 67-71; *C.A.*, 111 (1989) 190656z.
- 2483 Chang, J.Y., Knecht, R., Jenoe, P. and Vekemans, S.: Amino acid analysis at the femtomole level using the dimethylaminoazobenzene sulfonyl chloride precolumn derivatization method: potential and limitation. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 305-314; *C.A.*, 111 (1989) 228249j.
- 2484 Dupont, D.R., Keim, P.S., Chui, A.H., Bello, R., Bozzini, M.L. and Wilson, K.J.: A comprehensive approach to amino acid analysis. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 284-294; *C.A.*, 111 (1989) 228248h.
- 2485 Dyremark, A. and Ericsson, M.: An LCEC method for the analysis of synthetic amino acids in fruit juice. *Chromatographia*, 29 (1990) 51-53.
- 2486 Euerby, M.R.: Resolution of the neuroexcitatory N-methylaspartic acid (2-methylaminosuccinic acid) enantiomers by ligand-exchange thin-layer chromatography. *J. Chromatogr.*, 502 (1990) 226-229.

- 2487 Fürst, P., Pollack, L., Graser, T.A., Godel, H. and Stehle, P.: HPLC analysis of free amino acids in biological material - an appraisal of four pre-column derivatization methods. *J. Liq. Chromatogr.*, 12 (1989) 2733-2760.
- 2488 Gerstl, R. and Ranft, K.: (Determination of D,L-2-hydroxy-4-methylmercaptobutyric acid for the detection of Aliment supplements in mixed feeds.) *Landwirtsch. Forsch.*, 42 (1989) 6-11; *C.A.*, 111 (1989) 193171y.
- 2489 Gould, B.J., Walsh, D.P., Ah-Sing, E., Wright, J. and Stace, B.C.: Urinary carboxymethyl-lysine measurement as a potential indicator of diabetic complications. *Biochem. Soc. Trans.*, 17 (1989) 1087-1088; *C.A.*, 111 (1989) 192510w.
- 2490 Gunawan, S., Walton, N.Y. and Treiman, D.M.: High-performance liquid chromatographic determination of selected amino acids in rat brain by precolumn derivatization with phenylisothiocyanate. *J. Chromatogr.*, 503 (1990) 177-187.
- 2491 Guo, Y., Huang, Y. and Yao, Z.: (Analysis of total amino acids by pre-column derivatization with Fmoc). *Sepu*, 7 (1989) 219-221; *C.A.*, 111 (1989) 228268q.
- 2492 Hagen, S.R. and Augustin, J.: Determination of tryptophan in foods by isocratic reversed-phase high-performance liquid chromatography. *J. Micronutr. Anal.*, 5 (1989) 303-309; *C.A.*, 111 (1989) 230784y.
- 2493 Hagen, S.R., Frost, B. and Augustin, J.: Precolumn phenylisothiocyanate derivatization and liquid chromatography of amino acids in food. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 912-916.
- 2494 Haginaka, J. and Wakai, J.: Automated precolumn derivatization of amino acids with *ortho*-phthalaldehyde using a hollow-fibre membrane reactor. *J. Chromatogr.*, 502 (1990) 317-324.
- 2495 James, I.T., Perrett, D. and Thompson, P.W.: Rapid assay for hard tissue collagen cross-links using isocratic ion-pair reversed-phase liquid chromatography. *J. Chromatogr.*, 525 (1990) 43-57.
- 2496 Koshiishi, I. and Imanari, T.: Determination of homocitrullin residue in plasma proteins. *Anal. Sci.*, 5 (1989) 295-299; *C.A.*, 111 (1989) 228241a.
- 2497 Liebich, H.M., Bubeck, J.J., Pickert, A., Wahl, G. and Scheiter, A.: Hippuric acid and 3-carboxy-4-methyl-5-propyl-2-furanpropionic acid in serum and urine. Analytical approaches and clinical relevance in kidney diseases. *J. Chromatogr.*, 500 (1990) 615-627.
- 2498 Lippincott, S., Chesney, R.W., Friedman, A. Pityer, R., Barden, H. and Mazess, R.B.: Rapid determination of total hydroxyproline (HYP) in human urine by HPLC analysis of the phenylisothiocyanate (PITC)-derivative. *Bone*, 10 (1989) 265-268; *C.A.*, 112 (1990) 3795e.
- 2499 Lovell, G. and Corran, P.H.: Determination of L-thyroxine in reference serum preparations as the *o*-phthalaldehyde-N-acetylcysteine derivative by reversed-phase detection chromatography with electrochemical detection. *J. Chromatogr.*, 525 (1990) 287-296.
- 2500 Malencik, D.A., Zhao, Z. and Anderson, S.R.: Determination of dityrosine, phosphoserine by high-performance liquid chromatography. *Anal. Biochem.*, 184 (1990) 353-359.
- 2501 McGowan, C., Wiley, V.A. and Bates, R.P.: Application of methodology for RP-HPLC amino acid analysis to the measurement of hypoglycin A. *BioChromatography*, 4 (1989) 161-164; *C.A.*, 111 (1989) 190616m.
- 2502 Moitra, S., Mundhara, G.L. and Tiwari, J.S.: Studies of the sorption behavior of some aminoacids on chemically pretreated alumina in relation to chromatography. *Colloids Surf.*, 41 (1989) 311-326; *C.A.*, 112 (1990) 43413k.
- 2503 Morita, I., Kawamoto, M., Hattori, M., Eguchi, K., Sekiba, K. and Yoshida, H.: Determination of tryptophan and its metabolites in human plasma and serum by high-performance liquid chromatography with automated sample clean-up system. *J. Chromatogr.*, 526 (1990) 367-374.

- 2504 Nardini, M., Matarese, R.M., Pecci, L., Antonucci, A., Ricci, G. and Cavallini, D.: Detection of 2H-1,4-thiazine-5,6-dihydro-3-carboxylic acid (aminoethylcysteine ketimine) in the bovine brain. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1251-1256.
- 2505 Quershi, G.A. and Gokmen, S.: Determination of amino acids by liquid chromatography based on pre- and post-column derivatization. *Chim. Acta Turc.*, 15 (1987) 227-241; *C.A.*, 111 (1989) 224606a.
- 2506 Schuessler, U. and Ledl, F.: Direct determination of 2-amino-6-(2-formyl-5-hydroxymethyl-1-pyrrolyl)-hexanoic acid in hydrolyzed protein by reversed-phase HPLC. *Z. Lebensm.-Unters. Forsch.*, 189 (1989) 32-35; *C.A.*, 111 (1989) 193146u.
- 2507 Smith, A.J., Presley, J.M. and McIntire, W.: An evaluation of an automated high sensitivity amino acid analyzer based on the 9-fluorenylmethylchloroformate (FMOC) chemistry. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 255-265; *C.A.*, 111 (1989) 228188p.
- 2508 Stratford, M.R.L., Watfa, R.R., Murray, J.C. and Martin, S.G.: Determination of collagen and protein turnover by high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 383-395.
- 2509 Swartz, K.J., Matson, W.R., MacGarvey, U., Ryan, E.A. and Beal, M.F.: Measurement of kynurenic acid in mammalian brain extracts and cerebrospinal fluid by high-performance liquid chromatography with fluorometric and coulometric electrode array detection. *Anal. Biochem.*, 185 (1990) 363-376.
- 2510 Tabuchi, T., Okayama, A., Ogawa, Y., Miyajima, K., Hirata, I., Yoshida, T., Sugimoto, K. and Morimoto, K.: A new HPLC fluorometric method to monitor urinary delta-aminolevulinic acid ALA-U levels in workers exposed to lead. *Int. Arch. Occup. Environ. Health*, 61 (1989) 297-302; *C.A.*, 111 (1989) 188852s.
- 2511 Tsuruta, Y., Date, Y. and Kohashi, K.: Phthalimidylbenzenesulphonyl chlorides as fluorescence labelling reagents for amino acids in high-performance liquid chromatography. *J. Chromatogr.*, 502 (1990) 178-183.
- 2512 Van Eijk, H.M.H., Deutz, N.E.P., Wagenmakers, A.J.M. and Soeters, P.B.: 3-Methylhistidine determined in plasma by "high-performance" liquid chromatography. *Clin. Chem. (Winston-Salem)*, 36 (1990) 556-559.
- 2513 Ziegler, S.J. and Sticher, O.: HPLC of S-alk(en)yl-L-cystein derivatives in garlic including quantitative determination of (+)-S-allyl-L-cysteine sulfoxide (alliin). *Planta Med.*, 55 (1989) 372-378; *C.A.*, 112 (1990) 11976e.

For additional information see:

C.A., 111 (1989) 167513s, 170612d, 171491a, 228560d, 233519b;
112 (1990) 2158n, 3971j, 4050v, 15800r.

See also 2269, 2423, 2466, 2526, 2542, 2623, 2643, 2704, 2766.

18b. Peptides and peptidic and proteinous hormones

- 2514 Azumi, K., Yokosawa, H. and Ishii, S.: Halocyanamines: novel antimicrobial tetrapeptide-like substances isolated from the hemocytes of the solitary ascidian *Halocynthia roretzi*. *Biochemistry*, 29 (1990) 159-165.
- 2515 Bean, M.F., Pallante-Morell, S.L., Dulik, D.M. and Fenselau, C.: Protocol for liquid chromatography/mass spectrometry of glutathione conjugates using post-column solvent modification. *Anal. Chem.*, 62 (1990) 121-124.
- 2516 Brigstock, D.R., Heap, R.B., Barker, P.J. and Brown, K.D.: Purification and characterization of heparin-binding growth factors from porcine uterus. *Biochem. J.*, 266 (1990) 273-282.

- 2517 Burgoyne, R., Stacey, C., Young, P., Astephen, N. and Merion, M.: Analysis of peptides at low levels using a chromatography system optimized for narrow-bore columns. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 399-413; *C.A.*, 111 (1989) 228252e.
- 2518 Casteels, P., Ampe, C., Riviere, L., van Damme, J., Elicone, C., Fleming, M., Jacobs, F. and Tempst, P.: Isolation and characterization of abaecin, a major antibacterial response peptide in the honeybee (*Apis mellifera*). *Eur. J. Biochem.*, 187 (1990) 381-386.
- 2519 Chaiken, I., Fassina, G. and Shai, Y.: Analytical affinity chromatography, antisense peptides, and the synthetic design of recognition molecules. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (Protein Recognit. Immobilized Ligands) (1989) 2-23; *C.A.*, 111 (1989) 228465b.
- 2520 Cliffe, A.J., Revell, D. and Law, B.A.: A method for the reverse phase high performance liquid chromatography of peptides from Cheddar cheese. *Food Chem.*, 34 (1989) 147-160; *C.A.*, 111 (1989) 193143r.
- 2521 Domin, J., Yiangou, Y., Spokes, R.A., Aitken, A., Parmar, K.B., Chrysanthou, B.J. and Bloom, S.R.: The distribution, purification, and pharmacological action of an amphibian neuromedin U. *J. Biol. Chem.*, 264 (1989) 20881-20885.
- 2522 Engelhardt, H., Appelt, G. and Schweinheim, E.: Unexpected elution behaviour of peptides with various reversed-phase columns. *J. Chromatogr.*, 499 (1990) 165-175.
- 2523 Friedman, D.I. and Amidon, G.L.: Intestinal absorption mechanism of dipeptide angiotensin converting enzyme inhibitors of the lysyl-proline type: lisinopril and SQ 29,852. *J. Pharm. Sci.*, 78 (1989) 995-998.
- 2524 Henderson, D.E. and Mello, J.A.: Physicochemical studies of biologically active peptides by low-temperature reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 499 (1990) 79-88.
- 2525 Hodges, R.S., Burke, T.W.L. and Mant, C.T.: A novel approach to the purification of peptides by reversed-phase HPLC: sample displacement chromatography. In: Marshall, G.R. (Editor), *Pept. Chem. Biol., Proc. Am. Pept. Symp. 10th 1987*, ESCOM Sci. Pub., Leiden, 1988, pp. 226-228; *C.A.*, 111 (1989) 233607d.
- 2526 Hofsteenge, J., Stone, S.R., Donella-Deana, A. and Pinna, L.A.: The effect of substituting phosphotyrosine for sulphotyrosine on the activity of hirudin. *Eur. J. Biochem.*, 188 (1990) 55-59.
- 2527 Hwang, I.Y. and Elfarra, A.A.: Cysteine S-conjugates may act as kidney-selective prodrugs: formation of 6-mercaptopurine by the renal metabolism of S-(6-purinyl)-L-cysteine. *J. Pharmacol. Exp. Ther.*, 251 (1989) 448-454.
- 2528 Johnson, G.R. and Foster, C.J.: Partial characterization of a metalloendopeptidase activity produced by cultured endothelial cells that removes the COOH-terminal tripeptide from 125 I-atrial natriuretic factor. *Biochem. Biophys. Res. Commun.*, 167 (1990) 110-116.
- 2529 Kitagawa, K., Kawamoto, T., Futaki, S., Kiyama, S., Akita, T., Moritoki, H. and Kiso, Y.: Solution syntheses of two enkephalin-containing peptides, peptide E and dynorphin (1-24), using N^{in} -(2,4,6-trisopropylphenylsulfonyl) tryptophan. *Chem. Pharm. Bull.*, 37 (1989) 2631-2638.
- 2530 Kuroki, Y., Kanda, T., Kubota, I., Fujisawa, Y., Ikeda, T., Miura, A., Minamitake, Y. and Muneoka, Y.: A molluscan neuropeptide related to the crustacean hormone, RPCH. *Biochem. Biophys. Res. Commun.*, 167 (1990) 273-279.
- 2531 Lee, E.D., Henion, J.D. and Covey, T.R.: Microbore high-performance liquid chromatography-ion spray mass spectrometry for the determination of peptides. *J. Microcolumn Sep.*, 1 (1989) 14-18; *C.A.*, 111 (1989) 170512w.
- 2532 Liu, D., Dass, C., Wood, G. and Desiderio, D.M.: Opioid and tachykinin peptides, and their precursors and precursor-processing enzymes, in human cerebrospinal fluid. *J. Chromatogr.*, 500 (1990) 395-412.
- 2533 Lunte, S.M. and Wong, O.S.: Naphthalenedialdehyde-cyanide: a versatile fluorescent reagent for the LC analysis of peptides and other primary amines. *LC-GC*, 7 (1989) 908-916; *C.A.*, 112 (1990) 3690s.

- 2534 McLeod, A.N., Auf der Mauer, A. and Wood, S.P.: High-performance liquid chromatography of insulin. Accessibility and flexibility. *J. Chromatogr.*, 502 (1990) 325-336.
- 2535 Mega, T., Nakamura, N. and Ikenaka, T.: Modifications of substituted seryl and threonyl residues in phosphopeptides and a polysialoglycoprotein by β -elimination and nucleophile additions. *J. Biochem. (Tokyo)*, 107 (1990) 68-72.
- 2536 Menegatti, E., Scalia, S., Bortolotti, F., Ascenzi, P. and de Marco, A.: Controlled proteolysis of mouse epidermal growth factor. An RP-HPLC and $^1\text{H-NMR}$ study. *Int. J. Pept. Protein Res.*, 34 (1989) 161-165; *C.A.*, 111 (1989) 225411v.
- 2537 Moreau, T., Hoebeke, J., Lalmanach, G., Hattab, M. and Gauthier, F.: Simulation of the inhibitory cystatin surface by a synthetic peptide. *Biochem. Biophys. Res. Commun.*, 167 (1990) 117-122.
- 2538 Nagarajan, G.R., Boone, J.L.S., Stolzer, T.J. and Richieri, S.P.: Effects of pH and ion pairing agents on the reverse and its fragments. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 357-363; *C.A.*, 112 (1990) 3765v.
- 2539 Newcomb, R.: Comprehensive analysis of the peptide content of neuronal tissue. *LC-GC*, 7 (1989) 570-572; *C.A.*, 111 (1989) 170247p - a review with 16 refs.
- 2540 Svardal, A.M., Mansoor, M.A. and Ueland, P.M.: Determination of reduced, oxidized, and protein-bound glutathione in human plasma with precolumn derivatization with monobromobimane and liquid chromatography. *Anal. Biochem.*, 184 (1990) 338-346.
- 2541 Svensson, B., Clarke, A.J., Svendsen, I. and Moller, H.: Identification of carboxylic acid residues in glucoamylase G2 from *Aspergillus niger* that participate in catalysis and substrate binding. *Eur. J. Biochem.*, 188 (1990) 29-38.
- 2542 Szokan, G., Mezö, G., Hudecz, F., Majer, Z., Schön, I., Nyeki, O., Szirtes, T. and Döbling, R.: Racemization analyses of peptides and amino acid derivatives by chromatography with pre-column derivatization. *J. Liq. Chromatogr.*, 12 (1989) 2855-2875.
- 2543 Tamburini, P.P., Koehn, J.A., Gilligan, J.P., Charles, D., Palmesino, R.A., Sharif, R., McMartin, C., Eriton, M.D. and Miller, M.J.S.: Rat vascular tissue contains a neutral endopeptidase capable of degrading atrial natriuretic peptide. *J. Pharmacol. Exp. Ther.*, 251 (1989) 956-961.
- 2544 Tateyama, H., Hino, J., Minamino, N., Kangawa, K., Ogiwara, T. and Matsuo, H.: Characterization of immunoreactive brain natriuretic peptide in human cardiac atrium. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1080-1087.
- 2545 Voyksner, R.D. and Pack, T.W.: Semiquantitative method for high-molecular-weight neuropeptides by high-performance liquid chromatography/thermospray mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 897-903; *C.A.*, 111 (1989) 209253q.
- 2546 Watson, D.C., Yaguchi, M. and Lynn, K.R.: The amino acid sequence of chymopapain from *Carica papaya*. *Biochem. J.*, 266 (1990) 75-81.
- 2547 Wring, S.A., Hart, J.P. and Birch, B.J.: Determination of glutathione in human plasma using high-performance liquid chromatography with electrochemical detection with a carbon-epoxy resin composite electrode chemically modified with cobalt phthalocyanine. *Analyst (London)*, 114 (1989) 1571-1573.
- 2548 Yip, T.T. and Hutchens, T.W.: Development of high-performance immobilized metal affinity chromatography for the separation of synthetic peptides and proteolytic digestion products. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (1989) 45-66; *C.A.*, 111 (1989) 170324m.

For additional information see:

C.A., 111 (1989) 167522u, 170351t, 170363y, 170474k, 187677b, 188875b, 228459c, 228563g.

See also 2578, 2587, 2592, 2734, 2744, 2762, 2766, 2775, 3020.

18c. Elucidation of structure of proteins and enzymes

- 2549 Dong, M.W. and Tran, A.D.: Factors influencing the performance of peptide mapping by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 499 (1990) 125-139.
- 2550 Fukamizo, T., Goto, S., Torikata, T. and Araki, T.: State of binding subsites in Asp 101-modified lysozymes. *J. Biochem. (Tokyo)*, 107 (1990) 445-451.
- 2551 Kamei, K., Oshino, R. and Hara, S.: Amino acid sequence of glutathione S-transferase b from guinea pig liver. *J. Biochem. (Tokyo)*, 107 (1990) 111-117.
- 2552 Kubota, N., Orita, T., Hattori, K., Oh-eda, M., Ochi, N. and Yamazaki, T.: Structural characterization of natural and recombinant human granulocyte colony-stimulating factors. *J. Biochem. (Tokyo)*, 107 (1990) 486-492.
- 2553 Liu, S.-Y., Yoshizumi, K., Oda, N., Ohno, M., Tokunaga, F., Iwanaga, S. and Kihara, H.: Purification and amino acid sequence of basic protein II, a lysine-49-phospholipase A₂ with low activity, from *Trimeresurus flavoviridis* venom. *J. Biochem. (Tokyo)*, 107 (1990) 400-408.
- 2554 Mori, K., Fujii, R., Kida, N., Takahashi, H., Ohkubo, S., Fujino, M., Ohta, M. and Hayashi, K.: Complete primary structure of the human estrogen-responsive gene (pS2) product. *J. Biochem. (Tokyo)*, 107 (1990) 73-76.
- 2555 Nakano, Y., Itoh, M., Tanaka, E. and Kimura, K.: Identification of the reactive cysteinyl residue and ATP binding site in *Bacillus cereus* glutamine synthetase by chemical modification. *J. Biochem. (Tokyo)*, 107 (1990) 180-183.
- 2556 Ohsawa, F., Hirano, F. and Natori, S.: Purification and properties of a β -galactoside-binding lectin from neonatal marmoset. *J. Biochem. (Tokyo)*, 107 (1990) 431-434.
- 2557 Pernollet, J.C., Potier, B., Galle, A.M., Huet, J.C., Beauvais, F. and Sallantin, M.: 2D-HPLC separation, electrophoretic characterization and N-terminal sequences of oat seed prolamins. *Phytochemistry*, 28 (1989) 2565-2570; *C.A.*, 112 (1990) 2847m.
- 2558 Schlabach, T.D., Colburn, J.C., Mattaliano, R.J. and Yuen, S.: Meeting the challenge in peptide fragment purification for protein sequencing. In: Hugli, T.E. (Editor), *Tech. Protein. Chem.*, Academic, San Diego, 1989, pp. 497-505; *C.A.*, 111 (1989) 211274r.
- 2559 Sievert, H.-J.P., Wu, S.-L., Chloupek, R. and Hancock, W.S.: Automated evaluation of tryptic digest from recombinant human growth hormone using ultraviolet spectra and numeric peak information. *J. Chromatogr.*, 499 (1990) 221-234.
- 2560 Stone, K.L., LoPresti, M.B., Williams, N.D., Crawford, J.M., DeAngelis, R. and Williams, K.R.: Enzymic digestion of proteins and HPLC peptide isolation in the sub-nanomole range. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 377-391; *C.A.*, 111 (1989) 228251d.
- 2561 Wittmann-Liebold, B.: Moderne chemische Strukturanalyse von Proteinen und Peptiden: Strategien, Methoden, Einsatzmöglichkeiten. *Kontakte (Darmstadt)*, No. 3 (1989) 16-28 - a review with 60 refs.
- 2562 Yoshizaki, H., Mizoguchi, T., Arai, K., Shiratsuchi, M., Shidara, Y. and Maki, M.: Structure and properties of calphobindin II, an anticoagulant protein from human placenta. *J. Biochem. (Tokyo)*, 107 (1990) 43-50.

See also 2576, 2626.

19. PROTEINS

19a. General techniques

- 2563 Anspach, F.B., Johnston, A., Wirth, H.-J., Unger, K.K. and Hearn, M.T.W.: High-performance liquid chromatography of amino acids, peptides and proteins. XCV. Thermodynamic and kinetic investigations on rigid and soft affinity gels with varying particle and pore sizes: comparison of thermodynamic parameters and the adsorption behaviour of proteins evaluated from batch and frontal analysis experiments. *J. Chromatogr.*, 499 (1990) 103-124.
- 2564 Burkey, M.M. and Jorgenson, J.W.: Automated instrumentation for comprehensive two-dimensional high performance liquid chromatography of proteins. *Anal. Chem.*, 62 (1990) 161-167.
- 2565 Davankov, V.A., Kurganov, A.A. and Unger, K.K.: Reversed-phase high-performance liquid chromatography of proteins and polypeptides on polystyrene-coated silica supports. *J. Chromatogr.*, 500 (1990) 519-530.
- 2566 El Rassl, Z., de Ocampo, L.F. and Bacolod, M.D.: Binary and ternary salt gradients in hydrophobic-interaction chromatography of proteins. *J. Chromatogr.*, 499 (1990) 141-152.
- 2567 Engelhardt, H.: (HPLC of proteins). *GIT Fachz. Lab.*, 33 (1989) 699-705; *C.A.*, 111 (1989) 247204g.
- 2568 Gisach, D.J.: Rapid protein matrix reduction by a hydrophobic interaction-based solid phase extraction method. *BioTechniques*, 7 (1989) 562-564; *C.A.*, 111 (1989) 170528f.
- 2569 Glemza, A., Baskeviciute, B., Kadusevicius, V., Pesliakas, J. and Sudziuviene, O.: The influence of matrices on dye-protein interaction. In: Vijayalakshmi, M.A. and Bertrand, O. (Editors), *Protein-Dye Interact.*, [Proc. Int. Conf.] 1st 1988, Elsevier, Barking, 1989, pp. 107-114; *C.A.*, 111 (1989) 190628s.
- 2570 Goetz, H.: An evaluation of new microbore and high-speed columns for protein separations using HPLC and derivative spectroscopy. *BioChromatography*, 4 (1989) 159-160; *C.A.*, 111 (1989) 228243c.
- 2571 Hammond, P.M. and Scawen, M.D.: High-resolution fractionation of proteins in downstream processing. *J. Biotechnol.*, 11 (1989) 119-134; *C.A.*, 111 (1989) 190660w.
- 2572 Han, Y.: (Determination of protein molecular weight by fast protein liquid chromatography.) *Sepu*, 7 (1989) 186; *C.A.*, 111 (1989) 190649z.
- 2573 Hutchens, T.W. and Yip, T.-T.: Differential interaction of peptides and protein surface structures with free metal ions and surface-immobilized metal ions. *J. Chromatogr.*, 500 (1990) 531-542.
- 2574 Jagschies, G.: Chromatographic purification of proteins-support from physico-chemical research. *Ber. Bunsen-Ges. Phys. Chem.*, 93 (1989) 1004-1009; *C.A.*, 111 (1989) 211210s - a review with 10 refs.
- 2575 Jervis, L.: Fast analytical dye-ligand chromatography in process development. In: Vijayalakshmi, M.A. and Bertrand, O. (Editors), *Protein-Dye Interact.*, [Proc. Int. Conf.], 1st 1988, Elsevier, Barking, 1989, pp. 121-123; *C.A.*, 111 (1989) 228244d.
- 2576 Kalghatgi, K.: Micropellicular stationary phases for rapid protein analysis by high-performance liquid chromatography. *J. Chromatogr.*, 499 (1990) 267-278.
- 2577 Karger, B.L. and Blanco, R.: Protein conformational changes upon adsorption to chromatographic surfaces. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (1989) 141-148; *C.A.*, 111 (1989) 170327q.
- 2578 Kato, Y., Nakatani, S., Kitamura, T., Yamasaki, Y. and Hashimoto, T.: Reversed-phase high-performance liquid chromatography of proteins and peptides on a pellicular support based on hydrophilic resin. *J. Chromatogr.*, 502 (1990) 416-422.

- 2579 Khoklova, T.D., Nikitin, Yu.S. and Voroshilova, O.I.: (Macroporous silicas in immobilization and chromatography of biopolymers.) *Zh. Vses. Khim. O-va. im. D.I. Mendeleeva*, 34 (1989) 363-367; *C.A.*, 111 (1989) 190551m - a review with 23 refs.
- 2580 Kula, M.R., Walsdorf, A. and Cordes, A.: Affinity partitioning of proteins. *Ber. Bunsen-Ges. Phys. Chem.*, 93 (1989) 968-970; *C.A.*, 111 (1989) 228285t.
- 2581 Lesins, V. and Ruckenstein, E.: Chromatographic probing of protein-sorbent interactions. *J. Colloid Interface Sci.*, 132 (1989) 566-577; *C.A.*, 112 (1990) 12445t.
- 2582 Lin, S. and Karger, B.L.: Reversed-phase chromatographic behavior of proteins in different unfolded states. *J. Chromatogr.*, 499 (1990) 89-102.
- 2583 Mazzaroff, I., Varady, L., Mouchawar, G.A. and Regnier, F.E.: Thermodynamic model for electrostatic-interaction chromatography of proteins. *J. Chromatogr.*, 499 (1990) 63-77.
- 2584 McLoughlin, S.B. and Lowe, C.R.: Applications of triazinyl dyes in protein purification. *Rev. Prog. Color. Relat. Top.*, 18 (1988) 16-28; *C.A.*, 111 (1989) 190491s - a review with 163 refs.
- 2585 Mhatre, R., Krull, I.S. and Stuting, H.H.: Determination of biopolymer (protein) molecular weights by gradient elution, reversed-phase high-performance liquid chromatography with low-angle laser light scattering detection. *J. Chromatogr.*, 502 (1990) 21-46.
- 2586 Moriya, K., Tanizawa, K. and Kanaoka, Y.: Schiff base copper II chelate as a tool for immobilization of protein. *Chem. Pharm. Bull.*, 37 (1989) 2849-2851.
- 2587 Nice, E.C. and Simpson, R.J.: Micropreparative high-performance liquid chromatography of proteins and peptides. *J. Pharm. Biomed. Anal.*, 7 (1989) 1039-1053.
- 2588 Oscarsson, S. and Porath, J.: Protein chromatography with pyridine- and alkyl-thioether-based agarose adsorbents. *J. Chromatogr.*, 499 (1990) 235-247.
- 2589 Porath, J.: Electron-donor-acceptor chromatography EDAC for biomolecules in aqueous solutions. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (Protein Recognit. Immobilized Ligands), (1989) 101-122; *C.A.*, 111 (1989) 190493u - a review with 32 refs.
- 2590 Scopes, R.K.: The use of biospecific elution in purifying proteins from dye-ligand adsorbents. In: Vijayalakshmi, M.A. and Bertrand, O. (Editors), *Protein-Dye Interact., [Proc. Int. Conf.], 1st 1988*, Elsevier, Barking, 1989, pp. 97-106; *C.A.*, 111 (1989) 190627r.
- 2591 Shimizu, E., Kiguchi, S. and Yorifuji, T.: A simple approximation of weight-based protein measurement with HPLC apparatus. *Agric. Biol. Chem.*, 53 (1989) 2801-2803; *C.A.*, 112 (1990) 3789f.
- 2592 Southan, C.: The use of glass capillary tubes as disposable microbore columns for RP-HPLC of proteins and peptides. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 392-398; *C.A.*, 111 (1989) 228189q.
- 2593 Stewart, D.J., Hughes, P. and Lowe, C.R.: Affinity chromatography on triazine dyes immobilized on novel perfluorocarbon supports. *J. Biotechnol.*, 11 (1989) 253-265; *C.A.*, 111 (1989) 228276r.
- 2594 Ujhazy, A., Zsados, B. and Szejtli, J.: Gel chromatographic separation of proteins on cyclodextrin bead polymer stationary phases. *J. Liq. Chromatogr.*, 12 (1989) 2877-2885.
- 2595 Wajcman, H.: (Recent progress in the application of HPLC to protein separation). *Colloq. INSERM*, 175 (Biotechnol. Proteines Plasma) (1989) 137-143; *C.A.*, 112 (1990) 3672n - a review with 16 refs.
- 2596 Wilson, K.J.: Micro-level protein and peptide separations. *Trends Biochem. Sci. (Pers. Ed.)*, 14 (1989) 252-255; *C.A.*, 111 (1989) 190496x - a review with 19 refs.
- 2597 Yang, B.L., Goto, M. and Goto, S.: Enzyme purification by affinity chromatography combined with batchwise adsorption and columnwise elution. *J. Chem. Eng. Jpn.*, 22 (1989) 532-537; *C.A.*, 112 (1990) 3785b.

For additional information see:

C.A., 111 (1989) 170607f, 173147s, 173151p, 228558j.

See also 2198, 2216, 2519, 2780.

19b. Proteins of cells, viruses and subcellular particles

2598 Crooks, A.J., Lee, J.M., Dowsett, A.B. and Stephenson, J.R.: Purification and analysis of infectious virions and native non-structural antigens from cells infected with tick-borne encephalitis virus. *J. Chromatogr.*, 502 (1990) 59-68.

2599 Hayashi, Y., Matsui, H. and Tagaki, T.: Membrane protein molecular weight determined by low-angle laser light-scattering photometry coupled with high-performance gel chromatography. *Methods Enzymol.*, 172 (Biomembranes, Pt. S), (1989) 514-528; C.A., 111 (1989) 190903c.

2600 Kaiser, D.A., Goldschmidt-Clermont, P.J., Levine, B.A. and Pollard, T.D.: Characterization of renatured profilin purified by urea elution from poly-L-proline agarose columns. *Cell Motil. Cytoskeleton*, 14 (1989) 251-262; C.A., 112 (1990) 3798h.

See also 2362, 2672.

19c. Proteins synthesized by genetic manipulation

2601 Boschetti, E.: Applications of chromatography in downstream processing. *NATO ASI Ser., Ser. E*, 156 (1989) 401-418; C.A., 112 (1990) 5974m - a review with 8 refs.

2602 Diaz-Alonso, J.M., Kohnert, K.D., Keilacker, H., Ziegler, B. and Ziegler, M.: Purification of mouse monoclonal antibodies by chromatography on Sepharose 6 B. *Allerg. Immunol. (Leipzig)*, 35 (1989) 173-179; C.A., 111 (1989) 230210h.

2603 Minamitake, Y., Kodama, S., Katayama, T., Adachi, H., Tanaka, S. and Tsujimoto, M.: Structure of recombinant human interleukin 5 produced by Chinese hamster ovary cells. *J. Biochem. (Tokyo)*, 107 (1990) 292-297.

2604 Prickett, K.S., Amberg, D.C. and Hopp, T.P.: A calcium-dependent antibody for identification and purification of recombinant proteins. *BioTechniques*, 7 (1989) 580-589; C.A., 111 (1989) 228422k.

2605 Torres, A.R. and Peterson, E.A.: Purification of monoclonal antibodies by complex-displacement chromatography on CM-cellulose. *J. Chromatogr.*, 499 (1990) 47-54.

2606 Wu, S.-L., Hancock, W.S., Pavlu, B. and Gellerfors, P.: Application of high-performance hydrophobic-interaction chromatography to the characterization of recombinant DNA-derived human growth hormone. *J. Chromatogr.*, 500 (1990) 595-606.

See also 2651.

19d. Microbial and plant proteins

2607 Annan, R.S., Kim, C. and Martyn, J.: Measurement of D-tubocurarine chloride in human urine using solid-phase extraction and reversed-phase high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr.*, 526 (1990) 228-234.

2608 Cocchiara, R., Locorotondo, G., Parlato, A., Guarnotta, G., Ronchi, S., Albeggiani, G., Amoroso, S., Falgiani, P. and Geraci, D.: Purification of Par j I, a major allergen from *Parietaria judaica* pollen. *Int. Arch. Allergy Appl. Immunol.*, 90 (1989) 84-90; C.A., 111 (1989) 212961z.

- 2609 Hussain, A., Scanlon, M.G., Juliano, B.O. and Bushuk, W.: Discrimination of rice cultivars by polyacrylamide gel electrophoresis and high-performance liquid chromatography. *Cereal Chem.*, 66 (1989) 353-356; *C.A.*, 111 (1989) 211442u.
- 2610 Sanders, D.A., Gillece-Castro, B.L., Stock, A.M., Burlingame, A.L. and Koshland, D.E., Jr.: Identification of the site of phosphorylation of the chemotaxis response regulator protein, CheY. *J. Biol. Chem.*, 264 (1989) 21770-21778.
- 2611 Somlyo, B., Kovats, E., Keler, T. and Nowotny, A.: Column liquid chromatography of endotoxins. *J. Chromatogr.*, 525 (1990) 329-338.
- 2612 Yamakawa, Y., Sato, H. and Sato, Y.: Isolation of pertussis toxin subunit proteins by reverse-phase high-performance liquid chromatography and reconstitution of the holotoxin molecule. *Anal. Biochem.*, 185 (1990) 176-181.

For additional information see:

C.A., 111 (1989) 170613e, 211263m, 211271n.

19e. Proteins of blood, serum and blood cells

- 2613 Birkenmeier, G. and Kopperschlaeger, G.: Application of dyes in serum protein separation. *Colloq. INSERM*, 175 (1989) 217-228; *C.A.*, 112 (1990) 25410c - a review with 40 refs.
- 2614 Birkenmeier, G., Huse, K. and Kopperschlaeger, G.: The interaction of human α -fetoprotein with reactive dyes and the isolation of the protein by affinity chromatography. In: Vijayalakshmi, M.A. and Bertrand, O. (Editors), *Protein-Dye Interact.*, [Proc. Int. Conf.], 1st 1988, Elsevier, Barking, 1989, pp. 275-280; *C.A.*, 111 (1989) 190630m.
- 2615 Bouillon, R., Coopmans, W., Degroote, D.E.H., Radoux, D. and Eliard, P.H.: Immunoradiometric assay of parathyrin with polyclonal and monoclonal region-specific antibodies. *Clin. Chem. (Winston-Salem)*, 36 (1990) 271-276.
- 2616 Cianflone, K., Wu, H., Walsh, M., Baldu, A. and Sniderman, A.: Metabolic response of acylation stimulating protein to an oral fat load. *J. Lipid Res.*, 30 (1989) 1727-1733.
- 2617 Delaage, M., Mejan, O. and Bourgois, A.: Affinity chromatography and ion-exchange effects. The case of factor VIII-von Willebrandt factor complex. *Colloq. INSERM*, 175 (1989) 145-154; *C.A.*, 112 (1990) 62410y.
- 2618 Dodds, A.W. and Law, S.K.A.: The complement component C4 of mammals. *Biochem. J.*, 265 (1990) 495-502.
- 2619 Guiguet, M., Dethleux, M.C., Frigere, M.F., Bidan, Y., Bielefeld, P. and Mack, G.: Purification of the third component of rat complement using high performance liquid chromatography. *Colloq. INSERM*, 175 (1989) 253-259; *C.A.*, 111 (1989) 212971c.
- 2620 Harrison, P., Saundry, R.H. and Savidge, G.F.: The purification of the VIII/vWF complex using dextran sulphate Sepharose chromatography. *Colloq. INSERM*, 175 (1989) 279-285; *C.A.*, 112 (1990) 11822b.
- 2621 Haun, M., Incledon, B., Alles, P. and Wasi, S.: A rapid procedure for the purification of IgA1 and IgA2 subclasses from normal human serum using protein G and jackfruit lectin (jacalin) affinity chromatography. *Immunol. Lett.*, 22 (1989) 273-279; *C.A.*, 111 (1989) 230205k.
- 2622 Hocke, G., Kaffarnik, H., Münscher, G. and Steinmetz, A.: Purification of human serum amyloid A by anion-exchange fast protein liquid chromatography. *J. Chromatogr.*, 526 (1990) 203-209.
- 2623 Klein, M.L., Bartley, T.D., Davis, J.M., Whiteley, D.W. and Lu, H.S.: Isolation and structural characterization of three isoforms of recombinant consensus α interferon. *Arch. Biochem. Biophys.*, 276 (1990) 531-537.

- 2624 Koike, M., Akiyama, H. and Shimizu, T.: High-performance liquid chromatographic procedure for the determination of rat plasma concentrations of a new antibacterial agent, ($\pm$)-9-fluoro-6,7-dihydro-8-(4-hydroxy-1-piperidyl)-5-methyl-1-oxo-1H,5H-benzo[*i,j*]quinolizine-2-carboxylic acid, for topical use. *J. Chromatogr.*, 526 (1990) 235-239.
- 2625 Kosyak, A.V., Shukun, S.A., Poddubkova, T.N., Chikh, V.P. and Surovtsev, V.I.: (Human complement system components. Components C3 and C5: testing and isolation). *Biokhimiya (Moscow)*, 55 (1990) 276-284.
- 2626 Kumazaki, T. and Ishii, S.-i.: Disulfide bridge structure of ascidian trypsin inhibitor I: similarity to Kazal-type inhibitors. *J. Biochem. (Tokyo)*, 107 (1990) 414-419.
- 2627 Mbida, A., Kanoun, S. and Vijayalakshmi, M.A.: Purification of IgG subclass from human placenta by pseudo affinity chromatography. *Colloq. INSERM*, 175 (1989) 237-244; *C.A.*, 111 (1989) 212970b.
- 2628 More, J.E., Hitchcock, A.G., Price, S., Rott, J. and Harvey, M.J.: Recovery of albumin from cohn fraction IV using immobilized Procion Blue-HB: development of a full scale production process. In: Vijayalakshmi, M.A. and Bertrand, O. (Editors), *Protein-Dye Interact.*, [Proc. Int. Conf.], 1st 1988, Elsevier, Barking, 1989, pp. 265-274; *C.A.*, 111 (1989) 190629t.
- 2629 Naylor, S., Gan, L.-S., Day, B.W., Skipper, P.L. and Tannenbaum, S.R.: A rapid method for isolation of human hemoglobin benzo[*a*]pyrene diol epoxide derived adducts using high performance liquid chromatography. *Biomed. Chromatogr.*, 3 (1989) 266-268.
- 2630 Nishimura, K., Harada, K., Masuda, S. and Sugii, A.: High-performance liquid chromatography of serum albumins on an N-methylpyridinium polymer-based column. *J. Chromatogr.*, 525 (1990) 176-182.
- 2631 Otnad, E., Via, D.P., Sinn, H., Friedrich, E., Ziegler, R. and Dresel, H.A.: Binding characteristics of reduced hepatic receptors for acetylated low-density lipoprotein and maleylated bovine serum albumin. *Biochem. J.*, 265 (1990) 689-698.
- 2632 Plaigin, M., Lacoste-Bourgeacq, J.F., Mandaro, R. and Lemay, C.: Purification of plasma fractions by radial flow chromatography cartridges. *Colloq. INSERM*, 175 (1989) 169-175; *C.A.*, 112 (1990) 11821a.
- 2633 Sanchez-Corral, P., Anton, L.C., Alcolea, J.M., Marques, G., Sanchez, A. and Vivanco, F.: Separation of active and inactive forms of the third component of human complement C3, by fast protein liquid chromatography (FPLC). *J. Immunol. Methods*, 122 (1989) 105-113; *C.A.*, 111 (1989) 192681c.
- 2634 Silvestrini, B., Guglielmotti, A., Saso, L., Milanese, C., Melanitou, E., Grima, J. and Cheng, C.Y.: Development of an enzyme-linked immunosorbent assay with a monoclonal antibody prepared against α_1 -antitrypsin for diagnostic screening of inflammatory disorders. *Clin. Chem. (Winston-Salem)*, 36 (1990) 277-282.
- 2635 Stoltz, J.F., Rivat, C., Geschier, C., Colosetti, P. and Sertillanges, P.: Chromatographic purification of human albumin. Technical and economic aspects. *Colloq. INSERM*, 175 (1989) 191-200; *C.A.*, 112 (1990) 62411z.
- 2636 Suehiro, K., Kawabata, S., Miyata, T., Takeya, H., Takamatsu, J., Ogata, K., Kamiya, T., Saito, H., Niho, Y. and Iwanaga, S.: Blood clotting factor IX B_H Nagoya. Substitution of arginine 180 by tryptophan and its activation by α -chymotrypsin and rat mast cell chymase. *J. Biol. Chem.*, 264 (1989) 21257-21265.
- 2637 Te Booy, M.P.W.M., Faber, A., de Jong, E., Wolterink, E.P., Riethorst, W., Beugeling, T., Bantjes, A., Over, J. and König, B.W.: Large-scale purification of factor VIII by affinity chromatography: optimization of process parameters. *J. Chromatogr.*, 503 (1990) 103-114.

- 2638 Tedesco, J.L., Ohlson, S., Holmberg, A. and Rupp, R.: Evaluation of a new high performance liquid affinity chromatography system for rapid quantitation of IgG in cell culture media. *BioChromatography*, 4 (1989) 216-220; C.A., 111 (1989) 212966e.
- 2639 Tiusch, D., Allary, M., Saint-Blancard, J. and Boschetti, E.: Preparative purification of IgG from a human plasma alcohol precipitate. *Colloq. INSERM*, 175 (1989) 229-236; C.A., 111 (1989) 212969h.
- 2640 Van Oers, N.S.C., Boismenu, R., Cohen, B.L. and Murgita, R.A.: Analytical- and preparative-scale separation of molecular variants of α -fetoprotein by anion-exchange chromatography on monobeadTM resins. *J. Chromatogr.*, 525 (1990) 59-69.
- 2641 Watts, M.J., Dankert, J.R. and Morgan, B.P.: Isolation and characterization of a membrane-attack-complex inhibiting protein present in human serum and other biological fluids. *Biochem. J.*, 265 (1990) 471-477.

For additional information see:

C.A., 111 (1989) 169956n, 170610b, 172021j, 172029p, 190992f, 228284s,
228292t, 228601t;
112 (1990) 5931v, 11901b, 42549x.

See also 2261, 2263, 2445, 2605, 2667, 2754.

19f. Structural and muscle proteins

- 2642 Bessho, K., Tagawa, T. and Murata, M.: Purification of bone morphogenetic protein derived from bovine bone matrix. *Biochem. Biophys. Res. Commun.*, 165 (1989) 595-601.
- 2643 Kirsch, T. and van der Mark, K.: Isolation of bovine type X collagen and immunolocalization in growth-plate cartilage. *Biochem. J.*, 265 (1990) 453-459.
- 2644 Leszyk, J., Grabarek, Z., Gergely, J. and Collins, J.H.: Characterization of zero-length cross-links between rabbit skeletal muscle troponin C and troponin I: evidence for direct interaction between the inhibitory region of troponin I and the NH₂-terminal, regulatory domain of troponin C. *Biochemistry*, 29 (1990) 299-304.
- 2645 Villanueva, M.A., Ho, S.-C. and Wang, J.L.: Isolation and characterization of one isoform of actin from cultured soybean cells. *Arch. Biochem. Biophys.*, 277 (1990) 35-41.

19g. Protamines, histones and other chromosomal proteins

- 2646 Gohill, J., Pauls, J.D. and Fritzler, M.J.: Purification of histone H1 polypeptides by high-performance cation-exchange chromatography. *J. Chromatogr.*, 502 (1990) 47-57.
- 2647 vanfleteren, J.R., van Bun, S.M., de Baere, I. and van Beeumen, J.J.: The primary structure of a minor isoform (H1.2) of histone H1 from the nematode *Caenorhabditis elegans*. *Biochem. J.*, 265 (1990) 739-746.

19h. Chromoproteins and metalloproteins

- 2648 Ena, J.M., Castillo, H., Sanchez, L. and Calvo, M.: Isolation of human lactoferrin by affinity chromatography using insolubilized bovine β -lactoglobulin. *J. Chromatogr.*, 525 (1990) 442-446.
- 2649 Hidalgo, J., Dingman, A. and Garvey, J.S.: Preparative isolation of adult human liver metallothionein isoforms. *Rev. Esp. Fisiol.*, 45 (1989) 255-263; C.A., 112 (1990) 3769z.

- 2650 Hjerten, S. and Li, J.-P.: High-performance liquid chromatography of proteins on deformed non-porous agarose beads. Fast boronate affinity chromatography of haemoglobin at neutral pH. *J. Chromatogr.*, 500 (1990) 543-553.
- 2651 Johansson, C., Brodin, P., Grundström, T., Thullin, E., Forsén, S. and Drakenberg, T.: Biophysical studies of engineered mutant proteins based on calbindin D_{9k} modified in the pseudo EF-hand. *Eur. J. Biochem.*, 187 (1990) 455-460.
- 2652 Kobayashi, R., Hidaka, H. and Tashima, Y.: Purification, characterization, and partial sequence analysis of 32-kDa calcimedin from chicken gizzard. *Arch. Biochem. Biophys.*, 277 (1990) 203-210.
- 2653 Rahbar, S. and Asmerom, Y.: Rapid HPLC techniques for globin chain synthesis studies. *Hemoglobin*, 13 (1989) 475-487; *C.A.*, 112 (1990) 3777a.

For additional information see:

C.A., 111 (1989) 211273q, 228280n.

19i. *Proteins of glands, gland products, various zymogens (including milk proteins)*

- 2654 Baier, G., Wollensak, G., Mur, E., Redl, B., Stöffler, G. and Göttinger, W.: Analysis of human tear proteins by different high-performance liquid chromatographic techniques. *J. Chromatogr.*, 525 (1990) 319-328.
- 2655 Beyer, H.J. and Kessler, H.G.: (Determination of heat-induced whey protein denaturation by HPLC). *GIT-Suppl.*, No. 2 (1989) 22-24; *C.A.*, 111 (1989) 193264f.
- 2656 Blankenship, D.T., Brankamp, R.G., Manley, G.D. and Cardin, A.D.: Amino acid sequence of ghilanten: anticoagulant-antimetastatic principle of the south American leech, *Haementeria ghilianii*. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1384-1389.
- 2657 Christensen, T.M.I.E. and Munksgaard, L.: Quantitative fractionation of casein by precipitation or ion exchange chromatography. *Milchwissenschaft*, 44 (1989) 480-484; *C.A.*, 111 (1989) 230770r.
- 2658 Kisugi, J., Yamazaki, M., Ishii, Y., Tansho, S., Muramoto, K. and Kamiya, H.: Purification of a novel cytolytic protein from albumin gland of the sea hare, *Dolabella auricularia*. *Chem. Pharm. Bull.*, 37 (1989) 2773-2776.
- 2659 Loret, E.P., Manuelle, P., Rochat, H. and Granier, C.: Neurotoxins active on insects: amino acid sequences, chemical modifications, and secondary structure estimation by circular dichroism of toxins from the scorpion *Androctonus australis* Hector. *Biochemistry*, 29 (1990) 1492-1501.
- 2660 Oroszlan, P., Blanco, R., Lu, X.-M., Yarmush, D. and Karger, B.L.: Intrinsic fluorescence studies of the kinetic mechanism of unfolding of α -lactalbumin on weakly hydrophobic chromatographic surfaces. *J. Chromatogr.*, 500 (1990) 481-502.
- 2661 Sanogo, T., Paquet, D., Aubert, F. and Linden, G.: Purification of α_{s1} -casein by fast protein liquid chromatography. *J. Dairy Sci.*, 72 (1989) 2242-2246; *C.A.*, 112 (1990) 6232e.
- 2662 Shinomiya, T., Ohara, T., Wada, N., Omori, A. and Kamada, N.: Rat liver arginase suppresses mixed lymphocyte reaction. *J. Biochem. (Tokyo)*, 107 (1990) 435-439.

For additional information see:

C.A., 112 (1990) 6085j.

See also 2648.

19j. *Proteins of brain, cerebrospinal fluid and eye*

- 2663 Joniau, M., Coudijzer, K. and de Cuyper, M.: Reaction of α -tubulin with iodotyrosines catalyzed by tubulin:tyrosine ligase: carboxy-terminal labeling of tubulin with [125 I]monoiodotyrosine. *Anal. Biochem.*, 184 (1990) 325-329.
- 2664 Lacey, E. and Snowdon, K.L.: Isolation of mammalian brain tubulin by amino-activated gel chromatography. *J. Chromatogr.*, 525 (1990) 71-84.
- 2665 Landon, M., Smith, M.A., Clarke, K. and Kidd, M.: Immunoaffinity chromatographic purification of amyloid-related proteins from Alzheimer's disease brain tissue. *Prog. Clin. Biol. Res.*, 317 (1989) 903-912; *C.A.*, 111 (1989) 228443t.

19k. *Proteins of neoplastic tissue and transformed cells*

- 2666 Barbero, M.C., Garcia, M., Almena, A. and Alino, S.F.: HPLC analysis of supernatant soluble factors from tumor cell variants with defined metastatic status. *Cell. Mol. Biol.*, 35 (1989) 441-447; *C.A.*, 112 (1990) 3790z.

19l. *Specific binding and receptor proteins*

- 2667 Allary, M., Boschetti, E., Lafargue, D., Sene, C. and Saint Blancard, J.: Improved method for plasma fibronectin purification by double chromatographic fractionation on gelatin and heparin-trisacryl. *Colloq. INSERM*, 175 (1989) 61-66; *C.A.*, 112 (1990) 25433n.
- 2668 Fremont, S., Gueant, J.-L., Felden, F., Egloff, M., Gerard, A. and Nicolas, J.-P.: Dihydrotestosterone binding capacity of androgen-binding protein in tissue extract using high-performance size exclusion chromatography. *J. Chromatogr.*, 526 (1990) 186-193.
- 2669 Hu, W., Mazurier, J., Montreuil, J. and Spik, G.: Isolation and partial characterization of a lactotransferrin receptor from mouse intestinal brush border. *Biochemistry*, 29 (1990) 535-541.
- 2670 Ito, H., Yamamoto, H., Kimura, Y., Kambe, H., Okochi, T. and Kishimoto, S.: Affinity chromatography of human plasma gelsolin with polyphosphate compounds on immobilized Cibacron Blue F3GA. *J. Chromatogr.*, 526 (1990) 397-406.
- 2671 Johnsson, N. and Weber, K.: Structural analysis of p36, a Ca^{2+} /lipid-binding protein of the annexin family, by proteolysis and chemical fragmentation. *Eur. J. Biochem.*, 188 (1990) 1-7.
- 2672 Korner, M., Bellan, A.H., Brini, A.T. and Farrar, W.L.: Characterization of the human immunodeficiency virus type 1 enhancer binding proteins from the human T-cell line Jurkat. *Biochem. J.*, 265 (1990) 547-554.
- 2673 Leidenix, M.J., Jacoby, G.H., Henderson, T.A. and Young, K.D.: Separation of *Escherichia coli* penicillin-binding proteins into different membrane vesicles by agarose electrophoresis and sizing chromatography. *J. Bacteriol.*, 171 (1989) 5680-5686; *C.A.*, 111 (1989) 190958z.
- 2674 Mereau, A., Pieri, I., Gamby, C., Courty, J. and Barritault, D.: Purification of basic fibroblast growth factor receptors from bovine brain. *Biochimie*, 71 (1989) 865-871; *C.A.*, 111 (1989) 170369e.
- 2675 Miller, K.G., Field, C.M. and Alberts, B.M.: Actin-binding proteins from *Drosophila* embryos: a complex network of interacting proteins detected by F-actin affinity chromatography. *J. Cell Biol.*, 109 (6, Pt. 1) (1989) 2963-2976; *C.A.*, 111 (1989) 229631q.
- 2676 Nakajo, S., Sugiyura, M. and Inagami, T.: Native form of endothelin receptor in human placental membranes. *Biochem. Biophys. Res. Commun.*, 167 (1990) 280-286.
- 2677 Pang, I.H. and Sternweis, P.C.: Isolation of the α subunits of GTP-binding regulatory proteins by affinity chromatography with immobilized $\beta\gamma$ subunits. *Proc. Natl. Acad. Sci. U.S.A.*, 86 (1989) 203-208; *C.A.*, 112 (1990) 3778b.

- 2678 Raubenheimer, J. and van der Walt, L.A.: Multidimensional chromatographic separation of estrogen receptors. *Biomed. Chromatogr.*, 3 (1989) 255-261.
- 2679 Sairam, M.R.: Affinity chromatography of hormone receptors and antibodies on divinyl sulfonyl-sepharose. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (1989) 303-313; *C.A.*, 111 (1989) 211267r.
- 2680 Sviridov, O.V., Ermolenko, M.N., Pyshko, E.S., Karpova, T.V. and Strel'chyonok, O.A.: (Affinity chromatography of human blood serum thyroxine-binding proteins). *Biokhimiya (Moscow)*, 55 (1990) 329-337.
- 2681 Yannariello-Brown, J. and Madri, J.A.: A 48 kDa collagen-binding phosphoprotein isolated from bovine aortic endothelial cells interacts with the collagenous domain, but not the globular domain, of collagen type IV. *Biochem. J.*, 265 (1990) 383-392.

For additional information see:

C.A., 111 (1989) 170353v, 190669f, 209225g.

See also 2631, 3018.

19m. Urinary proteins

See 2350, 2641.

19n. Other proteins

- 2682 Dourges, M.A.J., Gulino, D., Courty, J., Badet, J., Barritault, D. and Jozefonvicz, J.: Affinity chromatography of fibroblast growth factors on substituted polystyrene. *J. Chromatogr.*, 526 (1990) 35-45.
- 2683 Gurley, L.R., London, J.E., Dethloff, L.A., Stavert, D.M. and Lehnert, B.E.: Analysis of proteins in bronchoalveolar lavage fluids during pulmonary edema resulting from nitrogen dioxide and cadmium exposure. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 479-489; *C.A.*, 111 (1989) 226681v.
- 2684 Vyas, K.P., Kari, P.H. and Pitzenberger, S.M.: Regioselectivity and stereoselectivity in the metabolism of HMG-CoA reductase inhibitors. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1155-1162.

For additional information see:

C.A., 111 (1989) 210185a.

See also 2756.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

20a. Oxidoreductases

- 2685 Abriola, D.P., MacKerell, A.D., Jr. and Pietruszko, R.: Correlation of loss of activity of human aldehyde dehydrogenase with reaction of bromoacetophenone with glutamic acid-268 and cysteine-302 residues. Partial-sites reactivity of aldehyde dehydrogenase. *Biochem. J.*, 266 (1990) 179-187.
- 2686 Britsch, L.: Purification of flavanone 3 β -hydroxylase from *Petunia hybrida*: antibody preparation and characterization of a chemogenetically defined mutant. *Arch. Biochem. Biophys.*, 276 (1990) 348-354.
- 2687 Fischer, D., Ebenau-Jehle, C. and Grisebach, H.: Phytoalexin synthesis in soybean: purification and characterization of NADPH:2'-hydroxydaidzein oxidoreductase from elicitor-challenged soybean cell cultures. *Arch. Biochem. Biophys.*, 276 (1990) 390-395.

- 2688 Halvorson, M., Greenway, D., Eberhart, D., Fitzgerald, K. and Parkinson, A.: Reconstitution of testosterone oxidation by purified rat cytochrome P450p (IIIA1). *Arch. Biochem. Biophys.*, 277 (1990) 166-180.
- 2689 Hatchikian, C.E., Traore, A.S., Fernandez, V.M. and Cammack, R.: Characterization of the nickel-iron periplasmic hydrogenase from *Desulfovibrio fructosovorans*. *Eur. J. Biochem.*, 187 (1990) 635-643.
- 2690 Heidlas, J. and Tressl, R.: Purification and properties of two oxidoreductases catalyzing the enantioselective reduction of diacetyl and other diketones from baker's yeast. *Eur. J. Biochem.*, 188 (1988) 165-174.
- 2691 Imaoka, S., Nagashima, K. and Funae, Y.: Characterization of three cytochrome P450s purified from renal microsomes of untreated male rats and comparison with human renal cytochrome P450. *Arch. Biochem. Biophys.*, 276 (1990) 473-480.
- 2692 Krook, M., Marekov, L. and Jörnvall, H.: Purification and structural characterization of placental NAD⁺-linked 15-hydroxyprostaglandin dehydrogenase. The primary structure reveals the enzyme to belong to the short-chain alcohol dehydrogenase family. *Biochemistry*, 29 (1990) 738-743.
- 2693 Nakagawa, M., Harada, T., Hara, A., Nakayama, T. and Sawada, H.: Purification and properties of multiple forms of dihydrodiol dehydrogenase from monkey liver. *Chem. Pharm. Bull.*, 37 (1989) 2852-2854.
- 2694 Neame, P.J. and Barber, M.J.: Conserved domains in molybdenum hydroxylases. The amino acid sequence of chicken hepatic sulfite oxidase. *J. Biol. Chem.*, 264 (1989) 20894-20901.
- 2695 Nozaki, M., Takeshige, K., Sumimoto, H. and Minakami, S.: Reconstitution of the partially purified membrane component of the superoxide-generating NADPH oxidase of pig neutrophils with phospholipid. *Eur. J. Biochem.*, 187 (1990) 335-340.
- 2696 Palmer, E.J., MacManus, J.P. and Mutus, B.: Inhibition of glutathione reductase by oncomodulin. *Arch. Biochem. Biophys.*, 277 (1990) 149-154.
- 2697 Porter, G.F., Savage, P.M. and McGuinness, E.T.: Affinity gel interactions of alcohol and polyol dehydrogenases: anomalous behaviour and structural correlations. *J. Chromatogr.*, 502 (1990) 69-78.
- 2698 Srivatsan, J., Weirich, M. and Agosin, M.: Cytochrome P-450-catalyzed formation of 20-hydroxy-ecdysone in larval housefly mitochondria. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1372-1377.
- 2699 Wrba, A., Jaenicke, R., Huber, R. and Stetter, K.O.: Lactate dehydrogenase from the extreme thermophile *Thermotoga maritima*. *Eur. J. Biochem.*, 188 (1990) 195-201.

See also 2375, 2732.

20b. *Transferases (excluding E.C. 2.7.-.-)*

- 2700 De Carolis, E. and Ibrahim, R.K.: Purification and kinetics of phenylpropanoid O-methyltransferase activities from *Brassica oleracea*. *Biochem. Cell Biol.*, 67 (1989) 763-769.
- 2701 Hoare, K. and Datta, K.: Characteristics of L-alanine-4,5-dioxovaleric acid transaminase: an alternate pathway of heme biosynthesis in yeast. *Arch. Biochem. Biophys.*, 277 (1990) 122-129.
- 2702 Holmquist, L.: High performance ion exchange chromatography of human plasma lecithin:cholesterol acyltransferase. *Biomed. Chromatogr.*, 3 (1989) 276-278.
- 2703 Kurihara, T., Ueda, M. and Tanaka, A.: Peroxisomal actooacetyl-CoA thiolase and 3-ketoacyl-CoA thiolase from an *n*-alkane-utilizing yeast, *Candida tropicalis*: purification and characterization. *J. Biochem. (Tokyo)*, 106 (1989) 474-478.
- 2704 Lash L.H., Elfarra, A.A., Rakiewicz-Nemeth, D. and Anders, M.W.: Bioactivation mechanism of cytotoxic homocysteine S-conjugates. *Arch. Biochem. Biophys.*, 276 (1990) 322-330.

- 2705 Maatsumoto, A., Nakajima, T. and Matsuda, K.: Mode of action of glycogen branching enzyme from *Neurospora crassa*. *J. Biochem. (Tokyo)*, 107 (1990) 118-122.
- 2706 Tan, T.M.C., Sit, K.H. and Wong, K.P.: UDP-Glucuronyltransferase activity toward harmol in human liver and human fetal liver cells in culture. *Anal. Biochem.*, 185 (1990) 44-50.
- 2707 Thomas, K.B., Zawilska, J. and Iuvone, P.M.: Arylalkylamine (serotonin) N-acetyltransferase assay using high-performance liquid chromatography with fluorescence or electrochemical detection of N-acetyltryptamine. *Anal. Biochem.*, 184 (1990) 228-234.
- 2708 Warren, M.J., Roessner, C.A., Santander, P.J. and Scott, A.I.: The *Escherichia coli* *cysG* gene encodes S-adenosylmethionine-dependent uroporphyrinogen III methylase. *Biochem. J.*, 265 (1990) 725-729.
- 2709 Yagi, T., Shounaka, M. and Yamamoto, S.: Distribution of aspartate aminotransferase activity in yeasts, and purification and characterization of mitochondrial and cytosolic isoenzymes from *Rhodotorula marina*. *J. Biochem. (Tokyo)*, 107 (1990) 151-159.

See also 2950.

20c. *Transferases transferring phosphorus-containing groups (E.C. 2.7.-.-)*

- 2710 Alcazar, A., Mendez, E., Martin, M.E. and Salinas, M.: Purification of a novel eIF-2 α protein kinase from calf brain. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1237-1244.
- 2711 Argoudelis, C.J.: Simple isocratic high-performance liquid chromatographic method for measuring pyridoxine kinase activity in crude biological extracts. *J. Chromatogr.*, 526 (1990) 25-33.
- 2712 Huse, K., Himmel, M., Gärtner, G., Kopperschlager, G. and Hofmann, E.: Use of an activated nylon membrane (Immunodyne) as an affinity adsorbent for the purification of phosphofructokinase and phosphoglycerate kinase from yeast. *J. Chromatogr.*, 502 (1990) 171-177.
- 2713 Le Grice, S.F.J. and Grüniger-Leitch, F.: Rapid purification of homodimer and heterodimer HIV-1 reverse transcriptase by metal chelate affinity chromatography. *Eur. J. Biochem.*, 187 (1990) 307-314.
- 2714 Lemaitre, R.N., King, W.C., MacDonald, D. and Glomset, J.A.: Distribution of distinct arachidonoyl-specific and non-specific isoenzymes of diacylglycerol kinase in baboon (*Papio cynocephalus*) tissues. *Biochem. J.*, 266 (1990) 291-299.
- 2715 Noda, M., Kunz, B.C., Tsai, S.-C., Adamik, R., Murtagh, J.J., Chen, H.-C., Halpern, J.L. and Moss, J.: Isolation and immunological properties of adenosine kinase. *Prepar. Biochem.*, 19 (1989) 351-361.
- 2716 Rotenberg, S.A., Krauss, R.S., Borner, C.M.B. and Weinstein, I.B.: Characterization of a specific form of protein kinase C overproduced by a C3H 10T^{1/2} cell line. *Biochem. J.*, 266 (1990) 173-178.
- 2717 Rush, J. and Konigsberg, W.H.: Rapid purification of overexpressed T4 DNA polymerase. *Prepar. Biochem.*, 19 (1989) 329-340.

20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*

- 2718 Anderson, D.J., Branum, E.L. and O'Brien, J.F.: Liver- and bone-derived isoenzymes of alkaline phosphatase in serum as determined by high-performance affinity chromatography. *Clin. Chem. (Winston-Salem)*, 36 (1990) 240-246.
- 2719 Baillyes, E.M., Ferguson, M.A.J., Colaco, C.A.L.S. and Luzio, J.P.: Inositol is a constituent of detergent-solubilized immunoprecipitated rat liver 5'-nucleotidase. *Biochem. J.*, 265 (1990) 907-909.

- 2720 Bruns, K. and Casillas, E.R.: Partial purification and characterization of an acetylcarnitine hydrolase from bovine epididymal spermatozoa. *Arch. Biochem. Biophys.*, 277 (1990) 1-7.
- 2721 Hauser, C.R. and Gray, H.B., Jr.: Precursor-product relationship of larger to smaller molecular forms of the BAL 31 nuclease from *Alteromonas espejiana*: preferential removal of duplex exonuclease relative to endonuclease activity by proteolysis. *Arch. Biochem. Biophys.*, 276 (1990) 451-459.
- 2722 Kamberov, E. and Ivanov, A.: Purification of phospholipase-C from *Bacillus cereus* by affinity chromatography on 2-(4-aminophenylsulphonyl)ethyl-cellulose. *J. Chromatogr.*, 525 (1990) 307-318.
- 2723 Komoda, T., Sato, M., Furiya, K., Sakagishi, Y. and Sekine, T.: Allelic and actopic polymorphisms in human placental alkaline phosphatases: sugar chain heterogeneities. *Clin. Chim. Acta*, 186 (1989) 203-210.
- 2724 Koshida, K. and Wahren, B.E.: Enzymatic heterogeneity of seminomas. *Clin. Chim. Acta*, 186 (1989) 255-264.
- 2725 Meador, III, J., Cannon, B., Cannistrato, V.J. and Kennell, D.: Purification and characterization of *Escherichia coli* RNase I. Comparisons with RNase M. *Eur. J. Biochem.*, 187 (1990) 549-553.
- 2726 Merryweather, A.T., Crampton, J.M. and Townson, H.: Purification and properties of an esterase from organophosphate-resistant strain of the mosquito *Culex quinquefasciatus*. *Biochem. J.*, 266 (1990) 83-90.
- 2727 Nikul'cheva, N.G., Soliternova, I.B. and Eliseeva, L.A.: (Purification and properties of rabbit liver triglyceride lipase). *Biokhimiya (Moscow)*, 55 (1990) 173-180.
- 2728 Stochaj, U., Dieckhoff, J., Mollenhauer, J., Cramer, M. and Mannherz, H.G.: Evidence for the direct interaction of chicken gizzard 5'-nucleotidase with laminin and fibronectin. *Biochim. Biophys. Acta*, 992 (1989) 385-392.
- 2729 Sugihara, A., Shimada, Y. and Tominaga, Y.: Separation and characterization of two molecular forms of *Geotrichum candidum* lipase. *J. Biochem. (Tokyo)*, 107 (1990) 426-430.

For additional information see:

C.A., 112 (1990) 3569j.

20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*

- 2730 Anand, L., Krishnamurthy, S. and Vithayathil, P.J.: Purification and properties of xylanase from the thermophilic fungus, *Humicola lanuginosa* (Griffon and Maublanc) Bunce. *Arch. Biochem. Biophys.*, 276 (1990) 546-553.
- 2731 Aoki, T., Araki, T. and Kitamikado, M.: Purification and characterization of a novel β -agarase from *Vibrio* sp. AP-2. *Eur. J. Biochem.*, 187 (1990) 461-465.
- 2732 Debeire, P., Priem, B., Strecker, G. and Vignon, M.: Purification and properties of an endo-1,4-xylanase excreted by a hydrolytic thermophilic anaerobe, *Clostridium thermolacticum*. A proposal for its action mechanism on larchwood 4-O-methylglucuronoxylan. *Eur. J. Biochem.*, 187 (1990) 573-580.
- 2733 Ito, Y., Yamada, H., Nakamura, S. and Imoto, T.: Purification, amino acid sequence, and some properties of rabbit kidney lysozyme. *J. Biochem. (Tokyo)*, 107 (1990) 236-241.
- 2734 Kikuchi, M., Taniyama, Y., Kanaya, S., Takao, T. and Shimonishi, Y.: Occurrence of S(1,2-dicarboxyethyl)-cysteine at position 77 in mutant human lysozyme secreted by *Saccharomyces cerevisiae*. *Eur. J. Biochem.*, 187 (1990) 315-320.
- 2735 Klein, R.D., Deibel, M.R., Sarcich, J.L., Zurcher-Neely, H.A., Reardon, I.M. and Heinrikson, R.L.: Purification and characterization of invertase from a novel industrial yeast, *Schwanniomyces occidentalis*. *Prepar. Biochem.*, 19 (1989) 293-319.

- 2736 Pisani, F.M., Rella, R., Raia, C.A., Rozzo, C., Nucci, R., Gambacorta, A., De Rosa, M. and Rossi, M.: Thermostable β -galactosidase from the archaeobacterium *Sulfolobus solfataricus*. Purification and properties. *Eur. J. Biochem.*, 187 (1990) 321-328.
- 2737 Szepeszy, L., Vida, L., Toth, M. and Laszlo, E.: Investigation of some α -amylases by high-performance hydrophobic-interaction chromatography. *J. Chromatogr.*, 499 (1990) 197-204.
- 2738 Tulsiani, D.R.P., Coleman, V.D. and Touster, O.: Asparagine-linked glycoprotein biosynthesis in rat brain: identification of glucosidase I, glucosidase II, and an endomannosidase (glucosyl mannosidase). *Arch. Biochem. Biophys.*, 277 (1990) 114-121.

For additional information see:

- C.A., 111 (1989) 227992j;
112 (1990) 3181b.

20f. Other hydrolases

- 2739 Abdus Sattar, A.K.M., Yamamoto, N., Yoshimoto, T. and Tsuru, D.: Purification and characterization of an extracellular prolyl endopeptidase from *Agaricus bisporus*. *J. Biochem. (Tokyo)*, 107 (1990) 256-261.
- 2740 Chantry, A., Gregson, N.A. and Glynn, P.: A novel metalloproteinase associated with brain myelin membranes. Isolation and characterization. *J. Biol. Chem.*, 264 (1989) 21603-21607.
- 2741 Chiczy, R.M. and Regnier, F.E.: Microenvironmental contributions to the chromatographic behavior of subtilisin in hydrophobic-interaction and reversed-phase chromatography. *J. Chromatogr.*, 500 (1990) 503-518.
- 2742 Crawford, C., Brown, N.R. and Willis, A.C.: Investigation of the structural basis of the interaction of calpain II with phospholipid and with carbohydrate. *Biochem. J.*, 265 (1990) 575-579.
- 2743 Dunaevsky, Ya.E. and Belozersky, M.A.: (The enzymes of the terminal step of proteolysis of the main storage protein from buckwheat seeds). *Biokhimiya (Moscow)*, 55 (1990) 3-11.
- 2744 Fortier, G. and Gagnon, J.: Kinetic study of nucleophile specificity in dipeptide synthesis catalyzed by clostridiopeptidase B. *Arch. Biochem. Biophys.*, 276 (1990) 317-321.
- 2745 Hou, K.C. and Zaniewski, R.: Purification of urokinase by combined cation exchanger and affinity chromatographic cartridges. *J. Chromatogr.*, 525 (1990) 297-306.
- 2746 Kinoshita, M., Hamakubo, T., Fukui, I., Murachi, T. and Toyohara, H.: Significant amount of multicatalytic proteinase identified on membrane from human erythrocyte. *J. Biochem. (Tokyo)*, 107 (1990) 440-444.
- 2747 Mantle, D., Lauffart, B., McDermott, J. and Gibson, A.: Characterization of aminopeptidases in human kidney soluble fraction. *Clin. Chim. Acta*, 187 (1990) 105-114.
- 2748 Mason, R.W.: Characterization of the active site of human multicatalytic proteinase. *Biochem. J.*, 265 (1990) 479-484.
- 2749 Miller, J.J. and Detwiler, T.C.: Characteristics of a thrombin inhibitor secreted by activated platelets. *Arch. Biochem. Biophys.*, 276 (1990) 364-368.
- 2750 Miller, R.P. and Farley, R.A.: β Subunit of $(Na^+ + K^+)$ -ATPase contains three disulfide bonds. *Biochemistry*, 29 (1990) 1524-1532.
- 2751 Ohashi, T., Ohno, T., Arata, J., Sugahara, K. and Kodama, H.: Characterization of prolidase I and II from erythrocytes of a control, a patient with prolidase deficiency and her mother. *Clin. Chim. Acta*, 187 (1990) 1-10.
- 2752 Oizumi, J. and Hayakawa, K.: Purification of the membrane protein enzyme lipamidase by affinity chromatography. *Biomed. Chromatogr.*, 3 (1989) 274-275.

- 2753 Scheel, E. and Schäfer, G.: Chemiosmotic energy conversion and the membrane ATPase of *Methanobolus tinadirus*. *Eur. J. Biochem.*, 187 (1990) 727-735.
- 2754 Suzuki, K., Kusumoto, H., Nishioka, J. and Komiyama, Y.: Bovine plasma protein C inhibitor with structural and functional homologous properties to human plasma protein C inhibitor. *J. Biochem. (Tokyo)*, 107 (1990) 381-388.
- 2755 Swadesh, J.K., Huang, I.-Y. and Budzynski, A.Z.: Purification and characterization of hementin, a fibrinogenolytic protease from the leech *Haementeria ghilianii*. *J. Chromatogr.*, 502 (1990) 359-369.
- 2756 Tavera, C., Guillemot, J.-C., Capdevielle, J., Ferrara, P., Leung-Tack, J. and Colle, A.: A rapid two-step purification of rat cystatin C, one major inhibitor of cysteine proteinases. *Prepar. Biochem.*, 19 (1989) 279-291.
- 2757 Yasuhara, T., Yokosawa, H. and Ishii, S.-i.: Purification and characterization of an aminopeptidase from sperm of the sea urchin, *Strongylocentrotus intermedius*. Ca^{2+} -dependent substrate specificity as a novel feature of the enzyme. *J. Biochem. (Tokyo)*, 107 (1990) 273-279.

For additional information see:

- C.A., 111 (1989) 227638y;
112 (1990) 25602s.

20g. Lyases

- 2758 Jawali, N.: The dimeric form of phosphoenolpyruvate carboxylase isolated from maize: physical and kinetic properties. *Arch. Biochem. Biophys.*, 277 (1990) 61-68.
- 2759 Lash, L.H., Nelson, R.M., van Dyke, R.A. and Anders, M.W.: Purification and characterization of human kidney cytosolic cysteine conjugate β -lyase activity. *Drug Metab. Disp.*, 18 (1990) 50-54.
- 2760 White, P.J., Young, J., Hunter, I.S., Nimmo, H.G. and Coggins, J.R.: The purification and characterization of 3-dehydroquinase from *Streptomyces coelicolor*. *Biochem. J.*, 265 (1990) 735-738.

20h. Isomerases

- 2761 Skarpeid, H.-J., Zimmer, T.-L., Shen, B. and Döhren, H.: The proline-activating activity of the multienzyme gramicidin S synthetase 2 can be recovered as a 115-kDa tryptic fragment. *Eur. J. Biochem.*, 187 (1990) 627-633.
- 2762 Vangrysperre, W., van Damme, J., Vandekerckhove, J., de Bruyne, C.K., Cornelis, R. and Kersters-Hilderson, H.: Localization of the essential histidine and carboxylate group in D-xylose isomerases. *Biochem. J.*, 265 (1990) 699-705.

See also 2804.

20i. Ligases

- 2763 Newton, R.P., Salvage, B.J. and Hakeem, N.A.: Cytidylate cyclase: development of assay and determination of kinetic properties of cytidine 3',5'-cyclic monophosphate-synthesizing enzyme. *Biochem. J.*, 265 (1990) 581-586.
- 2764 Wolfson, A.D., Motorin, Yu.A., Ribkinska, T.I. and Beresten, S.F.: Purification of mammalian tyrosyl-tRNA synthetase by high-performance liquid chromatography. *J. Chromatogr.*, 503 (1990) 277-281.

20j. Complex mixtures and incompletely identified enzymes

- 2765 Leontyevsky, L.A. and Golovleva, L.A.: (Extracellular lignin-decomposing enzymes of the fungus *Panus tigrinus*). *Biokhimiya (Moscow)*, 55 (1990) 423-431.

For additional information see:

C.A., 111 (1989) 169817t.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 2766 Alpert, A.J.: Hydrophilic-interaction chromatography for the separation of peptides, nucleic acids and other polar compounds. *J. Chromatogr.*, 499 (1990) 177-196.
- 2767 Assmann, B. and Haas, H.J.: Determination of creatinine-related urinary uracil excretion in children by high-performance liquid chromatography. *J. Chromatogr.*, 525 (1990) 277-285.
- 2768 Bai, Q.Y., Zelles, L., Scheunert, I. and Korte, F.: Determination of adenine nucleotides in soil by ion-paired reverse-phase high-performance liquid chromatography. *J. Microbiol. Methods*, 9 (1989) 345-351; C.A., 111 (1989) 172962s.
- 2769 Carpen, M.E., Poplack, D.G., Pizzo, P.A. and Balis, F.M.: High-performance liquid chromatographic method for analysis of 2',3'-dideoxyinosine in human body fluids. *J. Chromatogr.*, 526 (1990) 69-75.
- 2770 Cundy, K.C., Kohen, R. and Ames, B.N.: Determination of 8-hydroxydeoxyguanosine in human urine: a possible assay for *in vivo* oxidative DNA damage. *Basic Life Sci.*, 49 (Oxygen Radicals Biol. Med.) (1988) 479-482; C.A., 112 (1990) 3763t.
- 2771 Inaki, Y., Nagae, S., Miyamoto, T., Sugiura, Y. and Takemoto, K.: Nucleic acid analogs for high-performance liquid chromatography. *Polym. Sci. Technol. (Plenum)*, 38 (Appl. Bioacat. Polym. Mater.) (1988) 185-204; C.A., 111 (1989) 233438z.
- 2772 Kress, G. and Pentz, R.: A sensitive method for the determination of urinary mercapturic acids for use in biological monitoring. *Arch. Toxicol. Suppl.*, 13 (1989) 104-107; C.A., 111 (1989) 210160p.
- 2773 Mayernik, J.A., Conner, M.K. and Giam, C.S.: Methods for the isolation and detection of etheno adducts in nucleotide pools *in vivo* following exposure to ethyl carbamate. *J. Chromatogr.*, 526 (1990) 407-422.
- 2774 Mills, G.C.: Isolation and identification of S-adenosylmethionine from human urine. *Biochem. Med. Metab. Biol.*, 41 (1989) 217-223; C.A., 111 (1989) 170322j.
- 2775 Montaldo, P.G. and Cornaglia-Ferraris, P.: High-performance liquid chromatographic determination in human plasma of a new immunomodulatory agent with a peptidylhypoxanthine structure (RM 06). *J. Chromatogr.*, 525 (1990) 471-477.
- 2776 Niculescu-Duvaz, I. and Voiculescu, N.: Simultaneous analysis of bases, nucleosides, and nucleotides by HPLC. *Chromatogram*, 10 (1989) 7-9; C.A., 111 (1989) 228255h.
- 2777 Smith, R.E. and MacQuarrie, R.A.: Ion chromatographic determination of zwitterions in rat brain. *LC-GC*, 7 (1989) 775-777; C.A., 111 (1989) 211275s.
- 2778 Turcotte, J.G., Pivarnik, P.E., Shirali, S.S., Singh, H.K., Seghal, R.K., Macbride, D., Jang, N.-I. and Brown, P.R.: Preparative-scale high-performance liquid chromatographic separation and purification of 3'-azido-3'-deoxythymidine-5'-phosphate. *J. Chromatogr.*, 499 (1990) 55-61.
- 2779 Werner, G., Schneider, V. and Emmert, J.: Simultaneous determination of creatine, uric acid and creatinine by high-performance liquid chromatography with direct serum injection and multi-wavelength detection. *J. Chromatogr.*, 525 (1990) 265-275.

For additional information see:

C.A., 111 (1989) 173916s, 190883w, 190995j, 228478h, 230778z;
112 (1990) 5990p.

See also 2157, 2185, 2527.

21b. Nucleic acids, RNA

2780 Lindeberg, J., Srichaiyo, T. and Hjerten, S.: High-performance adsorption chromatography of transfer ribonucleic acids and proteins on 2- μ m spherical beads of hydroxyapatite. Influence of sodium chloride and magnesium ions on the resolution. *J. Chromatogr.*, 499 (1990) 153-163.

21c. Nucleic acids, DNA

2781 White, B.A. and Rosenzweig, S.: A reliable method for the purification of bacteriophage DNA. *BioTechniques*, 7 (1989) 694-696; C.A., 111 (1989) 211460y.

For additional information see:

C.A., 111 (1989) 170604c, 228559k, 228562f;
112 (1990) 4053y.

21d. Structural studies on RNA and RNA mapping

2782 Tanaka, K., Yazawa, K. and Nakajima, T.: High resolution chromatography of ribonucleosides and its application to RNA analysis. *Biomed. Chromatogr.*, 3 (1989) 246-250.

21e. Structural studies on DNA and DNA mapping

For additional information see:

C.A., 112 (1990) 3595q.

21f. Complex mixtures of nucleic acids and their fragments

See 2775.

22. ALKALOIDS

2783 Armstrong, D.W., Bertrand, G.L., Ward, K.D., Ward, T.J., Secor, H.V. and Seeman, J.I.: Evaluation of the effect of organic modifier and pH on retention and selectivity in reversed-phase liquid chromatographic separation of alkaloids on a cyclodextrin bonded phase. *Anal. Chem.*, 62 (1990) 332-338.

2784 Bouis, P., Taccard, G. and Boelsterli, U.A.: Determination of cocaine and norcocaine in plasma and cell cultures using high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 447-459.

2785 Cohen, M.L., Parli, C.J. and Fuller, R.W.: 5-Hydroxytryptamine₂ receptor antagonist activity of the acid metabolite (1-isopropyl dihydrolysergic acid) of the ergoline ester, sergolexole (LY281067). *J. Pharmacol. Exp. Ther.*, 251 (1989) 1006-1011.

2786 Knapp, S., Wardlow, M.L. and Thal, L.J.: Sensitive analysis of plasma physostigmine levels using dual-cell electrochemistry in the redox mode. *J. Chromatogr.*, 526 (1990) 97-107.

- 2787 Lau, S.W.J., Chow, D. and Feldman, S.: Simultaneous determination of physostigmine and tetrahydroaminoacridine in a transdermal permeation study by high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 87-95.
- 2788 Markin, R.S., Wadman, M.C., Bottjen, P.L., Haven, M.C. and Huth, J.A.: Short-column liquid chromatographic assay for caffeine and chloramphenicol in serum. *J. Chromatogr.*, 525 (1990) 464-470.
- 2789 O'Doherty, S., Cooke, M. and Roberts, D.J.: Enhancing the LC analysis of nicotine and its metabolites in urine using Meldrums acid as a complexing agent. *J. High Resolut. Chromatogr.*, 13 (1990) 74-77.
- 2790 Papp, E.: Fast separation of dihydro ergot alkaloids on an octylsilica column. *J. Chromatogr.*, 502 (1990) 241-242.
- 2791 Parikh, K.M., Doshi, V.J., Salunkhe, J.B. and Kanath, R.P.: High performance liquid chromatographic determination of vasicine and vasicinone from pharmaceutical formulations. *Indian Drugs*, 27 (1989) 64-66; *C.A.*, 112 (1990) 11998p.
- 2792 Parviainen, M.T., Puhakainen, E.V.J., Laatikainen, R., Savolainen, K., Heranen, J. and Barlow, R.D.: Nicotine metabolites in the urine of smokers. *J. Chromatogr.*, 525 (1990) 193-202.
- 2793 Sandberg, J.A. and Olsen, G.D.: Microassay for the simultaneous determination of cocaine, norcocaine, benzoylecgonine and benzoynorecgonine by high-performance liquid chromatography. *J. Chromatogr.*, 525 (1990) 113-121.
- 2794 Venn, R.F. and Michalkiewicz, A.: Fast reliable assay for morphine and its metabolites using high-performance liquid chromatography and native fluorescence detection. *J. Chromatogr.*, 525 (1990) 379-388.
- 2795 Venn, R.F., Hucks, D., McCarthy, J. and Wells, J.C.D.: A fast reliable assay for morphine and its metabolites. *Adv. Biosci. (Oxford)*, 75 (1989) 515-518; *C.A.*, 111 (1989) 166697t.
- 2796 Walker, J.S.: Phenylpropanolamine potentiates caffeine neurotoxicity in rats. *J. Pharm. Sci.*, 78 (1989) 986-989.
- 2797 Xie, F., Wang, H., Shu, H., Li, J., Jiang, J., Chang, J. and Hsieh, Y.: Separation and characterization of the metabolic products of lappaconitine in rat urine by high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 109-118.

For additional information see:

C.A., 112 (1990) 2125z, 3779c, 11973b, 42691n.

See also 3018.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 2798 Gratzfeld-Huesgen, A.: Simultaneous determination of porphyrins, aromatics, and olefins in crude oil by gel permeation chromatography and multiple signal detection. *LC-GC*, 7 (1989) 836-845; *C.A.*, 112 (1990) 59310k.
- 2799 Ho, J.W.: Effect of mobile phase pH, ionic strength, and buffer composition on the separation of porphyrins by high-performance liquid chromatography. *LC-GC*, 7 (1989) 348-352; *C.A.*, 112 (1990) 48016t.
- 2800 Nokel, A.Yu., Nizhnik, A.N. and Mironov, A.F.: (Study of the composition of hematorporphyrin derivatives). *Zh. Anal. Khim.*, 44 (1989) 1514-1517; *C.A.*, 111 (1989) 219359t.

For additional information see:

C.A., 111 (1989) 228287v.

See also 2708, 2791, 2910.

23b. *Bile pigments*

- 2801 Ma, J.: (Determination of bilirubin and its derivatives by high-pressure liquid chromatography). *Shenyang Yaoxueyuan Xuebao*, 6 (1989) 212-216; C.A., 111 (1989) 219346m - a review with 20 refs.
- 2802 Skar, V. and Sazerholt, H.: High-performance liquid chromatography of bilirubin conjugates in bile: effect of β -glucuronidase on the bile pigments. *Scand. J. Gastroenterol.*, 24 (1989) 657-665; C.A., 111 (1989) 211272p.

23c. *Indole derivatives*

- 2803 Auerbach, S.B., Minzenberg, M.J. and Wilkinson, L.O.: Extracellular serotonin and 5-hydroxyindoleacetic acid in the hypothalamus of the unanesthetized rat measured by *in vivo* dialysis coupled to high-performance liquid chromatography with electrochemical detection: dialysate serotonin reflects neuronal release. *Brain Res.*, 499 (1989) 281-290; C.A., 112 (1990) 1147w.
- 2804 Pawelek, J.M.: Dopachrome conversion factor functions as an isomerase. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1328-1333.

See also 2465, 2503, 2706, 2707.

23d. *Pyridine derivatives*

See 2150.

23e. *Other N-heterocyclic compounds*

- 2805 Hirata, Y., Minami, M., Naoi, M. and Nagatsu, T.: Studies on the uptake of N-methylisoquinolinium ion into rat striatal slices using high-performance liquid chromatography with fluorometric detection. *J. Chromatogr.*, 503 (1990) 189-195.
- 2806 Lafleur, A.L. and Nakagawa, Y.: Multimode size exclusion chromatography with poly(divinyl/benzene)/N-methylpyrrolidone for the characterization of coal-derived mixtures. *Fuel*, 68 (1989) 741-752; C.A., 112 (1990) 23333f.
- 2807 Mohseni, R.M. and Hurtubise, R.J.: Retention characteristics of several compound classes in reversed-phase liquid chromatograph with β -cyclodextrin as a mobile phase modifier. *J. Chromatogr.*, 499 (1990) 395-410.
- 2808 Rotschova, J. and Pospisil, J.: (Stabilizers derived from 2,2,6,6-tetramethylpiperidine and methods for their analytical determination in polymers). *Plaste Kautsch.*, 36 (1989) 289-293; C.A., 112 (1990) 57296m - a review with 49 refs.
- 2809 Urinov, E., Boimirzaev, A.S. and Rashidova, S.S.: (Chromatographic determination of N-vinylpyrrolidone content in a (poly(vinyl)pyrrolidone)-vinylpyrrolidone mixture). *Uzb. Khim. Zh.*, (1989) 29-31; C.A., 111 (1989) 215219z.

24. ORGANIC SULPHUR COMPOUNDS

- 2810 Aced, G., Möckel, H.J. and Nelsen, S.F.: Ion-pair chromatographic separation of onium salts on an ODS column. *J. Liq. Chromatogr.*, 12 (1989) 3201-3218.
- 2811 Khmel'nitskii, R.A. and Bondarenko, M.A.: (Instrumental methods for determination of lower mercaptans). *Zh. Anal. Khim.*, 44 (1989) 965-985; C.A., 111 (1989) 247199j - a review with 139 refs.

- 2812 Kokkonen, P., van der Greef, J., Niessen, W.M.A., Tjaden, U.R., Ten Hove, G.J. and van der Werken, G.: Separation and identification of intact gluconisolates using direct coupling of high-performance liquid chromatography with frit to fast-atom bombardment mass spectrometry. *Rapid Commun. Mass Spectrom.*, 3 (1989) 102-106; *C.A.*, 111 (1989) 170522z.
- 2813 Krause, R.T.: Liquid chromatographic-electrochemical determination of ethylenethiourea in foods by revised official method. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 975-979.
- 2814 Mancas, D., Leanca, M. and Chiriac, C.: Determination of 2-thiothiazolidine-4-carboxylic acid (TTCA) in urine by CGC. *J. High Resolut. Chromatogr.*, 12 (1989) 828-829.
- 2815 Semikin, V.V., Chikishev, Yu.G., Zeleneva, T.P. and Senyavin, M.M.: Chromatographic behavior of some derivatives of dithiocarbamic acids in reversed-phase HPLC. *Zh. Anal. Khim.*, 44 (1989) 120-126; *C.A.*, 112 (1990) 30055v.
- 2816 Tanaka, M., Shimizu, M. and Tsujimoto, Y.: High-performance liquid chromatographic determination of urinary mercapturic acids in rat. *J. Chromatogr.*, 526 (1990) 289-292.

For additional information see:

C.A., 111 (1989) 216867w;
112 (1990) 25311w.

See also 2113, 2504, 2513, 2515, 3121.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCLUDING SUGAR PHOSPHATES)

- 2817 Haky, J.E., Baird, D.M. and Falzone, S.: Reversed phase high performance liquid chromatographic analysis of triphenylphosphine in a reaction mixture. *Anal. Lett.*, 22 (1989) 2367-2651.
- 2818 Irth, H., Lamoree, M., de Jong, G.J., Brinkman, U.A.T., Frei, R.W., Kornfeldt, R.A. and Persson, L.: Determination of D-myo-1,2,6-inositol triphosphate by ion-pair reversed-phase liquid chromatography with post-column ligand exchange and fluorescence detection. *J. Chromatogr.*, 499 (1990) 617-625.
- 2819 Kabakoff, B.D., Doyle, J.W. and Kandutsch, A.A.: Relationships among dolichyl phosphate, glycoprotein synthesis, and cell culture growth. *Arch. Biochem. Biophys.*, 276 (1990) 382-389.
- 2820 Pullarkat, R.K., Pullarkat, P.S. and Morris, G.N.: Phosphorylated dolichols in aging. *Biochem. J.*, 265 (1990) 891-894.
- 2821 Sachinidis, A., Locher, R., Mengden, T. and Vetter, W.: Low-density lipoprotein elevates intracellular calcium and pH in vascular smooth muscle cells and fibroblasts without mediation of LDL receptor. *Biochem. Biophys. Res. Commun.*, 167 (1990) 353-359.
- 2822 Stephens, L.R. and Downes, C.P.: Produce-precursor relationship amongst inositol polyphosphates. Incorporation of [32 P]Pi into myo-inositol 1,3,4,6-tetrakisphosphate, myo-inositol 1,3,4,5-tetrakisphosphate, myo-inositol 3,4,5,6-tetrakisphosphate and myo-inositol 1,3,4,5,6-pentakisphosphate in intact avian erythrocytes. *Biochem. J.*, 265 (1990) 435-452.
- 2823 Verbesselt, R., Tjandra-Maga, T.B., de Schepper, J.P., Itakura, K. and Sugiro, T.: Determination of a platelet activating factor antagonist (CV-3988): methodology and clinical application. *J. Chromatogr.*, 526 (1990) 515-524.

For additional information see:

C.A., 112 (1990) 6191r, 15802t.

See also 2113, 2185, 2401, 2411.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 2824 Gehrig, C.C. and Fitzpatrick, T.M.: Determination of organic arsenicals in pesticide formulations by ion chromatography. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 994-996.
- 2825 Holak, W.: Liquid chromatographic-atomic absorption spectrophotometric method for determination of methyl mercury in seafood: collaborative study. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 926-930.
- 2826 Olson, G.J., Brinckman, F.E. and Blair, W.R.: Methods for the analysis for organometallic compounds in wastes. *ASTM Spec. Tech. Publ.*, 999 (1988) 130-145; *C.A.*, 111 (1989) 224533z - a review with 37 refs.
- 2827 Timerbaev, A.R., Vanifatova, N.G., Evstigerov, M.V., Salov, V.V. and Petrukhin, O.M.: (Liquid adsorption chromatography of metal chelates. Characteristics of the chromatographic behavior of nitroxyl-containing reagents). *Zh. Anal. Khim.*, 44 (1989) 834-841; *C.A.*, 112 (1990) 68801y.
- 2828 Whang, C.W. and Tsai, W.L.: Analysis of triorganotin compounds by high-performance liquid chromatography with reverse-pulse amperometric detection. *J. Chin. Chem. Soc. (Taipei)*, 36 (1989) 179-186; *C.A.*, 112 (1990) 69183k.

See also 2113, 2916, 3107.

26b. Boranes, silanes and related non-metallic compounds

- 2829 Grossed, C., Cantin, D. and Alary, J.: (Quantitative HPLC determination of antioxidants in pharmaceutical formulations with ultraviolet and electrochemical detections). *Analisis*, 17 (1989) 409-412; *C.A.*, 111 (1989) 239643j.

26c. Coordination compounds

- 2830 Blais, J.S. and Marshall, W.D.: Determination of ionic alkyllead compounds in water, soil and sediment by high-performance liquid chromatography-quartz tube atomic absorption spectrometry. *J. Anal. At. Spectrom.*, 4 (1989) 271-277; *C.A.*, 111 (1989) 224608c.
- 2831 Main, M.V. and Fritz, J.S.: Chromatographic determination of metal chelates of 2,6-diacetylpyridine bis(N-methylenepyridiniohydrazone). *Anal. Chim. Acta*, 229 (1990) 101-106.
- 2832 Maslowska, J. and Starzynski, S.: Studies on the separation of Al^{III} , Co^{III} , Cr^{III} complexes with acetylacetone by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 502 (1990) 221-225.
- 2833 Timerbaev, A.R. and Petrukhin, O.: (Liquid adsorption chromatography metal chelates. Stability and chromatographic properties of the chelates). *Zh. Anal. Khim.*, 44 (1989) 1424-1432; *C.A.*, 111 (1989) 240862m.

See also 2110, 2221, 3059, 3102, 3126.

27. VITAMINS AND VARIOUS GROWTH REGULATORS (NON-PEPTIDIC)

- 2834 Agarwal, V.K.: Liquid chromatographic determination of vitamin D in infant formula. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 1007-1009.

- 2835 Balschukat, D. and Kress, E.: Use of column switching for the determination of niacinamide in compound feed. *J. Chromatogr.*, 502 (1990) 79-85.
- 2836 Bhat, P.V. and Sundaresa, P.R.: High-performance liquid chromatography of vitamin A compounds. *Crit. Rev. Anal. Chem.*, 20 (1988) 197; *C.A.*, 111 (1989) 190510x - a review with 104 refs.
- 2837 Cody, M.K., Clark, G.B., Conway, B.O.B. and Crosby, N.T.: Identification of medicinal additives in animal feedingstuffs by high-performance liquid chromatography. *Analyst (London)*, 115 (1990) 1-8.
- 2838 Fernando, S.M. and Murphy, P.A.: HPLC determination of thiamin and riboflavin in soybeans and tofu. *J. Agric. Food Chem.*, 38 (1990) 163-167.
- 2839 Filimonov, V.N., Kolosova, I.F. and Balyatinskaya, L.N.: (Determination of fat-soluble vitamins in feed premixes by HPLC.) *Zh. Anal. Khim.*, 44 (1989) 1433-1436; *C.A.*, 111 (1989) 172574s.
- 2840 Hart, J.P. and Jordan, P.H.: Low-pressure liquid chromatography with electrochemical detection for the determination of vitamin A in multi-vitamin preparation. *Analyst (London)*, 114 (1989) 1633-1635.
- 2841 Hasselmann, C., Franck, D., Grimm, P., Diop, P.A. and Soules, C.: High-performance liquid chromatographic analysis of thiamin and riboflavin in dietetic foods. *J. Micronutr. Anal.*, 5 (1989) 269-279; *C.A.*, 111 (1989) 230782w.
- 2842 Horst, R.L., Koszewski, N.J. and Reinhardt, T.A.: 1α -Hydroxylation of 24-hydroxyvitamin D₂ represents a minor physiological pathway for the activation of vitamin D₂ in mammals. *Biochemistry*, 29 (1990) 578-582.
- 2843 Hou, W., Ji, H. and Wang, E.: Liquid chromatography of vitamin B⁶ with electrochemical detection using a carbon fibre electrode. *Anal. Chim. Acta*, 230 (1990) 207-211.
- 2844 Huang, X. and Li, Z.: (Determination of vitamin E in hen embryonic egg by HPLC.) *Sepu*, 7 (1989) 240-241; *C.A.*, 111 (1989) 193151s.
- 2845 Jansen, C.C. and de Kleijn, J.P.: The assay of cyanocobalamin in pharmaceutical preparations by solid-phase extraction and HPLC. *J. Chromatogr. Sci.*, 28 (1990) 42-45.
- 2846 Kaiser, S., Di Mascio, P., Murphy, M.E. and Sies, H.: Physical and chemical scavenging of singlet molecular oxygen by tocopherols. *Arch. Biochem. Biophys.*, 277 (1990) 101-108.
- 2847 Kennedy, J.F., Rivera, Z.S., Lloyd, L.L., Warner, F.P. and White, C.A.: The use of reversed-phase HPLC for the determination of purity and stability on storage of L-ascorbic acid samples. *J. Micronutr. Anal.*, 5 (1989) 281-289; *C.A.*, 111 (1989) 230783x.
- 2848 Kikuchi, M., Hanamizu, H., Narisawa, K. and Taka, K.: Assay of methylmalonyl CoA mutase with high-performance liquid chromatography. *Clin. Chim. Acta*, 184 (1989) 307-314.
- 2849 Klein, R., Thiery, R. and Tatischeff, I.: Dictyopterine, 6-(D-threo-1,2-dihydroxypropyl)-pterin, a new natural isomer of L-biopterine. Isolation from vegetative cells of *Dicystostelium discoideum* and identification. *Eur. J. Biochem.*, 187 (1990) 665-669.
- 2850 Kohashi, M., Takahashi, A. and Iwai, K.: Separation and determination of tetrahydrofolylpolyglutamates and their 5-methyl derivatives by reversed-phase high-performance liquid chromatography with electrochemical detection. *Pteridines*, 1 (1989) 99-102; *C.A.*, 112 (1990) 3766w.
- 2851 Matthey, M., Graf, H., Richter, G.H. and Flachowsky, G.: (Methodical recommendations for determination of plasma 25-OH D₃ vitamin by HPLC.) *Nahrung*, 33 (1989) 503-507; *C.A.*, 111 (1989) 190671a.
- 2852 Newcomb, S.A., Culling-Berglund, A.J. and Davis, T.P.: Endogenous levels of β -carotene in human buccal mucosa cells by reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 47-58.
- 2853 Ni, P., Caude, M. and Rosset, R.: (Automatic determination of folic acid in fortified foods by high-performance liquid chromatography.) *Analisis*, 17 (1989) 315-318; *C.A.*, 111 (1989) 193149x.

- 2854 Raju, N.R., Srilakshmi, P., Vinolia, G.A., Bhounsule, G.J. and Gaitonde, V.D.: Simultaneous determination of water soluble vitamins in pharmaceutical oral liquids by gradient high performance liquid chromatography-sample preparation with reserved-phase SepPak cartridge. *Indian Drugs*, 26 (1989) 697-701; C.A., 112 (1990) 62757s.
- 2855 Reddy G.S. and Tserng, K.: 24,25,28-Trihydroxyvitamin D₂ and 24,25,26-trihydroxyvitamin D₂: novel metabolites of vitamin D₂. *Biochemistry*, 29 (1990) 943-949.
- 2856 Reyes, E.S.P. and Subryan, L.: An improved method of simultaneous HPLC assay of riboflavin and thiamin in selected cereal products. *J. Food Compos. Anal.*, 2 (1989) 41-47; C.A., 111 (1989) 213435t.
- 2857 Sapers, G.M., Douglas, F.W., Jr., Ziolkowski, M.A., Miller, R.I. and Hicks, K.B.: Determination of ascorbic acid, dehydroascorbic acid and ascorbic acid-2-phosphate in infiltrated apple and potato tissue by high-performance liquid chromatography. *J. Chromatogr.*, 503 (1990) 431-436.
- 2858 Tonolo, G. and Marzo, S.: (Determination of natural tocopherols and added vitamin E in dietetic seed oils by HPLC). *Riv. Ital. Sostanze Grasse*, 66 (1989) 3-6; C.A., 112 (1990) 6204x.
- 2859 Weisz, O.A. and Schnaar, R.L.: Hepatocytes mediate coenzyme A transfer to specific carbohydrate-derivatized surfaces. *Biochem. Biophys. Res. Commun.*, 167 (1990) 67-73.
- 2860 Werner, E.R., Fuchs, D., Hausen, A., Reibnegger, G. and Wachter, H.: Extraction of neopterin from serum with bonded silica sorbents. *Biochem. Clin. Aspects Pteridines*, 5 (1989) 71-80; C.A., 111 (1989) 170515z.

For additional information see:

C.A., 111 (1989) 172577v, 190621j, 190659c, 214770k, 230765t, 230797e, 230801b;
112 (1990) 3771u, 4054z, 6210w, 25754t, 69162c.

See also 2330, 2385, 2905, 2908, 2912.

28. ANTIBIOTICS

- 2861 Agarwal, V.G.: High performance liquid chromatographic determination of gentamicin in milk. *J. Lig. Chromatogr.*, 12 (1989) 3265-3277.
- 2862 Cachet, T., van den Mooter, G., Hauchecorne, R., Vinckier, C. and Hoogmartens, J.: Decomposition kinetics of erythromycin A in acidic aqueous solutions. *Int. J. Pharm.*, 55 (1989) 59-65; C.A., 111 (1989) 219222t.
- 2863 Dine, T., Brunet, C., Luyckx, M., Cazin, M., Gosselin, P. and Cazin, J.L.: Rapid quantitative determination of epirubicin and its metabolites in plasma using high performance liquid chromatography and fluorescence detection. *Biomed. Chromatogr.*, 4 (1990) 20-23.
- 2864 Gause, G.F., Brazhnikova, M.G., Lomakina, N.N., Berdnikova, T.F., Fedorova, G.B., Tokareva, N.L., Borisova, V.N. and Batta, G.: Eremomycin - new glycopeptide antibiotic: chemical properties and structure. *J. Antibiot.*, 42 (1989) 1790-1799.
- 2865 Greiner, M. and Winkelmann, G.: Rapid identification and determination of herbicolin A and B by high-performance liquid chromatography. *J. Chromatogr.*, 502 (1990) 437-442.
- 2866 Griggs, D.J. and Wise, R.: A simple isocratic high-pressure liquid chromatographic assay. *J. Antimicrob. Chemother.*, 24 (1989) 437-445; C.A., 112 (1990) 100p.

- 2867 Hornedo-Nunez, A., Getek, T.A., Korfmacher, W.A. and Simenthal, F.: High-performance liquid chromatography of clindamycin and clindamycin phosphate with electrochemical detection. *J. Chromatogr.*, 503 (1990) 217-225.
- 2868 Kees, F., Raasch, W., Steger, M. and Grobecker, H.: High-performance liquid chromatographic assay for cefotiam and d3-cefotiam in human serum. *J. Chromatogr.*, 525 (1990) 484-489.
- 2869 Konishi, M., Nihio, M., Saitoh, K., Miyaki, T., Oki, T. and Kawaguchi, H.: Cispentacin, a new antifungal antibiotic. I. Production, isolation, physicochemical properties and structure. *J. Antibiot.*, 42 (1989) 1749-1755.
- 2870 Laakso, S., Scheinin, M. and Anttila, M.: Determination of erythromycin base and 2'-acetylerythromycin in human plasma using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr.*, 526 (1990) 475-486.
- 2871 Lee, M.D., Manning, J.K., Williams, D.R., Kuck, N.A., Testa, R.T. and Borders, D.B.: Calicheamicins, a novel family of antitumor antibiotics. 3. Isolation, purification and characterization of calicheamicins β_1^{Br} , γ_1^{Br} , α_2^I , α_3^I , β_1^I , γ_1^I and δ_1^I . *J. Antibiot.*, 42 (1989) 1070-1087.
- 2872 Maloney, J.A. and Awni, W.M.: High-performance liquid chromatographic determination of isepamicin in plasma, urine and dialysate. *J. Chromatogr.*, 526 (1990) 487-496.
- 2873 Mopper, B.: Liquid chromatographic determination of penicillin V potassium in tablets: collaborative study. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 883-884.
- 2874 Morgan, D., Cugier, P., Marello, B., Sarocka, C., Stroz, D. and Plasz, A.: Impurity profiling of clarithromycin using high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr.*, 502 (1990) 351-358.
- 2875 Nakajima, M., Okazaki, T., Iwado, S., Kinoshita, T. and Haneishi, T.: New diterpenoid antibiotics, spirocardins A and B. *J. Antibiot.*, 42 (1989) 1741-1748.
- 2876 Rice, P.D., Shao, Y.Y. and Bobbit, D.R.: Improvements in the determination of penicillin analogs by HPLC separation and laser-based polarimetric detection. *Talanta*, 36 (1989) 985-988; *C.A.*, 112 (1990) 42719c.
- 2877 Shoji, J., Sakazaki, R., Hattori, T., Matsumoto, K., Uotani, N. and Yoshida, T.: Isolation and characterization of agglomerins A, B, C and D. *J. Antibiot.*, 42 (1989) 1729-1733.
- 2878 Skinner, M.H., Hsieh, M., Torseth, J., Pauloin, D., Bhatia, G., Harkonen, S., Merigan, T.C. and Blaschke, T.F.: Pharmacokinetics of rifabutin. *Agents Chemother.*, 33 (1989) 1237-1241; *C.A.*, 111 (1989) 108467g.
- 2879 Stubbs, C., Skinner, M.F. and Kanfer, I.: Re-usability of solid-phase extraction columns during the HPLC analysis of some macrolide antibiotics in serum and urine. *Chromatographia*, 29 (1990) 31-34.
- 2880 Yurek, D.A., Kuo, M.S. and Li, G.P.: Assay of methylthiolincosaminide in fermentations by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 502 (1990) 184-188.

For additional information see:

C.A., 111 (1989) 186814g, 208620v, 208625a;
112 (1990) 25753s.

See also 3055.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

- 2881 Patel, B.M., Moye, H.A. and Weinberger, R.: Formation of fluorophores from nitrogenous pesticides by photolysis and reaction with OPA-2-mercaptoethanol for fluorescence detection in liquid chromatography. *J. Agric. Food Chem.*, 38 (1990) 126-134.

- 2882 Svoboda, L., Jandera, P. and Churacek, J.: Liquid chromatography of carbamate, urea, triazine, phenoxycarboxylic and organochlorine pesticides. *Collect. Czech. Chem. Commun.*, 54 (1989) 2120-2132; *C.A.*, 111 (1989) 227124c.

29a. Chlorinated insecticides

- 2883 Hogendoorn, E.A., van der Hoff, G.R. and van Zoonen, P.: Automated sample clean-up and fractionation of organochlorine pesticides and polychlorinated biphenyls in human milk using HP-HPLC with column-switching. *J. High Resolut. Chromatogr.*, 12 (1989) 784-789.

See also 2887.

29b. Phosphorus insecticides

- 2884 Husain, S., Alvi, S.N. and Rao, R.N.: High-performance liquid chromatographic separation and determination of quinalphos in technical and commercial formulations. *J. Chromatogr.*, 502 (1990) 208-213.
- 2885 Kumar, R.: Simultaneous determination of some organophosphorus pesticides by high performance liquid chromatography. *Biomed. Chromatogr.*, 3 (1989) 272-273.

29c. Carbamates

- 2886 Marvin, C.H., Brindle, I.D., Hall, C.D. and Chiba, M.: Development of an automated high-performance liquid chromatographic method for the on-line pre-concentration and determination of trace concentrations of pesticides in drinking water. *J. Chromatogr.*, 503 (1990) 167-176.

29d. Herbicides

- 2887 Bogus, E.R., Watschke, T.L. and Mumma, R.O.: Utilization of solid-phase extraction and reversed-phase and ion-pair chromatography in the analysis of seven agrochemicals in water. *J. Agric. Food Chem.*, 38 (1990) 142-144.
- 2888 Brown, R.W.: Synthesis and herbicidal activity of optically active methyl 3-hydroxy-4-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]pentanoates. *J. Agric. Food Chem.*, 38 (1990) 269-273.
- 2889 Durand, G. and Barcelo, D.: Determination of chlorotriazines and their photolysis products by liquid chromatography with photodiode-array and thermospray mass spectrometric detection. *J. Chromatogr.*, 502 (1990) 275-286.
- 2890 Gamble, D.S. and Khan, S.U.: Atrazine in organic soil: chemical speciation during heterogeneous catalysis. *J. Agric. Food Chem.*, 38 (1990) 297-308.
- 2891 Guinivan, R.A., Chang, W.M. and Prince, J.L.: Chlorimuron ethyl. *Anal. Methods Pestic. Plant Growth Regul.*, 16 (1988) 37-51; *C.A.*, 112 (1990) 2518e.
- 2892 Pastore, F., Lavagnini, I., Boaretto, A. and Magno, F.: Ion chromatographic determination of N-nitrosoglyphosate in a glyphosate matrix. *Anal. Chim. Acta*, 230 (1990) 29-34.
- 2893 Pommery, J., Mathieu, M., Mathieu, D. and Lhermitte, M.: High-performance liquid chromatographic determination of atrazine in human plasma. *J. Chromatogr.*, 526 (1990) 569-574.
- 2894 Powell, H.A., Kerby, N.W. and Rowell, P.: High-performance liquid chromatographic determination of the herbicide glyphosate and its metabolite (aminomethyl)phosphonic acid and their extraction from cyanobacteria. *J. Chromatogr.*, 502 (1990) 201-207.
- 2895 Schüssler, W.: Automatic measurement of bantazone and phenoxy acid herbicides by HPLC with three different detectors. *Chromatographia*, 29 (1990) 24-30.

- 2896 West, S.D., Langeland, K.A. and Laroche, F.B.: Residues of fluridone and a potential photoproduct (N-methylformamide) in water and hydrosol treated with the aquatic herbicide sonar. *J. Agric. Food Chem.*, 38 (1990) 315-319.
- 2897 Wigman, L.S., Thomson, M.L. and Wayne, R.S.: Single column ion chromatographic determination of anions in agriculturally useful quaternary compounds using indirect UV detection. *J. Liq. Chromatogr.*, 12 (1989) 3219-3229.

See also 2824.

29e. Fungicides

- 2898 Brandsteterova, E., Lehotay, J., Garaaj, J. and Leclercq, P.A.: High-performance liquid chromatographic determination of dibutyldithiocarbamate degradation products in cucumbers and soil. *J. Chromatogr.*, 502 (1990) 214-220.
- 2899 Engel, P., Dischinger, H.C., Jr. and Ullah, A.H.J.: High-performance liquid chromatography separation of nikkomycins X and Z. *Prepar. Biochem.*, 19 (1989) 321-328.
- 2900 Liu, C.-H., Mattern, G.C., Yu, X. and Rosen, J.D.: (Determination of benomyl by high-performance liquid chromatography/mass spectrometry/selected ion monitoring. *J. Agric. Food Chem.*, 38 (1990) 167-171.

29f. Other types of pesticides and various agrochemicals

- 2901 Irvine, G.B., Ennis, M. and Williams, C.H.: Visula detection of peptidase activity using fluorogenic substrates in a microtiter plate assay. *Anal. Biochem.*, 185 (1990) 304-307.
- 2902 Moye, H.A., Malagodi, M.H., Yoh, J., Deyrup, C.L., Chang, S.M., Leibee, G.L., Ku, C.C. and Wislocki, P.G.: Avermectin B_{1a} metabolism in celery: a residue study. *J. Agric. Food Chem.*, 38 (1990) 290-297.
- 2903 Werner, G.: (Analysis of plant-protection agents and pesticides). *DVGW-Schriftenr., Wasser*, 65 (1989) 57-74; *C.A.*, 111 (1989) 219025f.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 2904 Reynolds, S.L., Scotter, M.J. and Wood, R.: Determination of synthetic coloring matter in foodstuffs - collaborative trial. Part I: initial trial. *J. Assoc. Public Anal.*, 26 (1988) 7-25; *C.A.*, 111 (1989) 230760n.

For additional information see:

- C.A.*, 111 (1989) 230761p;
112 (1990) 48061d.

See also 2459.

30b. Chloroplast and other natural pigments

- 2905 Almela, L., Lopez-Roca, J.-M., Candela, M.E. and Alcazar, M.D.: Separation and determination of individual carotenoids in a *Capsicum* cultivar by normal-phase high-performance liquid chromatography. *J. Chromatogr.*, 502 (1990) 95-106.
- 2906 Czinkotai, B., Daood, H.G., Biacs, P.A. and Hajdu, F.: Separation and detection of paprika pigments by HPLC. *J. Liq. Chromatogr.*, 12 (1989) 2707-2717.

- 2907 Danehower, D.A. and Kelley, W.T.: Rapid extraction and high speed liquid chromatography of *Nicotiana tabacum* leaf pigments. *J. Chromatogr.*, 502 (1990) 431-436.
- 2908 De Leenheer, A.P., Lambert, W.E., de Bersaques, J.P. and Kint, A.H.: High-performance liquid chromatographic determination of etretinate and all-*trans*- and 13-*cis*-acitretin in human plasma. *J. Chromatogr.*, 500 (1990) 637-642.
- 2909 De las Rivas, J., Abadia, A. and Abadia, J.: A new reversed-phase HPLC method resolving all major higher plant phytosynthetic pigments. *Plant. Physiol.*, 91 (1989) 190-192; *C.A.*, 111 (1989) 190668e.
- 2910 Lützw, M. and Kleinig, H.: Chlorophyll-free chromoplasts from daffodil contain most of the enzymes for chlorophyll synthesis in a highly active form. *Arch. Biochem. Biophys.*, 277 (1990) 94-100.
- 2911 Minguez Mosquera, M.I., Montano Asquerino, A., Garrido Fernandez, J. and Gandul Rojas, B.: (Application of HPLC to the separation and identification of chlorophylls and carotenoids from olives.) *Grasas Aceites (Seville)*, 39 (1988) 363-366; *C.A.*, 111 (1989) 193156x.
- 2912 Pesek, C.A., Warthesen, J.J. and Taoukis, P.S.: A kinetic model for equilibration of isomeric β -carotenes. *J. Agric. Food Chem.*, 38 (1990) 41-45.
- 2913 Saleh, F.Y., Ong, W.A. and Chang, D.Y.: Structural features of aquatic fulvic acids. Analytical and preparative reversed-phase high-performance liquid chromatography separation with photodiode array detection. *Anal. Chem.*, 61 (1989) 2792-2800.
- 2914 Suemitsu, R., Horiuchi, K., Ohnishi, K. and Kubota, M.: High-performance liquid chromatographic determination of alterporriol D and E in fermentation of *Altenaria porri* (Ellis) ciferri. *J. Chromatogr.*, 503 (1990) 282-287.
- 2915 Susic, M. and Armstrong, L.G.: High-performance liquid chromatographic determination of humic acid in sodium aluminate solution. *J. Chromatogr.*, 502 (1990) 443-446.
- 2916 Taga, M., Tanaka, S. and Fukushima, M.: Measurement of copper complexing ability of humic acids by using diethylaminoethyl Sephadex A-25 column. *Anal. Sci.*, 5 (1989) 597-600; *C.A.*, 112 (1990) 43619g.

For additional information see:

- C.A.*, 111 (1989) 219347n, 239352p;
112 (1990) 3772v, 4902z, 6186t.

31. PLASTICS AND THEIR INTERMEDIATES

- 2917 Almdal, K.: Absolute molar mass distribution determination by size exclusion chromatography. Synthesis of narrow molar mass distribution polymers. Anionic polymerization under high vacuum conditions. A manual for synthesis of high molar mass polyisoprene, polybutadiene and polystyrene. *Risoe Natl. Lab., (Rep) Risoe-M*, Risoe-M-2787 (V. 2) (1989) 57 p.; *C.A.*, 111 (1989) 233773e.
- 2918 Almdal, K.: Absolute molar mass distribution determination by size exclusion chromatography. Synthesis of narrow molar mass distribution polymers. Characterization of the molar mass distribution of poly(2-hydroxyethyl methacrylate) by size exclusion chromatography with coupled refractive index and low angle laser light scattering detection. *Risoe Natl. Lab. (Rep.) Risoe-M*, Risoe-M-2787 (V. 1) (1989) 140 p.; *C.A.*, 111 (1989) 234052f.
- 2919 Begley, T.H. and Hollifield, H.C.: High-performance liquid chromatographic determination of migrating poly(ethylene terephthalate) oligomers in corn oil. *J. Agric. Food Chem.*, 38 (1990) 145-148.

- 2920 Belyakova, L.D., Platonova, N.P., Shevchenko, T.I., Svec, F. and Hradil, J.: Surface chemistry of methacrylate polymeric sorbents: chromatography and adsorption. *Pure Appl. Chem.*, 61 (1989) 1889-1896; *C.A.*, 112 (1990) 57653a - a review with 25 refs.
- 2921 Dmitrenko, A.V., Ulinskaya, N.N., Volkov, A.M. and Ivanchev, S.S.: (Structure of the grafted polymer layer and chromatographic properties of polymer-inorganic systems of aerosil-polystyrene type). *Vysokomol. Soedin., Ser. A*, 31 (1989) 1591-1596; *C.A.*, 112 (1990) 37267z.
- 2922 Estrin, Ya.I.: (Use of a Milikhrom chromatograph to study the molecular weight distribution of polymer). *Zavod. Lab.*, 55 (1989) 24-25; *C.A.*, 112 (1990) 36985v.
- 2923 Kuzaev, A.I., Sire, E.M., Sokolov, M.I. and Kamaeva, O.A.: (Determination of molecular weight characteristic of BSR rubber emulsion polymerization at the latex stage). *Kauch. Rezina*, (1989) 33-35; *C.A.*, 112 (1990) 22151h.
- 2924 Larroque, M. and Brun, S.: Dosage du méthylmethacrylate dans le serum par chromatographie liquide a haute performance. *J. Chromatogr.*, 525 (1990) 249-252.
- 2925 Maniar, M.L., Kalonia, D.S. and Simonelli, A.P.: Use of liquid chromatography and mass spectroscopy to select an oligomer representative of polyester hydrolysis pathways. *J. Pharm. Sci.*, 78 (1989) 858-862.
- 2926 Mori, S.: Separation of poly(styrene-vinyl acetate) block copolymers by liquid adsorption chromatography. *J. Chromatogr.*, 503 (1990) 411-419.
- 2927 Mourey, T.H., Miller, S.M., Ferrar, W.T. and Molaire, T.R.: Size-exclusion chromatography of poly(bis(trifluoroethoxy)phosphazene). *Macromolecules*, 22 (1989) 4286-4291; *C.A.*, 111 (1989) 21522u.
- 2928 Safieddine, A.M. and Hester, R.D.: Size exclusion chromatography of high-molecular-weight water-soluble polymers. *Polym. Prepr. (Am. Chem. Soc. Div. Polym. Chem.)*, 30 (1989) 369-370; *C.A.*, 112 (1990) 36989z.
- 2929 Singliar, M., Macho, V., Gasparovicova, E. and Stresinkova, D.: (Qualitative analysis for alkylphenylurethanes by gas chromatography and HPLC). *Chem. Prum.*, 39 (1989) 254-257; *C.A.*, 112 (1990) 30063w.
- 2930 Taromi, F.A. and Grubisic-Gallot, Z.: Characterization of star-shaped polystyrene samples by size exclusion chromatography. *Eur. Poly. J.*, 25 (1989) 1183-1187; *C.A.*, 112 (1990) 57168w.

For additional information see:

C.A., 111 (1989) 233946p, 234054h, 234055j, 234056k, 234816h, 234817j, 234818k;
112 (1990) 37171p, 48085q, 57969h.

See also 2152, 2229, 2808.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 2931 Cizmarik, J., Subert, J. and Vespalec, R.: High-performance liquid column chromatography in the pharmacopeial quality control of drugs. *Acta Fac. Pharm. Univ. Comenianae* 1988, 42 (1989) 59-85; *C.A.*, 112 (1990) 42683m - a review with 28 refs.
- 2932 Doyle, E., McDowall, R.D., Murkitt, G.S., Picot, V.S. and Rogers, S.J.: Automated sample handling for drug analysis from biological fluids using HPLC. *Anal. Proc. (London)*, 26 (1989) 294-295; *C.A.*, 111 (1989) 166622q - a review with 12 refs.

- 2933 Dyas, A.M., Robinson, M.L. and Fell, A.F.: New approaches to chiral resolution of drug enantiomers and related compounds by HPLC. *Anal. Proc. (London)*, 26 (1989) 346-348; *C.A.*, 112 (1990) 25740k - a review with 8 refs.
- 2934 Haginaka, J., Nishi, H., Shibukawa, A. and Nakagawa, T.: (High-performance liquid chromatography of drugs). *Bunseki*, (1989) 207-215; *C.A.*, 112 (1990) 42678p - a review with 86 refs.
- 2935 Kelly, M.T., Dadgar, D. and Smyth, M.R.: Some recent developments in high-performance liquid chromatography in biopharmaceutical analysis. *Anal. Proc. (London)*, 26 (1989) 341-343; *C.A.*, 112 (1990) 25739s - a review with 14 refs.
- 2936 Law, B.: Use of silica with reversed-phase type eluents for the analysis of basic drugs and metabolites. *TrAC*, 9 (1990) 31-36.
- 2937 McDowall, R.D., Doyle, E., Murkitt, G.S. and Picot, V.S.: Sample preparation for the HPLC analysis of drugs in biological fluids. *J. Pharm. Biomed. Anal.*, 7 (1989) 1087-1096 - a review with 25 refs.
- 2938 Radzik, D.M. and Lunte, S.M.: Application of liquid chromatography/electrochemistry in pharmaceutical and biochemical analysis: a critical review. *Crit. Rev. Anal. Chem.*, 20 (1989) 317-358; *C.A.*, 111 (1989) 19051ly - a review with 190 refs.
- 2939 Stevenson, D.: Approaches to method development for trace organic analysis. *J. Pharm. Biomed. Anal.*, 7 (1989) 1097-1101.
- 2940 Tamisier-Karolak, L., Farinotti, R., Bossant, M.J. and Dauphin, A.: (Importance and limits of new internal surface reverse phase supports for therapeutic drug monitoring.) *J. Pharm. Clin.*, 7 (1988) 543-581; *C.A.*, 111 (1989) 208614w.
- 2941 Thuaud, N., Seville, B., Deratani, A. and Lelievre, G.: Determination by high-performance liquid chromatography of the binding properties of charged β -cyclodextrin derivatives with drugs. *J. Chromatogr.*, 503 (1990) 453-458.
- 2942 Wainer, I.W.: The application of achiral/chiral coupled-column high-performance liquid chromatography to biomedical analysis. *J. Pharm. Biomed. Anal.*, 7 (1989) 1033-1038.
- 2943 Wong, S.H.Y.: Advances in liquid chromatography and related methodologies for therapeutic drug monitoring. *J. Pharm. Biomed. Anal.*, 7 (1989) 1011-1032 - a review with 94 refs.
- 2944 Zhang, L.: (Application of HPLC for drug analysis in body fluids). *Hebei Yixueyuan Xuebao*, 10 (1989) 123-128; *C.A.*, 111 (1989) 208537y - a review with 25 refs.

For additional information see:

C.A., 111 (1989) 166680g, 208641c, 208642d, 239635h;
112 (1990) 2168r, 4011h, 42682k.

See also 2187, 2218, 2249, 2268.

32b. Antirheumatics and antiinflammatory drugs

- 2945 Katagiri, Y., Noara, K., Ichikawa, N., Hayashibara, M. and Iwamoto, K.: High-performance liquid chromatographic procedure for the simultaneous determination of norfloxacin, fenbufen and felbinac in rat plasma. *Chem. Pharm. Bull.*, 37 (1989) 2858-2860.
- 2946 Larsen, C.: Quantitative determination of dextran-naproxen ester pro-drugs with varying molecular weights and degrees of substitution in biological media by means of high-performance size exclusion chromatography with fluorescence detection. *J. Pharm. Biomed. Anal.*, 7 (1989) 1173-1181.
- 2947 Rustum, A.M.: Measurement of ibuprofen in human whole blood by reversed-phase ion-paired high-performance liquid chromatography using a pH-stable polymeric column. *J. Chromatogr.*, 526 (1990) 246-253.

- 2948 Sane, R.T., Banavalikar, V.J., Joshi, M.D. and Nayak, V.: High-performance liquid chromatographic determination of ketoprofen in pharmaceuticals. *Indian Drugs*, 27 (1989) 54-57; *C.A.*, 112 (1990) 11995k.

see also 2218, 2439, 3026.

32c. Autonomic and cardiovascular drugs

- 2949 Boje, K.M. and Fung, H.-L.: Pharmacokinetic interaction between ethanol and nitrendipine in the rat: role of hypothermia as a contributing mechanism. *Drug Metab. Disp.*, 18 (1990) 127-129.
- 2950 Christ, D.D. and Walle, T.: Stereoselective sulfation of R,S-4'-hydroxypropranolol by canine hepatic cytosol and partially purified phenolsulfotransferases. *J. Pharmacol. Exp. Ther.*, 251 (1989) 949-955.
- 2951 Christ, D.D., Walle, U.K., Oatis, J.E., Jr. and Walle, T.: Pharmacokinetics and metabolism of the pharmacologically active 4'-hydroxylated metabolite of propranolol in the dog. *Drug Metab. Disp.*, 18 (1990) 1-4.
- 2952 Cook, C.S., Gwilt, P.R., Kowalski, K., Gupta, S., Oppermann, J. and Karim, A.: Pharmacokinetics of disopyramide in the dog. Importance of mono-N-dealkylated metabolite kinetics in assessing pharmacokinetic modeling of the parent drug. *Drug Metab. Disp.*, 18 (1990) 42-49.
- 2953 Dasgupta, A., Rosenzweig, I.B., Turgeon, J. and Raisys, V.A.: Encainide and metabolites analysis in serum or plasma using a reversed-phase high-performance liquid chromatographic technique. *J. Chromatogr.*, 526 (1990) 260-265.
- 2954 Enquist, M. and Hermansson, J.: Separation and quantitation of (R)- and (S)-atenolol in human plasma and urine using an α_1 -AGP column. *Chirality*, 1 (1989) 209-215; *C.A.*, 111 (1989) 186832m.
- 2955 Ginebault, P., Haddouche, A. and Bianchetti, G.: High-performance liquid and gas chromatographic methods for the determination of cicloprolol. *J. Chromatogr.*, 525 (1990) 359-367.
- 2956 Jajoo, H.K., Mayol, R.F., LaBudde, J.A. and Blair, I.A.: Structural characterization of urinary metabolites of the antiarrhythmic drug encainide in human subjects. *Drug Metab. Disp.*, 18 (1990) 28-35.
- 2957 Kountourellis, J.E. and Markopoulou, C.: A simultaneous assay by high performance liquid chromatography of bupropion hydrochloride and terbutaline sulphate in dosage forms. *J. Lig. Chromatogr.*, 12 (1989) 3279-3286.
- 2958 Lakings, D.B., Friis, J.M. and Bruns, M.B.: Determination of ditekiren, a renin inhibitor peptide, in monkey serum using high-performance liquid chromatography with solid-phase extraction. *J. Chromatogr.*, 526 (1990) 273-281.
- 2959 Li, X.Y., He, J.B., Tan, W.P., Yu, S.T., Huang, W., Sun, J.O., Yu, C.H. and Qian, F.: (Determination of flecainide concentration in blood plasma by HPLC and pharmacokinetic parameters in volunteers.) *Yaoxue Xuebao*, 24 (1989) 452-457; *C.A.*, 111 (1989) 208616y.
- 2960 Matuszewski, B.K., Constanzer, M.L., Hesse, G.A. and Bayne, W.F.: Development of direct stereoselective and nonstereoselective assays in biological fluids for the enantiomers of a thieno[2,3-b]thiopyran-2-sulfonamide, a topically effective carbonic anhydrase inhibitor. *J. Chromatogr.*, 526 (1990) 461-473.
- 2961 Mehvar, R.: Liquid chromatographic analysis of atenolol enantiomers in human plasma and urine. *J. Pharm. Sci.*, 78 (1989) 1035-1039.
- 2962 Neidlein, R., Volland, C. and Strein, K.: Pharmacokinetics of the partial β -agonist doxaminol in dog. *Arzneim.-Forsch.*, 40 (1990) 27-31.
- 2963 Neidlein, R., Volland, C. and Strein, K.: Biotransformation of the partial β -agonist doxaminol in dog. *Arzneim.-Forsch.*, 40 (1990) 130-136.
- 2964 Ou, C.-N., Rognerud, C.L., Duong, L.T. and Frawley, V.L.: Liquid-chromatographic determination of amiodarone and N-desethylamiodarone in serum. *Clin. Chem. (Winston-Salem)*, 36 (1990) 532-534.

- 2965 Persson, B.A., Balmer, K., Lagerström, P.-O and Schill, G.: Enantioselective determination of metoprolol in plasma by liquid chromatography on a silica-bonded α_1 -acid glycoprotein column. *J. Chromatogr.*, 500 (1990) 629-636.
- 2966 Piquette-Miller, M., Foster, R.T., Pasutto, F.M. and Jamali, F.: Stereospecific high-performance liquid chromatographic assay of acebutolol in human plasma and urine. *J. Chromatogr.*, 526 (1990) 129-137.
- 2967 Rudolph, M.: Direct enantiomeric separation of saterinone by high-performance liquid chromatography. *J. Chromatogr.*, 525 (1990) 161-168.
- 2968 Salem, M.A.S., Alaysi, H.N. and Badwan, A.A.: High pressure liquid chromatographic analysis and dissolution of famotidine in tablet formulation. *Anal. Lett.*, 22 (1989) 2501-2510.
- 2969 Sane, R.T., Ladage, K.D., Kothurkar, R.M., Tendolkar, R.V., Gangal, D.P. and Sonawane, K.K.: High-performance liquid chromatographic determination of pentoxifylline in pharmaceuticals. *Indian Drugs*, 27 (1989) 58-60; *C.A.*, 112 (1990) 11996m.
- 2970 Sastry, C.S.P., Suryanarayana, M.V., Tipirneni, A.S.R.P., Prasad, T.V.N., Murthy, D.V.N. and Reddy, K.N.: High-performance liquid chromatographic determination of triamterene and benzthiazide in combined dosage forms. *Indian Drugs*, 27 (1989) 191-194; *C.A.*, 112 (1990) 62774v.
- 2971 Schwende, F.J. and Lewis, R.C.: Determination of nicorandil in plasma using high-performance liquid chromatography with photoconductivity and ultraviolet detection. Application to pre-clinical pharmacokinetics in beagle dogs. *J. Chromatogr.*, 525 (1990) 151-160.
- 2972 Shimooka, K., Sawada, Y. and Tatamatsu, H.: Analysis of amlodipine in serum by a sensitive high-performance liquid chromatographic method with amperometric detection. *J. Pharm. Biomed. Anal.*, 7 (1989) 1267-1272.
- 2973 Singh, A.K., McArdle, C., Gordon, B., Ashraf, M. and Grandley, K.: Simultaneous analysis of furosemide and bumetanide in horse plasma using high performance liquid chromatography. *Biomed. Chromatogr.*, 3 (1989) 262-265.
- 2974 Steinhart, H. and Rudolph, M.: (Synthesis and analysis of metabolites of the β -blocker carazolol.) *Dtsch. Lebenam.-Rundsch.*, 85 (1989) 137-140; *C.A.*, 111 (1989) 166669k.
- 2975 Wang, Y. and Wu, L.: (Simultaneous determination of serum mexiletine, disopyramide, quinidine, lidocaine and its metabolite MEGX by high-performance liquid chromatography). *Zhonghua Yixue Jianyan Zazhi*, 12 (1989) 76-81; *C.A.*, 111 (1989) 108390b.
- 2976 Weil, A., Caldwell, J. and Strolin-Benedetti, M.: The metabolism and disposition of 14 C-fenofibrate in human volunteers. *Drug Metab. Disp.*, 18 (1990) 115-120.

For additional information see:

- C.A.*, 111 (1989) 186718d;
112 (1990) 133b, 11990e, 42552t, 62612r.

See also 2218, 2941, 2942.

32d. Central nervous system drugs

- 2977 Avramova, J.: Simultaneous determination of propyphenazone, paracetamol and caffeine in blood by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 7 (1989) 1221-1224.
- 2978 Barret, L., Jaussaud, P., Pautet, F., Guyot, J.L. and Daudon, M.: Metabolism of metapramine *in vitro* by chemical model systems and rat liver microsomes. *Arzneim.-Forsch.*, 39 (1989) 1574-1576.
- 2979 Bianchi, G., Caccia, S., Della Vedova, F. and Garattini, S.: Kinetics and brain uptake of S 9795, a new xanthine derivative, in rats. *Eur. J. Drug Metab.*, 14 (1989) 201-208.

- 2980 Blanchflower, W.J. and Kennedy, D.G.: A rapid screening procedure for the detection and quantification of clenbuterol in bovine urine by thermospray liquid chromatography/mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 935-936; *C.A.*, 111 (1989) 224731n.
- 2981 Bun, H., Duplan, V., Crevat-Pisano, P., Llurens, M. and Durand, A.: Rapid determination of the benzodiazepine antagonist flumazenil, Ro 15-1788, by high performance liquid chromatography. *Biomed. Chromatogr.*, 3 (1989) 269-271.
- 2982 Burton, L.C., Loftus, N.J., Vere, D.W. and Whelpton, R.: Determination of plasma nefopam by liquid chromatography and electrochemical detection. *J. Chromatogr.*, 526 (1990) 159-168.
- 2983 Carlucci, G.: High-performance liquid chromatographic assay for nizatidine, a new H₂ blocker, in human plasma and urine using disposable solid-phase extraction columns. *J. Chromatogr.*, 525 (1990) 490-494.
- 2984 Celardo, A., Passerini, P. and Bonati, M.: Placental transfer and tissue distribution of thiopental in the pregnant rat. *J. Pharmacokin. Biopharm.*, 17 (1989) 425-440.
- 2985 D'Aranno, V., Mancinelli, A., Borsini, F., Furio, M. and Meli, A.: Determination of nomifensine in rat brain by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr.*, 525 (1990) 234-239.
- 2986 De Vries, J.X., Rudi, J., Walter-Sack, I. and Conradi, R.: The determination of total and unbound midazolam in human plasma. A comparison of high performance liquid chromatography, gas chromatography and gas chromatography/mass spectrometry. *Biomed. Chromatogr.*, 4 (1990) 28-33.
- 2987 DeAngelis, R., Loebs, P., Maeher, R., Savarese, J. and Welch, R.: High-performance liquid chromatographic analysis of doxacurium, a new long-acting neuromuscular blocker. *J. Chromatogr.*, 525 (1990) 389-400.
- 2988 Degroodt, J.M., Wyhowski de Bukanski, B., Beernaert, H. and Courtheyn, D.: Clenbuterol residue analysis by HPLC-HPTLC in urine and animal tissues. *Z. Lebensm.-Unters. Forsch.*, 189 (1989) 128-131; *C.A.*, 111 (1989) 213437v.
- 2989 Echizen, H., Nakura, M. and Ishizaki, I.: Rapid and simple high-performance liquid chromatographic determination of saliva antipyrine for routine antipyrine test. *J. Chromatogr.*, 526 (1990) 296-299.
- 2990 Estus, S. and Blumer, J.L.: Critical period of phenytoin teratogenic action in the sea urchin *Arbacia punctulata* embryo. *J. Pharmacol. Exp. Ther.*, 251 (1989) 782-789.
- 2991 Fellegvari, I., Visy, J., Valko, K., Lang, T. and Simonyi, M.: Investigation of conformational diastereomers of 2,3-benzodiazepines by high performance liquid chromatography. *J. Liq. Chromatogr.*, 12 (1989) 2719-2732.
- 2992 Franklin, M.: Determination of plasma buspirone by high-performance liquid chromatography with coulometric detection. *J. Chromatogr.*, 526 (1990) 590-596.
- 2993 Gao, C.-X. and Krull, I.S.: Determination of enantiomeric drugs in physiological fluids using on-line solid phase derivatizations and reversed-phase liquid chromatography. *J. Pharm. Biomed. Anal.*, 7 (1989) 1183-1198.
- 2994 Gomita, Y., Nanba, M., Furuno, K., Eto, K. and Araki, Y.: Microdetermination of cimetidine in rat plasma by high-performance liquid chromatography. *Int. J. Pharm.*, 54 (1989) 89-93; *C.A.*, 111 (1989) 166696s.
- 2995 Hariharan, M., Kindt, E.K., VanNoord, T. and Tandon, R.: An improved sensitive assay for simultaneous determination of plasma haloperidol and reduced haloperidol levels by liquid chromatography using a coulometric detector. *Ther. Drug Monit.*, 11 (1989) 701-707; *C.A.*, 112 (1990) 106v.
- 2996 Inamura, Y., Shibasaki, S., Kubo, T., Kogure, M., Ishii, Y., Gotoh, Y., Asai, S. and Ishikawa, K.: High-performance liquid chromatographic determination of *cis*-N-(1-benzyl-2-methylpyrrolidin-3-yl)-5-chloro-2-methoxy-4-methylaminobenzamide in rat blood and brain using electrochemical detection. *J. Chromatogr.*, 526 (1990) 282-288.

- 2997 Johansson, M., Forsmo-Bruce, H., Alm, A.T. and Westerlund, D.: Effects of system peaks in a coupled column system for noscapine and its metabolites. *J. Pharm. Biomed. Anal.*, 7 (1989) 1055-1070.
- 2998 Kojima, K., Yamanaka, M., Nakanishi, Y. and Arakawa, S.: High-performance liquid chromatographic determination of etilefrine in human plasma using combined solid-phase and organic solvent extraction and electrochemical detection. *J. Chromatogr.*, 525 (1990) 210-217.
- 2999 Korfmacher, W.A., Getek, T.A., Hansen, E.B., Jr. and Cerniglia, C.E.: Direct analysis of microbial extracts containing metabolites of ethylenediamine-type antihistamines via high-performance liquid chromatography-thermospray mass spectrometry. *Anal. Biochem.*, 185 (1990) 136-142.
- 3000 Kountourellis, J.E. and Marcopoulou, C.: Rapid isocratic high-performance chromatographic determination of some hypnotics/sedatives in the presence of phenytoin in commercial formulations. *Indian Drugs*, 27 (1989) 66-70; *C.A.*, 112 (1990) 25803h.
- 3001 Kountourellis, J.E., Markopoulou, C. and Georgakopoulos, P.P.: Separation and simultaneous determination of bupropion and salbutamol in dosage forms by high-performance liquid chromatography. *J. Chromatogr.*, 502 (1990) 189-192.
- 3002 Loennechen, T. and Dahl, S.G.: High-performance liquid chromatography of levomepromazine (methotrimeprazine) and its main metabolites. *J. Chromatogr.*, 503 (1990) 205-215.
- 3003 Mancilla, J., Valdes, E. and Gil, L.: A novel isocratic HPLC method to separate and quantify acetanilide and its hydroxy aromatic derivatives: 2-, 3- or 4-hydroxyacetanilide (paracetamol or acetaminophen). *Eur. J. Drug Metab.*, 14 (1989) 241-244.
- 3004 Marini, D. and Balestrieri, F.: Characterization, purification and HPLC determination of famotidine. *Prod. Chim. Aerosol Sci.*, 30 (1989) 40-43; *C.A.*, 111 (1989) 180842t.
- 3005 Nagai, T. and Kamiyama, S.: Assay of the optical isomers of methamphetamine and amphetamine in rat urine using high-performance liquid chromatography with chiral cellulose-based columns. *J. Chromatogr.*, 525 (1990) 203-209.
- 3006 Nilsson, L.B.: Determination of remoxipride in plasma and urine by reversed-phase column liquid chromatography. *J. Chromatogr.*, 526 (1990) 139-150.
- 3007 Ogden, D.A., McKechnie, A.A., Bolton, D.A., Rankin, R.B., Dobbins, S.E. and Clements, J.A.: Determination of pipothiazine in human plasma by reversed-phase high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 7 (1989) 1273-1280.
- 3008 Okudzhava, V.M., Chankvetadze, B.G., Rukhadze, M.D. and Akhobadze, N.A.: (High-performance liquid-chromatographic assay of serum sodium ethaminal). *Farmatsiya (Moscow)*, 38 (1989) 58-60; *C.A.*, 111 (1989) 108416q.
- 3009 Perry, K.W. and Fuller, R.W.: Determination of brain concentrations of 8-hydroxy-2-(di-*n*-propylamino)tetralin by liquid chromatography with electrochemical detection. *Biochem. Pharmacol.*, 38 (1989) 3169-3173; *C.A.*, 112 (1990) 131z.
- 3010 Rustum, A.M. and Gutierrez, L.: Reversed-phase high-performance liquid chromatographic separation of the enantiomers of N-[4,4-di(3-methylthien-2-yl)-but-3-enyl]nipecotic acid on a Pirkle-type phenylglycine stationary phase. *J. Chromatogr.*, 503 (1990) 115-125.
- 3011 Sadana, G.S. and Potdar, A.J.: High performance liquid chromatographic determination of astemizole in pharmaceutical dosage forms. *Indian Drugs*, 27 (1989) 140-142; *C.A.*, 112 (1990) 62769x.
- 3012 Sakano, I. and Kokubo, S.: Simultaneous determination of acetaminophen, caffeine and dihydrocodeine phosphate in tablets by high-performance liquid chromatography. *Anal. Sci.*, 5 (1989) 623-624; *C.A.*, 112 (1990) 25790b.

- 3013 Sane, R.T., Bhate, V.R., Balakrishna, K., Joshi, M.D. and Nayak, V.G.: High-performance liquid chromatographic determination of alprazolam from pharmaceutical preparations. *Indian Drugs*, 26 (1989) 649-650; *C.A.*, 111 (1989) 219381u.
- 3014 Saville, B.A., Gray, M.R. and Tam, Y.K.: Evidence for lidocaine-induced enzyme inactivation. *J. Pharm. Sci.*, 78 (1989) 1003-1008.
- 3015 Selinger, K., Prevost, J. and Hill, H.M.: High-performance liquid chromatography method for the determination of diphenhydramine in human plasma. *J. Chromatogr.*, 526 (1990) 597-602.
- 3016 Thompson, R.D. and Carlson, M.: Determination of thymol in halothane anaesthetic preparations by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 7 (1989) 1199-1206.
- 3017 Treston, A.M. and Hooper, W.D.: Metabolic studies with phenobarbitone, primidone and their N-alkyl derivatives: quantification of substrates and metabolites using chemical ionization gas chromatography-mass spectrometry. *J. Chromatogr.*, 526 (1990) 59-68.
- 3018 Trnavská, Z.: Theophylline protein binding. *Arzneim.-Forsch.*, 40 (1990) 166-169.
- 3019 Uebel, R.A., Wium, C.A., Hawtrey, A.O. and Coetzee, J.: Electrochemical determination of 2,6-diisopropylphenol after high-performance liquid chromatography of extracts from serum. *J. Chromatogr.*, 526 (1990) 293-295.
- 3020 Verhoef, J. and van den Wildenberg, R.M.: Des-enkephalin- γ -endorphin (DE γ E): pharmacokinetics in dogs after intravenous and subcutaneous administration. *Eur. J. Drug Metab.*, 14 (1989) 229-234.
- 3021 Vogelgesang, J.: Determination of neuroleptic and carazolol in pork and beef. *Dtsch. Lebensm.-Rundsch.*, 85 (1989) 251-258; *C.A.*, 111 (1989) 193170x.
- 3022 Vree, T.B., Steegers-Theunissen, R.P.M., Baars, A.M. and Hekster, Y.A.: Direct high-performance liquid chromatographic analysis of *p*-hydroxyphenyl-phenylhydantoin glucuronide, the final metabolite of phenytoin, in human serum and urine. *J. Chromatogr.*, 526 (1990) 581-589.
- 3023 Wilhelm, D. and Kemper, A.: High-performance liquid chromatographic procedure for the determination of clozapine, haloperidol, droperidol and several benzodiazepines in plasma. *J. Chromatogr.*, 525 (1990) 218-224.
- 3024 Wuis, E.W., Janssen, M.G.A., Vree, T.B. and van der Kleijn, E.: Determination of a dantrolene metabolite, 5-(*p*-nitrophenyl)-2-furoic acid, in plasma and urine by high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 575-580.
- 3025 Xie, F., Wang, H.-C., Li, J.-H., Shu, H.-L., Jiang, J.-R., Chang, J.-P. and Hsieh, Y.-Y.: Studies on the metabolism of lappaconitine in humans. Identification of the metabolites of lappaconitine in human urine by high performance liquid chromatography. *Biomed. Chromatogr.*, 4 (1990) 43-46.
- 3026 Yu, H.-Y., Shen, Y.-Z., Sugiyama, Y. and Hanano, Y.: Drug interaction. Effects of salicylate on pharmacokinetics of valproic acid in rats. *Drug Metab. Disp.*, 18 (1990) 121-126.

For additional information see:

C.A., 111 (1989) 186715a, 224736t, 224739w;
112 (1990) 82j, 98u, 25779a.

See also 2786, 2787, 2942, 2975.

32e. Chemotherapeutics (except cytostatics and antibiotics)

- 3027 Al-Meshal, M.A.: Determination of ketoconazole in plasma and dosage forms by high-performance liquid chromatography and a microbiological method. *Anal. Lett.*, 22 (1989) 2249-2263.

- 3028 Hammiche, Y., Guermouche, M.H., Abdeddaim, K. and Brixi, M.D.: (HPLC analysis for pyrazinamide in rat serum). *Analusis*, 17 (1989) 394-397; *C.A.*, 111 (1989) 224740q.
- 3029 Hassanzadeh, M.K., Aweeka, F.T., Wu, S., Jacobson, M.A. and Gambertoglio, J.G.: Determination of phosphonoformic acid in human plasma and urine by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr.*, 525 (1990) 133-140.
- 3030 Horie, M., Saito, K., Hoshino, Y., Nose, N., Hamada, N. and Nakazawa, H.: Identification and determination of sulphamethazine and N⁴-acetylsulphamethazine in meat by high-performance liquid chromatography with photodiode-array detection. *J. Chromatogr.*, 502 (1990) 371-378.
- 3031 Jamaluddin, A.B.M., Sarwar, G., Rahim, M.A. and Rahman, M.K.: High-performance liquid chromatographic assay of rifampicin in human serum. *J. Chromatogr.*, 525 (1990) 495-497.
- 3032 Kaul, S., Dandekar, K.A. and Pittman, K.A.: Analytical method for the quantification of 2',3'-didehydro-3'-deoxythymidine, a new anti-human immunodeficiency virus (HIV) agent, by high-performance liquid chromatography (HPLC) and ultraviolet (UV) detection in rat and monkey plasma. *Pharm. Res.*, 6 (1989) 895-899; *C.A.*, 112 (1990) 104t.
- 3033 Klimowicz, A.: The comparison of some pharmacokinetic parameters of sulphadimethoxine estimated by high performance liquid chromatography and three spectrophotometric methods. *Eur. J. Drug Metab.*, 14 (1989) 181-186.
- 3034 Lacroix, C., Hoang, T.P., Wojciechowski, F., Duwoos, H., Nouveau, J. and Diquet, B.: Simultaneous quantification of zidovudine and its metabolites in serum and urine by high-performance liquid chromatography using a column-switching technique. *J. Chromatogr.*, 525 (1990) 240-245.
- 3035 Long, A.R., Short, C.R. and Barker, S.A.: Method for the isolation and liquid chromatographic determination of eight sulfonamides in milk. *J. Chromatogr.*, 502 (1990) 87-94.
- 3036 Rao, G.R., Murty, S.S.N., Rao, P.J. and Srivastava, C.M.R.: High-performance liquid chromatographic determination of morphazinamide hydrochloride. *Indian Drugs*, 26 (1989) 569-571; *C.A.*, 111 (1989) 239640f.
- 3037 Sane, R.T., Dhruaml, S.N., Mainkar, P.S., Nerurkar, V.R. and Patel, D.: Simultaneous determination of tetramisole and phenolphthalein from pharmaceutical preparations. *Indian Drugs*, 27 (1989) 61-63; *C.A.*, 112 (1990) 11997n.
- 3038 Sane, R.T., Mishra, P.D., Lagade, K.D. and Kothurkar, R.M.: High performance liquid chromatographic determination of chlorquinaldol from pharmaceutical preparations. *Indian Drugs*, 26 (1989) 701-703; *C.A.*, 112 (1990) 62758t.
- 3039 Seydel, J.K., Wempe, E.G., Rosenfeld, M., Jagannathan, R., Mahadevan, P.R. and Dhople, A.M.: *In vitro* and *in vivo* experiments with the new inhibitor of *Mycobacterium leprae* brodimoprim alone and in combination with dapsone. *Arzneim.-Forsch.*, 40 (1990) 69-75.
- 3040 Tjaden, U.R., Reeuwijk, H.J.E.M., van der Greef, J., Pattyn, G., de Bruijn, E.A. and van Oosterom, A.T.: Bioanalysis of suramin in human plasma by ion-pair high-performance liquid chromatography. *J. Chromatogr.*, 525 (1990) 141-149.
- 3041 Vree, T.B. and Shimoda, M.: High-performance liquid chromatography of sulphadimethoxine and its N₁-glucuronide, N₄-acetyl and N₄-acetyl-N₁-glucuronide metabolites in human plasma and urine. *J. Chromatogr.*, 526 (1990) 119-128.

For additional information see:

C.A., 111 (1989) 166686p, 166687q, 224714j, 230764s;
112 (1990) 25778d.

see also 2975, 3052.

32f. Cytostatics

- 3042 Bullen, W.W., Chang, T. and Whitfield, L.R.: High-performance liquid chromatographic assay for trimetrexate in human plasma. *J. Chromatogr.*, 526 (1990) 266-272.
- 3043 Ellenberger, T.E., Wright, J.E., Rosowsky, A. and Beverley, S.M.: Wild-type and drug-resistant *Leishmania major* hydrolyze methotrexate to N-10-methyl-4-deoxy-4-aminopteroate without accumulation of methotrexate polyglutamates. *J. Biol. Chem.*, 264 (1989) 15960-15966.
- 3044 Hais, I.M., Nobilis, M. and Kvasnickova, E.: Evaluation of photometric high-performance liquid chromatographic data for the determination of benflurone metabolites in biological materials with and without concomitant use of a standard. *J. Chromatogr.*, 500 (1990) 643-653.
- 3045 Itoh, M., Murata, H., Honda, M., Watanabe, Y. and Ochi, K.: Reaction of (R)-(-)-2-aminomethylpyrrolidine(1,1-cyclobutanedicarboxylato)platinum(II) with guanosine. *Chem. Pharm. Bull.*, 37 (1989) 2825-2827.
- 3046 Jehl, F., Debs, J., Herlin, C., Quoix, E., Gallion, C. and Monteil, H.: Determination of navelbine and desacetylnavelbine in biological fluids by high-performance liquid chromatography. *J. Chromatogr.*, 525 (1990) 225-233.
- 3047 Ko, R.J., Li, W.Y. and Koda, R.T.: Determination of the antimitotic agents N-desacetylcolchicine, demecolcine and colchicine in serum and urine. *J. Chromatogr.*, 525 (1990) 411-418.
- 3048 Polacek, J., Gustafsson, B., Brandin, S. and Ottersgard-Brorsson, A.: Determination of taumustine and its demethylated metabolites in plasma and urine. *J. Chromatogr.*, 526 (1990) 151-157.
- 3049 Reeuwijk, H.J.E.M., Tjaden, U.R., de Bruijn, E.A. and van der Greef, J.: Bioanalysis of (E)-5-(2-bromovinyl)-2'-deoxyuridine in plasma. *Chromatogram*, 10 (1989) 2-3; *C.A.*, 111 (1989) 166681h.
- 3050 Tatsuhara, T., Tabuchi, F., Yamane, M. and Hori, T.: Rapid and simple method for the determination of nimustine hydrochloride in human blood and brain by high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 507-514.
- 3051 Van Opstal, M.A.J., van der Horst, F.A.L., Holthuis, J.J.M., van Bennekom, W.P. and Bult, A.: Automated reversed-phase chromatographic analysis of etoposide and teniposide in plasma by using on-line surfactant-mediated sample clean-up and column-switching. *J. Chromatogr.*, 495 (1989) 139-151.

See also 2863, 2942.

32g. Other drug categories

- 3052 Dashbalyn, T., Prokof'eva, V.I., Egorov, A.E., Kashtanova, V.V. and Nechaeva, E.B.: (Application of thin-layer and high-performance liquid chromatography to the identification and separation of 10-acyl derivatives of phenothiazine). *Farmatsiya (Moscow)*, 38 (1989) 39-41; *C.A.*, 112 (1990) 62763r.
- 3053 Dixon, R., Fujitaki, J., Chrastil, L. and Lee, J.: AICA riboside: direct quantitation in ultrafiltrates of plasma by HPLC during pharmacokinetic studies in man. *Res. Commun. Chem. Pathol. Pharmacol.*, 65 (1989) 405-408; *C.A.*, 112 (1990) 96s.
- 3054 Krstulovic, A.M. and Vende, J.L.: Improved performance of the second generation α_1 -AGP columns: applications to the routine assay of plasma levels of alfuzosin hydrochloride. *Chirality*, 1 (1989) 243-245; *C.A.*, 111 (1989) 224715k.
- 3055 Lachno, D.R., Patel, N., Rose, M.L. and Yacoub, M.H.: Improved high-performance liquid chromatographic method for analysis of cyclosporin a using an automated sample processor. *J. Chromatogr.*, 525 (1990) 123-132.

- 3056 Lehmann, W.D. and Heinrich, H.C.: Ferrioxamine and its hexadentate iron-chelating metabolites in human post-Desferal urine studied by high-performance liquid chromatography and fast atom bombardment mass spectrometry. *Anal. Biochem.*, 184 (1990) 219-227.
- 3057 Lehr, K.H. and Damm, P.: Simultaneous determination of the sulphonylurea glimepiride and its metabolites in human serum and urine by high-performance liquid chromatography after pre-column derivatization. *J. Chromatogr.*, 526 (1990) 497-505.
- 3058 Lorusso, V., Poggesi, I., Dal Fiume, D.D. and Tirone, P.: High-performance liquid chromatographic assay of iomeprol in plasma and urine. *J. Chromatogr.*, 525 (1990) 401-409.
- 3059 Nakajima, K., Yasuda, T. and Nakazawa, H.: High-performance liquid chromatographic determination of zinc pyrithione in antidandruff preparations based on copper chelate formation. *J. Chromatogr.*, 502 (1990) 379-384.
- 3060 Nakaoka, M. and Hakusui, H.: Measurement of the formation of a thioglucuronide, a metabolite of malotilate, in rat hepatic microsomes by high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 254-259.
- 3061 Ostrovska, V., Pechova, A. and Svobodova, K.: Determination of metabolites of etophyllinclofibrate in plasma by reversed phase high performance liquid chromatography. *J. Liq. Chromatogr.*, 12 (1989) 2793-2799.
- 3062 Pue, M.A., Ransley, J.A., Writer, D.J., Dean, A.J., Franklin, E.R., Beattie, I.G. and Ross, D.A.: The disposition of SK&F L-94901, a selective thyromimetic in rat, dog and cynomolgus monkey. *Eur. J. Drug Metab.*, 14 (1989) 209-219.
- 3063 Shim, C.-K. and Hong, J.-S.: Inter- and intrasubject variations of ranitidine pharmacokinetics after oral administration to normal male subjects. *J. Pharm. Sci.*, 78 (1989) 990-994.
- 3064 Tjandra-Maga, B., Verbesselt, R., Scharpe, S., Verkerk, R., Lambert, W.E., van Liedekerke, B. and de Leenheer, A.: Comparison of cyclosporin A measurement in whole blood by six different methods. *J. Clin. Chem. Clin. Biochem.*, 28 (1990) 53-57.
- 3065 Tománková, H. and Sabartová, J.: Determination of azidothymidine and its degradation product thymine in pharmaceutical dosage forms by HPLC and HPTLC densitometry. *Chromatographia*, 29 (1990) 44-50.
- 3066 Weil, A., Caldwell, J., Guichard, J.P. and Picot, G.: Species differences in the chirality of the carbonyl reduction of [^{14}C] fenofibrate in laboratory animals and humans. *Chirality*, 1 (1989) 197-201; *C.A.*, 111 (1989) 208658p.
- 3067 Zarapkar, S.S., Salunkhe, U.B., Doshi, V.J. and Rele, R.V.: A high-performance liquid chromatographic method for the determination of glipizide in pharmaceutical preparations. *Indian Drugs*, 26 (1989) 564-566; *C.A.*, 111 (1989) 239638m.

For additional information see:

C.A., 111 (1989) 208639h, 224711f;
 112 (1990) 25359t, 25446u, 25749v, 25775a, 42211f, 42403v,
 42734d, 62734g.

See also 2314, 2434, 2514, 2837, 2908, 2945, 2973.

32h. Toxicological and forensic applications

- 3068 Blake, T.J.A. and Beattie, I.G.: Rapid structural analysis of the *in vitro* and *in vivo* metabolism of SKF 95448 by the combined use of thermospray liquid chromatography/mass spectrometry and liquid chromatography/tandem mass spectrometry. *Biomed. Environ. Mass Spectrom.* 1988, 18 (1989) 637-644; *C.A.*, 111 (1989) 166671e.

- 3069 Woolf, T.F.: Applications of HPLC coupled with radioactive flow detection in drug disposition studies. *LC-GC*, 7 (1989) 828-834; *C.A.*, 111 (1989) 208541v - a review with 15 refs.

For additional information see:

C.A., 111 (1989) 168820b.

See also 2510, 2793, 2924, 2973.

32i. Plant extracts

- 3070 Biedrich, P., Engelhardt, U.H. and Herzig, B.: Determination of quercetin-3-O-rutinoside in black tea by HPLC. *Z. Lebensm.-Unters. Forsch.*, 189 (1989) 149-150; *C.A.*, 111 (1989) 213438w.
- 3071 Carrasquer, A.M., Casals, I. and Alegre, L.: Semi-automated method for the determination of abscisic acid in crude plant extracts. *J. Chromatogr.*, 503 (1990) 459-465.
- 3072 Chen, G.L., Wang, H.J., Lin, H.C., Wu, W.L. and Chen, C.F.: A stability indicating method for geniposide by high-performance liquid chromatography/diode array detector. *Chung-hua Yao Hsueh Tsa Chih*, 41 (1989) 289-296; *C.A.*, 112 (1990) 62738m.
- 3073 Miething, H. and Holz, W.: Wirkstoffe in wässrigen Kamillenaufgüssen. 1. Mitteilung: Freisetzungskinetiken ausgewählter Flavonoidglycoside. *Pharmazie*, 44 (1989) 784-785.
- 3074 Perez, N., Matteo, C., Marcos, M.M., Cuervo, A.C. and Blunden, G.: (Quantitative determination of hecogenin and tigogenin in Venezuelan sisal, using high-resolution liquid chromatography.) *Rev. Latinoam. Quim.*, 20 (1989) 102-103; *C.A.*, 111 (1989) 190655y.
- 3075 Vennat, B., Pouget, M.P., Gross, D. and Pourrat, A.: (Preparations containing Hamamelis and horse chestnut extracts: simultaneous assay of hamamelitannin and esculoside). *J. Pharm. Belg.*, 44 (1989) 285-291; *C.A.*, 112 (1990) 11981c.
- 3076 Wagner, H., Bladt, S., Hartmann, U., Dally, A. and Berkulin, W.: Ginkgo biloba. DC- und HPLC-Analyse von Ginkgo-Extrakten und Ginkgo-Extrakte enthaltenden Phytopräparaten. *Dtsch. Apoth.-Ztg.*, 129 (1989) 2421-2429.

For additional information see:

C.A., 112 (1990) 25748u, 25772x.

See also 2452, 2470, 3025.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

See 2792.

33b. Complex mixtures and profiling (single compounds by cross-reference only)

- 3077 Stefansson, M. and Westerlund, D.: Reversed-phase ion-pair liquid chromatography of glucuronides. Retention and selectivity. *J. Chromatogr.*, 499 (1990) 411-421.

See also 2335, 2366, 2379, 2382, 2387, 2401, 2425, 2427, 2433, 2436, 2441, 2443, 2444, 2450, 2461, 2464, 2476, 2480, 2489, 2498, 2512, 2544, 2634, 2718, 2751, 2767, 2770, 2848, 2860.

34. FOOD ANALYSIS

34a. General papers and reviews

- 3078 Engelhardt, H.: (Applications of high-pressure liquid chromatography in food analysis.) *Lebensm.-Biotechnol.*, 6 (1989) 67-76; C.A., 111 (1989) 172525b - a review with 25 refs.
- 3079 Hibi, K. and Bounoshita, M.: (Use of high-performance liquid chromatography in the analysis of bitter compounds in food products.) *Shokuhin to Kagaku*, 31 (1989) 120-125; C.A., 111 (1989) 213399j - a review with 16 refs.

See also 2408, 2460.

34b. Complex mixtures (single compounds by cross-reference only)

- 3080 Bombardt, P.A., Rotwell, B.E., Closson, S.K. and Adams, W.J.: Quantitative determination of itazigrel in rodent diet by reversed-phase HPLC and evaluation of its stability at 20. *Biomed. Chromatogr.*, 3 (1989) 180-182; C.A., 111 (1989) 219370q.
- 3081 Mosandl, A., Kustermann, A., Hener, U. and Hagenauer-Hener, U.: (Analytical differentiation of natural and identical synthetic-lactones in foods.) *Dtsch. Lebensm.-Rundsch.*, 85 (1989) 205-209; C.A., 111 (1989) 172553j.

For additional information see:

C.A., 111 (1989) 172576u, 172578w, 230796d.

See also 2302, 2307, 2308, 2309, 2311, 2315, 2380, 2381, 2384, 2385, 2389, 2404, 2488, 2492, 2506, 2520, 2655, 2657, 2661, 2813, 2825, 2835, 2841, 2844, 2847, 2853, 2904, 2911, 2919, 3021, 3070, 3109, 3136, 3140, 3141.

35. ENVIRONMENTAL ANALYSIS

35a. General papers and reviews

- 3082 Harrison, R.M. and Rapsomanikis, S. (Editors): *Environmental Analysis Using Chromatography Interfaced with Atomic Spectroscopy*. Ellis Horwood Ltd., Chichester, 1989, 370 p.; C.A., 111 (1989) 224457c.

See also 2826, 2830.

35c. Water pollution (complex mixtures; single compounds by cross-reference only)

For additional information see:

C.A., 111 (1989) 247289p, 247290g.

See also 2886, 2895, 2903, 3111.

35d. Soil pollution (complex mixtures; single compounds by cross-reference only)

See 2457.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

- 3083 Funasaki, N., Hada, S. and Neya, S.: Gel filtration chromatographic study of micellar growth of polyethylene glycol dodecyl ether. *Bull. Chem. Soc. Jpn.*, 62 (1989) 2485-2491; *C.A.*, 111 (1989) 240570q.
- 3084 Okada, T.: Chromatographic oligomer separation of poly(oxyethylenes) on K^+ -form cation exchange resin. *Anal. Chem.*, 62 (1990) 327-331.
- 3085 Sasaki, H., Yokai, K. and Okumura, O.: The absorption behavior of dialkyl dimethyl ammonium chloride - nonionic surfactant binary systems. *J. Am. Oil Chem. Soc.*, 67 (1990) 64-67.
- 3086 Varughese, P., Gangoda, M.E. and Gilpin, R.K.: Reversed-phase liquid chromatographic studies of non-ionic surfactants and related solutes. *J. Chromatogr.*, 499 (1990) 469-475.

36b. Antioxidants and preservatives

- 3087 Arpino, P.J., Dilettato, D., Nguyen, K. and Bruchet, A.: Investigation of antioxidants and UV stabilizers from plastics. Part I. Comparison of HPLC and SFC; preliminary SFC/MS study. *J. High Resolut. Chromatogr.*, 13 (1990) 5-12.
- 3088 Dieffenbacher, A., Trisconi, M.J. and Durieux, B.: (Rapid determination of phenolic antioxidants in foods by HPLC.) *Mitt. Geb. Lebensmittelunters. Hyg.*, 80 (1989) 204-224; *C.A.*, 111 (1989) 172551g.

For additional information see:

C.A., 112 (1990) 6196w.

See also 3016.

36c. Various technical products

- 3089 Carbognani, L. and Izquierdo, A.: Preparative compound class separation of heavy oil vacuum residua by high performance liquid chromatography. *Fuel Sci. Technol. Int.*, 8 (1990) 1-15; *C.A.*, 112 (1990) 39421u.
- 3090 Koechne-Waechter, M.: (Contamination control in semiconductor manufacture by liquid chromatography). *GIT Fachz. Lab.*, 33 (1989) 873-875; *C.A.*, 111 (1989) 222732q - a review with 14 refs.
- 3091 Moliner, R., Ibarra, J.V. and Lagarza, M.D.: Size-exclusion chromatography of pyrolysis and hydropyrolysis tars from Spanish low-rank coal. *Fuel*, 68 (1989) 1487-1488; *C.A.*, 111 (1989) 236392r.
- 3092 Stein, J.: (Chromatographic characterization of reaction products of brown coal liquefaction in the carbon monoxide/water system). *Freiberg. Forschungsh. A*, 795 (1989) 23-37; *C.A.*, 112 (1990) 39551m.
- 3093 Zhang, Z.; Chen, S. and Cai, S.: (Analysis of rocks and minerals). *Fenxi Shiyanshi*, 7 (1989) 95-120; *C.A.*, 111 (1989) 224168j - a review with 145 refs.

For additional information see:

C.A., 112 (1990) 39091m.

See also 2402, 2798, 2806, 3103, 3144.

36d. Complex mixtures and unidentified compounds

- 3094 Nakanishi, S., Ando, K., Kawamoto, I., Yasuzawa, T., Sano, H. and Kase, H.: KS-504 compounds, novel inhibitors of Ca^{2+} and calmodulin-dependent cyclic nucleotide phosphodiesterase from *Mollisia ventosa*. *J. Antibiot.*, 42 (1989) 1775-1783.
- 3095 Oswald, G., Buschulte, T. and Schnabel, G.: (Simulation of an affinity chromatographic separation as purification step of a biotechnological production process). *Chem.-Ing.-Tech.*, 61 (1989) 723-725; *C.A.*, 112 (1990) 25592p.

See also 2149.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 3096 Hammer, D.A. and Lauffenburger, D.A.: Effects of nonspecific cell/surface interactions on cell affinity chromatographic separations. *Mater. Res. Soc. Symp. Proc.* 1987, 110 (Biomed. Mater. Devices) (1988) 739-745; *C.A.*, 112 (1990) 3776z.
- 3097 Ikuya, Y. (Editor): *Rapid Identification of Bacteria by Chromatography: Application of New Chromatographic Technology*. Saikon Publishing Co., Tokyo, 1988, 131 pp.; *C.A.*, 111 (1989) 211503q.
- 3098 Kennedy, R.T. and Oates, M.D.: Microcolumn separations and the analysis of single cells. *Science (Washington)*, 246 (1989) 57-63; *C.A.*, 111 (1989) 211494n.

38. INORGANIC COMPOUNDS

38a. Cations

- 3099 Alam, A.M.S. and Roy, B.: Application of cation exchange for separation of metals in mixed solvents (ethylene glycol-water-hydrochloric acid). *J. Bangladesh Acad. Sci.*, 13 (1989) 49-54; *C.A.*, 112 (1990) 15643s.
- 3100 Basova, E.M., Bol'shova, T.A., Ivanov, V.M. and Morozova, N.B.: (Determination of palladium, platinum and rhodium by HPLC after extractive preconcentration). *Zh. Anal. Khim.*, 44 (1989) 680-688; *C.A.*, 111 (1989) 247034b.
- 3101 Beauchemin, D., Siu, K.W.M., McLaren, J.W. and Berman, S.S.: Determination of arsenic species by high-performance liquid chromatography-inductively coupled plasma mass spectrometry. *J. Anal. At. Spectrom.*, 4 (1989) 285-289; *C.A.*, 111 (1989) 247008w.
- 3102 Bol'shova, T.A., Malykhin, A.Yu. and Basova, E.M.: (Separation and quantitative determination of 8-hydroxyquinoline complexes of molybdenum, tungsten, cobalt and chromium by adsorption high-performance liquid chromatography). *Vestn. Mosk. Univ., Ser. 2: Khim.*, 30 (1989) 172-175; *C.A.*, 112 (1990) 15725v.
- 3103 Bruins, J. and Maurer, W.: (Determination of unfixed ammonium in cement and cement mortar used for lining drinking water pipes). *ZKG Int., Ed. B*, 42 (1989) 464-467; *C.A.*, 111 (1989) 247111z.
- 3104 Campos, C.C. and Korkisch, J.: Anion-exchange behavior and separation of transition metals in cyclohexanone-methanol-hydrochloric acid media. *An. Acad. Bras. Cienc.*, 60 (1988) 35-40; *C.A.*, 111 (1989) 224263m.

- 3105 Das, N. and Das, J.: Selective separation of uranium(VI) on a chelating styrene-DVB based resin containing quinaldinic acid amide group. *Indian J. Chem., Sect. A*, 28A (1989) 150-153; C.A., 111 (1989) 246902w.
- 3106 Das, N.R. and Chattopadhyay, P.: Solvent and reversed phase extraction chromatographic separation of molybdenum and tungsten with quaternary ammonium salt reagent, Aliquat 336. *Bull. Chem. Soc. Jpn.*, 62 (1989) 3278-3282; C.A., 112 (1990) 27002v.
- 3107 Dilli, S., Haddad, P.R. and Htoon, A.K.: Further studies of diethyldithiocarbamate complexes by high-performance liquid chromatography. *J. Chromatogr.*, 500 (1990) 313-328.
- 3108 Dixit, C. and Khopkar, S.M.: Reversed phase extraction chromatographic separation of thallium(III) with tributyl phosphate as stationary phase. *Indian J. Chem. Sci.*, 1 (1987) 25-30; C.A., 112 (1990) 47734p.
- 3109 Durupthy, O., Durupthy, A. and Dubusc, M.: (Paper chromatography and electrophoresis of α -amino acids). *Bull. Union Physiciens*, 83 (1989) 622-628; C.A., 112 (1990) 6759g.
- 3110 Flieger, A.: (Distribution of some platinum group metals in the liquid anion-exchanger-Eriochrome Black T system studied by liquid-liquid extraction and extraction chromatography). *Chem. Anal. (Warsaw)*, 33 (1988) 411-417; C.A., 112 (1990) 47731k.
- 3111 Fuechtner, F.: Ion chromatographic determination of anionic impurities in the secondary water system of nuclear power stations. *Isotopenpraxis*, 25 (1989) 415-418; C.A., 111 (1989) 242160e.
- 3112 Gaikwad, A.G. and Khopkar, S.M.: Cation exchange chromatographic separation of scandium in mixed solvents. *J. Indian Counc. Chem.*, 3 (1987) 5-10; C.A., 112 (1990) 29911q.
- 3113 Grossmann, O., Popandopulo, Y.I. and Grazhulene, S.S.: Optimization of the extraction chromatography separation of large amounts of chromium(III) and cobalt(II) by experimental design. *Wiss. Ber.-Akad. Wiss. D.D.R. Zentralinst. Festkoerperphys. Werkstofforsch.*, 38 (1988) 20-27; C.A., 112 (1990) 15641q.
- 3114 Haddad, P.R. and Foley, R.C.: Modelling of cation retention in ion chromatography using fixed-site and dynamically coated ion-exchange columns. *J. Chromatogr.*, 500 (1990) 301-312.
- 3115 Jarvis, N.V.: Small-scale recovery and separation of actinides using tri-*n*-butyl phosphate extraction chromatography. *Solvent Extr. Ion Exch.*, 7 (1989) 1077-1088; C.A., 112 (1990) 64993q.
- 3116 Karandashev, V.K., Kuznetsov, R.A. and Grazulene, S.: (Use of coextraction and extraction suppression in extraction-chromatographic separation of elements). *Zh. Anal. Khim.*, 44 (1989) 624-628; C.A., 111 (1989) 246903x.
- 3117 Kawasaki, N., Ishigami, A., Tanimoto, T. and Tanaka, A.: Determination of iron(III) ion using ion chromatography with electrochemical detection and its application to the assay of the ferroxidase activity of ceruloplasmin. *J. Chromatogr.*, 503 (1990) 237-243.
- 3118 Kim, D.W. and Jung, J.H.: Influence of chelating agents in eluent on the separation factor of lithium isotopes. *J. Radioanal. Nucl. Chem.*, 130 (1989) 63-70; C.A., 111 (1989) 224268s.
- 3119 Kocjan, R. and Przeszlowski, S.: Retention of heavy metals and their separation of silica gel impregnated with a mixture of Aliquat 336 and nitroso-R-salt. *Chem. Anal. (Warsaw)*, 33 (1988) 419-425; C.A., 112 (1990) 15654w.
- 3120 Kozliti, E.A. and Kharitonow, O.V.: Algorithm for the separation of multicomponent systems by elution chromatography. *Zh. Fiz. Khim.*, 63 (1989) 1885-1890; C.A., 112 (1990) 15646v.
- 3121 Lih, J.-J., Yeh, K.-Y. and Liu, C.-Y.: Chromatographic separation by a new chelating resin containing β -hydroxydithiocinnamic acid. *Fresenius' Z. Anal. Chem.*, 336 (1990) 12-15.

- 3122 Malykhin, A.Yu., Bol'shova, T.A., Ermolenko, I.N., Bobrovich, I.B., Nikishkin, I.A. and Polyakova, L.A.: (Simultaneous determination of platinum-group metals by adsorption HPLC). *Zh. Anal. Khim.*, 44 (1989) 886-892; *C.A.*, 111 (1989) 224434t.
- 3123 Mende, A. and Baraniak, L.: (Fine separation of europium and gadolinium by ion exchange at solid strontium sulphate). *Isotopenpraxis*, 25 (1989) 464-466; *C.A.*, 112 (1990) 68806d.
- 3124 Müller, S.R., Simon, W., Widmer, H.M., Grolimund, K., Schomburg, G. and Kolla, P.: Separation of cations by open-tubular column liquid chromatography. *Anal. Chem.*, 61 (1989) 2747-2750.
- 3125 Nesterenko, P.N., Yuferova, I.B. and Kudryavtsev, G.V.: (Silica with chemically bonded amidoxime groups in high-performance liquid chromatography of transition metals). *Zh. Vses. Khim. O-va. im. D.I. Mendeleeva*, 34 (1989) 426-427; *C.A.*, 112 (1990) 47746u.
- 3126 Pikulikova, Z., Muchova, A. and Majer, J.: (Cation-exchange chromatography of metal ions in presence of chelate-forming agents with phosphono functional groups). *Acta Fac. Pharm. Univ. Comenianae*, 41 (1987) 83-109; *C.A.*, 111 (1989) 246894v.
- 3127 Schomburg, G., Kolla, P. and Laeubli, M.W.: Ion chromatography of alkali and alkaline earth metal ions. *Am. Lab. (Fairfield)*, 21 (1989) 92-101; *C.A.*, 112 (1990) 47732m.
- 3128 Si, K., Liu, F., Li, X., Zhao, F. and Tong, S.: (Preconcentration and separation of trace amount of copper(II) and nickel(II) with 4-(2-pyridyl-azo)resorcinol loaded on silica gel). *Beijing Daxue Xuebao, Ziran Kexueban*, 24 (1988) 680-686; *C.A.*, 112 (1990) 29912r.
- 3129 Suszczewski, W. and Rozycki, C.: (Complexometric determination of calcium in strontium and barium salts after extraction chromatographic separation). *Chem. Anal. (Warsaw)*, 33 (1989) 793-799; *C.A.*, 112 (1990) 69000y.
- 3130 Usami, S., Fukami, T., Kinoshita, E., Puri, B.K. and Satake, M.: Column chromatographic preconcentration of palladium with dimethyl glyoxime and acenaphthenequinone dioxime on naphthalene. *Anal. Chim. Acta*, 230 (1990) 17-22.
- 3131 Usami, S., Yamada, S., Puri, B.K. and Satake, M.: Chromatographic preconcentration of copper in environmental and biological samples using 2,4,6-tri(2-pyridyl)-1,3,6-triazinetetraphenylborate-naphthalene adsorbent. *Microchim. Acta*, 1 (1989) 263-270; *C.A.*, 112 (1990) 68927u.
- 3132 Yotsuyanagi, T., Hoshino, H. and Nakano, K.: On-column selective destruction and separation mode high-performance liquid chromatography for ultratrace amounts of metal ions. I. A new use of 2,2'-dihydroxyazobenzene derivatives as the reagents in spectrophotometry coupled with ion-pair reversed-phase HPLC for trace metal determination. *Kenkyu Hokoku-Asahi Garasu Kogyo Gijutsu Shoreikai*, 53 (1988) 317-325; *C.A.*, 112 (1990) 68815f.
- 3133 Yotsuyanagi, T., Hoshino, H. and Uehara, N.: On-column selective destruction and separation mode high-performance liquid chromatography for ultratrace amounts of metal ions II. Determination of aluminium by HPLC-fluorometric system with salicylaldehydebenzoylhydrazone. *Kenkyu Hokoku-Asahi Garasu Kogyo Gijutsu Shoreikai*, 53 (1988) 327-334; *C.A.*, 111 (1989) 246955r.
- 3134 Zhao, Y. and Fu, C.: Pre-column chelation liquid chromatographic separation and determination of trace V, Cu, Co, Pd, Fe and Ni. *Anal. Chim. Acta*, 230 (1990) 23-28.
- 3135 Zolotov, Yu.A., Spivakov, B.Ya., Maryutina, T.A., Baskov, V.L. and Pavlenko, I.V.: Partition countercurrent chromatography in inorganic analysis. *Fresenius' Z. Anal. Chem.*, 335 (1989) 938-944.

For additional information see:

C.A., 111 (1989) 224318h, 224350n, 246806t, 246898z;
112 (1990) 15660v, 15673b, 15727x, 29910p, 68975h, 68987p.

See also 2179, 2221, 2250, 2267, 2831, 2832, 3147, 3149.

38b. Anions

- 3136 Chan, T.L. and Tsai, W.C.: (Ion-chromatography analysis of nitrate in pineapple juice.) *Chung-kuo Nung Yeh Hua Hsueh Hui Chih*, 27 (1989) 187-195; C.A., 111 (1989) 193168c.
- 3137 Dannecker, W. and Selke, K.: Simultanbestimmung organischer und anorganischer Anionen aus verwitterten Natursteinoberflächen mittels Gradienten-Ionen-Chromatographie. *Fresenius' Z. Anal. Chem.*, 335 (1989) 966-970.
- 3138 Ito, K. and Sunahara, H.: Ion chromatography with ultraviolet and amperometric detection for iodide and thiocyanate in concentrated salt solutions. *J. Chromatogr.*, 502 (1990) 121-129.
- 3139 Ivask, J. and Penchuk, Ya.: Application of phenylalanine and tyrosine as eluent in ion chromatography. The effect of eluent concentration and pH on the retention time of inorganic anions. *Esti NSV Tead. Akad. Toim., Keem.*, 38 (1989) 125-129; C.A., 112 (1990) 47736r.
- 3140 Perez-Cerrada, M., Herrero-Villen, M.A. and Maquieira, A.: Sugar-rich food: determination of inorganic anions by ionic chromatography. *Food Chem.*, 34 (1989) 285-294; C.A., 111 (1989) 213456a.
- 3141 Perfetti, G.A., Joe, F.L., Jr. and Diachenko, G.W.: Liquid chromatographic determination of sulfite in grapes and selected grape products. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 903-906.
- 3142 Savage, J.C. and Gould, D.H.: Determination of sulfide in brain tissue and rumen fluid by ion-interaction reversed-phase high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 540-545.
- 3143 Shpigun, O.A., Obrezkov, O.N., Funk, J., Werner, G. and Zolotov, Yu.A.: (Ion-chromatographic analysis for sulphur-containing inorganic anions). *Vestn. Mosk. Univ., Ser. 2: Khim.*, 30 (1989) 273-277; C.A., 112 (1990) 15736z.
- 3144 Tusset, V. and Hancart, J.: Application of ion chromatography in steel industry. *Steel Res.*, 60 (1989) 241-245; C.A., 112 (1990) 29971j.

For additional information see:

C.A., 111 (1989) 224366x, 224406k, 246893u, 247175y, 247176z, 247178b;
112 (1990) 29913s, 68887f.

See also 3135, 3148.

38d. Volatile inorganic compounds

- 3145 Kim, H.-J.: Analysis of nitrogen dioxide in ambient air by ion-exclusion chromatography with electrochemical detection. *J. Chromatogr.*, 503 (1990) 466-470.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 3146 Farjo, K.: Application of the bidimensional ion exchange chromatography for the laboratory preparation of different nitrogen-15 enrichments and depletions. *Isotopenpraxis*, 25 (1989) 439-441; C.A., 112 (1990) 68807e.
- 3147 Kozlittin, E.A., Maslova, G.B., Nazarov, P.P. and Polyakova, N.I.: (Peculiarities of the use of EDTA for isolation of some radionuclides of rare earth elements by a chromatographic method). *Radiokhimiya*, 31 (1989) 98-104; C.A., 111 (1989) 220761z.

- 3148 Tananaev, I.G. and Dzyubenko, V.I.: (Anion-exchange purification of neptunium from plutonium). *Radiokhimiya*, 31 (1989) 104-107; *C.A.*, 111 (1989) 220762e.
- 3149 Van der Walt, T.N. and Coetzee, P.P.: Separation of lead-203 by ion-exchange chromatography on Chelex 100 after production of lead-203 by the $Pb(p,xn)$ ^{203}Bi - ^{203}Pb nuclear reaction. *Talanta*, 36 (1989) 451-455; *C.A.*, 111 (1989) 246905z.

For additional information see:

C.A., 111 (1989) 224267r.

See also 3118, 3123.

Gas Chromatography

1. REVIEWS AND BOOKS

- 943 Adamovics, J.A. (Editor): *Chromatographic Analysis of Pharmaceuticals*. M. Dekker, New York, 1989, 680 p.
- 944 Cortes, H.J. (Editor): *Multidimensional Chromatography. Techniques and Applications*. M. Dekker, New York, 1989, 392 p.
- 945 Descresce, R.P., Lifshitz, M.S., Mazura, A.C. and Tilson, J.E.: *Drug Testing in the Workplace*. Am. Soc. Clin. Pathologists, Washington, 1989.
- 946 Schreiber, H.P. and Lloyd, D.R.: Overview of inverse gas chromatography. *ACS Symp. Ser.*, 391 (1989) 1-10; *C.A.*, 112 (1990) 30028p - a review with 55 refs.
- 947 Yang, F. (Editor): *Microbore Column Chromatography: Unified Approach to Chromatography*. M. Dekker, New York, 1989, 424 p.

See also 975, 998, 1003, 1005, 1025, 1031, 1043, 1047, 1051, 1066, 1067, 1070, 1085, 1114, 1126, 1157, 1278, 1281, 1292, 1308, 1325.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 948 Dose, E.V. and Guiochon, G.: Normalized measure of overlap between non-gaussian chromatographic peaks. *Anal. Chem.*, 62 (1990) 174-181.
- 949 Li, J., Wang, S. and Jun, S.: Models of elution on a plateau method. *J. Chromatogr. Sci.*, 27 (1989) 596-599.
- 950 Meacham, R.I., Buffham, B.A. and Mason, G.: The role of viscosity in sorption-effect chromatography. *J. Chromatogr. Sci.*, 28 (1990) 34-41.

2b. Thermodynamics and theoretical relationships

- 951 Kalinichev, A.I.: (Theoretical description of dynamics of chromatographic peak broadening of the substance at nonlinear adsorption isotherm.) *Zh. Fiz. Khim.*, 63 (1989) 2984-2990.
- 952 Landau, V.V., Zabokritskii, M.P., Sterkhov, N.V. and Chizhkov, V.P.: (Work of partition in chromatography.) *Zh. Fiz. Khim.*, 63 (1989) 2978-2983.
- 953 Lebedev, Yu.Ya.: (Chromatographic systems with extreme characteristics of zone movement.) *Zh. Fiz. Khim.*, 63 (1989) 2991-2996.
- 954 Petrishchev, S.D. and Sakodyskii, K.I.: (Numerical calculation of central moments of any order of magnitude of peaks in gas chromatography.) *Zh. Fiz. Khim.*, 63 (1989) 2696-2701.

2c. Relationship between structure and chromatographic behaviour

- 955 Dowling, J.J., Evans, M.B. and Haken, J.K.: Gas-liquid chromatography in qualitative analysis. XVIII. Study of the dependence of the retention behaviour of polyoxyethylene glycol columns on the nature of the support and phase loading. *J. Chromatogr.*, 500 (1990) 355-365.
- 956 Gerasimenko, V.A. and Nabivach, V.M.: Relationships between gas chromatographic retention indices and molecular structure of aromatic hydrocarbons. *J. Chromatogr.*, 498 (1990) 357-366.

- 957 Hanai, T.: Effect of enthalpy on structure-relation correlation in capillary gas chromatography. *J. High Resolut. Chromatogr.*, 13 (1990) 178-181.
- 958 Naito, K., Sagara, T. and Takei, S.: Adsorption effect on the retention volume of hydrocarbons and dialkyl ethers in gas-liquid chromatography using a polar stationary phase and silica gel support. *J. Chromatogr.*, 503 (1990) 25-32.

See also 962, 1064, 1087.

2d. Measurement of physico-chemical and related values

- 959 Ali, L.H., Sulaiman, S.T. and Al-Tamer, M.Y.: Alumina-supported cobalt-molybdenum oxides as a catalyst for hydrosulfurization of organic sulfides and disulfides (Part-I) using stopped-flow gas chromatography. *Fuel Sci. Technol. Int.*, 7 (1989) 1209-1219; *C.A.*, 112 (1990) 23261f.
- 960 Ara, J. and Colenutt, B.A.: The use of Tenax GC as a porous polymer in gas chromatography. *J. Pure Appl. Sci.*, 8 (1989) 33-37; *C.A.*, 112 (1990) 8196v.
- 961 Cheong, W.J. and Carr, P.W.: Limiting activity coefficients and gas-liquid partition coefficients of alkylbenzenes in hydro-organic solvents. *J. Chromatogr.*, 500 (1990) 215-239.
- 962 Defayes, G., Fritz, D.F., Görner, T., Huber, G., de Reyff, C. and Kovats, E.: Organic solutes in paraffin solvents. Influence of the size of the solvent molecule on solution data. *J. Chromatogr.*, 500 (1990) 139-184.
- 963 Galin, M.: Gas-liquid chromatography study of the thermodynamics of interactions between linear polyethylene and nonpolar and polar solutes. *Polymer*, 30 (1989) 2074-2079; *C.A.*, 112 (1990) 36998b.
- 964 Golovnya, R.V., Grigoryeva, D.N. and Vasilyev, A.V.: Intra- and intermolecular interactions in the first members of homologous series of carbonyl containing compounds in gas chromatography. *J. High Resolut. Chromatogr.*, 13 (1990) 47-51.
- 965 Katsanos, N.A., Karaiskakis, G. and Vassilakos, C.: Adsorption, reaction, and desorption rate constants, measured simultaneously by gas chromatography. *Pure Appl. Chem.*, 61 (1989) 2057-2058; *C.A.*, 112 (1990) 12535x.
- 966 Kemnitz, E., Hass, D. and Kohne, A.: (Gas chromatographic determination of adsorption isotherm of C₁-Freons on chromium(III)-oxide.) *Z. Chem.*, 29 (1989) 348-350; *C.A.*, 112 (1990) 12386z.
- 967 Kovaleva, N.V., Levina, E.F., Nikitin, Yu.S. and Protonina, I.S.: (Investigation of heterogeneous surface of gamma-Fe₂O₃ by gas chromatographic method.) *Zh. Fiz. Khim.*, 63 (1989) 2667-2674.
- 968 Krestov, A.G., Blokhina, S.V., Kopytov, S.V. and Krestov, G.A.: (Specific features of the temperature dependence of the adsorption of *n*-alkenes and *n*-alkanols on nematic reentrant phase.) *Dokl. Akad. Nauk SSSR*, 307 (1989) 395-397.
- 969 Lavrenova, A.S., Zel'venskii, V.Yu., Bulatinskii, K.K. and Sakodynskii, K.I.: (Evaluation of accuracy indexes of procedures for measurements of the content of impurities in organic reagents by gas chromatography with an internal standard.) *Vysokochist. Veshchestva*, No. 5 (1988) 156-162; *C.A.*, 112 (1990) 30074a.
- 970 Ligner, G., Sidgi, M., Jagiello, J., Balard, H. and Papirer, E.: Characterization of specific interactions capacity of solid surfaces by adsorption of alkanes and alkenes. Part II: Adsorption on crystalline silica layer surfaces. *Chromatographia*, 29 (1990) 35-38.
- 971 Ma, F., Chao, B. and Ni, Z.: (Study of thermal stabilities of lanthanone acetates by using gas chromatography.) *Hangzhou Daxue Xuebao, Ziran Kexueban*, 16 (1989) 173-178; *C.A.*, 112 (1990) 90420v.

- 972 Nadiraze, N.A., Andronikashvili, T.G. and Berezkin, V.G.: (Effect of the carrier gas on characteristics of the distribution of hydrocarbon gases in the adsorbent-solid support system.) *Soobshch. Akad. Nauk Gruz. SSR*, 133 (1989) 573-576; *C.A.*, 112 (1990) 84830n.
- 973 Pomaville, R.M. and Poole, C.F.: Gas chromatographic study of the solution thermodynamics of organic solutes in tetraalkylammonium alkanesulfonate and perfluoroalkanesulfonate solvents. *J. Chromatogr.*, 499 (1990) 749-759.
- 974 Poole, S.K. and Poole, C.F.: Experimental protocol for the assessment of solvent strength and selectivity of liquid phases used in gas chromatography. *J. Chromatogr.*, 500 (1990) 329-348.
- 975 Schreiber, H.P.: Surface characterization by inverse gas phase chromatography. *Adv. Org. Coat. Sci. Technol. Ser.*, 11 (1989) 192-204; *C.A.*, 112 (1990) 38127r - a review with 33 refs.
- 976 Wojsz, R.: Adsorption mechanism and active carbon adsorption energy distribution function by gas chromatography. *Pol. J. Chem.*, 63 (1989) 223-231; *C.A.*, 112 (1990) 43342m.
- 977 Wu, F., Jiang, D. and Jiao, Q.: (Study of coke deposits on catalyst by temperature-programmed oxidation-gas chromatography.) *Shiyou Huagong*, 18 (1989) 557-561; *C.A.*, 112 (1990) 43516w.

See also 957.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 978 Duebelbeis, D.O.: A dual column reaction gas chromatographic system for the structural confirmation of chlorinated organics. Avail. *Univ. Microfilms Int.*, Order No. DA8904419, 1988, 128 p.; *C.A.*, 112 (1990) 48054d.
- 979 Hantos, K. and Leisztner, L.: (Control of analytical measurement methods by using the signal/noise ratio.) *Magy. Kem. Foly.*, 95 (1989) 110-114.
- 980 Johnson, G.L., Tipler, A. and Crowshaw, D.: The performance and application of a multidimensional system for gas chromatography. *J. High Resolut. Chromatogr.*, 2 (1990) 130-136.
- 981 PrevotEAU, D.: (A low-cost gas chromatograph.) *Bull. Union Physiciens*, 83 (1989) 701-702; *C.A.*, 112 (1990) 6772f.
- 982 Teply, J. and Pisa, L.: (Injector for gas chromatograph with check valve in mobile phase supply.) *Czech. CS* 254,820 (Cl. G01N30/24), 01 May 1989, Appl. 85/9,554, 19 Dec. 1985; 7 pp.; *C.A.*, 112 (1990) 69210s.
- 983 Zelikman, A.M., Davydenkov, A.K. and Lipavskii, V.N.: (Gas-chromatograph valve.) *U.S.S.R. SU* 1,516,971 (Cl. G01N30/24), 23 Oct. 1989, Appl. 4,368,550, 29 Jan. 1988; *C.A.*, 112 (1990) 48083n.
- 984 Zuder, G. and Weissensteiner, E.: Chromatograph device for gas analysis with temperature control. *Eur. Pat. Appl. EP* 330,650 (Cl. G01N30/30), 30 Aug. 1989, AT Appl. 88/412, 22 Feb. 1988; 7 pp.; *C.A.*, 112 (1990) 69026m.

See also 990, 1038, 1041, 1046, 1048, 1074, 1078, 1086.

3b. Detectors and detection reagents

- 985 Aue, W.A., Siu, K.W.M., Beauchemin, D. and Berman, S.S.: Hypercoulometry of the second kind in electron-capture detection. *J. Chromatogr.*, 500 (1990) 95-138.

- 986 Brina, R. and Pons, S.: Use of narrow gap line microelectrodes as sensitive and species selective gas chromatographic detectors. *Report*, Avail. NTIS, TR-116; Order No. AD-A200424, 1988, 24 p.; C.A., 112 (1990) 68728e.
- 987 Brina, R. and Pons, S.: The use of narrow gap microelectrodes as sensitive and species selective gas chromatographic detectors. *J. Electroanal. Chem. Interfacial Electrochem.*, C.A., 112 (1990) 68768t.
- 988 Brina, R., Pons, S. and Fleischmann, M.: Ultramicroelectrode sensors and detectors. Considerations of the stability, sensitivity, reproducibility, and mechanism of ion transport in gas-phase chromatography and in high-performance liquid-phase chromatography. *Report*, Avail. NTIS, TR-103; Order No. AD-A200306, 1988, 16 p.; C.A., 112 (1990) 68729f.
- 989 Carpio, R.A. and Lindt, E.: Use gas chromatography with HID for process gas impurities. *Semicond. Int.*, 12 (1989) 164-167; C.A., 112 (1990) 68759r.
- 990 Croteau, R.B. and Satterwhite, D.M.: Method for radio-capillary gas chromatography employing a modified oxidation-reduction train and flowthrough detector. *J. Chromatogr.*, 500 (1990) 349-354.
- 991 Fujii, T., Jimba, H. and Arimoto, H.: Mass spectrometric studies on the response mechanism of surface ionization detectors for gas chromatography. *Anal. Chem.*, 62 (1990) 107-111.
- 992 HJ Mohamad, A.: Element selective detection of halogenated compounds in capillary gas chromatography using helium microwave induced plasma spectrometry. Avail. *Univ. Microfilms Int.*, Order No. DA8908475, 1988, 137 p.; C.A., 112 (1990) 29917w.
- 993 Knapp, G., Leitner, E., Michaelis, M., Platzner, B. and Schalk, A.: Element specific GC-detection by plasma atomic emission spectroscopy - a powerful tool in environmental analysis. *Intern. J. Anal. Chem.*, 38 (1990) 369-378.
- 994 Montzka, S.A., Barkley, R.M. and Sievers, R.E.: Catalyzed redox occurring in a chemiluminescence detector used for environmental analysis. *S. Afr. J. Chem.*, 42, No. 2 (1989) 81-86; C.A., 112 (1990) 90918p.
- 995 Patterson, J.E.: Gas chromatographic detector based on enhancement of mercury 253.7-nm photoacoustic signals by hydrocarbons in a hydrogen carrier. *Anal. Chim. Acta*, 226 (1989) 99-107.
- 996 Popp, P., Leonhardt, J.W., Mothes, S. and Oppermann, G.: (Method of detection of gas chromatographically separated substances.) *Ger. (East)* DD 267,802 (Cl. G01N27/66), 10 May 1989, Appl. 309,874, 03 Dec. 1987; 3 pp.; C.A., 112 (1990) 69217z.
- 997 Wada, N.: (Hydrogen flame ionization detector for gas chromatograph.) *Jpn. Kokai Tokkyo Koho* JP 01,221,656 [89,221,656] (Cl. G01N27/62), 05 Sep. 1989, Appl. 88/47,281, 29 Feb. 1988; 4 pp.; C.A., 112 (1990) 90841h.
- 998 Westmoreland, D.G. and Rhodes, G.R.: Analytical techniques for trace organic compounds-II. Detectors for gas chromatography. *Pure Appl. Chem.*, 61 (1989) 1147-1160; C.A., 112 (1990) 69054u - a review with 41 refs.

See also 1052, 1069, 1071, 1075, 1078, 1203, 1321, 1354.

3c. Sorbents, carriers, column and layer performance, packing procedures

- 999 Abe, E. and Wasa, T.: (Thermally stable chiral stationary phases for amino acid enantiomeric analysis by gas chromatography.) *Chem. Express*, 4 (1989) 329-332; C.A., 112 (1990) 48012p.
- 1000 Aichholz, R.: Preparation of glass capillary columns coated with OH-terminated (3,3,3,-trifluoropropyl) methyl polysiloxane (PS 184.5). *J. High Resolut. Chromatogr.*, 13 (1990) 71-73.
- 1001 Alexandr, G.: Preparation of glass capillary columns coated with highly polar cyanoalkyl stationary phases. *J. High Resolut. Chromatogr.*, 13 (1990) 65-68.

- 1002 Bengård, A. and Blomberg, L.: Influence of adsorption at the gas-liquid interface on the determination of gas chromatographic retention indices on open-tubular columns coated with cyanosilicones. *J. Chromatogr.*, 502 (1990) 1-9.
- 1003 Blomberg, L.G.: Current aspects of stationary phase immobilization in open tubular column chromatography. *J. Microcolumn Sep.*, 2 (1990) 62-68 - a review with 95 refs.
- 1004 Bruner, F., Crescentini, G. and Mangani, F.: Gas-liquid-solid chromatography with coated graphitized carbon black. *Pure Appl. Chem.*, 61 (1989) 1997-2000; *C.A.*, 112 (1990) 12314z.
- 1005 Esin, M.S. and Vigdergauz, M.S.: (Use of high-temperature liquid crystals in gas chromatography for separation of polycyclic aromatic hydrocarbons.) *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 32, No. 6 (1989) 3-10; *C.A.*, 112 (1990) 47986x - a review with 51 refs.
- 1006 Grob, K. and Tschuor, R.: Optimal carrier gas velocities at high temperatures in capillary GC. *J. High Resolut. Chromatogr.*, 13 (1990) 193-194.
- 1007 Huang, A., Xu, X., Zhan, P. and Sun, Y.: (Preparation of cross-linking SE-30, SE-54 and cyanoethylmethylpolysiloxane glass capillary column with ozone as free radical initiator.) *Beijing Daxue Xuebao, Ziran Kexueban*, 25 (1989) 398-402; *C.A.*, 112 (1990) 84864b.
- 1008 Keibal, V.L., Nazarova, V.I. and Shcherbakova, K.D.: (An analytical and physicochemical application of packed capillary columns with different samples of graphitized carbon black.) *Vestn. Mosk. Univ., Ser. 2: Khim.*, 30 (1989) 283-252; *C.A.*, 112 (1990) 69142w.
- 1009 Kersten, B.R.: Characterization of the solvent properties of gas chromatographic stationary phases. *Avail. Univ. Microfilms Int.*, Order No. DA8910342, 1988, 382 p.; *C.A.*, 112 (1990) 26164n.
- 1010 Kotel'nikova, T.A. and Ageev, E.P.: (Modification of the surface of Polysorb 1 for gas-chromatographic determination of microimpurities of polar and nonpolar substances.) *Vysokochist. Veshchestva*, No. 5 (1989) 191-196; *C.A.*, 112 (1990) 48015s.
- 1011 Lobanova, S.A., Shcherbakova, O.A., Koifman, O.I. and Burmistrov, V.A.: (Structural selectivity of some polymorphous liquid crystals.) *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.*, 32, No. 6 (1989) 105-108; *C.A.*, 112 (1990) 69140u.
- 1012 Nadiradze, N.A., Andronikashvili, T.G. and Berezkin, V.G.: (Possibility of improving of the separation power in chromatographic analysis.) *Izv. Akad. Nauk Gruz. SSR, Ser. Khim.*, 15 (1989) 125-128; *C.A.*, 112 (1990) 90954x.
- 1013 Nayak, V.S.: New solid adsorbents for the separation of lower hydrocarbons and permanent gases. I. Ammonium tungstophosphate. *J. Chromatogr.*, 498 (1990) 349-356.
- 1014 Pitro, V. and Exnar, P.: (Surface modification of high-silica porous supports for gas chromatography.) *Czech. CS 259,182 (Cl. G01N30/48)*, 31 Mar. 1989, Appl. 86/5,228, 09 Jul. 1986; 7 pp.; *C.A.*, 112 (1990) 48087s.
- 1015 Qin, Y., Shao, L., Sun, K., Li, G. and Shen, Y.: (Study on polysiloxane liquid crystals as the stationary phase for glass capillary gas chromatography.) *Sepu*, 7 (1989) 325-330.
- 1016 Starobinets, G.L., Kulikov, V.I., Vinarskii, V.A. and Zhitkevich, N.V.: (Zinc chloride and its solutions in polyethylene glycol as new liquid stationary phases in gas chromatography.) *Dokl. Akad. Nauk BSSR*, 33 (1989) 819-822; *C.A.*, 112 (1990) 69113n.
- 1017 Takahata, Y., Yakabe, Y., Kato, M., Akiba, M. and Sudo, Y.: (Investigation of a wide diameter open tubular column for GC.) *Bunseki Kagaku*, 39 (1990) 163-166.

- 1018 Takeoka, G., Flath, R.A., Mon, T.R., BATTERY, R.G., Teranishi, R., Güntert, M., Lautamo, R. and Szejtli, J.: Further applications of permethylated β -cyclodextrin capillary gas chromatographic columns. *J. High Resolut. Chromatogr.*, 13 (1990) 202-206.
- 1019 Tarasevich, Yu.I., Zhukova, A.I., Bondarenko, S.V. and Nazarenko, A.V.: (Chromatographic properties of cetylpyridinium montmorillonite produced from its cesium-form.) *Ukr. Khim. Zh. (Russ. ed.)*, 55 (1989) 686-691; *C.A.*, 112 (1990) 48014r.
- 1020 Tien, L. and Fu, R.: (Preparation of wide-bore and thick-film apolar glass capillary columns.) *Sepu*, 7 (1989) 369-370.
- 1021 Todor, D.N., Tanase, A., David, V. and Baiulescu, G.E.: Thermoanalytical study of the complexes of palladium with α, α' -dipyridyl, *p*-nitrosodimethylaniline, and *p*-nitrosodiphenylamine. *Rev. Roum. Chim.*, 34 (1989) 877-881; *C.A.*, 112 (1990) 48010m.
- 1022 Tsitsishvili, G.V., Okropiridze, T.M., Laperashvili, L.Ya., Kvantaliani, L.K. and Chipashvili, D.S.: (Use of heat-stabilized clinoptilolite in gas-adsorption chromatography.) *Izv. Akad. Nauk Gruz. SSR, Ser. Khim.*, 15 (1989) 113-118; *C.A.*, 112 (1990) 90920h.
- 1023 Xiu, Z., Li, H., Wu, C. and Lu, P.: (Development and applications of a cross-linked glass capillary column with 25% cyanoethyl silicone.) *Sepu*, 7 (1989) 295-298.
- 1024 Xiu, Z., Li, H., Yu, W. and Lu, P.: (The preparation reproducibility and application of the methylphenylsilicone crosslinked glass capillary column.) *Sepu*, 7 (1989) 371-373.

See also 947, 960, 1035, 1036, 1045, 1059, 1060, 1073, 1082, 1101, 1146.

3d. Quantitative analysis

- 1025 Gemperline, P.J.: Mixture analysis using factor analysis I: calibration and quantitation. *J. Chemometrics*, 3 (1989) 549-568 - a review with 126 refs.
- 1026 Jin, X., Zeng, Q. and Huang, Y.: (Linear relation between separation factor and relative peak-height response factor of hydrocarbons on FID.) *Fenxi Huaxue*, 17 (1989) 355-357; *C.A.*, 112 (1990) 69080z.
- 1027 Medimagh, M.S. and Bouguerra, M.L.: Analyse en ligne par chromatographie en phase gazeuse des produits d'oxydation catalytique du naphthalene en milieu sels fondus a 425 C. *Analysis*, 7 (1989) 543-546.

See also 1074, 1113, 1299.

3e. Preparative scale chromatography

- 1028 Clark, P.J., Maroulis, P.J., Morris, V.J., Byerley, R.A. and Booker, J.: Automated preparative gas chromatograph. *Eur. Pat. Appl.* EP 330,095 (Cl. G01N30/84), 30 Aug. 1989, US Appl. 160,541, 25 Feb. 1988; 23 pp.; *C.A.*, 112 (1990) 9113c.

See also 1236.

3f. Programmed temperature, pressure, vapors, gradients

- 1029 Crandall, J.A., Bade, R.K. and Cook, R.L.: Current process analysis needs and the design of a programmed temperature process gas chromatograph. *Anal. Div.*, 23 (1989) 85-92; *C.A.*, 112 (1990) 30045s.
- 1030 Yin, H.F. and Sun, Y.L.: The achievement of reproducible temperature programmed retention indices in gas chromatography when using different columns and detectors. *Chromatographia*, 29 (1990) 39-43.

See also 1080.

3g. High performance procedures

- 1031 Hubball, J.: High-temperature gas chromatography: a status report. *LC-GC Int.*, 3, No. 2 (1990) 10-14 - a review with 28 refs.

4. SPECIAL TECHNIQUES

4a. Automation and computerization

- 1032 Doi, T.: (An analysis of chromatograms by personal computer.) *Gakujutsu Kiyo - Kochi Kogyo Koto Senmon Gakko*, 30 (1989) 17-24; *C.A.*, 112 (1990) 15764g.
- 1033 Ghaffari, S.: Microcomputer interfacing programs. *J. Chem. Educ.*, 66 (1989) 837-839.
- 1034 Papas, A.N. and Tougas, T.P.: Accuracy of peak deconvolution algorithms within chromatographic integrators. *Anal. Chem.*, 62 (1990) 234-239.
- 1035 Wu, Z. and Fan, Y.: (Optimization of column chromatographic conditions with a microcomputer - an orthogonal design and variance analysis evaluation.) *Sepu*, 7 (1989) 304-306.
- 1036 Xei, M., Zhou, C. and Wang, R.: (Computer optimization of packed gas-liquid chromatographic separation using modified simplex.) *Sepu*, 7 (1989) 361-364.

See also 1170, 1288, 1294.

4b. Combination of various chromatographic techniques

- 1037 Brorson, T., Sango, C. and Skarping, G.: Evaluation of chromatographic methods for the determination of isocyanates in air. *Intern. J. Anal. Chem.*, 38 (1990) 399-413.
- 1038 Fowles, I.A.: Application of balanced flow high oven temperature-cold on-column injection technique to fast solvent vapor elution and solute focusing in combined HPLC-HRGC. *J. High Resolut. Chromatogr.*, 13 (1990) 213-217.
- 1039 Jirovets, L., Nikiforov, A., Buchbauer, G. and Braun, D.: Correlation of TLC/densitometry-data with capillary GC-data of some main volatile components of essential oils. *Mikrochim. Acta*, 3 (1989) 1-6.
- 1040 Lukkari, P., Hannuksela, J., Mattinen, M.-L., Virolainen, M., Häkkinen, V.M.A. and Riekkola, M.-L.: Separation of metal complexes by on-line coupled LC-GC. *J. High Resolut. Chromatogr.*, 13 (1990) 170-172.
- 1041 Munari, F. and Grob, K.: Coupling HPLC to GC: why? how? with what instrumentation? *J. Chromatogr. Sci.*, 28 (1990) 61-66.
- 1042 Munari, F., Dugo, G. and Cotroneo, A.: Automated on-line HPLC-HRGC with gradient elution and multiple GC transfer applied to the characterization of citrus essential oils. *J. High Resolut. Chromatogr.*, 13 (1990) 56-61.
- 1043 Novotny, M.V.: Recent developments in analytical chromatography. *Science*, 246 (1989) 51.
- 1044 Vreuls, J.J., Cuppen, W.J.G.M., de Jong, G.J. and Brinkman, U.A.T.: Ethyl acetate for the desorption of a liquid chromatographic precolumn on-line into a gas chromatograph. *J. High Resolut. Chromatogr.*, 13 (1990) 157-161.

See also 943, 944, 947, 988, 1003, 1077, 1165, 1253, 1260, 1278, 1312, 1316.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 1045 Bowen, D.V. and Pullen, F.S.: Can aluminum-clad capillary columns be used for gas chromatography/mass spectrometry on a sector mass spectrometer? *Rapid Commun. Mass Spectrom.*, 3, No. 3 (1989) 67-68; *C.A.*, 112 (1990) 68767s.
- 1046 Evershed, R.P. and Prescott, M.C.: Facile connection of aluminum clad capillary GC columns to a magnetic sector mass spectrometer. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 503-506; *C.A.*, 112 (1990) 48053c.
- 1047 Goates, S.R. and Sin, C.H.: Supersonic jet spectroscopy coupled to chromatography for very high resolution chemical analysis. *Appl. Spectrosc. Rev.*, 25 (1989) 82-126; *C.A.*, 112 (1990) 47977v - a review with 103 refs.
- 1048 Gomory, A., Besenyi, I., Czira, G. and Tamas, J.: (The design and use of the microprocessor-controlled interface between a gas chromatograph and mass spectrometer by means of PDP-11-type of computer.) *Meres Autom.*, 37 (1989) 216-220; *C.A.*, 112 (1990) 68784v.
- 1049 Hogan, J.D. and Laude, D.A., Jr.: Suspended trapping procedure for alleviation of space charge effects in gas chromatography/Fourier-transform mass spectrometry. *Anal. Chem.*, 62 (1990) 530-535.
- 1050 Huang, E.C., Jackson, B.J., Markides, K.E. and Lee, M.L.: Application of capillary supercritical fluid chromatography-couple focusing mass spectrometry under negative ion chemical ionization conditions. *J. Microcolumn Sep.*, 2 (1990) 88-96.
- 1051 Smith, R.D., Wright, B.W. and Kalinoski, H.T.: Supercritical fluid chromatography-mass spectrometry (SFC-MS). *Prog. HPLC*, (1989) 111-155; *C.A.*, 112 (1990) 30027n - a review with 80 refs.
- 1052 Traitler, H. and Horman, I.: GC-MS; simultaneous quantification and identification by column effluent splitting. *J. High Resolut. Chromatogr.*, 13 (1990) 197-198.

See also 1173, 1179, 1215, 1258, 1264, 1316.

4e. Functional analysis

- 1053 Mukhamadiev, N.K., Khusainov, K.S., Rozhenko, I.N. and Sakodyskii, K.I.: (Group-structure method in gas-chromatographic identification of organic compounds.) *Dokl. Akad. Nauk UzSSR* No. 4 (1989) 42-44; *C.A.*, 112 (1990) 15783n.

4f. Trace analysis and preseparation techniques

- 1054 Hinshaw, J.V.: Purge-and-trap sampling systems. *LC-GC Int.*, 3, No. 2 (1990) 22-26.
- 1055 Maskarinec, M.P., Johnson, L.H. and Bayne, C.K.: Preparation of reference water and soil samples for performance evaluation of volatile organic analysis. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 823-827.
- 1056 Przyjazny, A. and Staszewski, R.: Evaluation of Polichrom A for preconcentration of volatile organic compounds from the atmosphere. *Anal. Chim. Acta*, 226 (1989) 213-219.
- 1057 Rudling, J.: Improvement of activated carbon for air sampling. *J. Chromatogr.*, 503 (1990) 33-40.
- 1058 Zhou, S.W., Malaiyandi, M. and Benoit, F.M.: An investigation on the concentration efficiencies of some macroreticular and ambersorb resins using radio-labelled organic contaminants commonly encountered in water. *Intern. J. Anal. Chem.*, 38 (1990) 439-471.

See also 993, 998, 1044, 1074, 1076, 1162, 1209, 1214, 1261, 1276, 1277, 1283, 1299.

4g. Separation of enantiomers

- 1059 Armstrong, D.W. and Jin, H.L.: Acylation effects on chiral recognition of racemic amines and alcohols by new polar and non-polar cyclodextrin derivative gas chromatographic phases. *J. Chromatogr.*, 502 (1990) 154-159.
- 1060 Armstrong, D.W., Li, W. and Pitha, J.: Reversing enantioselectivity in capillary gas chromatography with polar and nonpolar cyclodextrin derivative phases. *Anal. Chem.*, 62 (1990) 214-217.
- 1061 Boland, W., Koenig, W.A., Krebber, R. and Mueller, D.G.: Separation of enantiomeric algal pheromones and related hydrocarbons by gas-liquid chromatography on modified cyclodextrins as chiral stationary phases. Biosynthetic relevance of racemic by products. *Helv. Chim. Acta*, 72 (1989) 1288-1292; *C.A.*, 112 (1990) 52256p.
- 1062 Engel, K.H., Albrecht, W. and Heidlas, J.: Enantiodifferentiation of γ - and δ -lactones by gas chromatographic separation of diastereomeric (carbamoyloxy)carboxamide derivatives. *J. Agric. Food Chem.*, 38 (1990) 244-247; *C.A.*, 112 (1990) 55178a.
- 1063 Gasparrini, F., Misiti, D. and Villani, C.: Direct resolution of racemic compounds on chiral microbore columns by sub- and supercritical fluid chromatography. *J. High Resolut. Chromatogr.*, 13 (1990) 182-184.
- 1064 Guo, G. and Yu, A.: (Study on separation of diastereomers with capillary column gas chromatography.) *Youji Huaxue*, 9, No. 1 (1989) 31-37; *C.A.*, 112 (1990) 30026w.
- 1065 König, W.A., Lutz, S., Evers, P. and Knabe, J.: Enantiomer separation of chiral barbiturates and of α -lipoic acid by capillary gas chromatography with modified cyclodextrins as chiral stationary phases. *J. Chromatogr.*, 503 (1990) 256-259.
- 1066 Meyer, V.R.: (Chromatographic separation of enantiomers: The phenomenon of chirality.) *Pharm. Unserer Zeit*, 18 (1989) 140-145; *C.A.*, 112 (1990) 3668r - a review with 12 refs.
- 1067 Oi, N.: (Enantiomer separation by gas chromatography.) *Kikan Kagaku Sosetsu*, No. 6 (1989) 176-185; *C.A.*, 112 (1990) 20521e - a review with 40 refs.
- 1068 Roboz, J., Nieves, E. and Holland, J.F.: Separation and quantification by gas chromatography-mass spectrometry of arabinitol enantiomers to aid the differential diagnosis of disseminated candidiasis. *J. Chromatogr.*, 500 (1990) 413-426.

See also 999, 1104, 1136, 1242, 1243, 1244, 1249, 1252.

4h. Other special techniques

- 1069 Butey, J.L. and Kosman, J.J.: Multielement simulated distillation by capillary GC-MED. *J. Chromatogr. Sci.*, 28 (1990) 19-23.

4i. Supercritical fluid chromatography

- 1070 Bayona Termens, J.M.: (Modern trends in supercritical-fluid chromatography.) *Quim. Anal. (Barcelona)*, 7 (1988) 397-413; *C.A.*, 112 (1990) 47978w - a review with 64 refs.
- 1071 Chang, H.-C.K. and Taylor, L.T.: The performance of electron-capture detection after capillary supercritical fluid chromatography. *J. Chromatogr. Sci.*, 28 (1990) 29-33.
- 1072 Crow, J.A. and Foley, J.P.: Optimization of separations in supercritical fluid chromatography using a modified simplex algorithm and short capillary columns. *Anal. Chem.*, 62 (1990) 378-386.

- 1073 De Weerd, M., Dewaele, C., Verzele, M. and Sandra, P.: Packing material activity in packed capillary column SFC. *J. High Resolut. Chromatogr.*, 13 (1990) 40-46.
- 1074 Hawthorne, S.B., Miller, D.J. and Langenfeld, J.J.: Quantitative analysis using directly coupled supercritical fluid extraction-capillary gas chromatography (SFE-GC) with a conventional split/splitless injection port. *J. Chromatogr. Sci.*, 28 (1990) 2-8.
- 1075 Jinno, K., Mae, H. and Fujimoto, C.: Packed microcolumn SFC coupled with photodiode array UV detector and inductively coupled plasma detector (SFC-photodiode array UV-ICP). *J. High Resolut. Chromatogr.*, 13 (1990) 13-17.
- 1076 King, J.W.: Applications of capillary supercritical fluid chromatography-supercritical fluid extraction to natural products. *J. Chromatogr. Sci.*, 28 (1990) 9-14.
- 1077 Kithinji, J.P., Raynor, M.W., Egia, B., Davies, I.L., Bartle, K.D. and Clifford, A.A.: Analysis of coal tar polycyclic aromatic hydrocarbon LC-fractions by capillary SFC on a liquid crystalline stationary phase. *J. High Resolut. Chromatogr.*, 13 (1990) 27-33.
- 1078 Luffer, D.R., Galante, L.J., David, P.A., Novotny, M. and Hieftje, G.M.: Evaluation of a supercritical fluid chromatograph coupled to a surface-wave-sustained microwave-induced plasma detector. *Report*, Avail. NTIS, INDU/DC/GMH/TR-88-31; Order No. AD-A198728, 1988, 27 p.; *C.A.*, 112 (1990) 69092e.
- 1079 Manninen, P., Riekkola, M.-L., Holm, Y. and Hiltunen, R.: SFC in analysis of aromatic plants. *J. High Resolut. Chromatogr.*, 13 (1990) 167-169.
- 1080 Olesik, S.V. and Pekay, L.A.: A model for quantitatively describing linear velocity programming in capillary SFC. *Chromatographia*, 29 (1990) 69-75.
- 1081 Pawliszyn, J.: Inexpensive fluid delivery system for supercritical fluid extraction. *J. High Resolut. Chromatogr.*, 13 (1990) 199-202.
- 1082 Raynor, M.W., Moulder, R., Davies, I.L., Kithinji, J.P., Clifford, A.A., Bartle, K.D. and Hall, J.E.: Dual capillary column supercritical fluid chromatography. *J. High Resolut. Chromatogr.*, 13 (1990) 22-26.
- 1083 Steuer, W., Baumann, J. and Erni, F.: Separation of ionic drug substances by supercritical fluid chromatography. *J. Chromatogr.*, 500 (1990) 469-479.
- 1084 Veuthey, J.L., Janicot, J.L., Caude, M. and Rosset, R.: Comparative supercritical-fluid chromatographic properties of carbon dioxide and sulphur hexafluoride - ammonia mixtures with packed columns. *J. Chromatogr.*, 499 (1990) 637-643.
- 1085 Wright, B.W. and Smith, R.D.: Capillary supercritical fluid chromatography methods. *Chem. Anal.*, 101 (1989) 111-149; *C.A.*, 112 (1990) 47968t - a review with 75 refs.

See also 1050, 1051, 1063, 1126, 1153, 1162, 1198, 1236, 1316.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5a. Aliphatic hydrocarbons

- 1086 Figura, L.O. and Cammenga, H.K.: (In-line-stripping-cell for gas chromatography. Determination of dissolved gases in aqueous matrices.) *GIT Fachz. Lab.*, 33 (1989) 988-994; *C.A.*, 112 (1990) 25305x.
- 1087 Kuningas, K., Rang, S., Kailas, T., Korol, A. and Dubovik, M.: (Interlaboratory reproducibility of the retention characteristics of *n*-alkenes and *n*-alkynes on capillary columns.) *Eesti NSV Tead. Akad. Toim., Keem.*, 38 (1989) 159-166; *C.A.*, 112 (1990) 48046c.

- 1088 Pomonis, J.G., Hakk, H. and Fatland, C.L.: Synthetic methyl- and dimethylalkanes. Kovats indexes, carbon-13 NMR and mass spectra of some methylpentacosanes and 2,X-dimethylheptacosanes. *J. Chem. Ecol.*, 15 (1989) 2319-2333; *C.A.*, 112 (1990) 76662u.
- 1089 Shang, Z. and Zhang, Y.: (Determination of ethylene chloride in air.) *Lihua Jianyan, Huaxue Fence*, 25 (1989) 42; *C.A.*, 112 (1990) 83141b.
- 1090 Tambieva, N.A. and Vinnikov, Yu.Ya.: (Methane determination in natural waters and wastewaters by equilibrium vapor analysis.) *Gidrokhim. Mater.*, 105 (1989) 41-46; *C.A.*, 112 (1990) 83688k.

See also 995, 1002, 1008, 1012, 1013, 1022, 1026, 1061, 1291, 1294, 1297, 1298, 1329, 1334, 1335, 1338.

5b. Cyclic hydrocarbons

- 1091 Burse, V.W., Korver, M.P., Needham, L.L., Lapeza, C.R., Jr., Boozer, E.L., Head, S.L., Liddle, J.A. and Bayse, D.D.: Gas chromatographic determination of polychlorinated biphenyls (as Aroclor 1254) in serum: collaborative study. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 649-659.
- 1092 Huang, T., Wei, F., Xu, S., Gu, Y. and Qian, S.: (Application of GC/MS in the analysis of diaromatics, polyaromatics and polar compounds in light diesel oil from catalytic cracking.) *Shiyou Huagong*, 18 (1989) 328-333; *C.A.*, 112 (1990) 9610n.
- 1093 Speer, K., Steeg, E., Horstmann, P., Kühn, T. and Montag, A.: Determination and distribution of polycyclic aromatic hydrocarbons in native vegetable oils, smoked fish products, mussels and oysters, and bream from the river Elbe. *J. High Resolut. Chromatogr.*, 2 (1990) 104-105.

See also 961, 1019, 1077, 1082, 1294.

5c. Halogen derivatives

- 1094 Duebelbeis, D.O., Pieczonka, G., Kapila, S., Clevenger, T.E., Yanders, A.F. and Wilson, J.D.: Application of a dual column reaction chromatography system for confirmatory analysis of polychlorinated biphenyl congeners. *Chemosphere*, 19 (1989) 143-148; *C.A.*, 112 (1990) 2137e.
- 1095 Erm, A., Muks, E., Loiveke, I. and Heinvali, M.: (Capillary gas chromatography of methyl-substituted 6-chloro-6-methyl-2(E)-heptenes.) *Eesti NSV Tead. Akad. Toim., Keem.*, 38 (1989) 167-173; *C.A.*, 112 (1990) 90952v.
- 1096 Fuerst, P., Fuerst, C. and Groebel, W.: (Analysis of polychlorinated terpenes in fat-containing foods, for example fish [liver] oils.) *Dtsch. Lebensm.-Rundsch.*, 85 (1989) 273-279; *C.A.*, 112 (1990) 6195v.
- 1097 Kuznetsova, L.V.: (Determination of ethyl chloride in water.) *Gig. Sanit.*, No. 2 (1989) 49-50; *C.A.*, 110 (1989) 218679q.
- 1098 Mori, K., Nakamura, Y., Kaneko, M., Kan, T. and Watanabe, Y.: (Determination of Freons in household aerosol products.) *Eisei Kagaku*, 35 (1989) 347-353; *C.A.*, 112 (1990) 48080j.
- 1099 Patterson, D.G., Jr., Lapeza, C.R., Jr., Barnhart, E.R., Groce, D.F. and Burse, V.W.: Gas chromatographic/mass spectrometric analysis of human serum for non-ortho (coplanar) and ortho substituted polychlorinated biphenyls using isotope-dilution mass spectrometry. *Chemosphere*, 19 (1989) 127-134; *C.A.*, 112 (1990) 17420z.
- 1100 Rastogi, S.C.: A routine gas chromatographic method for the analysis of chlorofluorocarbons in aerosol cans. *Chromatographia*, 29 (1990) 13-15.

- 1101 Zhukova, A.I., Kozlova, V.S., Bondarenko, S.V. and Tarasevich, Yu.I.: (Use of organo-substituted montmorillonite for gas-chromatographic separation of chloro derivatives of toluene.) *Zh. Prikl. Khim. (Leningrad)*, 62 (1989) 1311-1315; C.A., 112 (1990) 55101v.

See also 978, 1055, 1248, 1299, 1338.

5d. Complex hydrocarbon mixtures

- 1102 Lopes Rodrigues, J., Vilhena, M.T. and Raposo, E.J.: (Gas phase chromatographic separation of aliphatic and aromatic hydrocarbons using OV-275 stationary phase at different temperatures.) *Bol. Soc. Port. Quím.*, 35 (1989) 23-24; C.A., 112 (1990) 8960q.
- 1103 Mukhopadhyay, P.K.: Characterization of amorphous and other organic matter types by microscopy and pyrolysis-gas chromatography. *Org. Geochem.*, 14 (1989) 269-284; C.A., 112 (1990) 9544u.

See also 1301, 1307.

6. ALCOHOLS

- 1104 Froehlich, O., Huffer, M. and Schreier, P.: Chirality evaluation of 2-heptanol and Z-4-hepten-2-ol from banana fruit using multidimensional gas chromatography. *Z. Naturforsch., C: Biosci.*, 44 (1989) 555-558; C.A., 112 (1990) 53987q.
- 1105 Ge, D., Wang, J., Guo, X. and He, B.: (A study of separation and measurement of methanol ethanol mixture with a new spherical macroporous polymer NKG-1.) *Sepu*, 7 (1989) 384-385.
- 1106 Piletz, J.E. and Halaris, A.: Gas-liquid chromatographic method for routine detection of plasma sulfo-, gluco-, and free 3-methoxy-4-hydroxyphenylglycol. *Clin. Chim. Acta*, 185 (1989) 165-176.
- 1107 Stolyarov, B.V. and Enikeeva, A.G.: (Chemical identification of oxygen-containing compounds trapped at the outlet of a chromatographic column.) *Vestn. Leningr. Univ., Ser. 4: Fiz., Khim.*, No. 3 (1989) 71-77; C.A., 112 (1990) 54391c.
- 1108 Unterweger, H. and Bandion, F.: (Determination and estimation of 1,3-propenediol content in wine.) *Vinohrad (Bratislava)*, 27 (1989) 210-212; C.A., 112 (1990) 19882s.

See also 1012, 1059, 1137, 1290, 1297, 1323.

7. PHENOLS

- 1109 Stuckenbruck, P. and Aquino Neto, F.R.: Behavior of phenolic compounds related to wood tar in quantitative HRGC with on-column injection. *J. High Resolut. Chromatogr.*, 13 (1990) 210-212.
- 1110 Tachochner, F., Pilz-Mittenburg, W., Benz, T., Brunner, H., Jaeger, W. and Hagenmaier, H.: Determination of chlorophenols in aqueous, solid, and gas samples by GC/ECD and GC/MS. *Z. Wasser Abwasser Forsch.*, 22 (1989) 267-271; C.A., 112 (1990) 9098q.
- 1111 Zubillaga, M.P. and Maerker, G.: Measurement of safrole and isosafrole in ham. *J. Food Sci.*, 54 (1989) 1475-1478; C.A., 112 (1990) 34508t.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8b. Aflatoxins and other mycotoxins

- 1112 Hu, X., Wang, L. and Cao, J.: (Determination of deoxynivalenol and nivalenol in cereals by gas chromatography.) *Huanjing Huaxue*, 8, No. 4 (1989) 1-5; C.A., 112 (1990) 53860t.
- 1113 Pawlosky, R.J., Mirocha, C.J., Wen, Y. and Abbas, H.K.: Use of deuterated internal standards for quantitation of T-2 and HT-2 toxins in human blood by tandem mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 807-812.
- 1114 Perkowski, J.: (Determination of mycotoxins of the trichothecene group by gas chromatography.) *Postępy Nauk Roln.*, 35, No. 4 (1988) 85-95; C.A., 112 (1990) 53796b - a review with 39 refs.

8c. Other compounds with heterocyclic oxygen (including tannins)

- 1115 Alexander, L.R., Maggio, V.L., Gill, J.B., Green, V.E., Turner, W.E., Patterson, D.G., Jr., Green, B.N., Gray, B.W., Guyan, S.A., et al.: Performance of the high sensitivity ion source for PCDD/PCDF analyses in human serum and adipose tissue. *Chemosphere*, 19 (1989) 241-248; C.A., 112 (1990) 17421s.
- 1116 Brown, R.S., Pettit, K. and Jones, P.W.: A GC/GC-MS study of 2,3,7,8-TCDD using a DB-225 column and selected decomposition monitoring as a quantitative tool. *Chemosphere*, 19 (1989) 171-176; C.A., 112 (1990) 2138f.
- 1117 Fuerst, P., Fuerst, C., Meemken, H.A. and Groebel, W.: (The determination of polychlorinated dibenzodioxins and dibenzofurans in breast milk.) *Z. Lebensm.-Unters. Forsch.*, 189 (1989) 338-345; C.A., 112 (1990) 53841n.
- 1118 Gonnord, M.F., Ignatiadis, I., Fraisse, D. and Becchi, M.: Incidence of the column reactivity on the PCDDs and PCDFs GC/MS quantitation. *Chemosphere*, 19 (1989) 189-194; C.A., 112 (1990) 2139g.
- 1119 Grainger, J., Reddy, V.V. and Patterson, D.G., Jr.: Analysis of tetra- through octachlorinated dibenzo-*p*-dioxins by gas chromatography/Fourier transform infrared spectroscopy. *Chemosphere*, 19 (1989) 249-254; C.A., 112 (1990) 50069f.
- 1120 Grochowalski, A., Gorski, L., Sliwka, I. and Lasa, J.: Gas chromatographic determination of polychlorinated dibenzo-*p*-dioxins in wood preservatives. *J. Chromatogr.*, 502 (1990) 160-166.
- 1121 Naikwadi, K.P. and Karasek, F.W.: Use of carbon-13 labelled dioxins mixture for analysis of polychlorinated dibenzo-*p*-dioxins in environmental samples by GC-MS. *Intern. J. Anal. Chem.*, 38 (1990) 329-342.
- 1122 Olie, K., Slot, P.C. and Wever, H.: Decomposition of octachlorodibenzofuran and formation of the HEPTA congeners during GLC analysis. *Chemosphere*, 19 (1989) 103-108; C.A., 112 (1990) 2136d.

See also 1327.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 1123 Frankel, E.N., Hu, M.-L. and Tappel, A.L.: Rapid headspace gas chromatography of hexanal as a measure of lipid peroxidation in biological samples. *Lipids*, 24 (1989) 976-981.
- 1124 Ichinose, T., Miller, M.G. and Shibamoto, T.: Gas chromatographic analysis of free and bound malonaldehyde in rat liver homogenates. *Lipids*, 24 (1989) 895-898.

- 1125 Mudd, A.: Fragmentation of 2-acylcyclohexane-1,3-dione ions and their 4-hydroxy derivatives. Elimination of water *via* an enol. *Rapid Commun. Mass Spectrom.*, 3 (1989) 197-202; *C.A.*, 112 (1990) 51683p.

See also 964, 1107, 1136, 1311, 1342.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 1126 Churms, S.C.: Recent developments in the chromatographic analysis of carbohydrates. *J. Chromatogr.*, 500 (1990) 555-583 - a review with 228 refs.
- 1127 Englmaier, P.: High resolution-GLC of carbohydrates as their dithioacetal-trimethylsilylates and -trifluoroacetates. *J. High Resolut. Chromatogr.*, 2 (1990) 121-125.
- 1128 Serra Bonvehi, J. and Bosch Callis, J.: (Determination of sugars in honey by gas chromatography.) *An. Quim., Ser. B*, 85 (1989) 38-46; *C.A.*, 112 (1990) 53828p.

See also 1068.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 1129 Ahmad, R., Ahmad, I. and Osman, S.M.: *Centratherum ritchiei* seed oil - characterization of vernolic acid. *Fat Sci. Technol.*, 91 (1989) 488-490.
- 1130 Ahmad, V.U., Rasheed, T. and Iqbal, S.: Characterization of some lipid components of *Cocculus hirsutus* by mass spectrometry. *Pak. J. Sci. Ind. Res.*, 32 (1989) 315; *C.A.*, 112 (1990) 4012j.
- 1131 Avorago, A. and Bier, D.M.: Contribution of 3-hydroxyisobutyrate to the measurement of 3-hydroxybutyrate in human plasma: comparison of enzymic and gas-liquid chromatography-mass spectrometry assays in normal and in diabetic subjects. *J. Lipid Res.*, 30 (1989) 1811-1817; *C.A.*, 112 (1990) 51513h.
- 1132 Begerow, J. and Angerer, J.: Improved method for the determination of urinary 2-ethoxyacetic acid by capillary gas chromatography with electron-capture detection. *Fresenius' Z. Anal. Chem.*, 266 (1990) 42-43.
- 1133 Brondz, I., Olsen, I. and Sjoestroem, M.: Gas chromatographic assessment of alcoholized fatty acids from yeasts: a new chemotaxonomic method. *J. Clin. Microbiol.*, 27 (1989) 2815-2819; *C.A.*, 112 (1990) 33305n.
- 1134 De Jong, C. and Badings, H.T.: Determination of free fatty acids in milk and cheese. Procedures for extraction, clean up, and capillary gas chromatographic analysis. *J. High Resolut. Chromatogr.*, 13 (1990) 94-98.
- 1135 Fourie, P.C. and Basson, D.S.: Application of a rapid transesterification method for identification of individual fatty acids by gas chromatography on three different nut oils. *J. Am. Oil Chem. Soc.*, 67 (1990) 18-20.
- 1136 Guichard, E., Kustermann, A. and Mosandl, A.: Chiral flavour compounds from apricots. Distribution of γ -lactone enantiomers and stereodifferentiation of dihydroactinidiolide using multi-dimensional gas chromatography *J. Chromatogr.*, 498 (1990) 396-401.
- 1137 Harvey, D.J.: Identification by gas chromatography/mass spectrometry of long-chain fatty acids and alcohols from hamster meibomian glands using picolinyl and nicotinate derivatives. *Biomed. Chromatogr.*, 3 (1989) 251-254.

- 1138 Kassahun, K., Burton, R. and Abbot, F.S.: Negative ion chemical ionization gas chromatography/mass spectrometry of valproic acid metabolites. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 918-926; *C.A.*, 112 (1990) 30142w.
- 1139 Kawamura, K. and Gagosian, R.B.: Atmospheric transport of soil-derived dicarboxylic acids over the North Pacific Ocean. *Naturwissenschaften*, 77 (1990) 25-27.
- 1140 Kroger, S.: Gas chromatographic determination of 2-ethylhexanoic acid in urine as its pentafluorobenzyl ester. *Analyst (London)*, 114 (1989) 1647-1648.
- 1141 Lee, H.-B., Peart, T.E. and Carron, J.M.: Gas chromatographic and mass spectrometric determination of some resin and fatty acids in pulp mill effluents as their pentafluorobenzyl ester derivatives. *J. Chromatogr.*, 498 (1990) 367-379.
- 1142 Link, W. and Spiteller, G.: Produkte der Dimerisierung ungesättigter Fettsäuren. I: Die Monomerfraktion der Dimerisierung reiner Ölsäure. *Fat Sci. Technol.*, 92 (1990) 19-25.
- 1143 Moss, C.W., Wallace, P.L. and Bosley, G.S.: Identification of monounsaturated fatty acids of *Aerococcus viridans* with dimethyl disulfide derivatives and combined gas chromatography-mass spectrometry. *J. Clin. Microbiol.*, 27 (1989) 2130-2132; *C.A.*, 112 (1990) 3959m.
- 1144 Rao, Y.N., Prasad, R.B.N. and Rao, S.V.: Positional distribution of fatty acids in oilseed phosphatidylcholines and phosphatidylethanolamines. *Fat Sci. Technol.*, 91 (1989) 482-484.
- 1145 Richardson, A.J., Calder, A.G., Stewart, C.S. and Smith, A.: Simultaneous determination of volatile and non-volatile acidic fermentation products of anaerobes by capillary gas chromatography. *Lett. Appl. Microbiol.*, 9 (1989) 5-8; *C.A.*, 112 (1990) 32786h.
- 1146 Ruseva, N.: Rapid gas-chromatographic analysis of the methyl esters of fatty acids on polyacrylonitrile. *Dokl. Bolg. Akad. Nauk*, 42, No. 4 (1989) 63-65; *C.A.*, 112 (1990) 69137y.
- 1147 Saito, S., Miki, T., Ito, H. and Kamoda, M.: (Trace components in sugars. VI. Quantitative determination of formic acid, acetic acid, propionic acid and butyric acid in sugars by GLC.) *Seito Gijutsu Kenkyu Kaishi*, No. 37 (1989) 61-67; *C.A.*, 112 (1990) 8930e.
- 1148 Schuerch, S. and Duebendorfer, G.: (Determination of nitrilotriacetic acid and ethylenediaminetetraacetic acid in water by gas chromatography.) *Mitt. Geb. Lebensmittelunters. Hyg.*, 80 (1989) 324-334; *C.A.*, 112 (1990) 83695k.
- 1149 Socolowsky, S., Hoehne, C. and Sandow, D.: (Gas chromatographic analysis of short-chain fatty acids and lactic acids in microbiological diagnostics.) *Z. Gesamte Hyg. Ihre Grenzgeb.*, 35 (1989) 596-598; *C.A.*, 112 (1990) 19505w.
- 1150 Wijesundera, R.C., Ratnayake, W.M.N. and Ackman, R.G.: Eicosapentaenoic acid geometrical isomer artifacts in heated fish oil esters. *J. Am. Oil Chem. Soc.*, 66 (1989) 1822-1830.

See also 1065, 1107, 1156, 1279, 1280, 1310, 1328, 1342.

11b. Prostaglandins

- 1151 Morrow, J.D., Harris, T.M. and Roberts, II, L.J.: Noncyclooxygenase oxidation formation of a series of novel prostaglandins: Analytical ramifications for measurement of eicosanoids. *Anal. Biochem.*, 184 (1990) 1-10.
- 1152 Scott, P.M., Lombaert, G.A., Pellaers, P., Bacler, S., Kanhere, S.R., Sun, W.F., Lau, P.Y. and Weber, D.: Application of capillary gas chromatography to a survey of wheat for five trichothecenes. *Food Addit. Contam.*, 6 (1989) 489-500; *C.A.*, 112 (1990) 6188v.

11c. Lipids and their constituents

- 1153 Demirbaker, M., and Blomberg, L.S.: Group separation of triacylglycerols on micropacked argentation columns using supercritical media as mobile phases. *J. Chromatogr. Sci.*, 28 (1990) 67-72.
- 1154 Kuksis, A., Myher, J.J. and Sandra, P.: Gas-liquid chromatographic profiling of plasma lipids using high-temperature-polarizable capillary columns. *J. Chromatogr.*, 500 (1990) 427-441.
- 1155 Wong, R. and Hwang, L.S.: (Capillary GC analysis of triglyceride composition of fats and oils.) *Shih P'in K'o Hsueh (Taipei)*, 16, No. 1 (1989) 1-13; *C.A.*, 112 (1990) 20004p.

See also 1157.

12. ORGANIC PEROXIDES

- 1156 Frank, H., Manfred, W. and Thiel, D.: Quantitative determination of fatty acid hydroperoxides *in vivo*. *Basic Life Sci.*, 49 (1988) 175-178; *C.A.*, 112 (1990) 2120u.
- 1157 Van Kuijk, F.J.G.M., Thomas, D.W., Stephens, R.J. and Dratz, E.A.: Detection of lipid peroxidation products in lipids and tissues by gas chromatography-mass spectrometry. *Basic Life Sci.*, 49 (1988) 179-183; *C.A.*, 112 (1990) 3650d - a review with 24 refs.

13. STEROIDS

13a. Pregnane and androstane derivatives

- 1158 Furuta, T., Kusano, K. and Kasuya, Y.: Simultaneous measurements of endogenous and deuterium-labelled tracer variants of androstenedione and testosterone by capillary gas chromatography-mass spectrometry. *J. Chromatogr.*, 525 (1990) 15-23.
- 1159 Garcia-Regueiro, J.A. and Diaz, I.: Evaluation of the contribution of skatole, indole, androstenone and androstenois to boar-taint in back fat of pigs by HPLC and capillary gas chromatography (CGC). *Meat Sci.*, 25 (1989) 307-316; *C.A.*, 112 (1990) 6320g.

See also 1278.

13b. Estrogens

- 1160 Lee, H.-M., Parish, E.J. and Bone, L.W.: The occurrence of estrone and estriol in *Trichostrongylus colubriformis* (Nematoda). *Lipids*, 24 (1989) 903-904.

See also 1278.

13c. Sterols

- 1161 Kojima, M., Ohnishi, M., Ito, S. and Fujino, Y.: Characterization of acylmono-, mono-, di-, tri- and tetraglycosylsterol and saponin in Adzuki bean (*Vigna angularis*) seeds. *Lipids*, 24 (1989) 849-853.
- 1162 Ong, C.P., Ong, H.M., Li, S.F.Y. and Lee, H.K.: The extraction of cholesterol from solid and liquid matrices using supercritical CO₂. *J. Microcolumn Sep.*, 2 (1990) 69-73.

- 1163 Park, P.S.W. and Addis, P.B.: Derivatization of 5 α -cholestane-3 β ,5,6 β -triol into trimethylsilyl ether sterol for GC analysis. *J. Am. Oil Chem. Soc.*, 66 (1989) 1632-1634.
- 1164 Serizawa, S., Nagai, T., Ito, M. and Sato, Y.: Simplified determination of serum cholesterol sulfate by gas-liquid chromatography combined with cyclohexylsilane-bonded phase column purification. *Arch. Dermatol. Res.*, 281 (1989) 411-416; *C.A.*, 112 (1990) 3797g.
- 1165 Walton, C.G., Ratnayake, W.M.N. and Ackman, R.G.: Total sterols in seafoods: Iatroscan TLC/FID versus the Kovacs GLC/FID method. *J. Food Sci.*, 54 (1989) 793-795; *C.A.*, 112 (1990) 34485h.

13d. Bile acids and alcohols

- 1166 Eguchi, T., Miyazaki, H. and Nakayama, F.: Simultaneous determination of keto and non-keto bile acids in human serum by gas chromatography with selected ion monitoring. *J. Chromatogr.*, 525 (1990) 25-42.
- 1167 Iida, T., Itoh, T., Hagiwara, K., Chang, F.C., Goto, J. and Nambara, T.: A method for the simultaneous analysis of unconjugated and glycine-conjugated bile acids by capillary gas-liquid chromatography. *Lipids*, 24 (1989) 1053-1055.
- 1168 Mao, D., Sha, S., Yao, S., Wang, Y. and Dong, C.: (Modified gas-liquid chromatography for determination of serum bile acids.) *Shanghai Dier Yike Daxue Xuebao*, 9 (1989) 239-241; *C.A.*, 112 (1990) 51524n.
- 1169 Stellaard, F., Langelaar, S.A., Kok, R.M. and Jakobs, C.: Determination of plasma bile acids by capillary gas-liquid chromatography-electron capture negative chemical ionization mass fragmentography. *J. Lipid Res.*, 30 (1989) 1647-1652; *C.A.*, 112 (1990) 3996w.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 1170 Craveiro, A.A. and Alencar, J.W.: (Identification of monoterpenes by library search computer programs: integrated spectral approach.) *Quim. Nova*, 11 (1988) 398-401; *C.A.*, 112 (1990) 30072y.
- 1171 Davies, N.W.: Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicone and Carbowax 20M phases. *J. Chromatogr.*, 503 (1990) 1-24.
- 1172 Speer, K.: (16-0-Methylcafestol - a new diterpene in coffee. Methods for its determination in crude and treated coffees.) *Z. Lebensm.-Unters. Forsch.*, 189 (1989) 326-330; *C.A.*, 112 (1990) 75404z.
- 1173 Woidich, A., Nikiforov, A. and Zbiral, E.: The use of N-alkylcarbamates in sample mapping of terpene alcohols. *Mikrochim. Acta*, 3 (1989) 117-124.
- 1174 Zhang, C., Sun, Q. and Nie, C.: (Study on GC/FTIR identification of some monoterpenes.) *Huaxue Xuebao*, 47 (1989) 389-392; *C.A.*, 112 (1990) 69151y.

See also 1018, 1272.

15b. Essential oils

- 1175 Lanças, F.M. and Cavicchioli, M.: Analysis of the essential oils of Brazilian citrus fruits by capillary gas chromatography. *J. High Resolut. Chromatogr.*, 13 (1990) 207-209.

- 1176 Pavlov, L., Bosnjak, N. and Zuber-Horgas, N.: (Gas chromatography in determination of *Mentha piperitae* L. oil quality.) *Prehrambeno-Tehnol. Biotehnol. Rev.*, 26 (1988) 121-125; *C.A.*, 112 (1990) 20036a.
- 1177 Pörschmann, J., Görig, A., Herzschiuh, R., Wettengel, R. and Günther, K.: Analyse und Aufarbeitung von Sulfitterpentinöl. *Chem. Techn. (Leipzig)*, 42 (1990) 27-31.
- 1178 Smit, Z., Karuza-Stojakovic, L. and Pekic, B.: (Proposal methods for determination of essential oil content in alcoholic extracts.) *Farm. Glas.*, 45 (1989) 287-292; *C.A.*, 112 (1990) 25761t.
- 1179 Wu, Q., Qiu, N. and Chen, Y.: (Analysis of Chinese herb preparations by GC/FTIR with Nicolet 55XC FTIR. II. Rapid determination of pharmaceutical essential oils.) *Fenxi Huaxue*, 17 (1989) 697-700; *C.A.*, 112 (1990) 42689t.

See also 1039, 1042, 1079.

16. NITRO AND NITROSO COMPOUNDS

- 1180 James, J.C., Pastel, M.J., Ledoux, K.A. and Baldoni, J.M.: Gas chromatographic/mass spectrometric (GC/MS) analysis of N-nitrosodiisopropanolamine at the nanogram-per-gram level in a gel formulation containing a nonsteroidal antiinflammatory agent. *Pharm. Res.*, 6 (1989) 892-894; *C.A.*, 112 (1990) 11992g.
- 1181 Song, J.: (Determination of mononitrotoluene (MNT) tautomers by gas chromatography.) *Fenxi Ceshi Tongbao*, 8, No. 3 (1989) 59-61; *C.A.*, 112 (1990) 39280x.
- 1182 Spriegelhalder, B., Kubacki, S.J. and Fischer, S.: A method for the determination of tobacco-specific nitrosamines (TSNA), nitrate, and nitrite in tobacco leaves and processed tobacco. *Beitr. Tabakforsch. Int.*, 14 (1989) 135-144; *C.A.*, 112 (1990) 31748k.
- 1183 Wigfield, Y.Y. and McLenaghan, C.C.: Nitrosamines in formulations of deet and EPTC. *Bull. Environ. Contam. Toxicol.*, 44 (1990) 13-18; *C.A.*, 112 (1990) 72168q.

See also 1256, 1300, 1345.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 1184 Bakthavachalam, J., Annan, R.S., Beland, F.A., Vouros, P. and Giese, R.W.: Selection of electrophoric derivatives of 1-aminopyrene and 2-aminofluorene for determination by gas chromatography with electron-capture negative-ion mass spectrometry. *J. Chromatogr.*, 500 (1990) 373-386.
- 1185 Macfarlane, R.G., Macleod, S.C., Midgley, J.M. and Watson, D.G.: Analysis of biogenic amines in bovine retina by gas chromatography-negative ion chemical ionization mass spectrometry. *J. Neurochem.*, 53 (1989) 1731-1736; *C.A.*, 112 (1990) 70118z.

See also 1059, 1305.

17d. *Other amine derivatives and amides (excluding peptides)*

1186 Leis, H.J.: Gas chromatography/negative-ion chemical ionization mass spectrometry of tertiary amines following Hofmann degradation. Application to the characterization of Loperamide. *Rapid Commun. Mass Spectrom.*, 3 (1989) 342-343; *C.A.*, 112 (1990) 130y.

1187 Li, Z. and Li, J.: (Determination of traces of diethylhydroxylamine inhibitor in C₅ fraction by gas chromatography.) *Hecheng Xiangjiao Gongye*, 12 (1989) 408-410; *C.A.*, 112 (1990) 79903j.

See also 1261.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. *Amino acids and their derivatives*

1188 Kapetanovic, I.M., Yonekawa, W.D. and Kupferberg, H.J.: Use of stable isotopes and gas chromatography-mass spectrometry in the study of different pools of neurotransmitter amino acids in brain slices. *J. Chromatogr.*, 500 (1990) 387-394.

See also 999.

19. PROTEINS

19e. *Proteins of blood, serum and blood cells*

See 1348.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. *Purines, pyrimidines, nucleosides, nucleotides*

1189 Gehrke, C.W. and Kuo, K.C.T. (Editors): *Chromatography and Modification of Nucleosides. Part A: Analytical Methods for Major and Modified Nucleosides - HPLC, GC, MS, NMR, UV and FT-IR.* *J. Chromatogr. Libr.*, Vol. 45A, Elsevier, Amsterdam, 1990.

See also 1272.

22. ALKALOIDS

1190 Hayes, M.J., Khemani, L., Bax, M. and Alkalay, D.: Quantitative determination of arecoline in plasma by gas chromatography chemical ionization mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 1005-1009; *C.A.*, 112 (1990) 113v.

1191 Mitsui, T., Hida, M. and Fujimura, Y.: Determination of alkaloidal narcotics by pyrolysis-gas chromatography. *J. Anal. Appl. Pyrolysis*, 17 (1989) 83-89.

- 1192 Ogden, M.W.: Gas chromatographic determination of nicotine in environmental tobacco smoke: collaborative study. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 1002-1006.
- 1193 Parviainen, M.T., Puhakainen, E.V.J., Laatikainen, R., Savolainen, K., Heranen, J. and Barlow, R.D.: Nicotine metabolites in the urine of smokers. *J. Chromatogr.*, 525 (1990) 193-202.
- 1194 Santos, V., Somanathan, R., Rivero Espejel, I.A., Guerrero Beltran, R. and Martinez, M.A.: GC-MS analysis of *Lupinus excubitus* var. *media* (Leguminosae). *Rev. Latinoam. Quim.*, 20 (1989) 86-87; *C.A.*, 112 (1990) 4567u.
- 1195 Yates, S.G., Petroski, R.J. and Powell, R.G.: Analysis of loline alkaloids in endophyte-infected tall fescue by capillary gas chromatography. *J. Agric. Food Chem.*, 38 (1990) 182-185.
- 1196 Zhang, Y., Jacob, P., III and Benowitz, N.L.: Determination of nornicotine in smokers' urine by gas chromatography following reductive alkylation to N'-propylnornicotine. *J. Chromatogr.*, 525 (1990) 349-357.

See also 1274, 1343.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 1197 Blum, W., Ramstein, P. and Eglinton, G.: Coupling of high temperature glass capillary columns to a mass spectrometer. GC/MS analysis of metalloporphyrins from Julia Creek oil shale samples. *J. High Resolut. Chromatogr.*, 13 (1990) 85-93.
- 1198 Wright, B.W. and Smith, R.D.: Supercritical fluid chromatography-mass spectrometry: a potentially useful technique for porphyrin analysis. *Org. Geochem.*, 14 (1989) 227-232; *C.A.*, 112 (1990) 23238d.

23c. Indole derivatives

See 1159.

24. ORGANIC SULPHUR COMPOUNDS

- 1199 Auger, J., Lecomte, C. and Thibout, E.: Leek odor analysis by gas chromatography and identification of the most active substance for the leek moth, *Acrolepiopsis assectella*. *J. Chem. Ecol.*, 15 (1989) 1847-1854; *C.A.*, 112 (1990) 33324t.
- 1200 Berents, A.D., Lakhman, L.N., Barinova, L.A., Larina, M.I., Kugucheva, E.E. and Polkovnikova, N.V.: (Determination of thiophene in pyrolytic benzene.) *Neftepererab. Neftekhim. (Moscow)*, No. 8 (1989) 24-26; *C.A.*, 112 (1990) 48074k.
- 1201 Mueller, B.: (Analytical evaluation of garlic preparations.) *Dtsch. Apoth. Ztg.*, 129 (1989) 2500-2504; *C.A.*, 112 (1990) 42708y.
- 1202 Rao, G.J., Sharma, V.K. and Chandra, H.: Quantitative analysis of thiocyanate in urine by head-space gas chromatography. *Curr. Sci.*, 58 (1989) 1103-1105; *C.A.*, 112 (1990) 71743m.
- 1203 Shearer, R.L., O'Neal, D.L., Rios, R. and Baker, M.D.: Analysis of sulfur compounds by capillary column gas chromatography with sulfur chemiluminescence detection. *J. Chromatogr. Sci.*, 28 (1990) 24-28.
- 1204 Velisek, J., Davidek, J., Michova, J. and Pokorny, J.: Rapid gas chromatographic determination of volatile degradation products of glucosinolates in rapeseed oil. *J. Chromatogr.*, 502 (1990) 167-170.

- 1205 Watts, S.F.: Separation and trapping of atmospheric sulfur compounds. *Environ. Technol. Lett.*, 10 (1989) 777-783; *C.A.*, 112 (1990) 61864n.
- 1206 Yasuhara, A. and Shibamoto, T.: Determination of physical and spectral data on thiazolidines for trace aldehyde analysis. *Agric. Biol. Chem.*, 53 (1989) 2273-2274; *C.A.*, 112 (1990) 6184r.

See also 1312.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCLUDING SUGAR PHOSPHATES)

- 1207 Kataoka, H., Sakiyama, N. and Makita, M.: Determination of 2-aminoethylphosphonic acid and its N-methyl derivative in animal tissues by gas chromatography with flame photometric detection. *Agric. Biol. Chem.*, 53 (1989) 2791-2796; *C.A.*, 112 (1990) 3788e.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 1208 Bloom, N.: Determination of picogram levels of methylmercury by aqueous phase ethylation, followed by cryogenic gas chromatography with cold vapor atomic fluorescence detection. *Can. J. Fish. Aquat. Sci.*, 46 (1989) 1131-1140; *C.A.*, 112 (1990) 42111y.
- 1209 Forsyth, D.S.: Purge and trap extract cleanup for gas chromatography of organolead compounds. *Sci. Total Environ.*, 89 (1989) 291-297; *C.A.*, 112 (1990) 75425g.
- 1210 Novotorova, L.G. and Novotorov, Yu.N.: (Analysis for lower alkyl derivatives of some Group II-VI elements by mass spectrometric and gas chromatographic methods.) *Vysokochist. Veshchestva*, No. 5 (1989) 15-26; *C.A.*, 112 (1990) 15-26.
- 1211 Ohhira, S. and Matsui, H.: Simultaneous determination of butyltin and phenyltin compounds in oysters by capillary gas chromatography. *J. Chromatogr.*, 525 (1990) 105-112.

26b. Boranes, silanes and related non-metallic compounds

- 1212 Krylov, V.A., Loginov, A.V., Shishov, V.N., Murskii, G.L., Filimonov, I.V., Durinov, I.Yu. and Gorshkov, Yu.V.: (Analysis of high-purity dimethyldichlorosilane.) *Vysokochist. Veshchestva*, No. 4 (1989) 217-219; *C.A.*, 112 (1990) 15799x.
- 1213 Zhou, Q., Jiang, L. and Ren, L.: (Determination of selenium in natural waters with 3,5-dibromo-o-phenylenediamine by gas chromatography.) *Huaxue Shiji*, 11 (1989) 295-297; *C.A.*, 112 (1990) 83741r.

26c. Coordination compounds

- 1214 Aggarwal, S.K., Konter, M., Wills, M.R., Savory, J. and Herold, D.A.: Determination of chromium in urine by stable isotope dilution gas chromatography/mass spectrometry using lithium bis(trifluoroethyl)dithiocarbamate as a chelating agent. *Anal. Chem.*, 62 (1990) 111-115.

See also 1040.

27. VITAMINS AND VARIOUS GROWTH REGULATORS (NON-PEPTIDIC)

- 1215 Banno, K., Matsuoka, M., Horimoto, S. and Kato, J.: Simultaneous determination of pantothenic acid and hopantenic acid in biological samples and natural products by gas chromatography-mass fragmentometry. *J. Chromatogr.*, 525 (1990) 255-264.
- 1216 Hall, K.C., Pearce, D.M.E. and Jackson, M.B.: A simplified method for determining 1-aminocyclopropane-1-carboxylic acid (ACC) in plant tissues using a mass selective detector. *Plant Growth Regul.*, 8 (1989) 297-307; *C.A.*, 112 (1990) 51692r.
- 1217 Ying, W., Li, Z., Xue, D., Chen, N. and Chen, Y.: (Micro-determination of vitamin E acetate in traditional Chinese medicine preparations by direct vaporization-gas chromatography.) *Sepu*, 7 (1989) 298-300.
- 1218 Yuji, K.: Analysis of bioactive compounds from plants. *Yukagaku*, 38 (1989) 823-831; *C.A.*, 112 (1990) 51496e.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. Chlorinated insecticides

- 1219 Hue, N.D., Dung, N.X., Huy, D.Q., Olie, K. and Willen, F.D.: (Determination of 2,3,7,8-TCDD by high-resolution gas chromatography.) *Tap Chi Hoa Hoc*, 27, No. 1 (1989) 25-29; *C.A.*, 112 (1990) 54222y.
- 1220 Masse, R., Lalanne, D., Messier, F. and Sylvestre, M.: Characterization of new bacterial transformation products of 1,1,1-trichloro-2,2-bis(4-chlorophenyl)ethane (DDT) by gas chromatography/mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 741-752; *C.A.*, 112 (1990) 50065b.
- 1221 Vinnikov, Yu.Ya., Boeva, L.V. and Ganin, G.I.: (Rapid gas-chromatographic determination of residual amounts of organochlorine pesticides in natural waters with microextraction concentration.) *Gidrokhim. Mater.*, 105 (1989) 20-28; *C.A.*, 112 (1990) 25296v.

See also 1288.

29b. Phosphorus insecticides

- 1222 Li, W.: (Determination of organophosphorus pesticide residues in water by gas chromatography.) *Huanjing Huaxue*, 8, No. 3 (1989) 54-56; *C.A.*, 112 (1990) 11675f.

See also 1288.

29c. Carbamates

See 1062.

29d. Herbicides

- 1223 Ahmad, I. and Crawford, G.: Trace residue analysis of the herbicide chlorsulfuron in soil by gas chromatography-electron capture detection. *J. Agric. Food Chem.*, 38 (1990) 138-141.
- 1224 Huang, Z. and Pen, S.: (Rapid determination of allethrin in mosquito coil and its raw material by gas chromatography.) *Sepu*, 7 (1989) 387-388.

- 1225 Kim, Y.J. and Kim, M.S.: Gas chromatography/mass spectrometry and gas chromatography/tandem mass spectrometry of some s-triazine pesticides. *Bull. Korean Chem. Soc.*, 10 (1989) 196-200; *C.A.*, 112 (1990) 50485p.
- 1226 Liu, H.H., Mattern, G.C., Singer, G.M. and Rosen, J.D.: Determination of daminozide in apples by gas chromatography/chemical ionization mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 984-986.
- 1227 Matter, L.: (Determination of protham and chloroprotham in potatoes by high-resolution capillary gas chromatography with ion-trap detection.) *GIT Fachz. Lab.*, 33 (1989) 1116; *C.A.*, 112 (1990) 53865y.
- 1228 Mattern, G.C., Singer, G.M., Louis, J., Robson, M. and Rosen, J.D.: Determination of linuron in potatoes using capillary column gas chromatography/mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 970-974.

29e. Fungicides

- 1229 Fedotov, A.S. and Metelev, V.V.: (Serocin and borcid toxicity for hydrobionts and method for their detection in water.) *Veterinariya*, No. 11 (1989) 67-68; *C.A.*, 112 (1990) 50291x.
- 1230 Mendes, M.C.S.: Evaluation and confirmation of acetylation-gas-liquid chromatographic method for the determination of triadimenol in foods. *J. Agric. Food Chem.*, 38 (1990) 174-178; *C.A.*, 112 (1990) 53818k.

29f. Other types of pesticides and various agrochemicals

- 1231 Torreti, L. and Simonella, A.: Multidimensional capillary GC of bromopropylate residues in honey: comparison of some identification procedures. *J. High Resolut. Chromatogr.*, 2 (1990) 142-145.
- 1232 Xin, Y., Qi, D., Han, H. and Xue, A.: (Residue determination of Comite in grapes.) *Nanjing Nongye Daxue Xuebao*, 11 (1988) 105-109; *C.A.*, 112 (1990) 75391t.

31. PLASTICS AND THEIR INTERMEDIATES

- 1233 Ding, J.K. and Liu, H.S.: A study of identification of trace rubber residues in marks from rubber-soled shoes and tires by Py-GC. *Forensic Sci. Int.*, 43 (1989) 45-50; *C.A.*, 112 (1990) 31777u.
- 1234 Frenkel, R.S. and Kirillova, T.I.: (Some structural features of nitrile rubbers.) *Kauch. Rezina*, No. 6 (1989) 44-45; *C.A.*, 112 (1990) 8588z.
- 1235 Haken, J.K., Harahap, N. and Burford, R.P.: Analysis of polycarbonate-polydimethylsiloxane block copolymers. *J. Chromatogr.*, 500 (1990) 367-371.
- 1236 Imahashi, T., Yamauchi, Y. and Saito, M.: (Fractionation and identification of additives in polyvinylchloride film by supercritical fluid extraction/semi-preparative supercritical fluid chromatography.) *Bunseki Kagaku*, 39 (1990) 79-85.
- 1237 Singliar, M., Macho, V. and Vojdasova, V.: (Some problems in the analysis of alkylphenylurethanes by gas chromatography.) *Petrochemia*, 28 (1988) 105-110; *C.A.*, 112 (1990) 30066z.
- 1238 Sukop, S., Kolar, L. and Vendra, R.: (Pyrolytic gas chromatography of some acrylate binders for leather finishes. I. Identification of polymers.) *Kozarstvi*, 39, No. 5 (1989) 82-86; *C.A.*, 112 (1990) 38542x.
- 1239 Sun, Y. and Yin, R.: (Analysis of volatiles in post curing of fluororubber by combined headspace gas chromatography-mass spectrometry.) *Hecheng Xiangjiao Gongye*, 12 (1989) 330-333; *C.A.*, 112 (1990) 57965d.
- 1240 Tsuge, S. and Ohtani, H.: (New developments in head space gas chromatography.) *Kagaku (Kyoto)*, 44 (1989) 212-213; *C.A.*, 112 (1990) 22150g.

- 1241 Xu, D., Cheng, G., Cao, S. and Hou, C.: (Curie point pyrolysis gas chromatographic-mass spectrometric identification of thermosetting polymers.) *Gaofenzi Cailiao Kexue Yu Gongcheng*, 4, No. 2 (1988) 28-32; *C.A.*, 112 (1990) 8198x.
- 1242 Xu, Z. and Li, A.: (Quantitative analysis of homopolymer blends and random copolymers of methyl methacrylate with *n*-butyl methacrylate by pyrolysis-gas chromatography.) *Sepu*, 7 (1989) 301-303.
- 1243 Xu, Z.: (Lecture on pyrolysis gas chromatography, Part 3. Applications to the polymer analysis and research.) *Sepu*, 7 (1989) 350-353.
- 1244 Xu, Z.: (Lecture of pyrolysis gas chromatography. Section 3. Application to the macromolecular analysis and investigation.) *Sepu*, 7 (1989) 271-276.
- 1245 Yamaguchi, S., Hirano, J. and Isoda, Y.: (Compositional analysis of acrylic coatings by pyrolysis-gas chromatography using multiple regression analysis.) *Kobunshi Ronbunshu*, 46 (1989) 493-498; *C.A.*, 112 (1990) 22370d.
- 1246 Yoo, Y.J.: Characterization of amine-substituted polydimethylsiloxanes by pyrolysis-high resolution capillary column gas chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA8906355, 1988, 375 p.; *C.A.*, 112 (1990) 78359z.
- 1247 Yoshida, R., Watanabe, Y. and Kazama, M.: (Determination of components in styrene copolymers by pyrolysis GC.) *Bunseki Kagaku*, 39 (1990) 183-187.

See also 963, 975, 1334.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 1248 Lopez-Avila, V., Dodhiwala, N.S., Milanes, J. and Beckert, W.F.: Evaluation of EPA method 8120 for determination of chlorinated hydrocarbons in environmental samples. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 593-602.

See also 943, 945.

32b. Antirheumatics and antiinflammatory drugs

- 1249 Shinohara, Y., Kirii, N., Tamaoki, H., Magara, H. and Baba, S.: Determination of the enantiomers of suprofen and [²H₃] suprofen in plasma by capillary gas chromatography-mass spectrometry. *J. Chromatogr.*, 525 (1990) 93-104.

See also 1180.

32c. Autonomic and cardiovascular drugs

- 1250 Czarny, R.J. and Hornbeck, C.L.: (Quantitation of methamphetamine and amphetamine in urine by capillary GC/MS. Part II. Derivatization with 4-carbethoxyhexafluorobutyl chloride.) *J. Anal. Toxicol.*, 13 (1989) 257-262; *C.A.*, 112 (1990) 17435z.
- 1251 Fuerst, P., Fuerst, C. and Groebel, W.: (Detection of the illegal use of salbutamol in the fattening of calves.) *Dtsch. Lebensm.-Rundsch.*, 85 (1989) 341-344; *C.A.*, 112 (1990) 71774x.
- 1252 Gross, A.S., Borstel, B. and Eichelbaum, M.: Determination of gallopamil in serum by gas chromatography-mass spectrometry. *J. Chromatogr.*, 525 (1990) 183-192.
- 1253 Guinebault, P., Haddouche, A. and Bianchetti, G.: High-performance liquid and gas chromatographic methods for the determination of cicloprolol. *J. Chromatogr.*, 525 (1990) 359-367.

- 1254 Huang, W.H., Lee, A.R. and Sung, K.S.: (Simultaneous analysis of acetaminophen, meprobamate and caffeine in oral muscle relaxant-analgesic tablets by capillary gas chromatography.) *Chung-hua Yao Hsueh Tsa Chih*, 41 (1989) 325-330; *C.A.*, 112 (1990) 42720w.
- 1255 Zavrzhnaya, T.A.: (Gas chromatographic determination of clofelin.) *Farmat-siya*, 38, No. 2 (1989) 33-34; *C.A.*, 112 (1990) 11986h.
- 1256 Zuccaro, P., Zuccaro, S.M., Pacifici, R., Pichini, S. and Boniforti, L.: Determination of isosorbide-5-mononitrate in human serum by gas chromatography-mass spectrometry. *J. Chromatogr.*, 525 (1990) 447-453.

See also 1083.

32d. Central nervous system drugs

- 1257 Basavaiah, K. and Ramappa, P.G.: GLC method for the determination of trifluoromazine hydrochloride. *Indian Drugs*, 26 (1989) 694-696; *C.A.*, 112 (1990) 42736f.
- 1258 Bjorkman, S., Aziz, N., Stein, D. and Stanski, D.R.: Determination of alfentanil in serum by radioimmunoassay or capillary column gas-liquid chromatography. A comparison of the assays. *Acta Pharm. Nord.*, 1 (1989) 211-220; *C.A.*, 112 (1990) 30159g.
- 1259 Davis, P.P., Choma, N., Crews, T. and Fukuda, E.: Quantitative determination of naltiazem in human plasma by gas chromatography/mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 978-982; *C.A.*, 112 (1990) 112u.
- 1260 De Vries, J.X., Rudi, J., Walter-Sack, I. and Conradi, R.: The determination of total and unbound midazolam in human plasma. A comparison of high performance liquid chromatography, gas chromatography and gas chromatography/mass spectrometry. *Biomed. Chromatogr.*, 4 (1990) 28-33.
- 1261 Nagata, K., Hayashi, T. and Naruse, H.: A study on effective derivatization for ultra-micro detection of dopamine by gas chromatography/electron capture negative ion chemical ionization mass spectrometry. *Anal. Lett.*, 22 (1989) 2281-2290.
- 1262 Pillai, G.K. and Rayanam, I.V.: Capillary column gas chromatographic determination of the ester glucuronides of diclofenac in human urine. *Indian Drugs*, 26 (1989) 579; *C.A.*, 112 (1990) 30144y.
- 1263 Siebert, H., Richter, K. and Zachiesche, M.: (Determination of phenobarbital, carbamazepine, primidone, and phenytoin in blood serum.) *Zentralbl. Pharm., Pharmakother., Pharmakother. Laboratoriumsdiagn.*, 128 (1989) 681-683; *C.A.*, 112 (1990) 30176k.
- 1264 Suzuki, S.: Studies on fentanyl and related compounds: II. Spectrometric discrimination of five monomethylated fentanyl isomers by gas chromatography/Fourier transform-infrared spectrometry. *Forensic Sci. Int.*, 43 (1989) 15-19; *C.A.*, 112 (1990) 50107s.

See also 1065, 1254, 1276.

32e. Chemotherapeutics (except cytostatics and antibiotics)

- 1265 Bondarev, V.S., Spiridonov, Yu.Ya. and Shestakov, V.G.: (Determination of chlorsulfuron.) *Khim. Sel'sk. Khoz.*, No. 9 (1989) 51-52; *C.A.*, 112 (1990) 2519f.
- 1266 Kmestak, S. and Dvorak, M.: Capillary gas chromatographic determination of sulphadimidine in pork tissues. *J. Chromatogr.*, 503 (1990) 260-265.
- 1267 Seefeld, F. and Dusing, M.: (Gas chromatographic determination of metalaxyl in plant material.) *Z. Gesamte Hyg. Ihre Grenzgeb.*, 35 (1989) 535-537; *C.A.*, 112 (1990) 6207a.

- 1268 Weber, L.: Rapid monitoring of chloramphenicol in microsamples of plasma by capillary gas chromatography with electron-capture detection. *J. Chromatogr.*, 525 (1990) 454-458.
- 1269 Ye, C. and Yang, Z.: (Determination of ethylparaben in Shengmai Yin oral liquid by gas chromatography.) *Zhongguo Zhongyao Zazhi*, 14 (1989) 539-540; *C.A.*, 112 (1990) 42712v.

32f. Cytostatics

- 1270 Markovic, S. and Kusec, Z.: (Gas chromatographic determination of cyclophosphamide in pharmaceutical preparations.) *Arh. Farm.*, 39 (1989) 23-28; *C.A.*, 112 (1990) 25793e.
- 1271 Weber, H., Marko, V., Kallay, Z., Trnovec, T. and Amlacher, R.: Determination of a cytostatic agent, bendamustine, in plasma using capillary gas chromatography with nitrogen-selective detection. *J. Chromatogr.*, 525 (1990) 459-463.

32g. Other drug categories

- 1272 Cen, L. and He, Z.: (Simultaneous determination of borneol and caffeine in Chinese prepared medicine Tan Kejing by programming temperature gas chromatography.) *Sepu*, 7 (1989) 367-368.
- 1273 Funke, P.T., Ivashkiv, E., Arnold, M.E. and Cohen, A.I.: Determination of pravastatin sodium and its major metabolites in human serum/plasma by capillary gas chromatography/negative ion chemical ionization mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 904-909; *C.A.*, 112 (1990) 30141v.
- 1274 Kompantseva, E.V., Tolkachev, O.N., Botezat-Belyi, Yu.K., Kudrin, S.A., Klochkov, S.V., Markova, O.M., Turbov, V.G., Ladanova, A.A. and Mochalova, N.L.: (Determination of glaucine by gas chromatography.) *Khim.-Farm. Zh.*, 23 (1989) 1116-1120; *C.A.*, 112 (1990) 62737k.

See also 1201, 1217.

32h. Toxicological and forensic applications

- 1275 Rendic, S.: Drug metabolism in identification of drugs misused in sport by gas chromatography-mass spectrometry. *Acta Pharm. Jugosl.*, 39 (1989) 173-180; *C.A.*, 112 (1990) 17437b - a review with 17 refs.
- 1276 Thompson, B.C., Kuzmack, J.M., Law, D.W. and Winslow, J.J.: Copolymeric solid-phase extraction for quantitating drugs of abuse in urine by wide-bore capillary gas chromatography. *LC-GC Int.*, 3, No. 2 (1990) 55-58.
- 1277 Vu-Duc, T., Vernay, A.: Evaluation of sample treatment procedures for the routine identification and determination at nanogram level of O⁶-monoacetylmorphine in urine by capillary GC and dual NPD-FID. *J. High Resolut. Chromatogr.*, 13 (1990) 162-166.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

- 1278 Robards, D. and Towers, P.: Chromatography as a reference technique for the determination of clinically important steroids. *Biomed. Chromatogr.*, 4 (1990) 1-19 - a review with 256 refs.

33b. Complex mixtures and profiling (single compounds by cross-reference only)

- 1279 Liebich, H.M. and Först, C.: Basic profiles of organic acids in urine. *J. Chromatogr.*, 525 (1990) 1-14.
- 1280 Liebich, H.M., Bubeck, J.I., Pickert, A., Wahl, G. and Scheiter, A.: Hippuric acid and 3-carboxy-4-methyl-5-propyl-2-furanpropionic acid in serum and urine. Analytical approaches and clinical relevance in kidney diseases. *J. Chromatogr.*, 500 (1990) 615-627.
- 1281 Yamaguchi, I.: (Gas chromatography/mass spectrometry of biochemical compounds.) *Yukagaku*, 38 (1989) 832-838; *C.A.*, 112 (1990) 18353h - a review with 60 refs.

See also 1091, 1106, 1131, 1137, 1149, 1154, 1196.

34. FOOD ANALYSIS

34b. Complex mixtures (single compounds by cross-reference only)

- 1282 Edwards, C.G. and Beelman, R.B.: Extraction and analysis of volatile compounds in white wines using Amberlite XAD-2 resin and capillary gas chromatography. *J. Agric. Food Chem.*, 38 (1990) 216-220.
- 1283 Hagman, G. and Roeraade, J.: Rapid determination of volatiles in alcoholic beverages by precolumn-backflush gas chromatography. *J. High Chromatogr.*, 2 (1990) 99-103.
- 1284 Horwood, J.F.: Headspace analysis of cheese. *Aust. J. Dairy Technol.*, 44 (1989) 91-96; *C.A.*, 112 (1990) 53863w.
- 1285 Inaba, A., Kubo, Y. and Nakamura, R.: Automated microcomputer system for measurement of oxygen uptake, carbon dioxide output, and ethylene evolution in fruits and vegetables. *Engel Gakkai Zasshi*, 58 (1989) 443-448; *C.A.*, 112 (1990) 34470z.
- 1286 Kawai, I. and Hara, T.: Evaluation of oxidation of vegetable oils by pyrolytic sulfurization gas chromatography. *Bull. Chem. Soc. Jpn.*, 62 (1989) 3007-3009; *C.A.*, 112 (1990) 58654p.
- 1287 Kwan, S. and Sporns, P.: Analysis of bee repellents in honey. *J. Apic. Res.*, 27 (1989) 162-168; *C.A.*, 112 (1990) 53869c.
- 1288 Lipinski, J. and Stan, H.J.: (Computer analysis of pesticides.) *GIT Fachz. Lab.*, 33 (1989) 1089-1092; *C.A.*, 112 (1990) 53864x.
- 1289 Murakami, A.A., Rader, S., Chicoye, E. and Goldstein, H.: Effect of hopping on the headspace volatile composition of beer. *J. Am. Soc. Brew. Chem.*, 47, No. 2 (1989) 35-42; *C.A.*, 111 (1989) 37893e.
- 1290 Nagaichuk, V.V., Kurganova, G.V. and Smirnov, V.A.: (Determination of 2-butanol.) *Pishch. Prom-st. (Moscow)*, No. 11 (1989) 52-53; *C.A.*, 112 (1990) 53621r.
- 1291 Ranfft, K., Gerstl, R. and Koische, G.: (Detection of wax treatment of apples.) *Z. Lebensm.-Unters. Forsch.*, 189 (1989) 331-337; *C.A.*, 112 (1990) 20035z.
- 1292 Rocha, T.E., Baliza, S.V., Soledade, L.E.B. and Guerra Filho, D.: (Gas chromatography of coking byproducts at CSN.) *Metalurgia (Sao Paulo)*, 45 (1989) 557-560; *C.A.*, 112 (1990) 59261v - a review with 14 refs.
- 1293 Tschiersky, H. and Baltes, W.: Curie-point pyrolysis and gas chromatography/mass spectrometry of 5,5'-oxydimethylenebis(2-furfural). *J. Anal. Appl. Pyrolysis*, 17 (1989) 91-93.

See also 1093, 1104, 1108, 1117, 1128, 1134, 1135, 1155, 1172, 1211, 1226, 1227, 1228, 1230, 1231, 1232, 1351.

35. ENVIRONMENTAL ANALYSIS

35a. General papers and reviews

See 993, 1184.

35b. Air pollution (complex mixtures; single compounds by cross-reference only)

- 1294 Einarson, O., Gorczak, J., Lundmark, B.-O. and Palmqvist, U.: Computerized capillary gas chromatographic analysis of hydrocarbon mixtures in workplace air. *J. Chromatogr.*, 498 (1990) 381-395.
- 1295 Li, Z. and Wang, G.: (Determination of acrylonitrile in air by gas chromatographic method.) *Shanghai Huanjing Kexue*, 8, No. 5 (1989) 26-27; *C.A.*, 112 (1990) 83137e.
- 1296 Lin, F. and Chen, C.: (The determination of isokitazine in air by gas chromatography with thermal desorption.) *Shanghai Yike Daxue Xuebao*, 15 (1988) 446-450; *C.A.*, 112 (1990) 11279e.
- 1297 Lipari, F.: Determination of individual hydrocarbons in automobile exhaust from gasoline-, methanol- and variable-fueled vehicles. *J. Chromatogr.*, 503 (1990) 51-68.
- 1298 Mazurek, M.A., Cass, G.R. and Simoneit, B.R.T.: Interpretation of high-resolution gas chromatography and high-resolution gas chromatography/mass spectrometry data acquired from atmospheric organic aerosol samples. *Aerosol Sci. Technol.*, 10 (1989) 408-420; *C.A.*, 112 (1990) 83151e.
- 1299 Müller, S. and Oehme, M.: Analysis of C₁- and C₂-halocarbons in ambient air from remote areas using stainless steel canister sampling, cold trap injection HRGC, and a static calibration technique. *J. High Resolut. Chromatogr.*, 13 (1990) 34-39.
- 1300 Nakanishi, M., Matsuura, A. and Watanabe, I.: (Determination of atmospheric peroxypropionyl nitrate (PPN) with a gas chromatograph-two ECD with a cryogenic collection method.) *Taiki Osen Gakkaishi*, 24 (1989) 152-159; *C.A.*, 112 (1990) 61860h.
- 1301 Teruya, Y., Kakimoto, T. and Izumiya, N.: (Determination of gasoline in automobile exhaust by gas chromatography.) *Kenkyu Hokoku Kurume Kogyo Daigaku*, No. 12 (1988) 43-49; *C.A.*, 112 (1990) 61859q.
- 1302 Zhiznevskii, V.M., Dudykevich, M.A. and Belitskaya, S.I.: (Identification of concentrations of noxious substances in the workplace air at a photopolymer plate plant.) *Gig. Sanit.*, No. 7 (1989) 27-29; *C.A.*, 112 (1990) 24917m.

See also 1037, 1056, 1089, 1110, 1205.

35c. Water pollution (complex mixtures; single compounds by cross-reference only)

- 1303 Bengtsson, G. and Annadotter, H.: Nitrate reduction in a groundwater microcosm determined by nitrogen-15 gas chromatography-mass spectrometry. *Appl. Environ. Microbiol.*, 55 (1989) 2861-2870; *C.A.*, 112 (1990) 18525r.
- 1304 Buchmiller, R.C.: Screening of ground water samples for volatile organic compounds using a portable gas chromatograph. *Ground Water Monit. Rev.*, 9 (1989) 126-130; *C.A.*, 112 (1990) 62258e.
- 1305 Dmitriev, M.T., Zaitseva, N.V., Malkov, V.Yu. and Ulanova, T.S.: (Gas-chromatographic identification of aromatic amines in water reservoirs.) *Gig. Sanit.*, No. 7 (1989) 29-31; *C.A.*, 112 (1990) 11677h.
- 1306 Foerst, C., Stieglitz, L., Roth, W. and Kuhnmuench, S.: Application of headspace analysis and aox-measurement to leachate from hazardous waste landfills. *Chemosphere*, 18 (1989) 1943-1954; *C.A.*, 111 (1989) 102419w.

- 1307 Kurashova, E.Kh., Musaev, I.A., Smirnov, M.V., Simanyuk, R.N., Mikaya, A.E., Ivanov, A.V. and Sanin, P.I.: (Hydrocarbons of Khariag crude oil). *Neftskhimiya*, 29 (1989) 616-627.
- 1308 Lyashenko, V.I. and Gakal, R.K.: (Gas-chromatographic analysis of contaminated water. Methods of sample preparation.) *Gig. Sanit.*, No. 8 (1989) 50-53; *C.A.*, 112 (1990) 62169b - a review with 31 refs.
- 1309 Stout, S.J., DaCunha, A.R., Boyd, J.E. and Devine, J.M.: Confirmation of phorate, terbufos, and their sulfoxides and sulfones in water by capillary gas chromatography/chemical ionization mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 987-991.
- 1310 Tulyupa, F.M. and Golovko, N.V.: (Extractive separation and gas-chromatographic determination of saturated carboxylic acids in waters.) *Khim. Tekhnol. Vody*, 11 (1989) 723-727; *C.A.*, 112 (1990) 42126g.
- 1311 Yasuhara, A., Dennis, K.J. and Shibamoto, T.: Development and validation of new analytical method for acrolein in air. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 749-751.

See also 1054, 1055, 1058, 1086, 1090, 1110, 1139, 1148, 1213, 1229.

35d. Soil pollution (complex mixtures; single compounds by cross-reference only)

- 1312 Brandsteterova, E., Lehotay, J., Garaj, J. and Leclercq, P.A.: High-performance liquid chromatographic determination of dibutyldithiocarbamate degradation products in cucumbers and soil. *J. Chromatogr.*, 502 (1990) 214-220.

See also 1055, 1121, 1219, 1223.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

- 1313 Liu, E., Yang, H. and Qi, Y.: (Structural analysis of polyoxyethylated aliphatic alcohol surfactants by pyrolysis gas chromatography.) *Sepu*, 7 (1989) 364-366.
- 1314 Saito, Y., Sato, T. and Anazawa, I.: The determination of the poly(oxyethylene) oligomer distribution in a non-ionic surfactant by means of thin-layer chromatography. *Bull. Chem. Soc. Jpn.*, 62 (1989) 3709-3710; *C.A.*, 112 (1990) 57165t.

36b. Antioxidants and preservatives

- 1315 Aoyama, M., Tachibana, M., Sasaki, F. and Hasebe, A.: (Analysis of antioxidants by capillary gas chromatography.) *Tokyo-to Suginami-ku Eisei Shikensho Nenpo*, No. 7 (1989) 51-54; *C.A.*, 112 (1990) 75409e.
- 1316 Arpino, P.J., Dilettato, D., Nguyen, K. and Bruchet, A.: Investigation of antioxidants and UV stabilizers from plastics. Part I: comparison of HPLC and SFC; preliminary SFC/MS study. *J. High Resolut. Chromatogr.*, 13 (1990) 5-12.
- 1317 Halkiewicz, J.: Micromethod for the determination of phthalate esters in parenteral solutions and inorganic salts in the preparation of such solutions and for environmental analysis. *Fresenius' Z. Anal. Chem.*, 335 (1989) 998-1001.
- 1318 Helaimia, F. and Messadi, D.: (Determination of linear programmed temperature gas chromatography retention indexes for some plasticizers.) *C. R. Acad. Sci., Ser. 2*, 309 (1989) 37-42; *C.A.*, 112 (1990) 78569t.

See also 1286.

36c. Various technical products

- 1319 Asanova, M. and Abdurazimov, B.K.: (Gas-liquid chromatographic analysis of the products of the synthesis of corn earworm pheromone.) *Rhim. Prir. Soedin.*, No. 4 (1989) 576-579; *C.A.*, 112 (1990) 50489t.
- 1320 Dart, P.J. and McCalley, D.V.: Determination of alkylketene dimer sizing agent products in paper by capillary gas chromatography. *Analyst (London)*, 115 (1990) 13-16.
- 1321 Di Sanzo, F.P.: Determination of oxygenates in gasoline range hydrocarbons by capillary column gas chromatography: a user's experience with an oxygen-specific detector (OFID). *J. Chromatogr. Sci.*, 28 (1990) 73-75.
- 1322 Dubrova, M.A., Shohedrina, M.M., Rudol'fi, T.A., Mindlin, L.O., Nikitina, M.A. and Bel'fer, A.G.: (Characterization of perfume quality by using vapor-phase gas chromatography.) *Fishch. Prom-st. (Moscow)*, No. 10 (1989) 64-65; *C.A.*, 112 (1990) 42214j.
- 1323 Evje, D.M., Figenschou, E. and Langseth-Manrique, K.: Determination of residual solvents and reagents in X-ray contrast media by automatic static headspace capillary gas chromatography. *J. High Resolut. Chromatogr.*, 2 (1990) 116-120.
- 1324 Ezheleva, A.E., Baturina, N.M., Krylov, V.A. and Chernyatin, A.K.: (Gas-chromatographic determination of impurities of methane, hydrogen, carbon monoxide, and carbon dioxide in high-purity monogermene by using selective electron capture sensitization.) *Vysokochist. Veshchestva*, No. 4 (1989) 231-235; *C.A.*, 112 (1990) 15714r.
- 1325 Feeney, M.J. and Jennings, W.J.: Advances in gas chromatographic analysis of thermally generated aromas. *ACS Symp. Ser.*, 409 (1989) 51-60; *C.A.*, 112 (1990) 19985c - a review with 11 refs.
- 1326 Gushchinskaya, M.M. and Posyl'nyi, V.Ya.: (Inverse gas chromatography of anthracites and products of their heat treatment.) *Rhim. Tverd. Topl. (Moscow)*, No. 5 (1989) 25-28; *C.A.*, 112 (1990) 23344k.
- 1327 Herzog, G., Dziwinski, E. and Kiedik, M.: (GC and GC/MS analysis of trioxane synthesis products.) *Przem. Chem.*, 68 (1989) 123-125; *C.A.*, 112 (1990) 20968f.
- 1328 Ivanov, A.F., Teplov, V.M. and Baiburina, E.G.: (Establishment of detection limits of impurities in high-purity (2-methyl-4-chlorophenoxy)acetic acid.) *Vysokochist. Veshchestva*, No. 3 (1989) 217-220; *C.A.*, 112 (1990) 90969f.
- 1329 Janz, S., Birkenfeld, T., Herzsuh, R. and Storch, H.: Gas chromatographic-mass spectrometric investigation of four commercial pristane (2,6,10,14-tetramethylpentadecane) preparations used for plasmacytoma induction in BALB/c mice. *Arch. Geschwulstforsch.*, 59 (1989) 415-422; *C.A.*, 112 (1990) 32003a.
- 1330 Mirzambetov, K. and Ostrovskii, V.: (Rapid methods (in cottonseed analysis).) *Khlopok*, No. 6 (1989) 38-40; *C.A.*, 112 (1990) 53868b.
- 1331 Mitra, G.D., Mohan, G. and Ghosh, S.K.: GC analysis of naphtha using packed column coupled in series. *Fuel Sci. Technol.*, 7, No. 4 (1988) 91-96; *C.A.*, 112 (1990) 9611p.
- 1332 Paryzkova, J. and Novakova, M.: (Method for determination of byproducts in 2,4-bis(4-hydroxy-3,5-bromo)propane by gas chromatography.) *Czech. CS 257,573 (Cl. G01N30/02)*, 15 Mar. 1989, Appl. 86/943, 12 Feb. 1986; 3 pp.; *C.A.*, 112 (1990) 15801s.
- 1333 Penton, Z.: (Composition of an industrial solvent.) *Lab. 2000*, 2, No. 9 (1988) 38-40; *C.A.*, 112 (1990) 48021r.
- 1334 Skrbic, B. and Ristic, M.: (Development of capillary gas chromatography for analysis of commercial heptane used in the production of polypropylene.) *Plast. Guma*, 9 (1989) 115-118; *C.A.*, 112 (1990) 57301j.

- 1335 Sokolova, I.M., Abryutina, N.N., Levshina, A.M., Shepeleva, T.V. and Petrov, A.A.: (Chemical typization of naphthenic crude oils.) *Neftekhimiya*, 29 (1989) 605-609.
- 1336 Timofeeva, A.N., Nikolaeva, N.M., Leont'eva, S.A. and Khodakovskaya, N.A.: (Gas-chromatographic evaluation of the composition of nonionic deemulsifiers.) *Khim. Tekhnol. Topl. Massel*, No. 11 (1989) 39-40; *C.A.*, 112 (1990) 59320p.
- 1337 Voyksner, R.D., Pack, T.W., Haney, C.A., Freeman, H.S. and Hsu, W.N.: Determination of the photodegradation products of Basic Yellow 2 by thermospray high-performance liquid chromatography and gas chromatography/mass spectrometry. *Biomed. Environ. Mass Spectrom.*, 18 (1989) 1079-1086; *C.A.*, 112 (1990) 79424d.

See also 1092, 1098, 1100, 1120, 1147, 1200.

36d. Complex mixtures and unidentified compounds

- 1338 Guo, Y.: (Gas chromatographic analysis of vent gases from the direct synthesis of phenyl chlorosilanes by fluidized-bed.) *Sepu*, 7 (1989) 385-386.
- 1339 Marek, E.M., Kochetova, N.F. and Rotter, O.F.: (Analysis for dimethoxybenzyl chloride by reaction gas-liquid chromatography.) *Khim.-Farm. Zh.*, 23 (1989) 358-361; *C.A.*, 112 (1990) 30077d.
- 1340 Proesch, J., Engewald, W., Thiele, W. and Sack, R.: (Gas-chromatographic examination of the exhaust air from an animal body waste salvage plant.) *Z. Gesamte Hyg. Ihre Grenzgeb.*, 35 (1989) 429-431; *C.A.*, 112 (1990) 11320m.
- 1341 Prokop, H.W., Zhou, H.J., Xu, S.Q., Wu, C.H. and Liu, C.C.: Analysis of the products from the electrochemical fluorination of octanoyl chloride. *J. Fluorine Chem.*, 43 (1989) 277-290; *C.A.*, 112 (1990) 30081a.
- 1342 Xue, L.I.: (The chromatographic analysis for water-formic acid-crotonaldehyde-crotonic acid system.) *Sepu*, 7 (1989) 388-389.

See also 1123.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

See 1130, 1133, 1143, 1145.

38. INORGANIC COMPOUNDS

38b. Anions

- 1344 Buchberger, W. and Huebauer, U.: Selective determination of bromide and iodide in serum and urine by gas chromatography. *Mikrochim. Acta*, 3 (1989) 137-142.
- 1345 Yang, X. and Deng, S.: (GC determination of nitrate nitrogen in water.) *Lihua Jianyan, Huaxue Fence*, 25 (1989) 287-288; *C.A.*, 112 (1990) 83711n.

38c. Permanent and rare gases

- 1346 Bardov, V.G., Kozyarin, I.P., Bryuzgina, T.S. and Kolomiets, E.N.: (Gas chromatographic technique for gaseous hydrogen detection in water.) *Gig. Tr. Prof. Zabol.*, No. 9 (1989) 57; *C.A.*, 112 (1990) 83681c.
- 1347 Guo, C.: (Experimental study on determination of the gas composition of mineral inclusions by the gas chromatography series connection system.) *Changchun Dizhi Xueyuan Xuebao*, 19 (1989) 99-104; *C.A.*, 112 (1990) 90767p.

- 1348 Johansson, M.B. and Wollmer, P.: Measurement of carboxyhemoglobin by spectrophotometry and gas chromatography. *Clin. Physiol.*, 9 (1989) 581-586; *C.A.*, 112 (1990) 71762s.
- 1349 Vasserman, I.I. and Nekhlyudov, I.B.: (Gas-chromatographic determination of trace impurities in high-purity gases with a heat conductivity detector.) *Vysokochist. Veshchestva*, No. 3 (1989) 171-175; *C.A.*, 112 (1990) 90620k.
- 1350 Zhang, Y. and Shang, Z.: (Determination of carbon monoxide in air.) *Lihua Jianyan, Huaxue Fence*, 24 (1988) 292-293; *C.A.*, 112 (1990) 83213b.

See also 995, 1013, 1022, 1324, 1353.

38d. Volatile inorganic compounds

- 1351 Demertzis, P.G. and Kontominas, M.G.: Water binding on egg powders by inverse gas chromatography: water cluster analysis. *Lebensm.-Wiss. Technol.*, 22 (1989) 228-232; *C.A.*, 112 (1990) 75490z.
- 1352 Glavin, G.G., Kvin, V.E., Beilin, Yu.A., Mindlin, B.E. and Nisel'son, L.A.: (Chromatographic-mass spectrometric identification of oxygen-containing impurities in germanium tetrachloride.) *Vysokochist. Veshchestva*, No. 2 (1989) 205-211; *C.A.*, 112 (1990) 29915u.
- 1353 Magnusson, T.: A method for equilibration chamber sampling and gas chromatographic analysis of the soil atmosphere. *Plant Soil*, 120 (1989) 39-47; *C.A.*, 112 (1990) 54214x.
- 1354 Peltonen, K.: The temperature optimization of electron-capture detection in gas chromatography. *LC-GC Int.*, 3, No. 3 (1990) 52-54.
- 1355 Ruan, Y. and Hu, G.: (Determination of carbon disulfide by gas chromatography.) *Zhonghua Laodong Weisheng Zhiyebing zazhi*, 7 (1989) 162-164; *C.A.*, 112 (1990) 47914x.
- 1356 Vorotyntsev, V.M., Drozdov, P.N., Nosyrev, S.A., Krylov, V.A. and Ezheleva, A.E.: (Preconcentration of a phosphine impurity in germane by diffusion through a polymeric membrane.) *Vysokochist. Veshchestva*, (1989) 236-239; *C.A.*, 112 (1990) 15715s.
- 1357 Wang, Y. and Jin, Y.: (Denitrification in the water-soil system - distribution of nitrogen compounds across a clay soil profile and determination of nitrous oxide.) *Huanjing Huaxue*, 8, No. 5 (1989) 13-20. *C.A.*, 112 (1990) 75798f.

See also 1303, 1324, 1338.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

See 990, 1113, 1121, 1303.

Planar Chromatography

1. REVIEWS AND BOOKS

- 462 Bodoga, P. and Marutoiu, C.: (Organic melts - a new type of mobile phase for thin-layer chromatography. I. Characterization and areas of application). *Rev. Chim. (Bucharest)*, 40 (1989) 717-719; *C.A.*, 112 (1990) 84767x - a review with 20 refs.
- 463 Dietzel, G., Kubisiak, H. and Stelzer, H.: A note on: development of 2-dimensional detectors for radio-thin-layer chromatography. *Recent Adv. Thin-Layer Chromatogr.*, [Proc. Chromatogr. Soc. Int. Symp.], 1987 (Publ. 1988) 117-123; *C.A.*, 112 (1990) 29873c - a review with 14 refs.
- 464 Luo, S.: (A survey of the determination of active constituents of Chinese traditional and herbal drugs). *Zhongcaoyao*, 20 (1989) 403-410; *C.A.*, 112 (1990) 42680h - a review without refs.
- 465 Rausch, R.: The importance of thin-layer chromatography as a rapid method for the control of optical purity. *Recent Adv. Thin-Layer Chromatogr.*, [Proc. Chromatogr. Soc. Int. Symp.], 1987 (Publ. 1988) 151-161; *C.A.*, 112 (1990) 47973r - a review with 40 refs.
- 466 Schuetz, H.: (The corrected R_f value. Determination and importance in toxicological analysis). *Pharm. Unserer Zeit*, 18 (1989) 161-168; *C.A.*, 112 (1990) 71697z - a review with 20 refs.
- 467 Wilson, I.D.: Contemporary thin-layer chromatography: an introduction. *Recent Adv. Thin-Layer Chromatogr.*, [Proc. Chromatogr. Soc. Int. Symp.], 1987 (Publ. 1988), 1-8; *C.A.*, 112 (1990) 47972q - a review with 20 refs.

See also 521, 571, 590, 644.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 468 Issaq, H.J. and Seburn, K.E.: Graphic presentation of binary mobile phase optimization in thin layer chromatography. *J. Liq. Chromatogr.*, 12 (1989) 3121-3128.

See also 492, 498.

2b. Thermodynamics and theoretical relationships

- 469 Salov, V.V., Timerbaev, A.R. and Petrukhin, O.M.: Solid-state dissociation of metal chelates and reagents in a silica gel stationary phase in TLC. III. Simulation model. *J. Planar Chromatogr.*, 3 (1990) 73-76.
- 470 Yang, S.: (Deduction of the distribution of solute according to binomial expansion in chromatography). *Huaxue Tongbao*, No. 2 (1989) 47-49; *C.A.*, 112 (1990) 15766j.

2c. Relationship between structure and chromatographic behaviour

- 471 De Voogt, P., van Zijl, G.A., Govers, H. and Brinkman, U.A.T.: Reversed-phase TLC and structure-activity relationships of polycyclic (hetero)aromatic hydrocarbons. *J. Planar Chromatogr.*, 3 (1990) 24-33.

- 472 Dimov, N., Mekenyan, O. and Banov, I.: Mathematical description of the chromatographic behaviour of isosorbide esters separated by thin layer chromatography. *Pharmazie*, 44 (1989) 771-773.

See also 470, 497, 619.

2d. *Measurement of physico-chemical and related values*

- 473 Gankina, E., Kurenbin, O., Kever, J., Malakhova, I., Belenkii, B., Lippmaa, E. and Pekhk, T.: Absolute method of determination of the molecular weight distribution of oligomers based on HPTLC and high resolution NMR. *J. Planar Chromatogr.*, 3 (1990) 68-72.

3. GENERAL TECHNIQUES

3a. *Apparatus and accessories*

- 474 Botz, L., Nyiredy, S. and Sticher, O.: A new solid phase sample application method and device for preparative planar chromatography. *J. Planar Chromatogr.*, 3 (1990) 10-14.
- 475 Campbell, B.D.: Scraping, collecting and eluting apparatus for thin-layer chromatographs. *U.S. Pat.* US 4,863,610 (Cl. 210-658; B01D11-02), 05 Sep. 1989, Appl. 60,402, 11 Jun. 1987; 11 pp.; *C.A.*, 112 (1990) 69213v.
- 476 Doherty, S.J. and Busch, K.L.: An optical CCD camera integrated with a secondary ion mass spectrometer for analysis of planar chromatograms. *J. Planar Chromatogr. - Mod. TLC*, 2 (1989) 149-151; *C.A.*, 112 (1990) 68782t.
- 477 Kochnev, I.V. and Rybal'chenko, I.V.: (Apparatus for scanning thin-layer chromatograms). *U.S.S.R. Pat.* SU 1,509,730 (Cl. G01N30/96), 23 Sep. 1989, Appl. 4,299,571, 06 Jul. 1987; *C.A.*, 111 (1989) 247283g.

3b. *Detectors and detection reagents*

- 478 Bladek, J. and Witkiewicz, Z.: (Chromatographic plate visualization method). *Pol. Pat.* PL 144,560 (Cl. G01N30/90), 30 Jun. 1988, Appl. 256,189, 11 Nov. 1985; 5 pp.; *C.A.*, 111 (1989) 247291h.
- 479 Dochinets, D.I., Petrenko, V.V. and Kubrak, E.A.: (Isatin hydrazone, a detection reagent for 3-ketosteroids). *Khim. Priir. Soedin.*, (1989) 728-729; *C.A.*, 112 (1990) 84294r.
- 480 Ebel, S. and Kang, J.S.: UV/VIS spectra and spectral libraries in TLC/HPTLC. 1. Background correction and normalization. *J. Planar Chromatogr.*, 3 (1990) 42-46.
- 481 Filthuth, H.: A note on digital autoradiograph: a new detector for radiochromatography and eletrophoresis. *Recent Adv. Thin-Layer Chromatogr.*, [Proc. Chromatogr. Soc. Int. Symp.], 1987 (Publ. 1988) 125-129; *C.A.*, 112 (1990) 48001j.
- 482 Imasaka, T., Tanaka, K. and Ishibashi, N.: Thin-layer chromatography with supersonic jet fluorometric detection. *Anal. Chem.*, 62 (1990) 374-378.
- 483 Kamat, S.S. and Kawale, G.B.: Some new observations on Fujiwara reaction. *J. Inst. Chem. (India)*, 60 (1988) 225-226; *C.A.*, 112 (1990) 69157e.
- 484 Kubis, A.J., Viswanadham, S.K., Sharakey, A.G. and Hercules, D.M.: (In situ analysis of compounds separated by thin-layer chromatography by laser mass spectrometry). *Microbeam Anal.*, 24 (1989) 365-365; *C.A.*, 112 (1990) 90882x.

- 485 Ling, B.L., Baeyens, W.R.G., Marysael, H., Stragier, K. and de Moerloose, P.: Influence of stationary phases and fluorescence dipping enhancers on the HPTLC densitometry of SBD-thiol derivatives. *J. Liq. Chromatogr.*, 12 (1989) 3135-3149.
- 486 Prekopa, A. and Liptak, J.: (Chromatographic separation and cultivation of antibiotics for detection). *Hung Teljes Pat.* HU 48,376 (Cl. G01N30/95), 29 May 1989, Appl. 87/3,571, 05 Aug. 1987: 6 pp.; *C.A.*, 112 (1990) 84331a.
- 487 Soper, S.A. and Kuwana, T.: Matrix-isolation luminescence of primary amines derivatized with naphthalenedialdehyde. *Appl. Spectrosc.*, 43 (1989) 883-886; *C.A.*, 112 (1990) 69195r.
- 488 Vasyuk, S.A. and Petrenko, V.A.: (Use of sodium 1,2-naphthoquinone-4-sulfonate as a detecting reagent). *Farmatsiya (Moscow)*, 38, No. 6 (1989) 72-74; *C.A.*, 112 (1990) 62766u.
- 489 Zhao, Y. and Ren, Y.: (Purification and identification of a new color reagent DBC-chlorophosphonazo for rare earths). *Huaxue Shiji*, 11 (1989) 78-81; *C.A.*, 112 (1990) 90575z.

See also 463, 476, 508, 597, 673.

3c. Sorbents, carriers, column and layer performance, packing procedures

- 490 Bold, A., Cruceanu, M., Bara, I., Popovici, E., Vasile, A. and Popa, A.: (Zeolitic molecular sieves used as stationary phases in thin layer chromatography. V. Alkaloid separations). *Rev. Chim. (Bucharest)*, 40 (1989) 137-139; *C.A.*, 112 (1990) 62733f.
- 491 Fischer, W., Hauck, H.E. and Jost, W.: New sorbents for thin-layer chromatography. *Recent Adv. Thin-Layer Chromatogr.*, [Proc. Chromatogr. Soc. Int. Symp.], 1987 (Publ. 1988) 139-149; *C.A.*, 112 (1990) 48011n.
- 492 Issaq, H.J.: Recent advances in multimodal thin-layer chromatography. *TRAC*, 9 (1990) 36-40; *C.A.*, 112 (1990) 115028p.
- 493 Jayaprakasam, R. and Sivakumar, R.: Thin-layer chromatography on cellulose. *Curr. Sci.*, 58 (1989) 980-982; *C.A.*, 112 (1990) 18398b.
- 494 Knyazev, A.S. and Belotserkovskii, G.M.: (Partition properties of active aluminum oxides in thin-layer chromatography). *Zh. Prikl. Khim. (Leningrad)*, 62 (1989) 1708-1711; *C.A.*, 112 (1990) 63526c.
- 495 Ovsepyan, G.S. and Mosikyan, S.M.: Influence of degree of carbonization on the chromatographic activity of amorphous calcium hydrometasilicate). *Arm. Khim. Zh.*, 41 (1988) 741-747; *C.A.*, 111 (1989) 247243u.
- 496 Petrovic, S.M., Perisic-Janjic, N.U., Popovic, M. and Kolarov, L.: Effect of the organic support on chromatographic retention. *J. Planar Chromatogr.*, 3 (1990) 61-64.
- 497 Ruane, R.J. and Wilson, I.D.: A note on: a preliminary investigation of the influence of the stationary phase in ion-pair thin-layer chromatography. *Recent Adv. Thin-Layer Chromatogr.*, [Proc. Chromatogr. Soc. Int. Symp.], 1987 (Publ. 1988) 171-176; *C.A.*, 112 (1990) 48035u.

See also 468, 630, 636, 702, 705, 706, 707.

3d. Quantitative analysis

See 507, 511, 693.

3e. Preparative scale chromatography

See 474, 478, 509.

3f. Programmed temperature, pressure, vapors, gradients

- 498 Touchstone, J.C., Sheikh, S.U., Fang, X.G., Alvarez, J.G. and Levin, S.S.: Subambient temperature in planar chromatography. *J. Liq. Chromatogr.*, 12 (1989) 3193-3199.

See also 500, 606, 630.

3g. High performance procedures

See 597, 693.

4. SPECIAL TECHNIQUES

4a. Automation and computerization

- 499 Gocan, S. and Liteanu, V.: A study on some performance indices used in thin layer chromatography. *Stud Univ. Babes-Bolyai, Chem.*, 33 (1988) 82-86; *C.A.*, 112 (1990) 90894c.
- 500 Markowski, W.: Computer-assisted selection of the optimum gradient programme in thin-layer chromatography. *J. Chromatogr.*, 485 (1989) 517-532.
- 501 Palamareva, M. and Kavrukova, I.: Microcomputer aided choice of mobile phases for thin-layer chromatographic separation of some E- and Z-oxazolones and related cinnamates on silica. *Izv. Khim.*, 21 (1988) 218-225; *C.A.*, 111 (1989) 247239x.
- 502 Qin-Sun, W. and Heng-Yan, W.: Computer-assisted comprehensive optimization of mobile phase selectivity in HPTLC. *J. Planar Chromatogr.*, 3 (1990) 15-19.

See also 556, 630, 673.

4b. Combination of various chromatographic techniques

- 503 Brenner, M., Moirandat, C. and Schlimme, R.: High-efficiency liquid extractor for isolation of a desired material from complex organic mixtures. *J. Chromatogr.*, 499 (1990) 721-748.
- 504 Nyiredy, S., Dallenbach-Toelke, K., Zogg, G.C. and Sticher, O.: Strategies of mobile phase transfer from thin-layer to medium-pressure liquid chromatography with silica as the stationary phase. *J. Chromatogr.*, 499 (1990) 453-462.

See also 519.

4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 505 Harada, K.: (Reconcentration of developed spot on TLC plate for TLC/mass spectrometry). *Iyo Masu Kenkyukai Koenshu*, 13 (1988) 107-110; *C.A.*, 112 (1990) 15765h.

See also 473.

4g. Separation of enantiomers

- 506 Duncan, J.D. and Armstrong, D.W.: Chiral mobile phase additives in reversed-phase TLC. *J. Planar Chromatogr.*, 3 (1990) 65-67.

See also 592.

4h. Other special techniques

- 507 Sarbu, C. and Marutoiu, C.: (Quantitative calculation of the information from the separation [data] and color of zones in thin-layer chromatography). *Rev. Chim. (Bucharest)*, 40 (1989) 140-143; *C.A.*, 112 (1990) 48030t.
- 508 Spurway, T.D., Wilson, I.D., Ruane, R.J. and Warrander, A.: OPLC of ^{14}C -radio-labelled paracetamol metabolites and beta-blockers with on- and off-line radioactivity detection and off-line identification using ^1H NMR. *J. Planar Chromatogr.*, 3 (1990) 51-54.
- 509 Traitler, H. and Studer, A.: Anticircular planar chromatography: analytical and preparative aspects. *Recent Adv. Thin-Layer Chromatogr.*, [Proc. Chromatogr. Soc. Int. Symp.], 1987 (Publ. 1988) 67-74; *C.A.*, 112 (1990) 48034x.

See also 547, 607.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5b. Cyclic hydrocarbons

- 510 Rod'ko, I.Ya. and Tarasova, N.P.: (Methods for determination of components of an impurity of polyaromatic hydrocarbons in products from electrothermal production of yellow phosphorus). *Izv. Vyssh. Uchbn. Zaved., Khim. Khim. Tekhnol.*, 32, No. 5 (1989) 48-50; *C.A.*, 112 (1990) 48075m.

See also 471, 697.

6. ALCOHOLS

- 511 Sherma, J. and Brubaker, K.: Quantitative thin-layer chromatographic determination of cinnamyl alcohol in perfumes. *J. Planar Chromatogr. - Mod. TLC*, 2 (1989) 392-393; *C.A.*, 112 (1990) 83837h.

7. PHENOLS

- 512 Rozylo, J.K. and Kolodziejczyk, H.: Application of TLC for the investigation of separation of some isomeric naphthalene derivatives. *J. Planar Chromatogr. - Mod. TLC*, 2 (1989) 161-162; *C.A.*, 112 (1990) 90945v.
- 513 Sayyah, S.M., Saad, S.M. and Elwy, E.: Characterization of resins prepared by polycondensation of Egyptian bagasse pith and bisphenol A. Part I. Physico-chemical properties of the molded shapes. *Acta Polym.*, 40 (1989) 667-672; *C.A.*, 111 (1989) 233805s.

See also 690, 695.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. *Flavonoids*

- 514 Fischer, D., Ebenau-Jehle, C. and Grisebach, H.: Phytoalexin synthesis and soybean: purification and characterization of NADPH:2'-hydroxydaidzein oxidoreductase from elicitor-challenged soybean cell cultures. *Arch. Biochem. Biophys.*, 276 (1990) 390-395.
- 515 Heimler, D., Mittempergher, L., Buzzini, P. and Boddi, V.: Quantitative HPTLC separation of flavonoid glycosides in the taxonomy of elm (*Ulmus* spp.). *Chromatographia*, 29 (1990) 16-20.

See also 686.

8b. *Aflatoxins and other mycotoxins*

- 516 Fulqueira, C.L. and de Bracalenti, B.J.C.: (Rapid and economical method for detecting aflatoxins and zearalenone in feed). *Rev. Microbiol.*, 20 (1989) 210-214; *C.A.*, 112 (1990) 97125m.

8c. *Other compounds with heterocyclic oxygen (including tannins)*

- 517 Borges, M.F.M. and Pinto, M.M.M.: Isocratic high performance liquid chromatography separation of esculetin and *cis/trans* isomers of caffeic acid. *J. Lig. Chromatogr.*, 12 (1989) 2345-2354.

See also 471, 504.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 518 Evstratova, K.I., Levitskaya, T.N., Malofis, S.L. and Malinovskaya, V.A.: (Method for control of vanillin production using thin-layer chromatography and spectrometry). *Zavod. Lab.*, 55 (1989) 11-15; *C.A.*, 112 (1990) 53817j.
- 519 Miksik, I., Struzinsky, R., Macek, K. and Deyl, Z.: Liquid column chromatography of oxo compounds as their 2,4-dinitrophenylhydrazones with special reference to glucose-glycin Maillard reaction products. *J. Chromatogr.*, 500 (1990) 607-613.

See also 504, 513, 688.

10. CARBOHYDRATES

10a. *Mono and oligosaccharides. Structural studies*

- 520 Aoki, T., Araki, T. and Kitamikado, M.: Purification and characterization of a novel β -agarase from *Vibrio* sp. AP-2. *Eur. J. Biochem.*, 187 (1990) 461-465.
- 521 Churms, S.C.: Recent developments in the chromatographic analysis of carbohydrates. (Review.) *J. Chromatogr.*, 500 (1990) 555-583.
- 522 Debeire, P., Priem, B., Strecker, G. and Vignon, M.: Purification and properties of an endo-1,4-xylanase excreted by a hydrolytic thermophilic anaerobe, *Clostridium thermolacticum*. A proposal for its action mechanism on larchwood 4-O-methylglucuronoxylan. *Eur. J. Biochem.*, 187 (1990) 573-580.

- 523 Takahashi, Y., Sakaguchi, F., Morimoto, K., Hashimoto, K., Funaba, T. and Hayauchi, Y.: (Physicochemical properties and stability of a carbose). *Iyakuhin Kenkyu*, 20 (1989) 763-783; *C.A.*, 112 (1990) 25451s.
- 524 Takeuchi, Y., Matsumoto, T., Ogata, E. and Shishiba, Y.: Isolation and characterization of proteoglycans synthesized by mouse osteoblastic cells in culture during the mineralization process. *Biochem. J.*, 266 (1990) 15-24.
- 525 Tulsiani, D.R.P., Coleman, V.D. and Touster, O.: Asparagine-linked glycoprotein biosynthesis in rat brain: identification of glucosidase I, glucosidase II, and an endomannosidase (glucosyl mannosidase). *Arch. Biochem. Biophys.*, 277 (1990) 114-121.

See also 472, 535, 682.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 526 Ioned, T.: Separation of homologues of methyl ester and 3-O-acetyl methyl ester derivatives of the corynomycolic acid fraction from *Corynebacterium pseudotuberculosis*. *J. Chromatogr.*, 481 (1989) 411-415.
- 527 Kawanishi, K., Yasufuku, J., Ishikawa, A. and Hashimoto, Y.: Long-chain alkyl ferulates in three varieties of *Ipomoea batatas* (L.) Lam. *J. Agric. Food Chem.*, 38 (1990) 105-108.
- 528 Mokhov, A.A., Leont'eva, L.B., Bardin, V.V., Krauklish, I.V. and Tselinskii, I.V.: (Method for determination of *o*-chlorobenzoic acid by paper chromatography). *U.S.S.R. Pat.* SU 1,509,728 (Cl. G01N30/90), 23 Sep. 1989, Appl. 4,359,124, 08 Jan. 1988; *C.A.*, 111 (1989) 247296p.
- 529 Sok, D.-E. and Kim, M.R.: Enzymatic formation of 9,16-dihydro(pero)xyoctadecatrienoic acid isomers from α -linolenic acid. *Arch. Biochem. Biophys.*, 277 (1990) 86-93.

See also 517, 697.

11b. Prostaglandins

See 537.

11c. Lipids and their constituents

- 530 Alvarez, J.G., Touchstone, J.C., Storey, B.T. and Grob, R.L.: Determination of sphingolipid sphingoid bases by HPTLC-fluorescence spectrodensitometry. *J. Liq. Chromatogr.*, 12 (1989) 3115-3120.
- 531 Asmis, R. and Jörg, A.: Calcium-ionophore-induced formation of platelet-activating factor and leukotrienes by horse eosinophils: a comparative study. *Eur. J. Biochem.*, 187 (1990) 475-480.
- 532 Battini, M.L., Careri, M., Casoli, A., Mangia, A. and Lugari, M.T.: Determination of N-alkanoyl 5-hydroxytryptamines (C-5-HT) in coffee beans by means of HPLC and TLC. *Ann. Chim. (Rome)*, 79 (1989) 369-377; *C.A.*, 112 (1990) 75407c.
- 533 Benfenati, E., Macconi, D., Noris, M., Icardi, G., Bettazzoli, L., de Bellis, G., Gavinelli, M., Rotondo, S. and Remuzzi, G.: Development of a mass spectrometric method to quantitate platelet activating factor in mouse urine. *J. Lipid Res.*, 30 (1989) 1977-1981.

- 534 Benfenati, E., de Bellis, G., Chen, S., Bettazzoli, L., Fanelli, R., Tacconi, M.T., Kirschner, G. and Pege, G.: A fast atom bombardment-mass spectrometric method to quantitative lysophosphatidylserine in rat brain. *J. Lipid Res.*, 30 (1989) 1983-1986.
- 535 Biffen, M. and Hanke, D.E.: Reduction in the level of intracellular myo-inositol in cultured soybean (*Glycine max.* cells inhibits cell division. *Biochem. J.*, 265 (1990) 809-814.
- 536 Boyle, L.E., Sklar, L.A. and Traynor-Kaplan, A.E.: An HPLC procedure for separating polyphosphoinositides on hydroxyapatite. *J. Lipid Res.*, 31 (1990) 157-159.
- 537 Brockman, M.J., Eiroa, A.M. and Marcus, A.J.: Albumin redirects platelet eicosanoid metabolism toward 12(S)-hydroxyeicosatetraenoic acid. *J. Lipid Res.*, 30 (1989) 1925-1932.
- 538 Churcher, Y., Allan, D. and Gomperts, B.D.: Relationship between arachidonate generation and exocytosis in permeabilized mast cells. *Biochem. J.*, 266 (1990) 157-163.
- 539 Cockcroft, S. and Stutchfield, J.: The receptors for ATP and fMetLeuPhe are independently coupled to phospholipases C and A₂ via G-protein(s). Relationship between phospholipase C and A₂ activation and exocytosis in HL60 cells and human neutrophils. *Biochem. J.*, 263 (1989) 715-723.
- 540 Dohmeyer, D.J., Corr, P.B. and Creer, M.H.: A sensitive, radiometric assay for lysophosphatidylcholine. *Anal. Biochem.*, 185 (1990) 36-43.
- 541 Ederveen, A.G.H., van Emst-de Vries, S.E., de Pont, J.J.H.M. and Willems, P.H.G.M.: The diacylglycerol kinase inhibitor, R59022, potentiates cholecystokinin-induced enzyme secretion from rabbit pancreatic acini. *Eur. J. Biochem.*, 188 (1990) 333-338.
- 542 Gu, X., Gu, T. and Yu, R.K.: Direct assay of glycosphingolipid glycosyltransferase activities on thin-layer chromatogram. *Anal. Biochem.*, 185 (1990) 151-155.
- 543 Hack, M.H. and Helmy, F.M.: Thin-layer chromatographic analysis of myelin lipids, their differential O-deacylation by primary alkylamines and their selective staining by thionine. A limited phylogenetic study. *J. Chromatogr.*, 525 (1990) 339-347.
- 544 Higgs, M.H., Sherma, J. and Fried, B.: Neutral lipids in the digestive gland-gonad complex of *Biomphalaria glabrata* snails, fed lettuce versus hen's egg yolk, determined by quantitative TLC-densitometry. *J. Planar Chromatogr.*, 3 (1990) 38-41.
- 545 Holmes, E.H. and Greene, T.G.: Characterization of two monoclonal antibodies specific for lacto-series type 1 chain Gal β 1 $\rightarrow$ 3GlcNAc-terminal structure. *Arch. Biochem. Biophys.*, 277 (1990) 181-188.
- 546 Kiguchi, K., Henning-Chubb, C.B. and Huberman, E.: Glycosphingolipid patterns of peripheral blood lymphocytes, monocytes, and granulocytes are cell specific. *J. Biochem. (Tokyo)*, 107 (1990) 8-14.
- 547 Kovacs, L., Pick, J. and Fucsek, J.: Lipid analysis by high-performance thin-layer chromatography. I. Microtechniques. *J. Planar Chromatogr. - Mod. TLC*, 2 (1989) 389-391; *C.A.*, 112 (1990) 73138s.
- 548 Kurosawa, M.: Inhibiting effects of azelastine on protein kinase C and diphosphoinositide kinase in rat mast cells. *Arzneim.-Forsch.*, 40 (1990) 162-165.
- 549 Kyogashima, M., Mulshine, J., Linnoila, R.I., Jensen, S., Magnani, J.L., Nudelman, E., Hakomori, S.-i. and Ginsburg, V.: Antibody 624H12, which detects lung cancer at early stages, recognizes a sugar sequence in the glycosphingolipid difucosylneolactonohexaosylceramide (V^{FucIII}FucnLc6Cer). *Arch. Biochem. Biophys.*, 275 (1989) 309-314.

- 550 Lendrath, G., Nasner, A. and Kraus, L.: Behaviour of vegetable phospholipids in thin-layer chromatography. Optimization of mobile phase, detection and direct evaluation. *J. Chromatogr.*, 502 (1990) 385-392.
- 551 Lo, S.-L. and Chang, E.L.: Purification and characterization of a liposomal-forming tetraether lipid fraction. *Biochem. Biophys. Res. Commun.*, 167 (1990) 238-243.
- 552 Miyake, R., Yokoyama, M. and Fukuzaki, H.: Inhibition of lysophospholipase by cholesterol in rabbit aorta. *Biochem. Biophys. Res. Commun.*, 167 (1990) 143-147.
- 553 Nakamura, K., Hashimoto, Y., Moriwaki, K., Yamakawa, T. and Suzuki, A.: Genetic regulation of GM4(NeuAc) expression in mouse erythrocytes. *J. Biochem. (Tokyo)*, 107 (1990) 3-7.
- 554 Nielsen, H.: Three-step one-dimensional thin-layer chromatographic separation of neutral lipids. *J. Chromatogr.*, 498 (1990) 423-427.
- 555 Okumura, K., Yamada, Y., Kondo, J., Kobayashi, N., Hashimoto, H. and Ito, T.: Quantitative determination of 1,2-diacylglycerol in thoracic aorta of the rat using Iatroscan TLC/FID: effect of norepinephrine. *Lipids*, 24 (1989) 982-984; *C.A.*, 112 (1990) 73114f.
- 556 Olsson, U., Kaufmann, P. and Herslöf, B.G.: TLC separation of lipids, optimized with chemometrical methods. *J. Planar Chromatogr.*, 3 (1990) 55-60.
- 557 Ritter, G., Boosfeld, E., Marakstein, E., Yu, R.K., Ren, S., Stallcup, W.B., Oettgen, H.F., Old, L.J. and Livingston, P.O.: Biochemical and serological characteristics of natural 9-O-acetyl GD3 from human melanoma and bovine buttermilk and chemically O-acetylated GD3. *Cancer Res.*, 50 (1990) 1403-1410.
- 558 Rustenbeck, I. and Lenzen, S.: Quantitation of hexadecylphosphocholine by high-performance thin-layer chromatography with densitometry. *J. Chromatogr.*, 525 (1990) 85-91.
- 559 Schick, P.K., Williams-Gartner, K. and He, X.: Lipid composition and metabolism in megakaryocytes at different stages of maturation. *J. Lipid Res.*, 31 (1990) 27-35.
- 560 Schweisguth, D.C., Quelle, F.W., Wachob, G. and Hammerstedt, R.H.: Isolation and characterization of brain lipids by solid phase extraction and thin layer chromatography. *Biochem. Educ.*, 17 (1989) 211-213; *C.A.*, 112 (1990) 97652f.
- 561 Singhal, A.K., Orntoft, T.F., Nudelman, E., Nance, S., Schibig, L., Stroud, M.R., Clausen, H. and Hakomori, S.: Profiles of Lewis^x-containing glycoproteins and glycolipids in sera of patients with adenocarcinoma. *Cancer Res.*, 50 (1990) 1375-1380.
- 562 Sundaram, K.S. and Lev, M.: Regulation of sulfotransferase activity by vitamin K in mouse brain. *Arch. Biochem. Biophys.*, 277 (1990) 109-113.
- 563 Taki, T., Nishiwaki, S., Ishii, K. and Handa, S.: A simple and specific assay of glycosyltransferase and glycosidase activities by an enzyme-linked immunosorbent assay method, and its application to assay of galactosyltransferase activity in sera from patients with cancer. *J. Biochem. (Tokyo)*, 107 (1990) 493-498.
- 564 Taki, T., Yamamoto, K., Takamatsu, M., Ishii, K., Myoga, A., Sekiguchi, K., Ikeda, I., Kurata, K., Nakayama, J., Handa, S. and Matsumoto, M.: Accumulation of gangliosides with N-acetylneuraminosyl(α 2-6)lactosamine structure in primary human hepatoma. *Cancer Res.*, 50 (1990) 1284-1290.
- 565 Tomono, Y., Abe, K. and Watanabe, K.: High-performance liquid affinity chromatography and *in situ* fluorescent labeling on thin-layer chromatography of glycosphingolipids. *Anal. Biochem.*, 184 (1990) 360-368.
- 566 Trinchera, M., Ghidoni, R., Greggia, L. and Tettamanti, G.: The N-acetylgalactosamine residue of exogenous GM₂ ganglioside is recycled for glycoconjugate biosynthesis in rat liver. *Biochem. J.*, 266 (1990) 103-106.

- 567 Vallari, D.S., Record, M. and Snyder, F.: Conversion of alkylacetyl glycerol to platelet-activating factor in HL-60 cells and subcellular localization of the mediator. *Arch. Biochem. Biophys.*, 276 (1990) 538-545.
- 568 Varela, I., Alvarez, J.F., Clemente, R., Ruiz-Albusac, J.M. and Mato, J.M.: Asymmetric distribution of the phosphatidylinositol-linked phospho-oligosaccharide that mimics insulin action in the plasma membrane. *Eur. J. Biochem.*, 188 (1990) 213-218.
- 569 Xu, Y., Jiang, X., Wang, Y., Guo, R. and Yu, L.: (Analysis of phospholipids in Wuzi Yanzhong Wan and its Chinese drugs). *Zhongcaoyao*, 20 (1989) 303-305; *C.A.*, 112 (1990) 25743p.
- 570 Yang, L.-Y., Kuksis, A. and Myher, J.J.: Lipolysis of menhaden oil triacylglycerols and the corresponding fatty acid alkyl esters by pancreatic lipase *in vitro*: a reexamination. *J. Lipid Res.*, 31 (1990) 137-148.

See also 588.

13. STEROIDS

- 571 Robards, K. and Towers, P.: Chromatography as a reference technique for the determination of clinically important steroids. *Biomed. Chromatogr.*, 4 (1990) 1-19 - a review with 255 refs.

13a. Pregnane and androstane derivatives

- 572 Lamparczyk, H., Ochocka, R.J., Zarzycki, P. and Zielinski, J.P.: Separation of steroids by reversed-phase HPTLC using various binary mobile phases. *J. Planar Chromatogr.*, 3 (1990) 34-37.

13b. Estrogens

See 572.

13c. Sterols

- 573 Florin-Christensen, M., Florin-Christensen, J., Garin, C., Isola, E., Brenner, R.R. and Rasmussen, L.: Inhibition of *Trypanosoma cruzi* growth and sterol biosynthesis by lovastatin. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1441-1445.
- 574 Fried, B., Duncan, M., Sherma, J. and Hoskin, G.P.: Thin layer chromatographic determination of sterols and other lipids in the hemolymph of *Biomphalaria glabrata* (Gastropoda). *J. Liq. Chromatogr.*, 12 (1989) 3151-3161.
- 575 Yan, J. and Bittman, R.: A facile synthesis of [14 C]epicholesterol. *J. Lipid Res.*, 31 (1990) 160-162.

See also 562.

13d. Bile acids and alcohols

- 576 Batta, A.K., Arora, R., Salen, G., Tint, G.S., Eskreis, D. and Katz, S.: Characterization of serum and urinary bile acids in patients with primary biliary cirrhosis by gas-liquid chromatography-mass spectrometry: effect of ursodeoxycholic acid treatment. *J. Lipid Res.*, 30 (1989) 1953-1962.

- 577 Kuipers, F., Radomska, A., Zimniak, P., Little, J.M., Havinga, R., Vonk, R.J. and Lester, R.: Defective biliary secretion of bile acid 3-O-glucuronides in rats with hereditary conjugated hyperbilirubinemia. *J. Lipid Res.*, 30 (1989) 1835-1845.
- 578 Ni, S., Chen, N. and Feng, W.: (Study of method of thin-layer chromatography of bile). *Shenyang Yaoxueyuan Xuebao*, 6 (1989) 197-198; *C.A.*, 111 (1989) 239622b.
- 579 Wang, J., Gao, Z., Li, Z., Wang, Z. and Liu, S.: (Thin-layer chromatography of free bile acid in bile). *Shaanxi Yixue Zazhi*, 18 (1989) 28-30; *C.A.*, 112 (1990) 51505g.

13f. Other steroids

See 682.

14. STEROID GLYCOSIDES AND SAPONINS

- 580 Xie, R., Lu, N., Liu, Q. and Deng, L.: (Separation of chemical constituents of diosgenin waste material and TLC densitometric determination of diosgenin). *Sichuan Daxue Xuebao, Ziran Kexueban*, 26 (1989) 208-211; *C.A.*, 112 (1990) 25758x.

See also 479, 683, 684.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 581 Endo, M., Mizutani, T., Moryasu, M., Ichmaru, M., Kato, A. and Hashimoto, Y.: (Determination of hinokitiol by HPLC or TLC). *Jpn. Kokai Tokyo Koho JP 01,185,441 [89,185,441]* (Cl. G01N30/88), 25 Jul. 1989, Appl. 88/11,627, 20 Jan. 1988; 4 pp.; *C.A.*, 112 (1990) 42753j.
- 582 Pachaly, P.: Salbeiblätter und Dreilappiger Salbei. *Dtsch. Apoth.-Ztg.*, 130 (1990) 169-170.
- 583 Wu, C., Yang, S. and Wu, S.: (Preparative separation of borneol and isoborneol in Bingpian by dry column chromatography). *Huaxi Yike Daxue Xuebao*, 20 (1989) 327-330; *C.A.*, 112 (1990) 25446u.

See also 615.

15b. Essential oils

See 582.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 584 Surgova, T.M., Sidorenko, M.V., Kofman, I.S. and Vinnitsky, V.B.: Determination of histamine in the presence of polyamines by spectrodensitometric TLC. *J. Planar Chromatogr.*, 3 (1990) 81-82.

See also 487, 695.

17b. *Catecholamines and their metabolites*

- 585 Yoshimura, T., Masubuchi, Y. and Okazaki, S.: (Separation and determination of catecholamines in food by electrochromatoscanning method). *Bunseki Kagaku*, 38 (1989) 613-617; *C.A.*, 112 (1990) 75395x.

17c. *Urea and guanidine derivatives*

- 586 Chmil, V.D. and Vasyagina, R.D.: (Thin-layer chromatography of carbamic acid derivatives). *Zh. Anal. Khim.*, 44 (1989) 757-759; *C.A.*, 112 (1990) 15781k.
587 Dukhovnaya, I.S. and Yun, E.M.: (Determination of diisocyanates by thin-layer chromatography). *Zh. Anal. Khim.*, 44 (1989) 1296-1301; *C.A.*, 112 (1990) 48073j.

17d. *Other amine derivatives and amides (excluding peptides)*

- 588 Hendrickson, H.S., Kotz, K.J. and Hendrickson, E.K.: Evaluation of fluorescent and colored phosphatidylcholine analogs as substrates for the assay of phospholipase A₂. *Anal. Biochem.*, 185 (1990) 80-83.

See also 676.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. *Amino acids and their derivatives*

- 589 Bhushan, R. and Ali, I.: Adsorption and partition of amino acids in a batch process compared to their TLC separation on silica gel. *J. Planar Chromatogr.*, 3 (1990) 85-87.
590 Bhushan, R. and Reddy, G.P.: Thin layer chromatography of dansyl and dinitrophenyl derivatives of amino acids. A review. *Biomed. Chromatogr.*, 3 (1989) 233-240 - a review with 47 refs.
591 Chiari, M., Chiesa, C., Righetti, P.G., Corti, M., Jain, T. and Shorr, R.: Kinetics of cysteine oxidation in immobilized pH gradient gels. *J. Chromatogr.*, 499 (1990) 699-711.
592 Euerby, M.R.: Resolution of the neuroexcitatory N-methylaspartic acid (2-methylaminosuccinic acid) enantiomers by ligand-exchange thin-layer chromatography. *J. Chromatogr.*, 502 (1990) 226-229.
593 Nesterova, O.V., Samylina, I.A. and Baratova, L.A.: Development of standardization methods for kukurbin from pumpkin seeds). *Farmatsiya (Moscow)*, 38, No. 6 (1989) 36-38; *C.A.*, 112 (1990) 84295s.
594 Prakasam, V.R.: An evaluation of butanol-acetic acid-water solvent system in ascending paper chromatography of amino acids. *J. Sci. Res. (Bhopal)*, 10 (1988) 103-105; *C.A.*, 112 (1990) 12467b.
595 Ueda, T., Kam, C.-M. and Powers, J.C.: The synthesis of arginylfluoroalkanes, their inhibition of trypsin and blood-coagulation serine proteinases and their anticoagulant activity. *Biochem. J.*, 265 (1990) 539-545.

See also 496, 506, 689.

18b. Peptides and peptidic and proteinous hormones

- 596 Coméra, C., Rothhut, B. and Russo-Marie, F.: Identification and characterization of phospholipase A₂ inhibitory proteins in human mononuclear cells. *Eur. J. Biochem.*, 188 (1990) 139-146.
- 597 Lin Ling, B., Baeyens, W.R.G. and Raemdonck, A.: Application of high-performance thin-layer chromatography-fluorescence densitometry to the simultaneous determination of reduced and oxidized glutathione. *J. Chromatogr.*, 502 (1990) 230-235.
- 598 Meador, J., III, Cannon, B., Cannistraro, V.J. and Kennell, D.: Purification and characterization of *Escherichia coli* RNase I. Comparisons with RNase M. *Eur. J. Biochem.*, 187 (1990) 549-553.
- 599 Yannariello-Brown, J. and Madri, J.A.: A 48 kDa collagen-binding phosphoprotein isolated from bovine aortic endothelial cells interacts with the collagenous domain, but not the globular domain, of collagen type IV. *Biochem. J.*, 265 (1990) 383-392.

See also 503.

19. PROTEINS

191. Specific binding and receptor proteins

- 600 Lewis, R.E., Cao, L., Perregaux, D. and Czech, M.P.: Threonine 1336 of the human insulin receptor is a major target for phosphorylation by protein kinase C. *Biochemistry*, 29 (1990) 1807-1813.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

20f. other hydrolases

- 601 Safarik, I.: Use of a densitometer for microdetermination of enzyme activity. *J. Planar Chromatogr.*, 3 (1990) 90-91.

20g. Lyases

- 602 Gibson, K.M., Lee, C.F., Kamali, V., Johnston, K., Beaudet, A.L., Craigen, W.J., Powell, B.R., Schwartz, R., Tsai, M.Y. and Tuchman, M.: 3-Hydroxy-3-methylglutaryl-CoA lyase deficiency as detected by radiochemical assay in cell extracts by thin-layer chromatography, and identification of three new cases. *Clin. Chem. (Winston-Salem)*, 36 (1990) 297-303.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 603 Chavdarova, V., Zlatanov, L., Koburova, K., Shkenderov, S.: (Densitometric determination of ATP and its degradation products in the preparation Phosphobion ampuls). *Izv. Durzh. Inst. Kontrol Lek. Sredstva*, 22 (1989) 24-29; *C.A.*, 112 (1990) 42704u.

- 604 Hamza, Y.E. and Kata, M.: (Influence of hydrotropic and complexing agents on the solubilization of allopurinol. *Pharm. Ind.*, 51 (1989) 1159-1162; *C.A.*, 112 (1990) 42387t.
- 605 Luke, W., Petry, H., Hoefler, K., Anders, A. and Anders, F.: Detection of RNA-dependent DNA polymerase activity in the *Xiphophorus* melanoma system. *Eur. J. Biochem.*, 188 (1990) 301-308.

22. ALKALOIDS

- 606 Cisowski, W. and Lamer-Zarawska, E.: Ergot alkaloid analysis by TLC with gradient elution. *J. Planar Chromatogr.*, 3 (1990) 47-50.
- 607 Fannin, F.F., Bush, L.P., Siegel, M.R. and Rowan, D.D.: Analysis of peramine in fungal endophyte-infected grasses by reversed-phase thin-layer chromatography. *J. Chromatogr.*, 503 (1990) 288-292.
- 608 Nguyen, K.C. and Nikolova, I.G.: (Quantitative determination of alkaloids in root bark of some species of *Rauwolfia* L. by thin-layer chromatography). *Rastit. Resur.*, 25 (1989) 594-599; *C.A.*, 112 (1990) 62744k.
- 609 Xie, D., Liu, L. and Xu, D.: (Quantitative determination of cantharidin injections). *Zhongguo Yiyao Gongye zazhi*, 20 (1989) 266-268; *C.A.*, 112 (1990) 42711u.

See also 490, 506, 665, 674, 675.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

See 641.

23e. Other N-heterocyclic compounds

- 610 Fernández, B.M. and Abasolo, M.I.: HPTLC for monitoring kinetics of the synthesis of quinoxaline derivatives. *J. Planar Chromatogr.*, 3 (1990) 20-23.
- 611 Valiotti, A.B., Ermakova, T.V., Pal'chevskii, V.V. and Churkina, S.I.: (Chromatographic separation of synthetic water-soluble cationic porphyrins). *Zh. Anal. Khim.*, 44 (1989) 169-171; *C.A.*, 112 (1990) 68554v.
- 612 Zhang, W., Chen, H. and Xiao, W.: (Determination of metronidazole in cream by TLC). *Zhongguo Yaoke Daxue Xuebao*, 20 (1989) 243-244; *C.A.*, 112 (1990) 25784c.

See also 471, 488, 637, 656, 667.

24. ORGANIC SULPHUR COMPOUNDS

- 613 Polyanskii, E.I., Butova, S.A., Malii, V.A. and Shenderovich, V.A.: (Analysis of the mixture of alkali metal alkylxanthates by a thin-layer chromatographic method). *Obogashch. Rud (Leningrad)*, 3 (1989) 32; *C.A.*, 112 (1990) 69200p.
- 614 Sibilev, A.V. and Dashbalyun, T.: (Development of unified chromatographic techniques for assessing the quality of phenothiazine derivatives). *Farmatsiya (Moscow)*, 38 (1989) 75; *C.A.*, 111 (1989) 239632e.

See also 471, 558.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCLUDING SUGAR PHOSPHATES)

- 615 Kabakoff, B.D., Doyle, J.W. and Kandutsch, A.A.: Relationships among dolichyl phosphate, glycoprotein synthesis, and cell culture growth. *Arch. Biochem. Biophys.*, 276 (1990) 382-389.
- 616 Pullarkat, R.K., Pullarkat, P.S. and Morris, G.N.: Phosphorylated dolichols in aging. *Biochem. J.*, 265 (1990) 891-894.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

- 617 Macias-Matos, C., Garica-Pino, C. and Arias-Verdes, J.A.: (Determination of organotin stabilizers in soy sauce in PVC bottles). *Rev. Agroquim. Tecnol. Aliment.*, 29 (1989) 281-284; *C.A.*, 112 (1990) 97126n.

26a. Organometallic compounds

- 618 Baba, T. and Sasaki, S.: (Rapid analysis of dibutyltin compounds and tricresyl phosphate in poly(vinyl chloride) by thin-layer chromatography). *Shokuhin Eiseigaku Zasshi*, 30 (1989) 321-323; *C.A.*, 112 (1990) 57395t.

26c. Coordination compounds

- 619 Timerbaev, A.R., Tsoi, I.G. and Petrukhin, O.M.: Correlation analysis in liquid chromatography of metal chelates. I. One-dimensional retention-structure models in reversed-phase thin-layer chromatography. *J. Chromatogr.*, 498 (1990) 337-348.

See also 469, 611.

27. VITAMINS AND VARIOUS GROWTH REGULATORS (NON-PEPTIDIC)

- 620 Gao, Z.: (Determination of pyridoxal in pyridoxine and its preparations). *Yaowu Fenxi Zazhi*, 9 (1989) 287-289; *C.A.*, 112 (1990) 42696t.
- 621 Jarzebinski, J. and Lugowska, E.: (Densitometric measurements in determining the active drug substances. XVI. Multicomponent drugs containing ascorbic acid). *Acta Pol. Pharm.*, 46 (1989) 160-164; *C.A.*, 112 (1990) 25799m.
- 622 Marciniak, B. and Stachowicz, M.: (Analysis of drug decomposition products. XLII. Determination of vitamins K in pharmaceutical preparations in the presence of the decomposition products). *Acta Pol. Pharm.*, 46 (1989) 138-145; *C.A.*, 112 (1990) 25767z.

See also 640.

28. ANTIBIOTICS

- 623 Anonymous: Antibiotic drugs: rifampin for injection. *United States Food and Drug Administration, Fed. Regist.* 18 Sep. 1989, 54, No. 179 (1989) 38374-38375; *C.A.*, 111 (1989) 239350m.
- 624 Chen, Q., Zhou, J., Duan, T. and Zhou, H.: (Preparative separation of cephalosporins by centrifugal TLC). *Kangshengsu*, 14 (1989) 161-167; *C.A.*, 112 (1990) 83945s.

- 625 Florent, J.C., Genot, A. and Monneret, C.: Synthesis and antitumor activity of new anthracyclines. *J. Antibiot.*, 42 (1989) 1823-1830.
- 626 Gause, G.F., Brazhnikova, M.G., Lomakina, N.N., Berdnikova, T.F., Fedorova, G.B., Tokareva, N.L., Borisova, V.N. and Batta, G.: Eremomycin - new glycopeptide antibiotic: chemical properties and structure. *J. Antibiot.*, 42 (1989) 1790-1799.
- 627 Lee, M.D., Manning, J.K., Williams, D.R., Kuck, N.A., Testa, R.T. and Borders, D.B.: Calichaemicins, a novel family of antitumor antibiotics. 3. Isolation, purification and characterization of calichaemicins β_1^{Br} , γ_1^{Br} , α_2^I , α_3^I , β_1^I , γ_1^I and δ_1^I . *J. Antibiot.*, 42 (1989) 1070-1087.
- 628 Moroi, R., Kinuno, K., Konno, T., Furukawa, M. and Kataoka, K.: (Physicochemical properties and stability of ofloxacin). *Kagaku Ryoho no Ryoiki*, 5 (1989) 1517-1528; C.A., 112 (1990) 62426h.
- 629 United States Food and Drug Administration: Antibiotic drugs; cephalexin hydrochloride monohydrate tablets. *Red. Regist.*, 54, No. 227 (28 Nov. 1989) 48859-48861; C.A., 112 (1990) 62407c.

See also 486.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

See 502.

29a. Chlorinated insecticides

- 630 De la Vigne, U. and Jänchen, D.: Determination of pesticides in water by HPTLC using automated multiple development (AMD). *J. Planar Chromatogr.*, 3 (1990) 6-9.

29b. Phosphorus insecticides

- 631 Funk, W., Cleres, L., Pitzer, H. and Donnevert, G.: Organophosphorus insecticides - quantitative HPTLC determination and characterization. *J. Planar Chromatogr. - Mod. TLC*, 2 (1989) 285-289; C.A., 112 (1990) 17680a.
- 632 Xiao, X.: (Determination of residual amount of 4049 in food without purification of the sample). *Lihua Jianyan, Huaxue Fence*, 24 (1988) 45; C.A., 112 (1990) 97096c.
- 633 Xie, B. and Zhou, Z.: (Rapid and simultaneous TLC determination of sulfur-containing organophosphorus pesticides and DDVP, Dipterex). *Lihua Jianyan, Huaxue Fence*, 25 (1989) 126; C.A., 112 (1990) 93755u.

29c. Carbamates

- 634 Gao, G.: (Rapid detection of carbofuran contamination) *Lihua Jianyan, Huaxue Fence*, 24 (1988) 102, 115; C.A., 112 (1990) 117440r.

29d. Herbicides

- 635 Lee, P.W., Forbis, A.D. and Franklin, L.: Bioaccumulation of cinmethylin in bluegill sunfish. *J. Agric. Food Chem.*, 38 (1990) 323-327.
- 636 Rathore, H.S. and Saxena, S.K.: A comparative study of some carboxylic acid herbicides by normal-phase, sequential, and ion-pair reversed phase thin-layer chromatography on calcium sulfate. *J. Planar Chromatogr. - Mod. TLC*, 2 (1989) 387-389; C.A., 112 (1990) 50494r.

29f. Other types of pesticides and various agrochemicals

See 607.

30. SYNTHETIC AND NATURAL DYES

See 505.

30a. Synthetic dyes

- 637 Feng, Y., Zhang, Z., Chen, Z. and Xia, C.: (Quantitative determination of N-phenylcarbazole from photoinduced reaction by TLC-UV spectrophotometry). *Shanxi Daxue Xuebao, Ziran Kexueban*, 12 (1989) 201-204; *C.A.*, 112 (1990) 59215w.
- 638 Yin, P., Qin, P. and Liu, Q.: (Separation of basic dyes by ion-pair reversed-phase thin-layer chromatography). *Sepu*, 7 (1989) 313-314; *C.A.*, 112 (1990) 79412y.

See also 666.

30b. Chloroplast and other natural pigments

- 639 Bounias, M. and Daurade, M.H.: (Quenching of the artifactual oxidation of chlorophyll pigments chromatographed on silica gel thin layers by impregnation of plates with glucose or thiosulfate). *Analysis*, 17 (1989) 401-404; *C.A.*, 112 (1990) 51486b - impregnation the plates with a mild reducing agent.
- 640 Kanasawud, P. and Crouzet, J.C.: Mechanism of formation of volatile compounds by thermal degradation of carotenoids in aqueous medium. 1. β -Carotene degradation. *J. Agric. Food Chem.*, 38 (1990) 237-243.
- 641 Lützw, M. and Kleinig, H.: Chlorophyll-free chromoplasts from daffodil contain most of the enzymes for chlorophyll synthesis in a highly active form. *Arch. Biochem. Biophys.*, 277 (1990) 94-100.

31. PLASTICS AND THEIR INTERMEDIATES

See 473, 617.

32. DRUG ANALYSIS*32a. Drug analysis, general techniques*

- 642 Amer, M.M., Khattab, F.I. and Hassan, N.Y.M.: Titrimetric determination of some pharmaceutical compounds using some halogenated derivatives. *Egypt. J. Pharm. Sci.*, 30 (1989) 207-220; *C.A.*, 112 (1990) 62754p.
- 643 Kulikov, V.A. and Kolkov, M.A.: (Use of thin-layer chromatography for separation and identification of components of drug mixtures). *Sovrem. Aspekty Teorii i Prakt. Farmatsii, L.*, (1988) 67-70; *C.A.*, 112 (1990) 12011y.
- 644 Singh, A.K., Granley, K., Ashraf, M. and Mishra, U.: Drug screening and confirmation by thin layer chromatography. *J. Planar Chromatogr.-Mod. TLC*, 2 (1989) 410-419; *C.A.*, 112 (1990) 111343q - a review with 110 refs.
- 645 Warren, B.: Thin-layer chromatography impurity tests-can they be improved? *Pharm. Ind.*, 51 (1989) 1057-1060; *C.A.*, 112 (1990) 84287r.

See also 674.

32b. *Antirheumatics and antiinflammatory drugs*

- 646 Chowdary, K.P.R. and Murty, A.S.R.: Studies on freeze-dried solid dispersions. *Indian J. Hosp. Pharm.*, 26 May-June (1989) 93-95; *C.A.*, 112 (1990) 11864s.
- 647 Giese, U.: Absorption and distribution of ibuprofen from a cream formulation after dermal administration to guinea pigs. *Arzneim.-Forsch.*, 40 (1990) 84-88.
- 648 Guneri, T.: The pH and light stability of isopropylaminophenazone in solutions. *Hacettepe Univ. Eczacilik Fak. Derg.*, 9 (1989) 27-33; *C.A.*, 112 (1990) 62453q.
- 649 Lachmann, G., Siegemund, B. and Kusche, W.: Pharmacokinetics and metabolism of 14 C-oxaceprol in beagle dogs after intramuscular and oral administration. *Arzneim.-Forsch.*, 40 (1990) 200-206.

See also 674.

32c. *Autonomic and cardiovascular drugs*

- 650 Cook, C.S., Gwilt, P.R., Kowalski, K., Gupta, S., Oppermann, J. and Karim, A.: Pharmacokinetics of disopyramide in the dog. Importance of mono-N-dealkylated metabolite kinetics in assessing pharmacokinetic modeling of the parent drug. *Drug Metab. Disp.*, 18 (1990) 42-49.
- 651 Neidlein, R., Volland, C. and Strein, K.: Biotransformation of the partial β -agonist doxaminol in dog. *Arzneim.-Forsch.*, 40 (1990) 130-136.
- 652 Sungur, S., Ersoy, L., Alpertunga, B. and Yalcin, M.S.: Thin layer chromatographic detection of some beta blockers after derivatization with dabsyl chloride. *Pharmazie*, 44 (1989) 864-865.
- 653 Weil, A., Caldwell, J. and Strolin-Benedetti, M.: The metabolism and disposition of 14 C-fenofibrate in human volunteers. *Drug Metab. Disp.*, 18 (1990) 115-120.
- 654 Zivanovic, L., Agatonovic, S. and Radulovic, D.: Determination of diuretics in pharmaceutical preparations using thin-layer chromatographic densitometry. *Pharmazie*, 44 (1989) 864.

See also 508, 580.

32d. *Central nervous system drugs*

- 655 Bezek, S., Kukan, M., Kállay, Z., Trnovec, T., tefek, M. and Piotrovskiy, L.B.: Disposition of ethimizol, a xanthine-related nootropic drug, in perfused rat liver and isolated hepatocytes. *Drug Metab. Disp.*, 18 (1990) 88-95.
- 656 Dashbalyn, T., Prokof'eva, V.I., Egorov, A.E., Kashtahova, V.V. and Nechaeva, E.B.: (Application of thin-layer and high-performance liquid chromatography to the identification and separation of 10-acyl derivatives of phenothiazine). *Farmatsiya (Moscow)*, 38, No. 6 (1989) 39-41; *C.A.*, 112 (1990) 62763r.
- 657 Sibilev, A.V., Prokof'eva, V.I., Arzamatsev, A.P., Lamm, M.M. and Kirpicheva, E.I.: (A unified procedure for assessing the quality of phenothiazine 10-alkyl derivatives by thin-layer chromatography). *Khim.-Farm. Zh.*, 23 (1989) 1267-1270; *C.A.*, 112 (1990) 62755q.
- 658 Tishchenkova, I.F. and Nazarov, U.Z.: (Estimation of the antihistamine agent diazoline in serum and urine by densitometric thin-layer chromatography). *Khim.-Farm. Zh.*, 23 (1989) 1393-1394; *C.A.*, 112 (1990) 48181t - pharmacokinetic study.

See also 674.

32e. Chemotherapeutics (except cytostatics and antibiotics)

- 659 Bellomonte, G., Franciosa, G. and Sanzini, E.: (Detection of sulfonamides in feed by high-performance TLC). *Riv. Soc. Ital. Sci. Aliment*, 18 (1989) 247-250; C.A., 112 (1990) 117461y.
- 660 Mohamed, A.-M.I.: Spectrophotometric determination of some benzene sulfonamides with tetracyanoquinodimethane. *J. Assoc. Off. Anal. Chem.*, 72 (1989) 885-889.
- 661 Semenido, Y.N. and Rodionova, R.A.: (Use of thin-layer chromatography in analysis of drug mixtures containing sulfanilamides). *Sovrem. Aspekty Teorii i Prakt. Farmatsii, L.*, (1988) 78-81; C.A., 112 (1990) 12012z.
- 662 Van Poucke, L., Rousseau, D., de Spiegeleer, B. and van Peteghem, C.: A rapid high-performance thin-layer chromatographic screening method for sulfonamides residues in animal muscle tissues. *Agric., Food Chem. Consum., Proc. Eur. Conf. Food Chem., 5th*, (1989) 438-442; C.A., 112 (1990) 75398a.

32f. Cytostatics

- 663 Bachrata, M., Stankovicova, M., Bezakova, Z. and Blesova, M.: A study of physicochemical properties of drug substances with cancerostatic effects. Part V. Study of the kinetics of acid hydrolysis. *Acta Fac. Pharm. Univ. Comeniana*, 42 (1988, Publ. 1989) 109-124; C.A., 112 (1990) 83954u.
- 664 Issaq, H.J., Seburn, K.E., Andrews, P. and Schaufelberger, D.E.: Multimodal thin layer chromatographic separation of bryostatins 1 and 2 from *Bugula neritina* extracts employing side-by-side silica gel/C₁₈ plates. *J. Liq. Chromatogr.*, 12 (1989) 3129-3134.
- 665 Kobayashi, J. and Oizumi, Y.: (Antitumor tetracyclic alkaloids from sea squirt (*Cystodites dellichiajei*)). *Jpn. Kokai Tokkyo Koho Pat.* JP 01,132,584 [89,132,584] (Cl. C07D471/04), 25 May 1989, Appl. 87/288,571, 17 Nov. 1987; 3 pp.; C.A., 112 (1990) 42550r.

See also 665.

32g. Other drug categories

- 666 Geeta, N., Veena, A.P. and Baggi, T.R.: Spectrophotometric determination of phenolphthalein in pharmaceutical products using 3-methylbenzothiazolinone-2-hydrazone. *Microchem. J.*, 40 (1989) 304-310; C.A., 112 (1990) 62761p.
- 667 Grabowska, I., Krackowska, A., Rajzer, D. and Smietanska, H.: (Studies on the stability of drugs derived from imidazoline. XI. Qualitative and quantitative evaluation of the stability of tetrahydrozoline hydrochloride in solutions). *Acta Pol. Pharm.*, 46 (1989) 61-69; C.A., 112 (1990) 25477e.
- 668 Han, S.D., Jiang, D.H. and Wang, J.L.: (Study on quality of gossypol recrystallized from chloroform). *Yaoxue Xuebao*, 24 (1989) 465-471; C.A., 112 (1990) 83944r.
- 669 Hiraki, Y., Nomaguchi, T., Kawazoe, T., Kumamoto, Y., Shinohara, T., Kusaba, K. and Syouji, Y.: (Effect of sunlight on the quality of ciprofloxacin preparation). *J. Jpn. Soc. Hosp. Pharm.*, 25 (1989) 1375-1377; C.A., 112 (1990) 42442g.
- 670 Marciniak, B. and Wojciechowska, B.: (Determination of 5-pyrazolinone derivatives in pharmaceutical preparations. I. Spectrophotometric determination of two-component preparations containing propyphenazone). *Acta Pol. Pharm.*, 46 (1989) 248-257; C.A., 112 (1990) 84319c.
- 671 Scotto di Tella, A. and Di Nunzio, C.: (Two-dimensional thin-layer chromatographic separation of eight antidepressants). *Riv. Tossicol.: Sper. Clin.*, 18 (1988) 93-99; C.A., 112 (1990) 91122m.

- 672 Tománková, H. and abartová, J.: Determination of azidothymidine and its degradation product thymine in pharmaceutical dosage forms by HPLC and HPTLC densitometry. *Chromatographia*, 29 (1990) 44-50.

See also 593, 612, 648.

32h. Toxicological and forensic applications

- 673 De Zeeuw, R.A., Witte, D.T. and Franke, J.-P.: Identification power of thin-layer chromatographic colour reactions and integration of colour codes in a database for computerized identification in systematic toxicological analysis. *J. Chromatogr.*, 500 (1990) 661-671.
- 674 Tanaka, T., Aramaki, S. and Momose, A.: Thin-layer chromatographic screening procedure for some drugs in horse plasma. *J. Chromatogr.*, 496 (1989) 407-415.

See also 466.

32i. Plant extracts

- 675 Abdullabekova, V.N., Sokolov, I.V., Sabirov, K.A. and Gunar, V.I.: (Determination of rutin during *Sophora japonica* treatment). *Khim.-Farm. Zh.*, 23 (1989) 1367-1370; *C.A.*, 112 (1990) 62743j.
- 676 Bauer, R. and Remiger, P.: *Echinacea*. Part V. TLC and HPLC analysis of alkanamides in *Echinacea* drugs. *Planta Med.*, 55 (1989) 367-371; *C.A.*, 112 (1990) 11975d.
- 677 Fu, X., He, Z., Liang, W. and Tu, G.: (Chemotaxonomic characteristics of *Yanhusuo*). *Yunnan Zhiwu Yanjiu*, 11 (1989) 193-197; *C.A.*, 112 (1990) 42315t.
- 678 Gaitonde, R.V. and Naik, P.L.: Estimation of quercetin in the leaves of *Maesa indica* wall by a new colorimetric method. *Curr. Sci.*, 58 (1989) 982-983; *C.A.*, 111 (1989) 239626f.
- 679 Harles, A., Turcanu, L., Ciuca, C. and Cosofret, V.V.: (Analytical studies for the quantitative determination of aesculin from *Aesculus hyppocastanum* Bark). *Rev. Chim. (Bucharest)*, 40 (1989) 436-440; *C.A.*, 112 (1990) 42690m.
- 680 Jia, Y., Yan, X. and Nie, K.: (Comparative analysis of ursolic acid in *Crataegus pinnatifida* Bge. var. *major* N. E. Br. before and after processing). *Zhongguo Zhongyao Zazhi*, 14 (1989) 466-468; *C.A.*, 112 (1990) 25750p.
- 681 Liu, T., Liang, W. and Tu, G.: (Quantitative determination of four components in *Codonopsis pilosula* by TLC-densitometry). *Yaowu Fenxi Zazhi*, 9 (1989) 227-230; *C.A.*, 112 (1990) 25744q.
- 682 Lu, Y., Yang, X. and Sha, D.: (Determination of Δ^7 -stigmastanol and fructose in Dangshen preparations). *Zhongguo Zhongyao Zazhi*, 14 (1989) 484-486; *C.A.*, 112 (1990) 25751q.
- 683 Rauwald, H.W. and Janssen, B.: TLC analysis of *Ruscus aculeatus* L. and HPLC determination of deglucoruscin and deglucoruscinate in crude drug and pharmaceutical formulations. *PZ Wiss.*, 1 (1988) 61-68; *C.A.*, 112 (1990) 25742n.
- 684 Sun, W., Sha, Z. and Yang, C.: (TLC-densitometric determination of sarsasapogenin in *Anemarrhena asphodeloides* Bge.). *Zhongguo Zhongyao Zazhi*, 14 (1989) 546-548; *C.A.*, 112 (1990) 25756v.
- 685 Valko, K., Olajos, S. and Cserhati, T.: Prediction of the high-performance liquid chromatographic retention behaviour of some benzodiazepine derivatives by thin-layer chromatography. *J. Chromatogr.*, 499 (1990) 361-371.
- 686 Wagner, H., Bladt, S., Hartmann, U., Dally, A. and Berkulin, W.: Ginkgo biloba. DC- und HPLC-Analyse von Ginkgo-Extrakten und Ginkgo-Extrakte enthaltenden Phytopräparaten. *Dtsch. Apoth.-Ztg.*, 129 (1989) 2421-2429.

See also 464, 504, 569, 582, 593.

33. CLINICO-CHEMICAL APPLICATIONS

33b. Complex mixtures and profiling (single compounds by cross-reference only)

- 687 Swahn, C.G., Persson, A. and Pauli, S.: Metabolism of the benzodiazepine antagonist ^{11}C -Ro 15-1788 after intravenous administration in man. *Hun. Psychopharmacol.*, 4 (1989) 297-301; C.A., 112 (1990) 91157b.

See also 549, 571, 576, 584, 602.

34. FOOD ANALYSIS

34b. Complex mixtures (single compounds by cross-reference only)

- 688 Lander, V. and Schreier, P.: (Rapid high-performance thin layer chromatographic test for the detection of asarone, acorenone, coumarin and thujone in vermouth and brandy). *Lebensmittelchem. Gerichtl. Chem.*, 43 (1989) 126-127; C.A., 112 (1990) 96867t.

See also 518, 532, 585, 617, 632, 634, 662, 692.

35. ENVIRONMENTAL ANALYSIS

35c. Water pollution (complex mixtures; single compounds by cross-reference only)

- 689 He, Y.: (Determination of amino acids in wastewater from manufacture of sodium glutamate by thin layer chromatography). *Sepu*, 7 (1989) 181-183; C.A., 111 (1989) 239211s.
- 690 Sharshakova, L.N. and Skidan, V.I.: (Method of determining aromatic hydroxy compounds in water by paper chromatography). *U.S.S.R. Pat.* SU 1,509,729 (Cl. G01N30/94), 23 Sep. 1989, Appl. 4,273,158; 30 Jun. 1988; C.A., 111 (1989) 247297q.

See also 630.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

- 691 Saito, Y., Sato, T. and Anazawa, I.: The determination of the poly(oxyethylene) oligomer distribution in a non-ionic surfactant by means of thin-layer chromatography. *Bull. Chem. Soc. Jpn.*, 62 (1989) 3709-3710; C.A., 112 (1990) 57165t.

36b. Antioxidants and preservatives

- 692 Gross, J. and Jungkunz, G.: (Detection of shellac on apples by thin-layer chromatography). *Lebensmittelchem. Gerichtl. Chem.*, 43 (1989) 55-56; C.A., 111 (1989) 213421k.
- 693 Tomankova, H. and Pinkasova, M.: (TLC (HPTLC)-spectrodensitometry in pharmaceutical analysis. III. HPTLC-spectrodensitometric simultaneous determination of parabens in pharmaceutical preparations). *Cesk. Farm.*, 38 (1989) 299-303; C.A., 112 (1990) 42715y.

36c. Various technical products

- 694 Brahm, R., Ziegenbalg, W. and Renger, B.: TLC-spectrodensitometric determination of Polidocanol (Thesit) in different pharmaceutical preparations. *J. Planar Chromatogr.*, 3 (1990) 77-78.
- 695 Davidkova, P., Kopecek, J. and Gasparic, J.: Identification of developing agents by thin-layer chromatography. *J. Inf. Rec. Mater.*, 17 (1989) 117-124; C.A., 112 (1990) 90956z.
- 696 Gubernatorova, L.V.: (Streamlined determination of rosin by thin-layer chromatography). *U.S.S.R. Pat.* SU 1,492,269 (Cl. G01N30/90), 07 Jul. 1989, Appl. 4,283,129; 14 Jul. 1987; C.A., 112 (1990) 8921c.
- 697 Iyengar, D.R. and McCarthy, T.J.: Trends in adsorption of end-functionalized polystyrenes by thin layer chromatography. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 30 (1989) 154-155; C.A., 112 (1990) 8136a.

See also 513, 587.

38. INORGANIC COMPOUNDS

38a. Cations

- 698 Ajmal, M., Mohammad, A., Fatima, N. and Ahmad, J.: Aqueous alkali halides as impregnant and eluent in TLC of heavy metals: quantitative separation of Al, Ti, Fe, An, and Cd. *J. Liq. Chromatogr.*, 12 (1989) 3163-3191.
- 699 Gaibakyan, D.S., Karakhanyan, R.P. and Khachatryan, A.G.: (Study of complex formation of rhenium, molybdenum, vanadium, and tungsten in systems containing thiocyanic acid and separation of microquantities of these elements by paper chromatography). *Zh. Anal. Khim.*, 44 (1989) 754-756; C.A., 111 (1989) 246904u.
- 700 Grane, N., Martin, M.L., Santiago, J.M. and Berenguer, V.: (Separation and identification of frequent and nonfrequent iron-group cations by ascending paper chromatography). *Afinidad*, 46, No. 419 (1989) 11-16; C.A., 111 (1989) 24892t.
- 701 Ishida, K., Uchida, Y., Ninomiya, S. and Osawa, M.: Thin-layer chromatographic behaviour and separation of rare earths in silica gel - aqueous alkali metal chloride systems. *Fresenius' Z. Anal. Chem.*, 336 (1990) 419-422.
- 702 Jain, A., Singh, O.V. and Tandon, S.N.: Reversed-phase TLC behavior of lanthanides and some other elements using mono-(2-ethylhexyl) acid phosphate as an impregnant. *J. Planar Chromatogr.*, 3 (1990) 79-81.
- 703 Ma, G.: (Spectrophotometric determination of gold in electrolytic copper anode muds). *Yefin Fenxi*, 8 (1988) 52-53; C.A., 112 (1990) 47840v.
- 704 Medanic, Z., Eskinja, I., Turina, S. and Kovacicek, F.: (Zinc determination in aluminium alloys by thin-layer chromatography with anodic sampling). *Kem. Ind.*, 38 (1989) 305-314; C.A., 112 (1990) 29991r.

- 705 Misra, A.K. and Rajput, R.P.S.: Chromatography of some 's', 'p', and 'd' block metals on Zr(IV) phosphoantimonate thin layers in buffered EDTA solutions. *J. Planar Chromatogr.*, 3 (1990) 83-85.
- 706 Shimizu, T., Ohtomo, T. and Shimizu, T.: TLC of inorganic ions on polyethyleneimine cellulose in mixed sulfuric acid and dioxane media. *J. Planar Chromatogr.*, 3 (1990) 88-89.
- 707 Skvorc, I., Zambeli, N., Iskric, S. and Hadziija, O.: Thin-layer chromatography of some metals on silica gel layers impregnated with salicylic, syringic and *o*-phthalic acid. *J. Chromatogr.*, 498 (1990) 429.
- 708 Wang, Y. and Zeng, J.: (Separation and the extraction mechanism of rare earth elements by reversed-phase paper chromatography). *Zhongguo Xitu Xuebao*, 7 (1989) 53-57; *C.A.*, 112 (1990) 68798c.

See also 484, 489.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 709 Al-Janabi, M.A.A.: *In vitro* preparation of indium-113m (calcium)-phytate colloid for liver scintigraphy. *J. Labelled Compd. Radiopharm.*, 27 (1989) 1137-1141; *C.A.*, 112 (1990) 62592j.

See also 481, 508.

Electrophoresis

1. REVIEWS AND BOOKS

- 805 Pickering, M.V.: Capillary electrophoresis: proteins, problems, and promises. *LC-GC*, 7 (1989) 752-756; *C.A.*, 111 (1989) 211200p - a review with no refs.
- 806 Shiba, K.: *Graphic Electrophoresis, 1: Cellulose Acetate Membrane.* Ishiyaku Publishers, Tokyo, 1988, 174 p.

See also 829, 833, 876, 879, 892, 1020.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 807 Robert, G.O.: Analysis of electrophoresis performance. Final report, 21 Feb. 1985 - 30 Jun. 1988. *NASA Contract Rep.*, NASA-CR-183508, NAS1.26:183508, RAI-88-EL2-4, 1988, 102 p.; *C.A.*, 112 (1990) 47667u.
- 808 Westrop, R.: Capillary electrophoresis is here. *Lab. Pract.*, 38 (1989) 13-14; *C.A.*, 112 (1990) 69057x.

2b. Thermodynamics and theoretical relationships

- 809 McManigill, D. and Swedberg, S.A.: Factors affecting plate height in high performance zonal capillary electrophoreses (HPZCE). In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 468-478; *C.A.*, 111 (1989) 211367y.

See also 838.

2d. Measurement of physico-chemical and related values

- 810 Sallantin, M., Huet, J.-C., Demartean, C. and Pernollet, J.-C.: Reassessment of commercially available molecular weight standards for peptide sodium dodecyl sulfate-polyacrylamide gel electrophoresis using electroblotting and microsequencing. *Electrophoresis (Weinheim)*, 11 (1990) 34-36.
- 811 Sluyterman, L.A.A. and Kooistra, C.: Isotachophoretic phenomena on ion exchangers. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (1989) 123-130; *C.A.*, 111 (1989) 219975r.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 812 Nieman, E.: Gel electrophoretic system. *Eur. Pat. Appl.* EP 334,615 (Cl. G01N27/26), 27 Sep. 1989, ZA Appl. 88/2,019, 22 Mar. 1988; 23 p.; *C.A.*, 111 (1989) 224635j.
- 813 Ogawa, M. and Makino, Y.: Electrophoresis medium membrane, pulsed field electrophoresis method and apparatus. *Eur. Pat. Appl.* EP 320,937 (Cl. G01N27/26), 21 Jun. 1989, JP Appl. 87/320,708, 18 Dec. 1987; 17 pp.; *C.A.*, 112 (1990) 4064c.

- 814 Paul, A.E., Ferl, R.J., Boudreaux, G.S. and Miesel, M.W.: A simple optoelectronic device for controlling an electrophoresis apparatus. *Rev. Sci. Instrum.*, 60 (1989) 3072-3073; *C.A.*, 111 (1989) 240884v.
- 815 Spiker, S.: A vertical submarine electrophoresis apparatus for polyacrylamide minigels. *Anal. Biochem.*, 185 (1990) 270-273.

See also 819.

3b. Detection procedures and detectors

- 816 Cheng, Y.F. and Vo-Dinh, T.: Paper electrophoresis with surface-enhanced fluorimetric detection. *Anal. Chim. Acta*, 229 (1990) 295-297.
- 817 Filthuth, H.: Digital autoradiography - a new detector for radiochromatography and electrophoresis. In: Baillie, T.A., Jones, J.R. (Editors), *Synth. Appl. Isot. Labelled Cpd. 1988, Proc. Int. Symp. 1988*, Elsevier, Amsterdam, 1989, pp. 315-321; *C.A.*, 111 (1989) 211222x.
- 818 Filthuth, H.: A note on digital autoradiography: a new detector for radiochromatography and electrophoresis. In: Dallas, F.A.A. (Editor), *Recent Adv. Thin-Layer Chromatogr. 1987*, Plenum, New York, 1988, pp. 125-129; *C.A.*, 112 (1990) 48001j.
- 819 Hakim, B.H., Betrencourt, C. and Volochine, B.: Fully automated electrophoretic light-scattering device. *Rev. Sci. Instrum.*, 60 (1989) 3421-3424; *C.A.*, 112 (1990) 63468k.
- 820 Hogan, B.L. and Yeung, E.S.: Indirect fluorometric detection of tryptic digests separated by capillary zone electrophoresis. *J. Chromatogr. Sci.*, 28 (1990) 15-18.
- 821 Jay, G.D., Culp, D.J. and Jahneke, M.R.: Silver staining of extensively glycosylated proteins on sodium dodecyl sulfate-polyacrylamide gels: enhancement by carbohydrate-binding dyes. *Anal. Biochem.*, 185 (1990) 324-330.
- 822 Nakamura, S.: Sample encapsulation to facilitate application in gel electrophoresis. *Jpn. Kokai Tokkyo Koho JP 01,126,537 [89,126,537]* (Cl. G01N27/26), 18 May 1989, Appl. 87/284,650, 11 Nov. 1987; 3 pp.; *C.A.*, 112 (1990) 4063b.
- 823 Roth, R.I. and Levin, J.: Amplification of chromogenic staining of proteases within electrophoretic gels. *J. Biochem. Biophys. Methods*, 19 (1989) 129-141; *C.A.*, 111 (1989) 210822b.

See also 881, 888, 891, 893, 909, 943, 1031, 1086.

3c. Electrophoresis in stabilized media

- 824 Boschetti, E.: Polyacrylamide derivatives to the service of bioseparations. *J. Biochem. Biophys. Methods*, 19 (1989) 21-36; *C.A.*, 111 (1989) 228466c.
- 825 Kambara, H. and Nishikawa, T.: Method and apparatus for processing of used electrophoresis gels for reuse. *Ger. Offen. DE 3,821,454* (Cl. B01D57/02), 09 Feb. 1989, JP Appl. 87/155,447, 24 Jun. 1987; 6 pp.; *C.A.*, 111 (1989) 228580k.
- 826 Kozulic, B. and Mosbach, K.: Polymers containing N-acryloyltrimethylaminomethane (NAT) and their use as gels for electrophoresis. *PCT Int. Pat. Appl. WO 88 09,981* (Cl. G08F220/58), 15 Dec. 1988, GB Appl. 87/13,854, 13 Jun. 1987; 35 pp.; *C.A.*, 111 (1989) 211535b.
- 827 Peisker, K.: Fabric-reinforced ultrathin polyacrylamide gels for sodium dodecyl sulfate-polyacrylamide gel electrophoresis and isoelectric focusing. *Electrophoresis (Weinheim)*, 11 (1990) 152-155.
- 828 Righetti, P.G., Chiari, M., Casale, E., Chiesa, C., Jain, T. and Shorr, R.: HydroLink gel electrophoresis (HLGE). I. Matrix characterization. *J. Biochem. Biophys. Methods*, 19 (1989) 37-49; *C.A.*, 111 (1989) 211362t.

- 829 Righetti, P.G.: Of matrices and men. *J. Biochem. Biophys. Methods*, 19 (1989) 1-20; *C.A.*, 111 (1989) 170260n - a review with 79 refs.
- 830 Terai, F., Yukawa, K. and Suefuji, M.: Method of producing gradient gel medium for electrophoresis. *Eur. Pat. Appl.* EP 293,010 (Cl. G01N27/26), 30 Nov. 1988, *JF Appl.* 87/12i,504, 29 May 1987; 9 pp.; *C.A.*, 111 (1989) 211536c.

See also 882, 895, 1021, 1055, 1056, 1061, 1083.

4. SPECIAL TECHNIQUES

4a. Combination of electrophoretic techniques with chromatography

See 1086.

4b. Preparative and continuous procedures

See 980.

4c. Isoelectric focusing

- 832 Vaidya, N.R., Gothoskar, B.P. and Banerji, A.P.: Column isoelectric focusing in natural pH gradients generated by biological buffers. *Electrophoresis (Weinheim)*, 11 (1990) 156-161.

See also 834, 847, 1080.

4d. Isotachophoresis

See 811, 869, 873, 1072, 1082, 1084, 1085.

4e. Two dimensional electrophoresis

See 875, 886, 892, 894, 914, 915, 916, 922, 935, 937, 938, 954, 980, 1000.

4f. Affinity electrophoresis

- 833 Shimura, K. and Kasai, K.: (Two-dimensional affinophoresis). *Seibutsu Butsuri Kagaku*, 33 (1989) 171-176; *C.A.*, 112 (1990) 3664m - a review with 19 refs.

4g. Other special techniques

- 834 Altland, K.: IPGMAKER: A program for IBM-compatible personal computers to create and test recipes for immobilized pH gradients. *Electrophoresis (Weinheim)*, 11 (1990) 140-147.
- 835 Babich, Ya.A., Dukhin, S.S. and Tarovskii, A.A.: (Electrophoresis of the second kind in strong field). *Dokl. Akad. Nauk Ukr. SSR, Ser. B: Geol. Khim. Biol. Nauki*, (1989) 30-33; *C.A.*, 111 (1989) 240821x.
- 836 Brenner, N. and Osborne, J.C., Jr.: Separation of zone formation from electroosmotic impulse in tubular electrophoretic systems. *US. Pat.* US 4,859,301 (Cl.204-180.1; G01N27-28), 22 Aug. 1989, *Appl.* 238,676, 30 Aug. 1988; 5 p.; *C.A.*, 111 (1989) 224632f.
- 837 Inoue, S.: Supercritical fluid extraction in electrophoresis. *Jpn. Kokai Tokkyo Koho JP* 01 47,428 [89 47,428] (Cl. B01D57/02), 21 Feb. 1989, *Appl.* 87/205,368, 19 Aug. 1987; 5 pp.; *C.A.*, 111 (1989) 228584q.

- 838 Jansson, M., Emmer, A. and Roeraade, J.: Some design considerations in miniaturized electrokinetic separation systems. *J. High Resolut. Chromatogr.*, 12 (1989) 797-801.
- 839 Lux, J.A., Yin, H. and Schomburg, G.: Influence of polymer coating of capillary surfaces on migration behavior in micellar electrokinetic capillary chromatography. *J. High Resolut. Chromatogr.*, 13 (1990) 145-147.
- 840 Morris, M.D., Shick, R.A., Peck, K. and Parus, S.J.: Optoelectric electrophoresis analysis systems. *PCT Int. Appl. WO 89 01,620* (Cl. G01N21/00), 23 Feb. 1989, US Appl. 85,538, 13 Aug. 1987; 24 pp.; C.A., 112 (1990) 4025r.
- 841 Righetti, P.G., Ettori, C., Chafey, P. and Wahrmann, J.P.: Enantiomer resolution in immobilized pH gradient gels via inclusion of a chiral discriminator. *Electrophoresis (Weinheim)*, 11 (1990) 1-4.

See also 805, 809, 820, 871, 896, 956, 960, 1074.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 842 Al-Hakim, A. and Linhardt, R.J.: Isolation and recovery of acidic oligosaccharides from polyacrylamide gels by semi-dry electrotransfer. *Electrophoresis (Weinheim)*, 11 (1990) 23-28.
- 843 Scherz, H.: Thin-layer electrophoretic separation of monosaccharides, oligosaccharides and related compounds on reverse phase silica gel. *Electrophoresis (Weinheim)*, 11 (1990) 18-22.

See also 1023.

10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 844 Lyon, M. and Gallagher, J.T.: A general method for the detection and mapping of submicrogram quantities of glycosaminoglycan oligosaccharides on polyacrylamide gels by sequential staining with azure A and ammoniacal silver. *Anal. Biochem.*, 185 (1990) 63-70.
- 845 Min, H.: Electrophoretic analysis of glycosaminoglycans and vitreous hyaluronate. *Diss. Abstr. Int. B*, 50 (1989) 547; C.A., 111 (1989) 212384p.
- 846 Rykova, V.I., Kropotova, S.A., Kaledin, V.I. and Semenova, L.A.: (A comparative study of glycosaminoglycan composition in plasma membranes of normal and transformed mouse liver cells). *Biokhimiya (Moscow)*, 55 (1990) 439-444.

10c. Glycoproteins and their components

- 847 Bhattacharyya, L. and Brewer, C.F.: Isoelectric focusing studies of concanavalin A and the lentil lectin. *J. Chromatogr.*, 502 (1990) 131-142.
- 848 Kennel, S.J., Foote, L.J., Falcioni, R., Sonnenberg, A., Stringer, C.D., Crouse, C. and Hemler, M.E.: Analysis of the tumor-associated antigen TSP-180. Identity with $\alpha^6\beta^4$ in the integrin superfamily. *J. Biol. Chem.*, 264 (1989) 15515-15521.
- 849 Muramoto, K., Tanaka, H., Kimura, S., Kubo, K., Yoshihara, S., Yokoyama, M., Yoshida, Y. and Endo, M.: Quantitative determination of sulfated glycopeptide by two-dimensional electrophoresis on cellulose acetate membrane. *J. Biochem. Biophys. Methods*, 19 (1989) 75-82; C.A., 111 (1989) 170449f.

- 850 Parmelee, D.C., Benjamin, T., Niu, C.H. and Thorgeirsson, S.S.: Direct sequence analysis of rat liver membrane glycoproteins separated by 2D-PAGE. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 42-50; *C.A.*, 111 (1989) 211464c.
- 851 Scherberich, J.E., Birk, H.W. and Schoeppe, W.: Tubule-derived membrane glycoproteins in the urine of patients (including those with AIDS) as analysed by radioimmunoblotting. *J. Clin. Chem. Clin. Biochem.*, 28 (1990) 31-35.
- 852 Shimamura, M., Ohara, T., Imazawa, M. and Miyamoto, K.: Purification of myelin-associated glycoprotein from calf brain using high-performance liquid chromatography. *J. Chromatogr.*, 526 (1990) 535-539.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

See 1069.

11c. Lipids and their constituents

- 853 Trusova, V.M., Mezentsev, V.V., Ksenzenko, S.M., Tikhonov, A.N. and Molchanov, M.I.: (Phospholipids of thylakoid membranes of chloroplasts devoid of photo-system II). *Biokhimiya (Moscow)*, 55 (1990) 480-488.

11d. Lipoproteins and their constituents

- 854 Applebaum-Bowden, D., McLean, P., Steinmetz, A., Fontana, D., Matthys, C., Warnick, G.R., Cheung, M., Albers, J.J. and Hazzard, W.R.: Lipoprotein, apolipoprotein, and lipolytic enzyme changes following estrogen administration in postmenopausal women. *J. Lipid Res.*, 30 (1989) 1895-1906.
- 855 Bouma, M.E., Beucler, I., Pessah, M., Heinzmann, C., Lusi, A.J., Naim, H.Y., Ducastelle, T., Leluyer, B., Schmitz, J., Infante, R. and Aggerbeck, L.P.: Description of two patients with abetalipoproteinemia: synthesis of a normal-sized apolipoprotein B-48 in intestinal organ culture. *J. Lipid Res.*, 31 (1990) 1-15.
- 856 Dashti, N., Smith, E.A. and Alaupovic, P.: Increased production of apolipoprotein B and its lipoproteins by oleic acid in Caco-2 cells. *J. Lipid Res.*, 31 (1990) 113-123.
- 857 Dergunov, A.D., Shuvaev, V.V. and Perova, N.V.: (The dynamic behaviour of human plasma very low density lipoprotein apoproteins and regulation of lipolysis). *Biokhimiya (Moscow)*, 55 (1990) 134-146.
- 858 Ferrell, R.E., Sepehrnia, B., Kamboh, M.I. and VandeBerg, J.L.: Highly polymorphic apolipoprotein A-IV locus in the baboon. *J. Lipid Res.*, 31 (1990) 131-135.
- 859 France, D.S., Hughes, T.E., Miserendino, R., Spirito, J.A., Babiak, J., Eskesen, J.B., Tapparelli, C. and Paterniti, J.R., Jr.: Nonimmunochemical quantitation of mammalian apolipoprotein A-I in whole serum or plasma by nonreducing gel electrophoresis. *J. Lipid Res.*, 30 (1989) 1997-2004.
- 860 Holmquist, L.: Identification and quantification of apolipoprotein D in normal human urine. *Electrophoresis (Weinheim)*, 11 (1990) 93-94.
- 861 Huang, L.-S., Ripps, M.E. and Breslow, J.L.: Molecular basis of five apolipoprotein B gene polymorphisms in noncoding regions. *J. Lipid Res.*, 31 (1990) 71-77.
- 862 Junker, L.H. and Davis, R.A.: Receptor-mediated uptake of low density lipoprotein stimulates bile acid synthesis by cultured rat hepatocytes. *J. Lipid Res.*, 30 (1989) 1933-1941.

- 863 Le Boeuf, R.C., Doolittle, M.H., Montcalm, A., Martin, D.C., Reue, K. and Lusis, A.J.: Phenotypic characterization of the ATH-1 gene controlling high density lipoprotein levels and susceptibility to atherosclerosis. *J. Lipid Res.*, 31 (1990) 91-101.
- 864 Mailly, F., Davignon, J. and Nestruck, A.C.: Analytical isoelectric focusing with immobilized pH gradients of human apolipoprotein E from very low density lipoproteins and total plasma. *J. Lipid Res.*, 31 (1990) 149-155.
- 865 Smit, M., de Knijff, P., van der Kooij-Meijls, E., Groenendijk, C., van den Maagdenberg, A.M.J.M., Gevers Leuven, J.A., Stalenhoef, A.F.H. et al.: Genetic heterogeneity in familial dysbetalipoproteinemia. The E2(Lys 146-Gln) variant results in a dominant mode of inheritance. *J. Lipid Res.*, 31 (1990) 45-53.

See also 919.

13. STEROIDS

13c. Sterols

- 866 Salcedo, J., Delgado, A. and Gonzales-Caballero, F.: The effect of pH on the electrical properties of the cholesterol/water interface. *J. Colloid Interface Sci.*, 133 (1989) 278-281; *C.A.*, 111 (1989) 240909g.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 867 Chiari, M., Manzocchi, A. and Righetti, P.G.: Formation of a cysteine-acrylamide adduct in isoelectric focusing gels. *J. Chromatogr.*, 500 (1990) 697-704.
- 868 Durupthy, O., Durupthy, A. and Dubusc, M.: (Paper chromatography and electrophoresis of α -amino acids). *Bull. Union Physiciens*, 83 (1989) 622-628; *C.A.*, 112 (1990) 6759g.
- 869 Mizobuchi, N., Ageta, T. and Kodama, H.: Determination of N-monoacetylcystathionine in biological samples using isotachophoresis. *J. Chromatogr.*, 526 (1990) 550-555.

See also 841.

18b. Peptides and peptidic and proteinous hormones

- 870 Fischer, W.H., Vaughan, J., Karr, D., McClintock, R., Spiess, J., Rivier, J. and Vale, W.: Recovery of biological activity, blotting and sequence analysis of protein hormones after SDS-PAGE. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 36-41; *C.A.*, 111 (1989) 167499s.
- 871 Hecht, R.I., Coleman, J.F., Jr., Morris, J.C., Stover, F.S. and Demarest, C.: Micropreparative capillary zone electrophoresis of recombinant human interleukin-3. *Prepar. Biochem.*, 19 (1989) 363-366.
- 872 Kalmokoff, M.L., Karnauchow, T.M. and Jarell, K.F.: Conserved N-terminal sequences in the flagellins of archaebacteria. *Biochem. Biophys. Res. Commun.*, 167 (1990) 154-160.
- 873 Stehle, P., Monter, B., Bahsitta, W.-P. and Fürst, P.: Analytical control of protease catalyzed synthesis of dipeptides - capillary isotachophoresis vs. reversed phase high performance liquid chromatography. *J. Liq. Chromatogr.*, 12 (1989) 2821-2838.

- 874 Urios, P. and Cittanova, N.: Adaptation of fluorescence polarization immunoassay to the assay of macromolecules. *Anal. Biochem.*, 185 (1990) 308-312.

See also 945, 976.

18c. Elucidation of structure of proteins and enzymes

- 875 Bauw, G., van Damme, J., Fuype, M., Vandekerckhove, J., Gesser, B., Ratz, G.P., Lauridsen, J.B. and Celis, J.E.: Protein-electroblotting and -microsequencing strategies in generating protein data bases from two-dimensional gels. *Proc. Natl. Acad. Sci. U.S.A.*, 86 (1989) 7701-7705; *C.A.*, 112 (1990) 3878j.
- 876 Hirano, H.: (Microsequence analysis of proteins separated by gel electrophoresis). *Seibutsu Butsuri Kagaku*, 33 (1989) 165-170; *C.A.*, 112 (1990) 3663k - a review with 24 refs.
- 877 Kikuta, Y., Kusunose, E., Okumoto, T., Kubota, I. and Kusunose, M.: Purification and characterization of two forms of cytochrome P-450 with ω -hydroxylase activities toward prostaglandin A and fatty acids from rabbit liver microsomes. *J. Biochem. (Tokyo)*, 107 (1990) 280-286.
- 878 Moskalenko, A.A., Erokhin, Yu.E. and Kuznetsova, N.Yu.: (Effect of trypsin on aggregated photosystem II enriched particles). *Biokhimiya (Moscow)*, 55 (1990) 240-246.
- 879 Wittmann-Liebold, B.: Moderne chemische Strukturanalyse von Proteinen und Peptiden: Strategien, Methoden, Einsatzmöglichkeiten. *Kontakte (Darmstadt)*, No.3 (1989) 16-28 - a review with 60 refs.
- 880 Yates, J.R., Shabanowitz, J., Griffin, P.R., Zhu, N.Z. and Hunt, D.F.: Sequence analysis of protein C-terminal proteolytic fragments, protein isolated from 2D-gels, and murine calbindin: new methodology. In: Hugli, T.E. (Editor), *Tech. Protein Chem.*, Academic, San Diego, 1989, pp. 168-175; *C.A.*, 111 (1989) 228475e.

See also 820, 900, 969.

19. PROTEINS

19a. General techniques

- 881 Ali, R. and Sayeed, S.A.: A novel dye for staining electrophoretically resolved proteins. In: Zaidi, Z.H., *Protein Struct.-Funct. Relat., Proc. Int. Symp.*, Elsevier, Amsterdam, 1988, pp. 15-26; *C.A.*, 111 (1989) 170444a.
- 882 Alvarez, L. and Nieto, A.: Electrophoresis of native proteins on agarose gels containing volatile buffers. *Anal. Lett.*, 22 (1989) 2737-2745.
- 883 Bier, M.: The lasting and the ephemeral virtues of electrophoresis. *UCLA Symp. Mol. Cell. Biol., New Ser.*, 80 (1989) 237-240; *C.A.*, 111 (1989) 170447d.
- 884 Christy, K.G., Jr., LaTart, D.B. and Osterhoudt, H.W.: Modifications for SDS-PAGE of proteins. *BioTechniques*, 7 (1989) 692-693; *C.A.*, 111 (1989) 228380v.
- 885 Desvaux, F.X., David, B. and Peltre, G.: Multiple successive immunoprinting: a fast blotting technique of a single agarose isoelectric focusing gel. *Electrophoresis (Weinheim)*, 11 (1990) 37-41.
- 886 Goldfarb, M.F.: Two-dimensional electrophoresis with silver staining: a sensitive method for analyzing complex protein mixtures. *Lect. Notes Coastal Estuarine Stud.*, 25 (Immunochem. Approaches Coastal, Estuarine Oceanogr. Quest.) (1988) 130-133; *C.A.*, 111 (1989) 228385a.

- 887 Hayes, J.D. and Stockman, P.K.: Electrophoresis of proteins and nucleic acids: I. Theory. *Br. Med. J.*, 299 (1989) 843-846; *C.A.*, 112 (1990) 3662j - review with 16 refs.
- 888 Helbing, A.R. and Jansen, J.W.: Application of colloidal gold staining in protein electrophoresis: an improvement? *Tijdschr. Ned. Ver. Klin. Chem.*, 14 (1989) 165-169; *C.A.*, 111 (1989) 228302w.
- 889 Inczedy-Marcsek, M., Lindner, E., Haessler, R., Zwack-Megele, G., Roisen, F. and Yorke, G.: Extraction of proteins within ultrathin-layer polyacrylamide electrophoresis (SDS-PAGE) and isoelectric focusing (PAGIF) of cryostat sections and tissue culture specimens. *Acta Histochem., Suppl.*, 36 (1988) 377-394; *C.A.*, 111 (1989) 170445b.
- 890 Jacobson, G. and Karsnäs, P.: Important parameters in semi-dry electrophoretic transfer. *Electrophoresis (Weinheim)*, 11 (1990) 46-52.
- 891 Neuhoﬀ, V., Stamm, R., Pardowitz, I., Arold, N., Ehrhardt, W. and Taube, D.: Essential problems in quantification of proteins following colloidal staining with Coomassie Brilliant Blue dyes in polyacrylamide gels, and their solution. *Electrophoresis (Weinheim)*, 11 (1990) 101-117.
- 892 Ohashi, M., Toda, T., Fujita, T. and Yoshida, M.: (Two-dimensional electrophoresis of native proteins on cellulose acetate membranes). *Seibutsu Butsuri Kagaku*, 33 (1989) 177-180; *C.A.*, 112 (1990) 3665n - a review with 12 refs.
- 893 Pinon, J.-M., Puygauthier-Toubas, D., Lèpan, H., Marx, C., Bonhomme, A., Boulant, J., Geers, R. and Dupont, H.: Rapid detection of proteins by enzyme-linked immunofiltration assay after transfer onto nitrocellulose membranes. *Electrophoresis (Weinheim)*, 11 (1990) 41-45.
- 894 Rabilloud, T., Gianazza, E., Catto, N. and Righetti, P.G.: Amidodisulfobetaines, a family of detergents with improved solubilization properties: application for isoelectric focusing under denaturing conditions. *Anal. Biochem.*, 185 (1990) 94-102.
- 895 Sugihara, M. and Ogawa, M.: Acrylamide polymer-containing media for gel electrophoresis of proteins. *Jpn. Kokai Tokkyo Koho JP 01 21,346* [89 21,346] (Cl. G01N27/26), 24 Jan. 1989, Appl. 87/177,979, 16 Jul. 1987, 9 pp.; *C.A.*, 111 (1989) 228581m.
- 896 Swedberg, S.A.: Characterization of protein behavior in high-performance capillary electrophoresis using a novel capillary system. *Anal. Biochem.*, 185 (1990) 51-56.

See also 810.

19b. Proteins of cells, viruses and subcellular particles

- 897 Berdinskikh, N.K., Korol', D.R. and Livshits, V.I.: (The cell membrane protein modification during the N-nitrosodiethylamine induced hepatocarcinogenesis). *Biokhimiya (Moscow)*, 55 (1990) 432-438.
- 898 Crooks, A.J., Lee, J.M., Dowsett, A.B. and Stephenson, J.R.: Purification and analysis of infectious virions and native non-structural antigens from cells infected with tick-borne encephalitis virus. *J. Chromatogr.*, 502 (1990) 59-68.
- 899 Prescott, M. and McLennan, A.G.: Synthesis and applications of 8-azido photoaffinity analogs of P^1, P^3 -bis(5'-adenosyl)triphosphate and P^1, P^4 -bis(5'-adenosyl)tetraphosphate. *Anal. Biochem.*, 184 (1990) 330-337.
- 900 Walpita, P., Connor, J.D. and Pfeifer, D.: Protein fingerprinting: a novel virus identification system. *J. Virol. Methods*, 25 (1989) 315-324; *C.A.*, 112 (1990) 3883g.
- 901 Warlow, R.S. and Bernard, C.C.A.: Improved detection of lymphocyte membrane proteins in purified form and as a crude mixture using native and denaturing polyacrylamide gel electrophoresis by optimisation of Coomassie Brilliant Blue and silver staining. *Electrophoresis (Weinheim)*, 11 (1990) 53-60.

- 902 Yokota, H. and Yuasa, A.: Increase of a form of UDP-glucuronyltransferase glucuronizing various phenolic xenobiotics and the corresponding translatable mRNA in 3-methylcholanthrene-treated rat liver. *J. Biochem. (Tokyo)*, 107 (1990) 92-96.
- 903 Zu, Y., Kohno, M., Kubota, I., Nishida, E., Hanaoka, M. and Namba, Y.: Characterization of interleukin 2 stimulated 65-kilodalton phosphoprotein in human T cells. *Biochemistry*, 29 (1990) 1055-1062.

See also 964, 979.

19c. Proteins synthesized by genetic manipulation

- 904 Nishitani, H., Kobayashi, H., Ohtsubo, M. and Nishimoto, T.: Cloning of *Xenopus* RCC1 cDNA, a homolog of the human RCC1 gene: complementation of tsBN2 mutation and identification of the product. *J. Biochem. (Tokyo)*, 107 (1990) 228-235.
- 905 Ohnishi, K., Nakazima, A., Matasubara, K and Kiritani, K.: Cloning and nucleotide sequences of livB and livC, the structural genes encoding binding proteins of the high-affinity branched-chain amino acid transport in *Salmonella typhimurium*. *J. Biochem. (Tokyo)*, 107 (1990) 202-208.
- 906 Righetti, P.G., Wenisch, E., Jungbauer, A., Katinger, H. and Faupel, M.: Preparative purification of human monoclonal antibody isoforms in a multi-compartment electrolyser with immobilized membranes. *J. Chromatogr.*, 500 (1990) 681-696.
- 907 Shing, Y., Folkman, J., Weisz, P.B., Joullie, M.M. and Ewing, W.R.: Affinity of fibroblast growth factors for β -cyclodextrin tetradecasulfate. *Anal. Biochem.*, 185 (1990) 108-111.

19d. Microbial and plant proteins

- 908 Bramanti, T.E. and Holt, S.C.: Iron-regulated outer membrane proteins in the periodontopathic bacterium, *Bacteroides gingivalis*. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1146-1154.
- 909 Clements, R.L.: Alternative methods for fixing and staining gliadins in polyacrylamide gels. *Electrophoresis (Weinheim)*, 11 (1990) 123-128.
- 910 Fukumori, F., Takeuchi, N., Hagiwara, T., Ohbayashi, H., Endo, T., Kochibe, N., Nagata, Y. and Kobata, A.: Primary structure of a fucose-specific lectin obtained from a mushroom, *Aleuria aurantia*. *J. Biochem. (Tokyo)*, 107 (1990) 190-196.
- 911 Gueguen, J., Bollecker, S., Schwenke, K.D. and Raab, B.: Effect of succinylation on some physicochemical and functional properties of the 12S storage protein from rapeseed (*Brassica napus* L.). *J. Agric. Food Chem.*, 38 (1990) 61-69.
- 912 Hussain, A., Scanlon, M.G., Juliano, B.O. and Bushuk, W.: Discrimination of rice cultivars by polyacrylamide gel electrophoresis and high-performance liquid chromatography. *Cereal Chem.*, 66 (1989) 353-356; *C.A.*, 111 (1989) 211442u.
- 913 Odintsova, T.I. and Egorov, T.A.: (N-Terminal sequences of ω -gliadins of *Aegilops longissima*: on the origin of polyploid wheat genomes). *Biokhimiya (Moscow)*, 55 (1990) 509-516.
- 914 Patton, W.F., Phanak, M.R. and Jacobson, B.S.: Analysis of plasma membrane protein changes in *Dictyostelium discoideum* during concanavalin A induced receptor redistribution using two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 11 (1990) 79-85.
- 915 Roggero, P. and Pennazio, S.: Two-dimensional polyacrylamide gel electrophoresis of extracellular soybean pathogenesis-related proteins using PhastSystem. *Electrophoresis (Weinheim)*, 11 (1990) 86-90.

- 916 Yoon, H.H. and Kim, S.H.: Two-dimensional electrophoretic analysis of rice proteins. *Han'guk Nonghwa Hakhoechi*, 32 (1989) 85-90; *C.A.*, 111 (1989) 230986r.

See also 1004.

19e. *Proteins of blood, serum and blood cells*

- 917 Cianflone, K., Wu, H., Walsh, M., Baldu, A. and Sniderman, A.: Metabolic response of acylation stimulating protein to an oral fat load. *J. Lipid Res.*, 30 (1989) 1727-1733.
- 918 Fernandez, N., Labeta, M., Kurpisz, M., Nakatsuji, T., Sachs, J. and Festenstein, H.: Major histocompatibility complex (MHC) protein analysis by optimised two-dimensional electrophoretic methods. *Electrophoresis (Weinheim)*, 11 (1990) 167-174.
- 919 Hussain, M.M. and Zannis, V.I.: Intracellular modification of human apolipoprotein AII (ApoAII) and sites of ApoAII mRNA synthesis: comparison of ApoAII with ApoCII and ApoCIII isoproteins. *Biochemistry*, 29 (1990) 209-217.
- 920 Klein, M.L., Bartley, T.D., Davis, J.M., Whiteley, D.W. and Lu, H.S.: Isolation and structural characterization of three isoforms of recombinant consensus α interferon. *Arch. Biochem. Biophys.*, 276 (1990) 531-537.
- 921 Monokovic, D.D. and Tracy, P.B.: Activation of human factor V by factor Xa and thrombin. *Biochemistry*, 29 (1990) 1118-1128.
- 922 Polotskaya, A.V. and Adler, V.V.: (Regulation by glucocorticoids of proliferative and protein-synthesizing reactions in human peripheral blood lymphocytes). *Biokhimiya (Moscow)*, 55 (1990) 102-108.
- 923 Silvestrini, B., Guglielmotti, A., Saso, L., Milanese, C., Melanitou, E., Grima, J. and Cheng, C.Y.: Development of an enzyme-linked immunosorbent assay with a monoclonal antibody prepared against α_1 -antitrypsin for diagnostic screening of inflammatory disorders. *Clin. Chem. (Winston-Salem)*, 36 (1990) 277-282.
- 924 Sinosich, M.J.: Molecular characterization of pregnancy-associated plasma protein-A by electrophoresis. *Electrophoresis (Weinheim)*, 11 (1990) 70-78.
- 925 Suzuki, Y., Yoshida, K., Ichimiya, T., Yamamoto, T. and Sinahara, H.: Trypsin inhibitors in guinea pig plasma: isolation and characterization of contrapsin and two isoforms of α -1-antiproteinase and acute phase response of four major trypsin inhibitors. *J. Biochem. (Tokyo)*, 107 (1990) 173-179.
- 926 Tahir, M.A., Kimes, D.R. and Sarwar, M.: An alternative buffer system for group specific Gc component. *Sci. Int. (Lahore)*, 1 (1989) 281-282; *C.A.*, 111 (1989) 210198g.
- 927 Thymann, M., Svensmark, O., Masumba, G., Brokso, H. and Skibsby, L.B.: Haptoglobin subtype determination by isoelectric focusing in agarose gel: application to paternity testing and presentation of a new α_2 -variant. *Electrophoresis (Weinheim)*, 11 (1990) 61-65.

See also 874, 973, 1013.

19f. *Structural and muscle proteins*

- 928 Brennan, M.: Changes in the cross-linking of collagen from rat tail tendons due to diabetes. *J. Biol. Chem.*, 264 (1989) 20953-20960.
- 929 Chang, H.W., Bock, E. and Bonilla, E.: Dystrophin in electric organ of *Torpedo californica* homologous to that in human muscle. *J. Biol. Chem.*, 264 (1989) 20831-20834.
- 930 Fujii, T., Maachino, K., Andoh, H., Satoh, T. and Kondo, Y.: Calcium-dependent control of caldesmon-actin interaction by S100 protein. *J. Biochem. (Tokyo)*, 107 (1990) 133-137.

- 931 Grabarek, Z. and Gergely, J.: Zero-length crosslinking procedure with the use of active esters. *Anal. Biochem.*, 185 (1990) 131-135.
- 932 Hironi, M., Tanaka, R. and Watanabe, Y.: *Tetrahymena* actin: copolymerization with skeletal muscle actin and interactions with muscle actin-binding proteins. *J. Biochem. (Tokyo)*, 107 (1990) 32-36.
- 933 Kobayashi, R., Tashima, Y. and Yoshioka, N.: The 30 kDa abnormal protein in avian dystrophic muscle is indistinguishable from carbonic anhydrase III by physicochemical, immunological, and enzymological criteria. *J. Biochem. (Tokyo)*, 107 (1990) 56-60.
- 934 Kobayashi, R., Toyoshima, I., Masamune, O. and Tashima, Y.: Detection and isolation of a 30 kDa abnormal protein in avian dystrophic muscle. *J. Biochem. (Tokyo)*, 107 (1990) 51-55.
- 935 Kojima, T., Sano, K., Hirabayashi, T. and Obinata, T.: Characterization of C-protein isoforms expressed in developing, denervated, and dystrophic chicken skeletal muscles by two-dimensional gel electrophoresis. *J. Biochem. (Tokyo)*, 107 (1990) 470-475.
- 936 Oshishi, I., Morikawa, Y. and Baba, T.: ADP-ribosylation of nonmuscle actin by component I of botulinum C₂ toxin inactivates the ability to interact with unmodified actin. *J. Biochem. (Tokyo)*, 107 (1990) 420-425.
- 937 Pulyaeva, E.V., Kovalae, L.I., Tsvetkova, M.N., Shishkin, S.S. and Boldyrev, N.I.: (A two-dimensional map of human heart left ventricular proteins). *Biokhimiya (Moscow)*, 55 (1990) 489-498.
- 938 Sakai, Y., Okamoto, H., Mogami, K., Yamada, T. and Hotta, Y.: Actin with tumor-related mutation is antimorphic in *Drosophila* muscle: two distinct modes of myofibrillar disruption by antimorphic actins. *J. Biochem. (Tokyo)*, 107 (1990) 499-505.
- 939 Singer, H.A., Oren, J.W. and Benscoter, H.A.: Myosin light chain phosphorylation in ³²P-labeled rabbit aorta stimulated by phorbol 12,13-dibutyrate and phenylephrine. *J. Biol. Chem.*, 264 (1989) 21215-21222.
- 940 Villanueva, M.A., Ho, S.-C. and Wang, J.L.: Isolation and characterization of one isoform of actin from cultured soybean cells. *Arch. Biochem. Biophys.*, 277 (1990) 35-41.
- 941 Wilkinson, M.J., Cohen, R.L. and Shuman, M.A.: A nonradioactive assay for type IV collagen degradation. *Anal. Biochem.*, 185 (1990) 294-296.
- 19g. *Protamines, histones and other chromosomal proteins*
- 942 Baci, P.C. and Durham, J.P.: A procedure for the extraction and high resolution two-dimensional gel electrophoresis of total nuclear phosphoproteins from isotopically purified nuclei. *Electrophoresis (Weinheim)*, 11 (1990) 162-166.
- 943 Csordas, A., Pedrini, M. and Grunicke, H.: Suitability of staining techniques for the detection and quantitation of nonhistone high mobility group proteins. *Electrophoresis (Weinheim)*, 11 (1990) 118-123.
- 944 Jackson, V.: *In vivo* studies on the dynamics of histone-DNA interaction: evidence for nucleosome dissolution during replication and transcription and a low level of dissolution independent of both. *Biochemistry*, 29 (1990) 719-731.
- 945 Vanfleteren, J.R., van Bun, S.M., de Baere, I. and van Beeumen, J.J.: The primary structure of a minor isoform (H1.2) of histone H1 from the nematode *Caenorhabditis elegans*. *Biochem. J.*, 265 (1990) 739-746.
- 19h. *Chromoproteins and metalloproteins*
- 946 Bonfils, C. and Combalbert, J.: Resolution of rabbit liver cytochrome P450 3b and 3c (P450 IIC3 and IIIA4) from microsomal membrane by two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 11 (1990) 182-186.

- 947 Hinkamp, S. and Dunemann, L.: Electrophoretic techniques in element species analysis. Comparison between flow-through and flat-bed techniques. *Anal. Chim. Acta*, 230 (1990) 1-7.
- 948 Kobayashi, R., Hidaka, H. and Tashima, Y.: Purification, characterization, and partial sequence analysis of 32-kDa calcimedin from chicken gizzard. *Arch. Biochem. Biophys.*, 277 (1990) 203-210.
- 949 Shayiq, R.M. and Avadhani, N.G.: A phenobarbital-inducible hepatic mitochondrial cytochrome P-450 immunochemically related to microsomal P-450b. *Biochemistry*, 29 (1990) 866-873.

19i. *Proteins of glands, gland products, various zymogens (including milk proteins)*

- 950 Yosida, H., Ono, M., Manabe, T. and Okuyama, T.: (Analysis of human seminal proteins by micro two dimensional electrophoresis. II.) *Seibutsu Butsuri Kagaku*, 33 (1989) 119-124; *C.A.*, 111 (1989) 211365w.

19j. *Proteins of brain, cerebrospinal fluid and eye*

- 951 Elizarov, S.M. and Simirsky, V.N.: (A CAMP-independent protein kinase form amphibian eye lens: identification, distribution and phosphorylation substrates). *Biokhimiya (Moscow)*, 55 (1990) 554-563.
- 952 Fujii, T., Nakamura, A., Ogoma, Y., Kondo, Y. and Arai, T.: Selective purification of microtubule-associated proteins 1 and 2 from rat brain using poly(L-aspartic acid). *Anal. Biochem.*, 184 (1990) 268-273.
- 953 Lacey, E. and Snowdon, K.L.: Isolation of mammalian brain tubulin by amino-activated gel chromatography. *J. Chromatogr.*, 525 (1990) 71-84.
- 954 Mattila, K., Pirttilä, T. and Frey, H.: Horizontal two-dimensional electrophoresis of cerebrospinal fluid proteins with immobilized pH gradients in the first dimension. *Electrophoresis (Weinheim)*, 11 (1990) 91-92.
- 955 Semple-Rowland, S.L. and Ulshafer, R.J.: Analysis of proteins in developing RD (retinal degeneration) chick retina using two-dimensional gel electrophoresis. *Exp. Eye Res.*, 49 (1989) 665-675; *C.A.*, 111 (1989) 228397f.
- 956 Wallingford, R.A.: Development of capillary zone electrophoresis for the analysis of neuronal microenvironments. Avail. *Univ. Microfilms Int.*, order No. DA8910079; *C.A.*, 111 (1989) 228398g.

19k. *Proteins of neoplastic tissue and transformed cells*

- 957 Prasad, S.C., Thraves, P.J., Bhatia, K.G., Smulson, M.E. and Dritschilo, A.: Enhanced poly(adenosine diphosphate ribose) polymerase activity and gene expression in Ewing's sarcoma cells. *Cancer Res.*, 50 (1990) 38-43.

See also 848.

19l. *Specific binding and receptor proteins*

- 958 Barbry, P., Chassande, O., Marsault, R., Lazdunski, M. and Frelin, C.: [³H]Phenamil binding protein of the renal epithelium Na⁺ channel. Purification, affinity labeling, and functional reconstitution. *Biochemistry*, 29 (1990) 1039-1045.
- 959 Bresnick, E.H., Dalman, F.C. and Pratt, W.B.: Direct stoichiometric evidence that the untransformed M_r 300 000, 9S, glucocorticoid receptor is a core unit derived from a larger heteromeric complex. *Biochemistry*, 29 (1990) 520-527.
- 960 Chan, K.-F.J. and Chen, W.H.: High-performance capillary electrophoresis of calmodulin. *Electrophoresis (Weinheim)*, 11 (1990) 15-18.

- 961 Hu, W., Mazurier, J., Montreuil, J. and Spik, G.: Isolation and partial characterization of a lactotransferrin receptor from mouse intestinal brush border. *Biochemistry*, 29 (1990) 535-541.
- 962 Ito, H., Yamamoto, H., Kimura, Y., Kambe, H., Okochi, T. and Kishimoto, S.: Affinity chromatography of human plasma gelsolin with polyphosphate compounds on immobilized Cibacron Blue F3GA. *J. Chromatogr.*, 526 (1990) 397-406.
- 963 Kadowaki, H., Symanski, L.A., Rys-Sikora, K.E. and Koff, R.S.: Lactosylceramide molecular species specificity of rat liver CMP-N-acetylneuraminat:lactosylceramide sialyltransferase. *J. Lipid Res.*, 30 (1989) 1789-1797.
- 964 Korner, M., Bellan, A.H., Brini, A.T. and Farrar, W.L.: Characterization of the human immunodeficiency virus type 1 enhancer binding proteins from the human T-cell line Jurkat. *Biochem. J.*, 265 (1990) 547-554.
- 965 Lin, S.M., Hopfer, S.M., Brennan, S.M. and Sunderman, F.W., Jr.: Protein blotting method for detection of nickel-binding proteins. *Res. Commun. Chem. Pathol. Pharmacol.*, 65 (1989) 275-288; *C.A.*, 112 (1990) 3986t.
- 966 Miedel, M.C., Hulmes, J.D. and Pan, Y.E.: Limited proteolysis of recombinant human soluble interleukin-2 receptor. Identification of an interleukin-2 binding core. *J. Biol. Chem.*, 264 (1989) 21097-21105.
- 967 Mohammad, F. and Farber, E.: *In situ* detection of DNA-binding proteins in herpes simplex virus type 1-infected cells. *Acta Virol.*, 33 (1989) 401-409; *C.A.*, 112 (1990) 3879K.
- 968 Pandian, M.R., Morgan, C., Nelson, J.C. and Fisher, D.A.: Differentiating various abnormalities of thyroxin binding to serum proteins by radioelectrophoresis of thyroxin and immunoassay of binding proteins. *Clin. Chem. (Winston-Salem)*, 36 (1990) 457-461.
- 969 Sunahara, R.K., Murphy, B.J. and Tuana, B.S.: A sensitive and rapid method for identification and characterization of low abundance receptors. *Anal. Biochem.*, 185 (1990) 143-146.
- 970 Yazawa, M., Matsuzawa, F. and Yagi, K.: Inter-domain interaction and the structural flexibility of calmodulin in the connecting region of the terminal two domains. *J. Biochem. (Tokyo)*, 107 (1990) 287-291.

See also 976.

19m. Urinary proteins

- 971 Beilby, J.P. and O'Leary, B.A.: α_1 -Acid glycoprotein decreases recovery of total protein in urine when trichloroacetic acid is used to precipitate the proteins. *Clin. Chem. (Winston-Salem)*, 36 (1990) 565-567.
- 972 Boege, F., Koehler, B. and Liebermann, F.: Identification and quantification of Bence-Jones proteinuria by automated nephelometric screening. *J. Clin. Chem. Clin. Biochem.*, 28 (1990) 37-42.

19n. Other proteins

- 973 Chao, S., Chai, K.X., Chao, L. and Chao, J.: Molecular cloning and primary structure of rat α_1 -antitrypsin. *Biochemistry*, 29 (1990) 323-329.
- 974 Huet, G., Balduyck, M., Mizon, C., Colombel, J.F. and Mizon, J.: Fecal α_1 -antichymotrypsin in inflammatory bowel diseases. *Clin. Chem. (Winston-Salem)*, 36 (1990) 576.
- 975 Pietron, W., Gieryng, R. and Pirog, Z.: (Electrophoretic characteristics of soluble proteins of *Fasciola hepatica* (Linnaeus, 1758)). *Wiad. Parazytol.*, 34 (1988) 263-277; *C.A.*, 112 (1990) 3877h.
- 976 Yannariello-Brown, J. and Madri, J.A.: A 48 kDa collagen-binding phosphoprotein isolated from bovine aortic endothelial cells interacts with the collagenous domain, but not the globular domain, of collagen type IV. *Biochem. J.*, 265 (1990) 383-392.

See also 1014.

20. ENZYMES AND ENZYME ACTIVITY ESTIMATION

- 977 Chippindale, P.: A high pH discontinuous buffer system for resolution of isozymes in starch gel electrophoresis. *Stain Technol.*, 64 (1989) 61-64; C.A., 112 (1990) 3098e.

20a. Oxidoreductases

- 978 Iwata, N., Inazu, N. and Satoh, T.: Immunological and enzymological localization of carbonyl reductase in ovary and liver of various species. *J. Biochem. (Tokyo)*, 107 (1990) 209-212.
- 979 Kim, S.G. and Novak, R.F.: Induction of rat hepatic P450IIE1 (CYP 2E1) by pyridine: evidence for a role of protein synthesis in the absence of transcriptional activation. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1072-1079.
- 980 Kocha, T., Fukuda, T., Isobe, T. and Okuyama, T.: Large-scale purification of bovine brain lactate dehydrogenase by affinity chromatography on immobilized colchicine. *J. Biochem. (Tokyo)*, 107 (1990) 138-143.
- 981 Loseva, L.P., Shatilov, V.R. and Kretovich, W.L.: (An immunological analysis of glutamate dehydrogenases from unicellular green algae). *Biokhimiya (Moscow)*, 55 (1990) 408-415.
- 982 Masson, P., Arciero, D.M., Hooper, A.B. and Balny, C.: Electrophoresis at elevated hydrostatic pressure of the multiheme hydroxylamine oxidoreductase. *Electrophoresis (Weinheim)*, 11 (1990) 128-133.
- 983 Tahir, M.A. and Day, J.R.: Alternative buffer system for lactate dehydrogenase (LDH) electrophoresis utilized in the identification of menstrual blood. *Sci. Int. (Lahore)*, 1 (1989) 215-218; C.A., 111 (1989) 168817f.

See also 1019.

20b. Transferases (excluding E.C. 2.7.-.-)

- 984 Chang, L., Chuang, L., Tsai, C., Tu, C.D. and Tam, M.F.: Characterization of glutathione S-transferases from day-old chick livers. *Biochemistry*, 29 (1990) 744-750.
- 985 Inglese, J., Johnson, D.L., Shiau, A., Smith, J.M. and Benkovic, S.J.: Subcloning, characterization, and affinity labeling of *Escherichia coli* glycylamide ribonucleotide transformylase. *Biochemistry*, 29 (1990) 1436-1443.
- 986 Oshino, R., Kamei, K., Nishioka, M. and Shin, M.: Purification and characterization of glutathione S-transferases from guinea pig liver. *J. Biochem. (Tokyo)*, 107 (1990) 105-110.

20c. Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)

- 987 Alcázar, A., Méndez, E., Martín, M.E. and Salinas, M.: Purification of a novel eIF-2 α protein kinase from calf brain. *Biochem. Biophys. Res. Commun.*, 166 (1990) 1237-1244.
- 988 Crabb, J.W., Harris, W.R., Johnson, C.M., Sotiroidis, T.G., Kuhn, C.C. and Heilmeyer, L.M.G., Jr.: Electrophoretic purification of the alpha and beta subunits of phosphorylase kinase and evidence in support of the deduced amino acid sequences. *Electrophoresis (Weinheim)*, 11 (1990) 133-140.
- 989 Kanemitsu, F. and Okigaki, T.: Clinical application of subforms of creatine kinase MM and macro creatine kinases. *J. Chromatogr.*, 526 (1990) 423-438.

- 990 Le Grice, S.F.J. and Grüniger-Leitch, F.: Rapid purification of homodimer and heterodimer HIV-1 reverse transcriptase by metal chelate affinity chromatography. *Eur. J. Biochem.*, 187 (1990) 307-314.
- 991 Linz, U., Delling, U. and Rübbsamen-Waigmann, H.: Systematic studies on parameters influencing the performance of the polymerase chain reaction. *J. Clin. Chem. Clin. Biochem.*, 28 (1990) 5-13.
- 992 Megow, D. and Jacobasch, G.: Microheterogeneity of human erythrocyte pyruvate kinase: application of immunological visualization techniques after isoelectric focusing in ultrathin gels. *Electrophoresis (Weinheim)*, 11 (1990) 65-69.
- 993 Miller, R.M., Sies, H., Park, E.-M. and Thomas, J.A.: Phosphorylase and creatine kinase modification by thiol-disulfide exchange and by xanthine oxidase-initiated S-thiolation. *Arch. Biochem. Biophys.*, 276 (1990) 355-363.
- 994 Noda, M., Kunz, B.C., Tsai, S.-C., Adamik, R., Murtagh, J.J., Chen, H.-C., Halpern, J.L. and Moss, J.: Isolation and immunological properties of adenosine kinase. *Prepar. Biochem.*, 19 (1989) 351-361.
- 995 Okada, M. and Nakagawa, H.: A protein tyrosine kinase involved in regulation of pp60^{c-src} function. *J. Biol. Chem.*, 264 (1989) 20886-20893.
- 996 Panteghini, M., Bonora, R. and Pagani, F.: A two-tube immunochemical method for determination of CK-MB isoenzyme in serum evaluated. *Clin. Chem. (Winston-Salem)*, 36 (1990) 550-553.
- 997 Reppin, R.: Determination of creatine kinase and lactate dehydrogenase isoenzymes by cellulose acetate strip electrophoresis. *Z. Med. Laboratoriums-diagn.*, 30 (1989) 304-307; *C.A.*, 112 (1990) 3105e.
- 998 Rush, J. and Konigsberg, W.H.: Rapid purification of overexpressed T4 DNA polymerase. *Prepar. Biochem.*, 19 (1989) 329-340.
- 999 Sprott, S., Hammond, K.D. and Savage, N.: Electrophoretic separation of protein kinases: altered mobility with different crosslinking agents in the presence of certain detergents. *Electrophoresis (Weinheim)*, 11 (1990) 29-33.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 1000 Abou-Haila, A. and Fain-Maurel, M.-A.: Micro two-dimensional gel electrophoresis of serum and testis esterases from different strains of mice. *Electrophoresis (Weinheim)*, 11 (1990) 175-181.
- 1001 Fried, S.K. and Zechner, R.: Cachectin/tumor necrosis factor decreases human adipose tissue lipoprotein lipase mRNA levels, synthesis, and activity. *J. Lipid Res.*, 30 (1989) 1917-1923.
- 1002 Jobb, E.A. and Callahan, J.W.: Biosynthesis of sphingomyelinase in normal and Niemann-Pick fibroblasts. *Biochem. Cell Biol.*, 67 (1989) 801-807.
- 1003 Kern, P.A., Martin, R.A., Carty, J., Goldberg, I.J. and Ong, J.M.: Identification of lipoprotein lipase immunoreactive protein in pre- and postheparin plasma from normal subjects and patients with type I hyperlipoproteinemia. *J. Lipid Res.*, 31 (1990) 17-26.
- 1004 Krupyanko, V.I., Kudryavtseva, A.I., Trushkina, A.G., Ezhov, V.A. and Nikolaeva, V.M.: (Isolation and characterization of two multiple forms of acid phosphatase from the fungus *Penicillium brevicompactum*). *Biokhimiya (Moscow)*, 55 (1990) 395-403.
- 1005 Kubicz, A., Szalewicz, A., Fawcett, J.S. and Chrambach, A.: Electrofocusing of acid phosphatases from frog liver, using an immobilized pH gradient. *Electrophoresis (Weinheim)*, 11 (1990) 147-151.
- 1006 Matsukage, A. and Nishimoto, Y.: Coprecipitation of 3'-5' exonuclease and DNA polymerase γ with anti-DNA polymerase γ antibody. *J. Biochem. (Tokyo)*, 107 (1990) 213-216.
- 1007 Parthasarathy, L., Ramesh, T.G., Devi, C.S.S. and Parthasarathy, R.: Specific staining of myo-inositol-1-phosphatase on polyacrylamide gels after electrophoresis. *J. Biosci.*, 14 (1989) 249-253; *C.A.*, 112 (1990) 3137s.

See also 991, 1019.

20e. Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)

- 1008 Anand, L., Krishnamurthy, S. and Vithayathil, P.J.: Purification and properties of xylanase from the thermophilic fungus, *Humicola lanuginosa* (Griffon and Maublanc) Bunce. *Arch. Biochem. Biophys.*, 276 (1990) 546-553.
- 1009 Kataeva, I.A., Golovchenko, N.P., Dzhavakhishvili, T.D., Tsoi, T.V., Boronin, A.M. and Akimenko, V.K.: (Purification and some properties of β -glucosidases encoded by the bgl gene of *Clostridium thermocellum* and its 3'-deletion mutant). *Biokhimiya (Moscow)*, 55 (1990) 299-307.
- 1010 Klein, R.D., Deibel, M.R., Sarcich, J.L., Zurcher-Neely, H.A., Reardon, I.M. and Heinrikson, R.L.: Purification and characterization of invertase from a novel industrial yeast, *Schwanniomyces occidentalis*. *Prepar. Biochem.*, 19 (1989) 293-319.

See also 855.

20f. Other hydrolases

- 1011 Aronovich, E.L., Akhunov, V.S. and Krasnopol'skaya, K.D.: (Isolation and properties of hexosaminidase A and activator protein from human kidney). *Biokhimiya (Moscow)*, 55 (1990) 43-51.
- 1012 Hervatin, F., Benkouka, F., Robert, J.C., Peranzi, G. and Soumarmon, A.: The ontogeny of rat gastric H^+/K^+ -ATPase. *Biochim. Biophys. Acta*, 985 (1989) 320-324.
- 1013 Suzuki, K., Kusumoto, H., Nishioka, J. and Komiyama, Y.: Bovine plasma protein C inhibitor with structural and functional homologous properties to human plasma protein C inhibitor. *J. Biochem. (Tokyo)*, 107 (1990) 381-388.
- 1014 Tavera, C., Guillemont, J.-C., Capdevielle, J., Ferrara, P., Leung-Tack, J. and Collé, A.: A rapid two-step purification of rat cystatin C, one major inhibitor of cysteine proteinases. *Prepar. Biochem.*, 19 (1989) 279-291.
- 1015 Yamauchi, T., Nagahama, M., Watanabe, T., Ishizuka, Y., Hori, H. and Murakami, K.: Site-directed mutagenesis of human prorenin. Substitution of three arginine residues in the propeptide with glutamine residues yields active prorenin. *J. Biochem. (Tokyo)*, 107 (1990) 27-31.
- 1016 Yokoyama, K., Hisabori, T. and Yoshida, M.: The reconstituted $\alpha_3\beta_3$ complex of the thermostable F_1 -ATPase. *J. Biol. Chem.*, 264 (1989) 21837-21841.

See also 823, 855.

20g. Lyases

- 1017 Stanley, B.A., Pegg, A.E. and Holm, I.: Site of pyruvate formation and processing of mammalian S-adenosylmethionine decarboxylase proenzyme. *J. Biol. Chem.*, 264 (1989) 21073-21079.

20i. Ligases

- 1018 Bobkova, E.V., Volfson, A.D., Ankilova, V.N. and Lavrik, O.I.: (Separation and comparative analysis of subunits of phenylalanyl-tRNA-synthetases from *Escherichia coli* MRE-600 and *Thermus thermophilus* HB8). *Biokhimiya (Moscow)*, 55 (1990) 525-533.

20j. Complex mixtures and incompletely identified enzymes

- 1019 Feng, W., He, G., Ye, Z., Liu, H. and Zhong, S.: (Isoenzyme analyses in the digestive system of the giant panda by polyacrylamide gel disc electrophoresis). *Sichuan Daxue Xuebao, Ziran Kexueban*, 26 (1989) 212-216; *C.A.*, 112 (1990) 3210k.
- 1020 Shirley, M.W.: Enzyme characterization of the Eimeriidae. *Colloq. INRA*, 49 (1989) 111-124; *C.A.*, 112 (1990) 4105s - a review with 15 refs.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

21a. Purines, pyrimidines, nucleosides, nucleotides

- 1021 Guttman, A., Cohen, A.S., Heiger, D.N. and Karger, B.L.: Analytical and micropreparative ultrahigh resolution of oligonucleotides by polyacrylamide gel high-performance capillary electrophoresis. *Anal. Chem.*, 62 (1990) 137-141.
- 1022 Mishra, A.P. and Yadava, K.L.: Study of mercury(II), beryllium(II) and copper(II)-cytosine complexes by paper electrophoresis. *Vijnana Parishad Anusandhan Patrika*, 32 (1989) 129-135; *C.A.*, 112 (1990) 64065p.
- 1023 Schuddeemat, J., van Leeuwen, C.C.M., van den Broek, P.J.A. and Steveninck, J.: Mechanism of transport-associated phosphorylation of 2-deoxy-D-glucose in the yeast *Kluyveromyces marxianus*: characterization of the phosphoryl source. *Biochim. Biophys. Acta*, 986 (1989) 290-294.

See also 1073.

21b. Nucleic acids, RNA

- 1024 Amaya, Y., Nakano, A., Ito, K. and Mori, M.: Isolation of a yeast gene, SRH1, that encodes a homologue of the 54K subunit of mammalian signal recognition particle. *J. Biochem. (Tokyo)*, 107 (1990) 457-463.
- 1025 Hajnsdorf, E. and Expert-Bezancon, A.: An electrophoretic method for the purification of RNA regions involved in protein crosslinking. *Anal. Biochem.*, 185 (1990) 103-107.
- 1026 Kawamura, T., Shigesada, K., Yanagisawa, S. and Izui, K.: Phosphoenolpyruvate carboxylase prevalent in maize roots: Isolation of a cDNA clone and its use for analyses of the gene and gene expression. *J. Biochem. (Tokyo)*, 107 (1990) 165-168.
- 1027 Matsumoto, T., Nishikawa, K., Hori, H., Ohta, T., Miura, K.-i. and Watanabe, K.: Recognition sites of tRNA by a thermostable tRNA(guanosine-2'-)-methyltransferase from *Thermus thermophilus* HB27. *J. Biochem. (Tokyo)*, 107 (1990) 331-338.
- 1028 Miyauchi, T., Kanekura, T., Yamaoka, A., Ozawa, M., Miyazawa, S. and Muramatsu, T.: Basigin, a new, broadly distributed member of the immunoglobulin superfamily, has strong homology with both the immunoglobulin V domain and the β -chain of major histocompatibility complex class II antigen. *J. Biochem. (Tokyo)*, 107 (1990) 316-323.
- 1029 Oberbäumer, I. and Speth, C.: Detection of RNA on Northern blots by negative staining with aurointricarboxylic acid. *Anal. Biochem.*, 185 (1990) 77-79.
- 1030 Pallansch, L., Beswick, H., Talian, J. and Zelenka, P.: Use of an RNA folding algorithm to choose regions for amplification by the polymerase chain reaction. *Anal. Biochem.*, 185 (1990) 57-62.

- 1031 Pollard-Knight, D., Read, C.A., Downes, M.J., Howard, L.A., Leadbetter, M.R., Pheby, S.A., McNaughton, E., Syms, A. and Brady, M.A.W.: Nonradioactive nucleic acid detection by enhanced chemiluminescence using probes directly labeled with horseradish peroxidase. *Anal. Biochem.*, 185 (1990) 84-89.
- 1032 Reilly, J.D., Freeman, S.K., Melhem, R.F., Kierzek, R., Caruthers, M.H., Edmonds, M. and Munns, T.W.: Antibodies specific for branched ribonucleic acids. *Anal. Biochem.*, 185 (1990) 125-130.

See also 887, 1001, 1053.

21c. Nucleic acids, DNA

- 1033 Aakerman, B., Jonsson, M., Norden, B. and Lalande, M.: Orientational dynamics of T2 DNA during agarose gel electrophoresis: influence of gel concentration and electric field strength. *Biopolymers*, 28 (1989) 1541-1571; *C.A.*, 111 (1989) 170451a.
- 1034 Adeli, K. and Ogbonna, G.: Rapid purification of human DNA from whole blood for potential application in clinical chemistry laboratories. *Clin. Chem. (Winston-Salem)*, 36 (1990) 261-264.
- 1035 Chuprina, V.P.: Critical comments on the one recent DNA bending model and its fit to electrophoretic data. *J. Biomol. Struct. Dyn.*, 7 (1989) 35-39; *C.A.*, 111 (1989) 228382x.
- 1036 Flick, M.B., Warters, R.L., Yasui, L.S. and Krisch, R.E.: Measurement of radiation-induced DNA damage using gel electrophoresis or neutral filter elution shows an increased frequency of DNA strand breaks after exposure to pH 9.6. *Radiat. Res.*, 119 (1989) 452-465; *C.A.*, 111 (1989) 228026j.
- 1037 Gaubatz, J.W. and Flores, S.C.: Purification of eucaryotic extrachromosomal circular DNAs using exonuclease III. *Anal. Biochem.*, 184 (1990) 305-310.
- 1038 Guo, Q., Lu, M., Seeman, N.C. and Kallenbach, N.R.: Drug binding by branched DNA molecules: analysis by chemical footprinting of intercalation into an immobile junction. *Biochemistry*, 29 (1990) 570-578.
- 1039 Heller, C. and Pohl, F.M.: A systematic study of field inversion gel electrophoresis. *Nucleic Acids Res.*, 17 (1989) 5989-6003; *C.A.*, 111 (1989) 170450z.
- 1040 Henry, J.M., Raina, A.K. and Ridgway, R.L.: Isolation of high-molecular-weight DNA from insects. *Anal. Biochem.*, 185 (1990) 147-150.
- 1041 Holmes, D.L. and Stellwagen, N.C.: The electric field dependence of DNA mobilities in agarose gels: a reinvestigation. *Electrophoresis (Weinheim)*, 11 (1990) 5-15.
- 1042 Holmes, D.L. and Stellwagen, N.C.: Electrophoresis of DNA in oriented agarose gels. *J. Biomol. Struct. Dyn.*, 7 (1989) 311-327; *C.A.*, 112 (1990) 3887m.
- 1043 Ishihara, H. and Shikita, M.: Electroblotting of double-stranded DNA for hybridization experiments: DNA transfer is complete within 10 minutes after pulsed-field gel electrophoresis. *Anal. Biochem.*, 184 (1990) 207-212.
- 1044 Jiang, X., Trujillo, J.M., Dao, D. and Liang, J.C.: Studies of BCR and ABL gene rearrangements in chronic myelogenous leukemia patients by conventional and pulsed-field gel electrophoresis using gel inserts. *Cancer Genet. Cytogenet.*, 42 (1989) 287-294; *C.A.*, 112 (1990) 3895n.
- 1045 Lamperti, E.D., Villa-Komaroff, L.: Generation of deletion subclones for sequencing by partial digestion with restriction endonucleases. *Anal. Biochem.*, 185 (1990) 187-193.
- 1046 Lockett, T.J.: A bacteriophage λ DNA purification procedure suitable for the analysis of DNA from either large or multiple small lysates. *Anal. Biochem.*, 185 (1990) 230-234.

- 1047 Lumpkin, O.: One-dimensional translational motion of a two-spring chain with strong nonlinear drag: a possible model for time-dependent DNA gel electrophoresis. *Phys. Rev. A: Gen. Phys.*, 40 (1989) 2634-2642; *C.A.*, 111 (1989) 228488m.
- 1048 Muskhelishvili, G.D., Karseladze, M.V. and Prangashvili, D.A.: (Incorporation of exogenous precursors into nucleic acids of the extremely thermophilic acidophilic archaeobacterium *Sulfolobus acidocaldarius*). *Biokhimiya (Moscow)*, 55 (1990) 517-524.
- 1049 Pollard-Knight, D., Simmonds, A.C., Schaap, A.P., Akhavan, H. and Brady, M.A.W.: Nonradioactive DNA detection on Southern blots by enzymatically triggered chemiluminescence. *Anal. Biochem.*, 185 (1990) 353-358.
- 1050 Polman, J.K. and Larkin, J.M.: Purification of DNA from agarose gels. *Biotechnol. Tech.*, 3 (1989) 329-332; *C.A.*, 111 (1989) 228395d.
- 1051 Prior, T.W., Friedman, K.J., Highsmith, W.E., Jr., Perry, T.R. and Silverman, L.M.: Molecular probe protocol for determining carrier status in Duchenne and Becker muscular dystrophies. *Clin. Chem. (Winston-Salem)*, 36 (1990) 441-445.
- 1052 Sagara, Y., Sawae, H., Kimura, A., Sagara-Nakano, Y., Morohashi, K.-i., Miyoshi, K.-i. and Horiuchi, T.: Structural organization of the bovine adrenodoxin gene. *J. Biochem. (Tokyo)*, 107 (1990) 77-83.
- 1053 Schlechte, H., Kunde, D. and Jacobasch, K.H.: Determination of nucleic acid fractions in neoplastic and preneoplastic tissues by gel electrophoresis. *Ger. (East)* DD 265,239 (Cl. GolN33/561), 22 Feb. 1989, Appl. 306,781, 09 Sep. 1987; 5 pp.; *C.A.*, 112 (1990) 5712z.
- 1054 Serwer, P. and Hayes, S.J.: A new mode of rotating gel electrophoresis for fractionating linear and circular duplex DNA: the effects of electrophoresis during the gel's rotation. *Appl. Theor. Electrophor.*, 1 (1989) 95-98; *C.A.*, 112 (1990) 3893k.
- 1055 Smith, C.L., Ewing, C.M., Mellon, M.T., Kane, S.E., Jain, T. and Shorr, R.G.L.: HydroLink gel electrophoresis (HLGE). II. Applications of a new polymer matrix to dsDNA analysis. *J. Biochem. Biophys. Methods*, 19 (1989) 51-64; *C.A.*, 111 (1989) 211363u.
- 1056 Smith, C.L., Ewing, C.M., Mellon, M.T., Shorr, R.G.L. and Jain, T.: HydroLink gel electrophoresis (HLGE). III. High DNA loading capacity and recovery of dsDNA. *J. Biochem. Biophys. Methods*, 19 (1989) 65-73; *C.A.*, 111 (1989) 211364v.
- 1057 Sobral, B.W.S. and Atherly, A.G.: Pulse time and agarose concentration affect the electrophoretic mobility of cccDNA during electrophoresis in CHEF and in FIGE. *Nucleic Acids Res.*, 17 (1989) 7359-7369; *C.A.*, 111 (1989) 211377b.
- 1058 Sorimachi, H., Emori, Y., Kawasaki, H., Suzuki, K. and Inoue, Y.: Molecular cloning and characterization of cDNAs coding for apolysialoglycoproteins in cherry salmon (*Oncorhynchus masou*) eggs. *J. Biochem. (Tokyo)*, 107 (1990) 61-67.
- 1059 Stürzl, M. and Roth, W.K.: "Run-off" synthesis and application of defined single-stranded DNA hybridization probes. *Anal. Biochem.*, 185 (1990) 164-169.
- 1060 Takenaka, S.: (Analysis of DNA replication terminus (ter) using gel electrophoresis). *Bunseki*, (1989) 727-731; *C.A.*, 111 (1989) 226059y - a review with 12 refs.
- 1061 Weber, J., Scheid, W. and Traut, H.: Enhancement of bleomycin-induced DNA damage by exposing isolated DNA to high concentrations of the calcium antagonist verapamil. *Arzneim.-Forsch.*, 39 (1989) 1550-1554.
- 1062 Weidner, M.F., Millard, J.T. and Hopkins, P.B.: Determination at single nucleotide resolution of the sequence specificity of DNA interstrand cross-linking agents in DNA fragments. *J. Am. Chem. Soc.*, 111 (1989) 9270-9282; *C.A.*, 112 (1990) 3896p.

See also 865, 887, 1024, 1026, 1031.

21d. Structural studies on RNA and RNA mapping

See 1032.

21e. Structural studies on DNA and DNA mapping

- 1063 Arpagaus, M., Kott, M., Vatsis, K.P., Bartels, C.F., La Du, B.N. and Lockridge, O.: Structure of the gene for human butyrylcholinesterase. Evidence for a single copy. *Biochemistry*, 29 (1990) 124-131.
- 1064 Blocher, D., Einspinner, M. and Zajackowski, J.: CHEF electrophoresis, a sensitive technique for the determination of DNA double-strand breaks. *Int. J. Radiat. Biol.*, 56 (1989) 437-448; *C.A.*, 111 (1989) 228028m.
- 1065 Levine, J.D. and Cech, C.L.: Low frequency restriction enzymes in pulsed field electrophoresis. *Bio/Technology*, 7 (1989) 1033-1036; *C.A.*, 111 (1989) 228173e.
- 1066 Sugiura, S., Tanaka, M., Masamune, Y. and Yamaguchi, K.: DNA binding properties of purified replication initiator protein (Rep) encoded by plasmid pSC101. *J. Biochem. (Tokyo)*, 107 (1990) 369-376.
- 1067 Tokunaga, K., Kay, P.H., Christiansen, F.T., Saueracker, G. and Dawkins, R.L.: Comparative mapping of the human major histocompatibility complex in different racial groups by pulsed field gel electrophoresis. *Hum. Immunol.*, 26 (1989) 99-106; *C.A.*, 111 (1989) 226319h.

See also 861, 865.

21f. Complex mixtures of nucleic acids and their fragments

- 1068 Nolan-Sorden, N.L., Lesiak, K., Bayard, B., Torrence, P.F. and Silverman, R.H.: Photochemical crosslinking in oligonucleotide-protein complexes between a bromine-substituted 2-5A analog and 2-5A-dependent RNase by ultraviolet lamp or laser. *Anal. Biochem.*, 184 (1990) 298-304.

See also 865.

24. ORGANIC SULPHUR COMPOUNDS

- 1069 Abe, S., Endo, M. and Sone, T.: Isotachopheretic separation and detection of thiophene-mono, di- and tricarboxylic acids. *Fresenius' Z. Anal. Chem.*, 336 (1990) 40-41.
- 1070 Khmel'nitskii, R.A. and Bondarenko, M.A.: (Instrumental methods for determination of lower mercaptans). *Zh. Anal. Khim.*, 44 (1989) 965-985; *C.A.*, 111 (1989) 247199j.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCLUDING SUGAR PHOSPHATES)

See 1023.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26c. Coordination compounds

- 1071 Koch, S. and Ackermann, G.: (Electrophoretic investigations of the complexation of iron(III) with polyphenols). *Z. Chem.*, 29 (1989) 185-186; *C.A.*, 111 (1989) 220232c.

28. ANTIBIOTICS

- 1072 Holaka, W. and Gwozdz, E.: Determination of colistin concentration by isotachophoresis. *J. Liq. Chromatogr.*, 12 (1989) 2761-2767.
- 1073 Kimura, K.-i., Morinaga, T., Miyata, N. and Kawanishi, G.: Nucleotide sequence of SN-07 chromophore binding site. *J. Antibiot.*, 42 (1989) 1838-1844.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29d. Herbicides

- 1074 Foret, F., Sustacek, V. and Bocek, P.: Separation of some triazine herbicides and their solvolytic products by capillary zone electrophoresis. *Electrophoresis (Weinheim)*, 11 (1990) 95-97.

32. DRUG ANALYSIS

32d. Central nervous system drugs

- 1075 Jokl, V., Mokra-Pospichalova, J. and Vitkovic, B.: (Capillary isotachophoresis of tricyclic antidepressive and antihistamine agents: determination of the individual drugs and separation of amitriptyline and nortriptyline). *Cesk. Farm.*, 38 (1989) 294-298; *C.A.*, 112 (1990) 62751k.

32h. Toxicological and forensic applications

- 1076 Choots, A.C., Verheggen, T.P.E.M., de Vries, P.M.J.M. and Everaerts, F.M.: Ultraviolet-absorbing organic anions in uremic serum separated by capillary zone electrophoresis, and quantification of hippuric acid. *Clin. Chem. (Winston-Salem)*, 36 (1990) 435-440.

See also 926, 927.

33. CLINICO-CHEMICAL APPLICATIONS

33b. Complex mixtures and profiling (single compounds by cross-reference only)

- See 851, 854, 855, 859, 860, 865, 923, 924, 954, 968, 972, 983, 996, 1003, 1034, 1051, 1076.

34. FOOD ANALYSIS

34b. Complex mixtures (single compounds by cross-reference only)

See 947.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36d. Complex mixtures and unidentified compounds

1077 Imae, T., Otani, W. and Oka, K.: Electrophoretic light scattering of surfactant micelle colloids. *J. Phys. Chem.*, 94 (1990) 853-855; *C.A.*, 112 (1990) 63159k.

1078 Van Kemenade, A.W.C. and Everaerts, F.M.: Electrophoresis in processing of swine excrement. *PT-Procestechn.*, 44 (1989) 54-57; *C.A.*, 111 (1989) 213576q.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

1079 Kennedy, R.T., Oates, M.D., Cooper, B.R., Nickerson, B. and Jorgenson, J.W.: Microcolumn separations and the analysis of single cells. *Science (Washington)*, 246 (1989) 57-63; *C.A.*, 111 (1989) 211494n.

38. INORGANIC COMPOUNDS

38a. Cations

1080 Ito, R., Hirata, Y., Katsurada, K. and Shigesada, H.H.: (Separation of rare-earth ions by electrophoretic focusing). *Kagaku Kogaku Ronbunshu*, 15 (1989) 690-694; *C.A.*, 112 (1990) 47739u.

1081 Kaneko, K., Kaneko, H., Hirabayashi, M. and Ihara, H.: (Electrophoretic determination of metals in superconductor oxides). *Jpn. Kokai Tokkyo Koho Pat. JP 01,152,353* (89,152,353) (Cl. G01N27/26), 14 Jun. 1989, Appl. 87/311,197, 09 Dec. 1987; 5 p.; *C.A.*, 111 (1989) 247180w.

1082 Kaniansky, D., Zelensky, I., Valaskova, I., Marak, J. and Zelenska, V.: Isotachophoretic separation of alkali and alkaline earth metal cations in water-polyethylene glycol mixtures. *J. Chromatogr.*, 502 (1990) 143-153.

1083 Yang, J.Y., Yans, M.H. and Lin, S.M.: Effect of wet decomposition methods on the determination of cobalt, copper, selenium, and zinc in biological samples using electrophoresis. *Anal. Chem.*, 62 (1990) 146-150.

See also 1022.

38b. Anions

1084 Kojima, K.: (Electrolytic solutions for isotachophoresis). *Jpn. Kokai Tokkyo Koho JP 01 09,353* (89 09,353) (Cl. G01N27/26), 12 Jan. 1989, Appl. 87/164,562, 30 Jun. 1987; 4 p.; *C.A.*, 112 (1990) 69036q.

- 1085 Taga, M., Kan, M., Komatsu, F., Tanaka, S. and Yoshida, H.: Capillary isotachophoretic determination of phosphate after enrichment on membrane filter as ion pair of molybdophosphate with bis(2-(5-chloro-2-pyridylazo)-5-diethylaminophenolato)cobalt(III). *Anal. Sci.*, 5 (1989) 219-220; *C.A.*, 111 (1989) 246968x.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 1086 Smith, I. and Furst, V.: Computer analysis of radiochromatograms and gels: qualitative and quantitative methods with the AMBIS system. *J. Planar Chromatogr. - Mod. TLC*, 2 (1989) 233-237; *C.A.*, 111 (1989) 211225a.

See also 818.