

Author Index

- Ageta, T., see Mizobuchi, N. 321
- Aherne, G.W., see Burton, N.K. 253
- Akira, K., see Baba, S. 31
- Ascalone, V.
- and Dal Bo', L.
Chromatographic selectivity and maintenance of column efficiency during the high-performance liquid chromatographic analysis of progabide and its acid metabolite 412
- Baba, S.
- , Akira, K., Sasaki, Y. and Furuta, T.
Synchronized accumulating radioisotope detector equipped with a mutual anti-coincidence guard for radio-gas chromatography 31
- Badcock, N.R.
- , O'Reilly, D.A. and Pinnock, C.B.
Liquid chromatographic determination of retinol and α -tocopherol in human buccal mucosal cells 290
- Bayne, W.F., see Demetriades, J.L. 225
- Berthou, F., see Lucas, D. 57
- Bettendorff, L.
- , Grandfils, C., De Rycker, C. and Schoffeniels, E.
Determination of thiamine and its phosphate esters in human blood serum at femtomole levels 297
- Beving, H.
- , Kristensson, J., Malmgren, R., Olsson, U., Schmidt, S. and Svensson, J.
Determination of platelet-megakaryocyte regeneration time in painters occupationally exposed to organic solvents 67
- Black, A.E.
- and Chang, T.
Determination of 1-(4-methoxybenzoyl)-5-oxo-2-pyrrolidinepropanoic acid, a potential nootropic agent, in human plasma and urine by reversed-phase high-performance liquid chromatography 361
- Blocka, K.L., see Richardson, C.J. 382
- Bo', L. Dal, see Ascalone, V. 412
- Bonakdar, M., see Wang, J. 343
- Borbe, H.O., see Terlinden, R. 372
- Borchert, M.E., see Frost, S.K. 314
- Brandenberger, H., see Hattori, H. 135
- Bühning, K.U.
- and Garbe, A.
Determination of the new β -blocker bisoprolol and of metoprolol, atenolol and propranolol in plasma and urine by high-performance liquid chromatography 215
- Burton, N.K.
- and Aherne, G.W.
Sensitive measurement of glutathione using isocratic high-performance liquid chromatography with fluorescence detection 253
- Buu, N.T., see Racz, K. 117
- Chang, T., see Black, A.E. 361
- Chichester, C.O., see Sassa, Jr., W.C. 258
- Coutts, R.T., see Singh, N.N. 331
- Cramers, C.A., see De Jong, A.P.J.M. 19
- Dal Bo', L., see Ascalone, V. 412
- Damm, P., see Lehr, K.H. 355
- Debinski, W., see Racz, K. 117
- De Jong, A.P.J.M.
- , Kok, R.M., Cramers, C.A. and Wadman, S.K.
Determination of acidic catecholamine metabolites in plasma and cerebrospinal fluid using gas chromatography-negative-ion mass spectrometry 19
- Demetriades, J.L.
- , Souder, P.R., Entwistle, L.A., Vincek, W.C., Musson, D.G. and Bayne, W.F.
High-performance liquid chromatographic determination of cilastatin in biological fluids 225
- De Rycker, C., see Bettendorff, L. 297
- Dethloff, L.A.
- , Gilmore, L.B. and Hook, G.E.R.
Separation of pulmonary surfactant phospholipids by high-performance liquid chromatography 79
- De Vries, J.X.
- , Walter-Sack, I., Van de Loo, A. and Kocher, J.
Determination of benzarone in human plasma and urine by high-performance liquid chromatography and gas chromatography-mass spectrometry. Identification of the conjugates 167
- Dobó, R., see Kovács, L. 308
- Dreisbach, J.H.
- and Veca, A.
Rapid method for the analysis of urinary pyrrole-2-carboxylic acid using reversed-phase high-performance liquid chromatography 280
- Entwistle, L.A., see Demetriades, J.L. 225

- Evans, W.E., see Riley, C.A. 199
- Floch, H.H., see Lucas, D. 57
- Friis, J.M.
- and Lakings, D.B.
High-performance liquid chromatographic method for the determination of cefpimizole in tissue 399
- Frost, S.K.
- , Borchert, M.E. and Thorsteinsdottir, S.
Rapid and sensitive thin-layer chromatographic assay procedure for measuring xanthine dehydrogenase activity from tissue extracts 314
- Furuta, T., see Baba, S. 31
- Garbe, A., see Bühring, K.U. 215
- Gillen, L.E., see Holdiness, M.R. 99
- Gilmore, L.B., see Dethloff, L.A. 79
- Goldman, M.E.
- , Weber, R.J., Newman, A.H., Rice, K.C., Skolnick, P. and Paul, S.M.
High-performance fast affinity chromatographic purification of anti-benzodiazepine antibodies 264
- Grandfils, C., see Bettendorff, L. 297
- Haber, M.
- , Kavallaris, M. and Stewart, B.W.
Two-stage incorporation of thymidine triphosphate into mammalian DNA as indicated by chromatography on benzoylated DEAE-cellulose 127
- Hada, T., see Yamamoto, T. 270
- Haeffner, E.W.
- , Holl, A. and Schroeter, D.
Preparation of two plasma membrane fractions from ascites tumour cells by gel chromatography on Sephacryl S-1000 107
- Harrison, E.F., see Holdiness, M.R. 99
- Hashimoto, T., see Kataoka, H. 242
- Hattori, H.
- , Suzuki, O. and Brandenberger, H.
Positive- and negative-ion mass spectrometry of butyrophonones 135
- Henry, J.A., see Zoer, J. 189
- Higashino, K., see Yamamoto, T. 270
- Holdiness, M.R.
- , Morgan, Jr., L.R., Gillen, L.E. and Harrison, E.F.
High-performance liquid chromatographic determination of N-acetylcysteine in human serum following acetaminophen overdose 99
- Holl, A., see Haeffner, E.W. 107
- Holland, M.L., see Yancey, M. 3
- Hook, G.E.R., see Dethloff, L.A. 79
- Hutchings, A.
- , Spragg, B.P. and Routledge, P.A.
High-performance liquid chromatographic assay of amiodarone and desethylamiodarone in plasma 389
- Inger, F., see Lacroix, C. 275
- Inoue, K., see Kohashi, M. 303
- Iwai, K., see Kohashi, M. 303
- Jakobsen, P.
- and Waldorff, S.
Determination of digoxin, digoxigenin and dihydrodigoxigenin in urine by extraction, derivatization and high-performance liquid chromatography 349
- Jamali, F., see Singh, N.N. 331
- James, C.A.
- and Rogers, H.J.
Estimation of mesna and dimesna in plasma and urine by high-performance liquid chromatography with electrochemical detection 394
- Jandke, J.
- and Spitteller, G.
Dipeptide analysis in human urine 39
- Johnson, V.A.
- and Wilson, J.T.
Flurbiprofen analysis in plasma and breast milk by high-performance liquid chromatography 367
- Jong, A.P.J.M. de, see De Jong, A.P.J.M. 19
- Juskom, W.J., see Shum, L. 207
- Kataoka, H.
- , Yamamoto, H., Sumida, Y., Hashimoto, T. and Makita, M.
Gas chromatographic determination of hypotaurine 242
- Kavallaris, M., see Haber, M. 127
- Kelly, J.G., see Kinney, C.D. 377
- Kinney, C.D.
- and Kelly, J.G.
Estimation of concentrations of diltiazem in plasma using normal-phase column liquid chromatography with ultraviolet detection 377
- Kocher, J., see De Vries, J.X. 167
- Kodama, H., see Mizobuchi, N. 321
- Kohashi, M.
- , Inoue, K., Sotobayashi, H. and Iwai, K.
Microdetermination of folate monoglutamates in serum by liquid chromatography with electrochemical detection 303
- Kok, R.M., see De Jong, A.P.J.M. 19
- Kovács, L.
- , Zalka, A., Dobó, R. and Pucsok, J.
One-dimensional thin-layer chromatographic separation of lipids into four-

- teen fractions by two successive developments on the same plate 308
- Kristensson, J., see Beving, H. 67
- Kuchel, O., see Racz, K. 117
- Labadarios, D.
- , Moodie, I.M. and Shephard, G.S.
Gas chromatography of amino acids in urine: identification and removal of creatinine as the major interfering substance 237
- Lacroix, C.
- , Inger, F., Menager, S. and Lafont, O.
Détermination simultanée de l'acide hippurique et des acides *o*-, *m*-, *p*-méthylhippuriques urinaires par chromatographie liquide 275
- Lafont, O., see Lacroix, C. 275
- Lakings, D.B., see Friis, J.M. 399
- Lehr, K.H.
- and Damm, P.
Simultaneous determination of fenbendazole and its two metabolites and two triclofenbendazole metabolites in plasma by high-performance liquid chromatography 355
- Limratana, N., see Speek, A.J. 284
- Loo, A. van de, see De Vries, J.X. 167
- Lucas, D.
- , Ménez, J.F., Berthou, F., Pennec, Y. and Floch, H.H.
Determination of free acetaldehyde in blood as the dinitrophenylhydrazine derivative by high-performance liquid chromatography 57
- Makita, M., see Kataoka, H. 242
- Malmgren, R., see Beving, H. 67
- Maurer, H.
- and Pflieger, K.
Identification and differentiation of beta-blockers and their metabolites in urine by computerized gas chromatography-mass spectrometry 147
- Menager, S., see Lacroix, C. 275
- Ménez, J.F., see Lucas, D. 57
- Mizobuchi, N.
- , Ageta, T., Sasaki, K. and Kodama, H.
Isotachopheric analyses of cystine, homocystine and cystathionine in urines from patients with inborn errors of metabolism 321
- Moodie, I.M., see Labadarios, D. 237
- Morgan, Jr., L.R., see Holdiness, M.R. 99
- Moriwaki, Y., see Yamamoto, T. 270
- Musson, D.G., see Demetriades, J.L. 225
- Mutschler, E., see Schweitzer, C. 405
- Newman, A.H., see Goldman, M.E. 264
- Novotny, M., see Yancey, M. 3, 47
- Olsson, U., see Beving, H. 67
- O'Reilly, D.A., see Badcock, N.R. 290
- Palmer, J.L.
Novel method of sample preparation for the determination of paracetamol in plasma by high-performance liquid chromatography with electrochemical detection 338
- Pasutto, F.M., see Singh, N.N. 331
- Paul, S.M., see Goldman, M.E. 264
- Pehal, F., see Verner, P. 232
- Pennec, Y., see Lucas, D. 57
- Pflieger, K., see Maurer, H. 147
- Pinnock, C.B., see Badcock, N.R. 290
- Pucsok, J., see Kovács, L. 308
- Racz, K.
- , Kuchel, O., Debinski, W. and Buu, N.T.
Improved liquid chromatographic determination of dopamine- β -hydroxylase activity in tissues and plasma 117
- Rice, K.C., see Goldman, M.E. 264
- Richardson, C.J.
- , Ross, S.G., Blocka, K.L. and Verbeeck, R.K.
High-performance liquid chromatographic analysis of piroxicam and its major metabolite 5'-hydroxypiroxicam in human plasma and urine 382
- Riley, C.A.
- and Evans, W.E.
Simultaneous analysis of antipyrine and lorazepam by high-performance liquid chromatography 199
- Rodgers, R.L., see Sessa, Jr., W.C. 258
- Rogers, H.J., see James, C.A. 394
- Ross, S.G., see Richardson, C.J. 382
- Routledge, P.A., see Hutchings, A. 389
- Russell, A.S., see Singh, N.N. 331
- Russell, B.W., see Smuts, H.E.M. 326
- Rycker, C. De, see Bettendorff, L. 297
- Salari, H.
Solid-phase extraction and reversed-phase high-performance liquid chromatographic technique for isolation and estimation of platelet activating factor in plasma 89
- Saowakontha, S., see Speek, A.J. 284
- Sasaki, K., see Mizobuchi, N. 321
- Sasaki, Y., see Baba, S. 31
- Schmidt, S., see Beving, H. 67
- Schoffeniels, E., see Bettendorff, L. 297
- Schreurs, W.H.P., see Speek, A.J. 284
- Schroeter, D., see Haeflner, E.W. 107
- Schweitzer, C.
- , Spann, H. and Mutschler, E.
Fluorimetric determination of fluvox-

- amine or clovoxamine in human plasma after thin-layer chromatographic or high-performance liquid chromatographic separation 405
- Sessa, Jr., W.C.
- , Rodgers, R.L. and Chichester, C.O. Quantification of collagen synthesis by reversed-phase high-performance liquid chromatography utilizing 4-chloro-7-nitrobenzofurazan derivatives 258
- Shephard, G.S., see Labadarios, D. 237
- Shum, L.
- and Jusko, W.J. Reversed-phase high-performance liquid chromatographic determination of verofylline in rat serum 207
- Singh, N.N.
- , Jamali, F., Pasutto, F.M., Coutts, R.T. and Russell, A.S. Stereoselective gas chromatographic analysis of etodolac enantiomers in human plasma and urine 331
- Skolnick, P., see Goldman, M.E. 264
- Smuts, H.E.M.
- and Russell, B.W. Two isotachophoretically separated components in human cerebrospinal fluid identified as folic acid and uric acid 326
- Sotobayashi, H., see Kohashi, M. 303
- Souder, P.R., see Demetriades, J.L. 225
- Spahn, H., see Schweitzer, C. 405
- Speck, A.J.
- , Wongkham, C., Limratana, N., Saowakontha, S. and Schreurs, W.H.P. Microdetermination of vitamin A in human plasma using high-performance liquid chromatography with fluorescence detection 284
- Spiteller, G., see Jandke, J. 39
- Spragg, B.P., see Hutchings, A. 389
- Stewart, B.W., see Haber, M. 127
- Stuart, R., see Yancey, M. 3, 47
- Sumida, Y., see Kataoka, H. 242
- Suzuki, O., see Hattori, H. 135
- Svensson, J., see Beving, H. 67
- Takahashi, S., see Yamamoto, T. 270
- Terlinden, R.
- and Borbe, H.O. Determination of amitriptylinoxide and its major metabolites amitriptyline and nortriptyline in plasma by high-performance liquid chromatography 372
- Thomson, S.B.
- and Tucker, D.J. Analysis of homocysteine in human urine using high-performance liquid chromatography with electrochemical detection 247
- Thorsteinsdottir, S., see Frost, S.K. 314
- Timm, U., see Zell, M. 175
- Tucker, D.J., see Thomson, S.B. 247
- Van de Loo, A., see De Vries, J.X. 167
- Veca, A., see Dreisbach, J.H. 280
- Verbeeck, R.K., see Richardson, C.J. 382
- Verner, P.
- and Pehal, F. Ion-exchange extractive alkylation of organic acids 232
- Vincek, W.C., see Demetriades, J.L. 225
- Virgili, P., see Zoer, J. 189
- Vries, J.X. de, see De Vries, J.X. 167
- Wadman, S.K., see De Jong, A.P.J.M. 19
- Waldorff, S., see Jakobsen, P. 349
- Walter-Sack, I., see De Vries, J.X. 167
- Wang, J.
- and Bonakdar, M. Liquid chromatography of reserpine and rescinnamine using electrochemical detection 343
- Weber, R.J., see Goldman, M.E. 264
- Wiesler, D., see Yancey, M. 3, 47
- Wilson, J.T., see Johnson, V.A. 367
- Wongkham, C., see Speck, A.J. 284
- Yamamoto, H., see Kataoka, H. 242
- Yamamoto, T.
- , Moriwaki, Y., Takahashi, S., Hada, T. and Higashino, K. Separation of hypoxanthine and xanthine from pyrazinamide and its metabolites in plasma and urine by high-performance liquid chromatography 270
- Yancey, M.
- , Holland, M.L., Stuart, R., Wiesler, D. and Novotny, M. Urinary profiles of organic acids and volatile metabolites during the starvation process in rats 3
- Yancey, M.
- , Stuart, R., Wiesler, D. and Novotny, M. Quantitative alterations in the metabolism of carbonyl compounds due to diet-induced lipid peroxidation in rats 47
- Zalka, A., see Kovács, L. 308
- Zell, M.
- and Timm, U. Highly sensitive assay of a benzodiazepine antagonist in plasma by capillary gas chromatography with nitrogen-selective detection 175

Zoer, J.

—, Virgili, P. and Henry, J.A.

High-performance liquid chromatographic assay for morphine with electro-

chemical detection using an unmodified silica column with a non-aqueous ionic eluent 189