

CHERN CLASSES OF VECTOR BUNDLES WITH HOLOMORPHIC CONNECTIONS ON A COMPLETE SMOOTH COMPLEX VARIETY

HÉLÈNE ESNAULT & V. SRINIVAS

Introduction

Let X be a complete smooth variety over the complex field \mathbf{C} , X_{an} the associated complex manifold, and \mathcal{E} a holomorphic vector bundle (locally free sheaf) on X_{an} with a holomorphic connection $\nabla: \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{O}_{X_{an}}} \Omega_{X_{an}}^1$, where $\Omega_{X_{an}}^1$ is the sheaf of holomorphic 1-forms on X_{an} . It is well known that \mathcal{E} has vanishing Chern classes in $H(X_{an}, \mathbf{Q})$, so that the integral Chern classes are torsion.

Recall that the i th Deligne complex $\mathcal{D}(i) = \mathcal{D}(i)X_{an}$ is defined by

$$0 \rightarrow \mathbf{Z}(i) \rightarrow \mathcal{O}_{X_{an}} \xrightarrow{d} \Omega_{X_{an}}^1 \xrightarrow{d} \dots \xrightarrow{d} \Omega_{X_{an}}^{i-1} \rightarrow 0,$$

where $\mathbf{Z}(i)$ is the subsheaf of abelian groups of the constant sheaf \mathbf{C} on X_{an} generated by $(2\pi\sqrt{-1})^i \mathbf{Z}$. The Deligne-Beilinson cohomology group (see [4] and the references given there) $H_{\mathcal{D}}^j(X_{an}, i)$ is defined to be the j th hypercohomology of $\mathcal{D}(i)$. Then there is an exact sequence

$$0 \rightarrow J^i(X) \rightarrow H_{\mathcal{D}}^{2i}(X_{an}, i) \xrightarrow{\rho} Hg^i(X_{an}) \rightarrow 0,$$

where $Hg^i(X_{an}) \subset H^{2i}(X, \mathbf{Z}(i))$ is the subspace of classes of Hodge type (i, i) (i.e., which maps to $F^i H^{2i}(X_{an}, \mathbf{C})$ in $H^{2i}(X_{an}, \mathbf{C})$, where F denotes the Hodge filtration), and $J^i(X)$ is the i th intermediate Jacobian of X , defined by

$$J^i(X) = H^{2i-1}(X_{an}, \mathbf{C}) / \{ \text{im} H^{2i-1}(X_{an}, \mathbf{Z}(i)) + F^i H^{2i-1}(X_{an}, \mathbf{C}) \}.$$

The topological Chern class $c_i(\mathcal{E}) \in Hg^i(X_{an}) \subset H^{2i}(X_{an}, \mathbf{Z}(i))$ is the image under ρ of the "refined" Chern class with values in Deligne-Beilinson cohomology,

$$c_i^{\mathcal{D}}(\mathcal{E}) \in H_{\mathcal{D}}^{2i}(X_{an}, i).$$

If \mathcal{E} has a connection ∇ , then $c_i(\mathcal{E})$ is the torsion, so for some integer $N > 0$, $Nc_i(\mathcal{E}) \in J^i(X)$.

For $i = 1$, $J^1(X) = \text{Pic}^0(X)$, the Picard variety of X , and it is a consequence of Hodge theory and GAGA that every element of $\text{Pic}^0(X)$ is the class of an invertible sheaf \mathcal{L} with an integrable connection.

For $i = 2$, Bloch [2] shows that the elements of $H^4_{\mathcal{D}}(X_{an}, 2)$, which are second Chern classes $c_2^{\mathcal{D}}(\mathcal{E})$ for locally free \mathcal{E} with an integrable connection, form a countable set. More precisely, he defines a countable subgroup $\Delta \subset C$ using the *dilogarithm* function, and shows that

$$Nc_2^{\mathcal{D}}(\mathcal{E}) \in \text{im}(H^3(X_{an}, \Delta) \rightarrow H^3(X_{an}, \mathbf{C}) \rightarrow J^2(X)),$$

where N is the exponent of $c_2(\mathcal{E})$ in $H^4_{\mathcal{D}}(X_{an}, \mathbf{Z}(2))$. He also comments on the relationships between his results and a conjecture of Cheeger and Simons, in the light of which he conjectures that $c_i^{\mathcal{D}}(\mathcal{E})$ is the *torsion* for all $i > 1$ for any locally free sheaf \mathcal{E} with integrable connection.

Our aim in this note is to prove the following result.

Theorem. *Let X be a smooth complete variety over \mathbf{C} . Then for any $i > 1$, the set*

$$\{c_i^{\mathcal{D}}(\mathcal{E}) \in H^{2i}_{\mathcal{D}}(X_{an}, i) \mid \mathcal{E} \text{ has a holomorphic connection}\}$$

is countable.

Note that we do not require the connections to be integrable.

1. Proof of the Theorem

We begin by noting that by GAGA,

(i) if \mathcal{E} is a locally free $\mathcal{O}_{X_{an}}$ -module of finite rank (i.e., a holomorphic vector bundle), then there is a locally free \mathcal{O}_X -module \mathcal{E}_0 , unique up to isomorphism, such that \mathcal{E} is the associated analytic sheaf;

(ii) if $\mathcal{E}, \mathcal{E}_0$ are as in (i), and ∇ is a holomorphic connection on \mathcal{E} , then there is an algebraic connection ∇_0 on \mathcal{E}_0 , unique up to isomorphism, such that the associated analytic connection on $(\mathcal{E}_0)_{an} \simeq \mathcal{E}$ is ∇ .

One way to see (ii) is as follows: if X is any smooth algebraic variety, and \mathcal{F} a locally free \mathcal{O}_X -module of finite rank, then consider the sheaf of algebraic 1-jets of the locally free \mathcal{O}_X -module \mathcal{F} , defined by

$$\mathcal{J}^1(\mathcal{F}) = p_*^2(p_1^* \mathcal{F} \otimes_{\mathcal{O}_{X \times X}} \mathcal{O}_{X \times X} / \mathcal{I}_{\Delta}^2),$$

where \mathcal{I}_{Δ} is the ideal sheaf of the diagonal on $X \times X$, and $p_i : X \times X \rightarrow X$

are the projections. The natural exact sequence (the *jet sequence*)

$$(*) \quad 0 \rightarrow \mathcal{F} \otimes_{\mathcal{O}_X} \Omega_{X/C}^1 \rightarrow \mathcal{J}^1(\mathcal{F}) \rightarrow \mathcal{F} \rightarrow 0 \quad \dots$$

obtained as p_{2*} of the sequence by tensoring

$$0 \rightarrow \mathcal{F}_\Delta / \mathcal{F}_\Delta^2 \rightarrow \mathcal{O}_{X \times X} / \mathcal{F}_\Delta^2 \rightarrow \mathcal{O}_\Delta \rightarrow 0$$

with $p_1^* \mathcal{F}$, yields an extension class

$$A(\mathcal{F}) \in \text{Ext}_X^1(\mathcal{F}, \mathcal{F} \otimes_{\mathcal{O}_X} \Omega_{X/C}^1) \simeq H^1(X, \mathcal{E}nd_{\mathcal{O}_X}(\mathcal{F}) \otimes_{\mathcal{O}_X} \Omega_{X/C}^1),$$

the *Atiyah class* of \mathcal{F} , whose vanishing is a necessary and sufficient condition for \mathcal{F} to have an algebraic connection (see [1]). In fact connections on \mathcal{F} are naturally in bijection with splittings of the jet sequence.

There is a corresponding Atiyah class for the existence of a holomorphic connection on \mathcal{F}_{an} , which lies in

$$H^1(X_{an}, \mathcal{E}nd_{\mathcal{O}_{X_{an}}}(\mathcal{F}_{an}) \otimes_{\mathcal{O}_{X_{an}}} \Omega_{X_{an}}^1)$$

where $\Omega_{X_{an}}^1$ is the sheaf of holomorphic 1-forms on X_{an} . Further, the jet sequence for \mathcal{F}_{an} is the sequence of holomorphic sheaves associated to the algebraic jet sequence, so $A(\mathcal{F}) \mapsto A(\mathcal{F}_{an})$ under the natural map on cohomology groups. By GAGA, if X is complete, then the map on cohomology is an isomorphism, and therefore in this case, if $A(\mathcal{F}_{an})$ vanishes, so does $A(\mathcal{F})$.

Hence, in (ii), we see that \mathcal{E}_0 has some algebraic connection ∇' . Now $\sigma = \nabla'_{an} - \nabla$ is a holomorphic section

$$\sigma \in H^0(X_{an}, \mathcal{E}nd_{\mathcal{O}_{X_{an}}}(\mathcal{E}) \otimes_{\mathcal{O}_{X_{an}}} \Omega_{X_{an}}^1).$$

Again by GAGA, any holomorphic section σ as above is of the form $\sigma = \tau_{an}$, where τ is an algebraic section

$$\tau \in H^0(X, \mathcal{E}nd_{\mathcal{O}_X}(\mathcal{E}_0) \otimes_{\mathcal{O}_X} \Omega_{X/C}^1);$$

now $\nabla_0 = \nabla' - \tau$ is an algebraic connection on \mathcal{E}_0 such that $(\nabla_0)_{an} = \nabla$.

The Atiyah class $A(\mathcal{F})$ is also related to the topological Chern classes $c_i(\mathcal{F}_{an}) \in Hg^i(X_{an}) \subset H^{2i}(X_{an}, \mathbf{Z}(i))$, as follows (see [1]—this relationship will be exploited in the proof of the Theorem). If X is any smooth algebraic variety over \mathbf{C} , and \mathcal{F} is locally free of finite rank on X , then the exterior product of differentials and composition of endomorphisms induces a map of sheaves

$$(\mathcal{E}nd_{\mathcal{O}_X}(\mathcal{F}) \otimes_{\mathcal{O}_X} \Omega_{X/C}^1)^{\otimes i} \rightarrow \mathcal{E}nd_{\mathcal{O}_X}(\mathcal{F}) \otimes_{\mathcal{O}_X} \Omega_{X/C}^i,$$

and hence a map on cohomology

$$\psi_i : H^1(X, \mathcal{E}nd_{\mathcal{O}_X}(\mathcal{F}) \otimes_{\mathcal{O}_X} \Omega_{X/\mathbb{C}}^1)^{\otimes i} \rightarrow H^i(X, \mathcal{E}nd_{\mathcal{O}_X}(\mathcal{F}) \otimes_{\mathcal{O}_X} \Omega_{X/\mathbb{C}}^i).$$

Let

$$\begin{aligned} M_i(\mathcal{F}) &= \psi_i(A(\mathcal{F})^{\otimes i}) = \psi_i(A(\mathcal{F}) \otimes \cdots \otimes A(\mathcal{F})) \\ &\in H^i(X, \mathcal{E}nd_{\mathcal{O}_X}(\mathcal{F}) \otimes_{\mathcal{O}_X} \Omega_{X/\mathbb{C}}^i), \end{aligned}$$

and let

$$N_i(\mathcal{F}) = \text{tr}(M_i(\mathcal{F})) \in H^i(X, \Omega_{X/\mathbb{C}}^i),$$

where ‘tr’ is the map on cohomology induced by the trace on the sheaf of endomorphisms. The classes $N_i(\mathcal{F})$ are the *Newton classes* of \mathcal{F} , where $N_i(\mathcal{F})$ is a polynomial (with integral coefficients) in the Chern classes

$$c_j(\mathcal{F}) \in H^j(X, \Omega_{X/\mathbb{C}}^j)$$

for $j \leq i$, such that $c_i(\mathcal{F})$ has a nonzero coefficient (in terms of the splitting principle, the Newton class N_i is the sum of the i th powers of the ‘Chern roots’). Conversely, the i th Chern class is a polynomial (with rational coefficients) in the Newton class $N_j(\mathcal{F})$ for $j \leq i$.

In particular, if the Atiyah class $A(\mathcal{F})$ vanishes (i.e., if \mathcal{F} has an algebraic connection), the Chern classes with values in $H^i(X, \Omega_{X/\mathbb{C}}^i)$ vanish.

If X is smooth and complete over \mathbb{C} , the topological Chern class $c_i(\mathcal{F}_{an})$ is compatible with the Chern class $c_i(\mathcal{F}) \in H^i(X, \Omega_{X/\mathbb{C}}^i)$ in the following way: Hodge theory and GAGA yield maps (the latter two are isomorphisms)

$$\begin{aligned} Hg^i(X_{an}) \rightarrow F^i H^{2i}(X, \mathbb{C}) \cap \bar{F}^i H^{2i}(X, \mathbb{C}) &\cong H^i(X_{an}, \Omega_{X_{an}}^i) \\ &\cong H^i(X, \Omega_{X/\mathbb{C}}^i) \end{aligned}$$

under which $c_i(\mathcal{F}_{an})$ maps to $c_i(\mathcal{F})$. Hence for smooth and complete X , $c_i(\mathcal{F}) = 0 \Leftrightarrow c_i(\mathcal{F}_{an})_{\mathbb{Q}} = 0$, where $c_i(\mathcal{F}_{an}) \mapsto c_i(\mathcal{F}_{an})_{\mathbb{Q}} \in H_g^i(X_{an}) \otimes \mathbb{Q} \subset H^{2i}(X_{an}, \mathbb{Q}(i))$.

More generally, if k is a field, $f : X \rightarrow S$ a smooth morphism of smooth k -varieties, and \mathcal{F} a locally free \mathcal{O}_X -module of finite rank, then one has the notion of an algebraic connection on \mathcal{F} relative to S , which is a map of sheaves

$$\mathcal{F} \rightarrow \mathcal{F} \otimes_{\mathcal{O}_X} \Omega_{X/S}^1$$

satisfying the Leibniz rule. There is a Atiyah class

$$A_S(\mathcal{F}) \in H^1(X, \mathcal{E}nd_{\mathcal{O}_X}(\mathcal{F}) \otimes_{\mathcal{O}_X} \Omega_{X/S}^1),$$

constructed using the ideal of the diagonal in $X \times_S X$, whose vanishing is equivalent to the existence of an algebraic connection on \mathcal{F} relative to S . This is compatible with the ‘global’ Atiyah class $A_k(X)$ (the obstruction to the existence of a connection relative to $\text{Spec } k$), in the sense that $A(\mathcal{F}) \mapsto A_S(\mathcal{F})$ under the map induced by the sheaf map

$$\Omega_{X/k}^1 \rightarrow \Omega_{X/S}^1.$$

Further, it is compatible with base change $S' \rightarrow S$, where S' is a smooth k -variety.

The following result is the main step in the proof.

Proposition 1. (Rigidity). *Let X be a smooth complete variety over \mathbb{C} , and Y a smooth connected variety over \mathbb{C} . Let \mathcal{E} be a locally free $\mathcal{O}_{X \times Y}$ -module of finite rank on $X \times Y$ which has a connection relative to the projection $p_2: X \times Y \rightarrow Y$. Then for any $i > 1$, the mapping*

$$c(i): Y \rightarrow H_{\mathcal{D}}^{2i}(X_{an}, i), \quad y \mapsto c_i^{\mathcal{E}}((\mathcal{E} \otimes \mathbb{C}(y))_{an}),$$

is constant.

Proof. To simplify the notation, we drop the subscript ‘an.’ Since any two points of Y lie on the image of a morphism from a connected smooth affine curve, we are reduced to the case where Y is an affine curve.

The map $c(i)$ has the following alternative description. One has the ‘algebraic’ Chern class $c_i^{CH}(\epsilon) = \xi_i \in CH^i(X \times Y)$, the Chow group of codimension i algebraic cycles on $X \times Y$ (see [5]), for example. The Chern class $c_i^{CH}(\mathcal{E} \otimes \mathbb{C}(y)) \in CH^i(X)$ is the image of ξ_i under the natural map

$$i_y^*: CH^i(X \times Y) \rightarrow CH^i(X),$$

where $i_y: X \rightarrow Y \times X$ is $i_y(x) = (x, y)$. The map $c(i)$ is then given by

$$c(i)(y) = Cl_{\mathcal{D}}(i_y^*(\xi_i)),$$

where

$$Cl_{\mathcal{D}}: CH^i(X) \rightarrow H_{\mathcal{D}}^{2i}(X, i)$$

is the cycle class map with values in Deligne-Beilinson cohomology. If we fix a base point $y_0 \in Y$, then the algebraic cycle $i_y^*(\xi_i) - i_{y_0}^*(\xi_i)$ is (co)homologous to 0 on X , and

$$c(i)(y) - c(i)(y_0) = Cl_{\mathcal{D}}(i_y^*(\xi_i)) - Cl_{\mathcal{D}}(i_{y_0}^*(\xi_i)) \in J^i(X),$$

the i th intermediate Jacobian of X ; one property of the cycle class in Deligne-Beilinson cohomology is that this element of $J^i(X)$ is the image of $i_y^*(\xi_i) - i_{y_0}^*(\xi_i)$ under the Abel-Jacobi mapping.

Let \bar{Y} be the projective smooth curve associated to Y , and let

$$\bar{\xi}_i \in CH^i(X \times \bar{Y})$$

be a preimage of ξ_i under the restriction map

$$CH^i(X \times \bar{Y}) \rightarrow CH^i(X \times Y).$$

Choose an algebraic cycle $\sum_j n_j Z_j$ representing ξ_i , and take $\bar{\xi}_i$ to be the class of $\sum_j n_j \bar{Z}_j$, where \bar{Z}_j is the Zariski closure Z_j . Then the Abel-Jacobi map gives a map from zero cycles of degree 0 on \bar{Y} to $J^i(X)$, by

$$\theta : \sum_j ((y_j) - (y_0)) \mapsto Cl_{\mathcal{O}} \left(\sum_j (i_y^*(\bar{\xi}_i) - i_{y_0}^*(\bar{\xi}_i)) \right) \in J^i(X),$$

whose value on $(y) - (y_0)$ is $c(i)(y) - c(i)(y_0)$ for $y \in Y$. The mapping θ clearly factors through the Jacobian of \bar{Y} , since $Cl_{\mathcal{O}}$ is well defined on rational equivalence classes, and so there is an induced mapping

$$[\bar{\xi}_i] : J(\bar{Y}) \rightarrow J^i(X).$$

We are reduced to proving this map is *constant*.

The mapping $[\bar{\xi}_i] : J(\bar{Y}) \rightarrow J^i(X)$ induced by the class $\bar{\xi}_i \in CH^i(X \times \bar{Y})$ is related to the topological cycle class of $\bar{\xi}_i$ in $H^{2i}(X \times \bar{Y}, \mathbf{Z}(i))$ in the following way (see Part One of the article [3] of Clemens and Griffiths). There is a Künneth component $\eta_i \in H^{2i-1}(X, \mathbf{Z}(i)) \otimes H^1(\bar{Y}, \mathbf{Z})$ of this topological cycle class (this Künneth component in fact depends only on ξ_i); its image in $H^{2i}(X \times \bar{Y}, \mathbf{C})$ lies in $F^i \cap \bar{F}^i$, where F^i is the Hodge filtration on $H^{2i}(X \times \bar{Y}, \mathbf{C})$. Under the isomorphism (Hodge theory)

$$F^i \cap \bar{F}^i \simeq H^i(X \times \bar{Y}, \Omega_{X \times \bar{Y}/\mathbf{C}}^i),$$

η_i is mapped to an element in the subspace

$$H^i(X, \Omega_{X/\mathbf{C}}^{i-1}) \otimes_{\mathbf{C}} H^0(\bar{Y}, \Omega_{\bar{Y}}^1) \oplus H^{i-1}(X, \Omega_{X/\mathbf{C}}^i) \otimes_{\mathbf{C}} H^1(\bar{Y}, \mathcal{O}_{\bar{Y}}),$$

and these two summands are the complex conjugates of each other. Hence we may write image $(\eta_i) = \mu_i + \bar{\mu}_i$ with

$$\mu_i \in H^i(X, \Omega_{X/\mathbf{C}}^{i-1}) \otimes_{\mathbf{C}} H^0(\bar{Y}, \Omega_{\bar{Y}}^1) \simeq \text{Hom}_{\mathbf{C}}(H^1(\bar{Y}, \mathcal{O}_{\bar{Y}}), H^i(X, \Omega_{X/\mathbf{C}}^{i-1})),$$

and $\bar{\mu}_i$ is the complex conjugate of μ_i , since their sum is a real cohomology class. Similarly we may regard η_i as an element of

$$\text{Hom}_{\mathbf{Z}}(H^1(\bar{Y}, \mathbf{Z}(1)), H^{2i-1}(X, \mathbf{Z}(i))).$$

This homomorphism is the mapping on lattices inducing the Abel-Jacobi map $[\bar{\xi}_i]: J(\bar{Y}) \rightarrow J^{2i-1}(X)$; the mapping μ_i , composed with the inclusion

$$H^i(X, \Omega_{X/C}^{i-1}) \simeq F^{i-1} \cap \bar{F}^i \hookrightarrow H^{2i-1}(X, \mathbf{C})/F^i H^{2i}(X, \mathbf{C}),$$

is the corresponding map of \mathbf{C} -vector spaces.

The upshot of this is that we are reduced to proving that $\mu_i = 0$.

Since $A_Y(\mathcal{E}) = 0$,

$$\begin{aligned} A(\mathcal{E}) &\in \ker(H^1(X \times Y, \mathcal{E}nd(\mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/C}^1) \\ &\rightarrow H^1(X \times Y, \mathcal{E}nd(\mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/Y}^1)). \end{aligned}$$

Now the natural map

$$\Omega_{X \times Y/C}^1 \rightarrow \Omega_{X \times Y/Y}^1$$

induces an isomorphism

$$p_1^* \Omega_{X/C}^1 \simeq \Omega_{X \times Y/Y}^1,$$

and similarly there is an isomorphism

$$p_2^* \Omega_{Y/C}^1 \simeq \Omega_{X \times Y/Y}^1.$$

This leads to a direct sum decomposition

$$\Omega_{X \times Y}^1 \simeq p_1^* \Omega_{X/C}^1 \oplus p_2^* \Omega_{Y/C}^1;$$

there is a similar decomposition on $X \times \bar{Y}$. This yields a direct sum decomposition

$$\begin{aligned} &H^1(X \times Y, \mathcal{E}nd(\mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/C}^1) \\ &= H^1(X \times Y, \mathcal{E}nd(\mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} p_1^* \Omega_{X/C}^1) \\ &\quad \oplus H^1(X \times Y, \mathcal{E}nd(\mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} p_2^* \Omega_{Y/C}^1) \end{aligned}$$

such that the Atiyah class $A(\mathcal{E})$ has components $A_Y(\mathcal{E}) = 0$ and $A_X(\mathcal{E})$ in the respective summands. Hence $A(\mathcal{E}) = A_X(\mathcal{E})$ lies in the subgroup

$$H^1(X \times Y, \mathcal{E}nd(\mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} p_2^* \Omega_{Y/C}^1).$$

Since Y is a curve, $\Omega_{Y/C}^i = 0$ for $i > 1$. Thus

$$M_i(\mathcal{E}) \in H^i(X \times Y, \mathcal{E}nd(\mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} \Omega_{X \times Y/C}^i)$$

lies in the subspace

$$H^i(X \times Y, \mathcal{E}nd(\mathcal{E}) \otimes_{\mathcal{O}_{X \times Y}} p_2^* \Omega_{Y/C}^i) = 0 \quad \text{for } i > 1.$$

Hence the Newton classes $N_i(\mathcal{E})$ vanish for $i > 1$. This implies that the Chern class

$$c_i(\mathcal{E}) \in H^i(X \times Y, \Omega_{X \times Y/C}^i)$$

is a rational multiple of $N_1(\mathcal{E})^i$, where

$$N_1(\mathcal{E}) \in H^1(X \times Y, \Omega_{X \times Y/C}^1).$$

But again $N_1(\mathcal{E})$ lies in the subspace

$$H^1(X \times Y, p_2^* \Omega_{Y/C}^1),$$

so

$$N_1(\mathcal{E})^i \in H^i(X \times Y, p_2^* \Omega_{Y/C}^i) = 0 \quad \text{for } i > 1.$$

We observe that the restriction map

$$H^i(X \times \bar{Y}, \Omega_{X \times \bar{Y}/C}^i) \rightarrow H^i(X \times Y, \Omega_{X \times Y/C}^i)$$

respects the decompositions

$$\begin{aligned} H^i(X \times \bar{Y}, \Omega_{X \times \bar{Y}/C}^i) &= H^i(X \times \bar{Y}, p_1^* \Omega_{X/C}^i) \\ &\quad \oplus H^i(X \times \bar{Y}, p_1^* \Omega_{X/C}^{i-1} \otimes_{\mathcal{O}_{X \times \bar{Y}}} p_2^* \Omega_{\bar{Y}/C}^1), \\ H^i(X \times Y, \Omega_{X \times Y/C}^i) &= H^i(X \times \bar{Y}, p_1^* \Omega_{X/C}^i) \\ &\quad \oplus H^i(X \times Y, p_1^* \Omega_{X/C}^{i-1} \otimes_{\mathcal{O}_{X \times Y}} p_2^* \Omega_{Y/C}^1). \end{aligned}$$

The summands

$$H^i(X \times \bar{Y}, p_1^* \Omega_{X/C}^i), \quad H^i(X \times \bar{Y}, p_1^* \Omega_{X/C}^{i-1} \otimes_{\mathcal{O}_{X \times \bar{Y}}} p_2^* \Omega_{\bar{Y}/C}^1)$$

further decompose respectively as

$$\begin{aligned} H^i(X, \Omega_{X/C}^i) \otimes H^0(\bar{Y}, \mathcal{O}_{\bar{Y}}) \oplus H^{i-1}(X, \Omega_{X/C}^i) \otimes H^1(\bar{Y}, \mathcal{O}_{\bar{Y}}), \\ H^i(X, \Omega_{X/C}^{i-1} \otimes H^0(\bar{Y}, \Omega_{\bar{Y}/C}^1) \oplus H^{i-1}(X, \Omega_{X/C}^{i-1}) \otimes H^1(\bar{Y}, \Omega_{\bar{Y}/C}^1). \end{aligned}$$

Thus

$$Cl(\bar{\xi}_i) \in H^i(X \times Y, \Omega_{X \times Y/C}^i)$$

is a sum of four components, two of which are μ_i and $\bar{\mu}_i$; in particular μ_i is the component in

$$H^i(X, \Omega_{X/C}^{i-1}) \otimes H^0(\bar{Y}, \Omega_{\bar{Y}/C}^1).$$

The restriction map

$$H^i(X \times \bar{Y}, p_1^* \Omega_{X/C}^{i-1} \otimes_{\mathcal{O}_{X \times \bar{Y}}} p_2^* \Omega_{\bar{Y}/C}^1) \rightarrow H^i(X \times Y, p_1^* \Omega_{X/C}^{i-1} \otimes p_2^* \Omega_{Y/C}^1)$$

is injective on the summand

$$H^i(X, \Omega_{X/C}^{i-1}) \otimes H^0(\bar{Y}, \Omega_{\bar{Y}/C}^1)$$

of the domain, and vanishes on the other summand. Since the restriction of $Cl(\bar{\xi}_i)$ vanishes for $i > 1$, μ_i restricts to 0 on $X \times Y$, i.e., $\mu_i = 0$, as desired. q.e.d.

We now give a short alternative proof using the construction of the Deligne-Beilinson cohomology $H_{\mathcal{D}}^j(i)$ of open smooth complex varieties (see [4]) but not using [3].

Alternative proof. As before, we reduce our proof to the case where Y is a smooth, connected affine curve, so that the Chern classes $c_i(\mathcal{E}) \in H^i(X \times Y, \Omega_{X \times Y}^i)$ vanish for $i \geq 2$. There is an exact sequence

$$\begin{aligned} 0 \rightarrow H^{2i-1}(X \times Y, \mathbf{C}/\mathbf{Z}(i))/F^i H^{2i-1}(X \times Y) \\ \rightarrow H_{\mathcal{D}}^{2i}(X \times Y, i) \rightarrow F_{\mathbf{Z}(i)}^i H^{2i}(X \times Y) \rightarrow 0, \end{aligned}$$

where

$$F_{\mathbf{Z}(l)}^j H^i := \{\omega \in F^j H^i \text{ such that image of } \omega \text{ vanishes in } H^i(\mathbf{C}/\mathbf{Z}(l))\}.$$

One has an exact sequence

$$0 \rightarrow F_{\mathbf{Z}(i)}^{i+1} H^{2i}(X \times Y) \rightarrow F_{\mathbf{Z}(i)}^i H^{2i}(X \times Y) \rightarrow H^i(X \times \bar{Y}, \Omega_{X \times \bar{Y}}^i(\log D)),$$

where \bar{Y} is as above, $D = X \times \{\infty\}$ with $\{\infty\} := \bar{Y} - Y$, and

$$\begin{aligned} H^i(X \times \bar{Y}, \Omega_{X \times \bar{Y}}^i(\log D)) \\ = H^i(X, \Omega_X^i) \otimes H^0(\bar{Y}, \mathcal{O}_{\bar{Y}}) \oplus H^i(X, \Omega_X^{i-1}) \otimes H^0(\bar{Y}, \Omega_{\bar{Y}}^1(\log\{\infty\})) \\ \oplus H^{i-1}(X, \Omega_X^i) \otimes H^1(\bar{Y}, \mathcal{O}_{\bar{Y}}). \end{aligned}$$

(Since Y is an affine curve, $H^1(\bar{Y}, \Omega_{\bar{Y}}^1(\log\{\infty\})) = 0$.) In this decomposition the image of $c_i^{\mathcal{D}}(\mathcal{E}) \in H_{\mathcal{D}}^{2i}(X \times Y, i)$ is written as $a_{i,i} + a_{i-1,i} + a_{i,i-1}$. From the vanishing image of $c_i^{\mathcal{D}}(\mathcal{E})$ in

$$\begin{aligned} H^i(X \times Y, \Omega_{X \times Y}^i) = H^i(X, \Omega_X^i) \otimes H^0(Y, \mathcal{O}_Y) \\ \oplus H^i(X, \Omega_X^{i-1}) \otimes H^0(Y, \Omega_Y^1), \end{aligned}$$

and from the injectivities of $H^0(\bar{Y}, \mathcal{O}_{\bar{Y}})$ and $H^0(\bar{Y}, \Omega_{\bar{Y}}^1(\log\{\infty\}))$ respectively in $H^0(Y, \mathcal{O}_Y)$ and $H^0(Y, \Omega_Y^1)$, it follows that $a_{i,i} = a_{i-1,i} = 0$.

As $a_{i-1,i} = a_{i,i-1}$, where the dual space to $H^1(\bar{Y}, \mathcal{O}_{\bar{Y}})$ is $H^0(\bar{Y}, \Omega_{\bar{Y}}^1)$ in $H^0(\bar{Y}, \Omega_{\bar{Y}}^1(\log\{\infty\}))$, one obtains that $c_i^{\mathcal{D}}(\mathcal{E})$ maps to $F_{\mathbf{Z}(i)}^{i+1}H^{2i}(X \times Y)$. As $F_{\mathbf{Z}(i)}^{i+1}H^{2i}(X \times \bar{Y}) = 0$, one has $F_{\mathbf{Z}(i)}^{i+1}H^{2i}(X \times Y) \hookrightarrow F_{\mathbf{Z}(i-1)}^iH^{2i-1}(X \times \{\infty\}) = 0$ via the Gysin sequence. Therefore $c_i^{\mathcal{D}}(\mathcal{E})$ comes from a class $\gamma_i \in H^{2i-1}(X \times Y, \mathbf{C}/\mathbf{Z}(i))$, with $\gamma_i = \alpha_i + \beta_i$, $\alpha_i \in H^{2i-1}(X, \mathbf{C}/\mathbf{Z}(i)) \otimes H^0(Y, \mathbf{Z})$, $\beta_i \in H^{2i-2}(X, \mathbf{C}/\mathbf{Z}(i)) \otimes H^1(Y, \mathbf{Z})$, and one has

$$\begin{aligned} c_i^{\mathcal{D}}(\mathcal{E}|_{X \times \{y\}}) &= c_i^{\mathcal{D}}(\mathcal{E})|_{X \times \{y\}} \text{ via the morphism} \\ &H_{\mathcal{D}}^{2i}(X \times Y, i) \rightarrow H_{\mathcal{D}}^{2i}(X \times \{y\}, i) \\ &= \text{image}(\gamma_i|_{X \times \{y\}}) \text{ via the morphism} \\ &H^{2i-1}(X \times Y, \mathbf{C}/\mathbf{Z}(i)) \rightarrow H_{\mathcal{D}}^{2i}(X \times Y, i) \\ &= \text{image}(\alpha_i|_{X \times \{y\}}) \text{ via the morphism} \\ &H^{2i-1}(X \times \{y\}, \mathbf{C}/\mathbf{Z}(i)) \rightarrow H_{\mathcal{D}}^{2i}(X \times \{y\}, i). \end{aligned}$$

The class of α_i is constant as desired. q.e.d.

The proof of the Theorem is now completed by a routine argument. Let $k \subset \mathbf{C}$ be a countable algebraically closed field of definition for X . Let X_0 be a model of X over k , i.e., a smooth complete k -variety with $X_0 \times_{\text{Spec } k} \text{Spec } \mathbf{C} = X$. First note that, up to isomorphism, there is only a countable number of locally free \mathcal{O}_{X_0} -modules \mathcal{E}_0 which have an algebraic connection over k . This is because there are in fact only countably many locally free \mathcal{O}_{X_0} -modules up to isomorphism over k (cover X_0 by infinitely many affine $\text{Spec } A_i$; there are only countably many projective A_i -modules up to isomorphism for each i , and only countably many possibilities for transition matrices).

Each locally free sheaf \mathcal{E}_0 defined over k and carrying a connection yields a locally free \mathcal{O}_X -module \mathcal{E} by extension of scalars. Clearly there are only countably many classes $c_i^{\mathcal{D}}(\mathcal{E})$ with \mathcal{E} of this special form.

By the rigidity result, it then suffices to prove that if \mathcal{F} is any locally free \mathcal{O}_X -module with a connection, there exist the following:

- (i) a connected smooth variety Y_0 defined over k , and the corresponding complex variety $Y = (Y_0)_{\mathbf{C}}$,
- (ii) a locally free $\mathcal{O}_{X_0 \times Y_0}$ -module \mathcal{E}_0 with a connection ∇_0 relative to Y_0 , and the corresponding objects \mathcal{E}, ∇ over \mathbf{C} , and
- (iii) a closed point $y \in Y$ such that $(\mathcal{E}, \nabla) \otimes \mathbf{C}(y)$ is isomorphic to the given locally free sheaf \mathcal{F} with its given connection.

Given this data, the Chern class $c_i^{\mathcal{G}}(\mathcal{F})$ equals $c_i^{\mathcal{G}}(\mathcal{E})$, where $y_0 \in Y_0$ is a closed point, regarded in a natural way as a \mathbf{C} -point of Y , and $\mathcal{E} = \mathcal{G} \otimes \mathbf{C}(y_0)$. Then $\mathcal{E}_0 = \mathcal{G}_0 \otimes k(y_0)$ is a locally free sheaf defined over k with a connection, and $(\mathcal{E}_0)_{\mathbf{C}} = \mathcal{E}$. This would prove the Theorem.

To make the claimed construction, note that \mathcal{F} and its given connection are defined over a finitely generated k -subalgebra K of \mathbf{C} . Let \mathcal{F}_K and ∇_K be corresponding objects over K . Let Y_0 be a smooth k -variety with function field K ; then X_K is the generic fiber of the proper and smooth morphism $X_0 \times_k Y_0 \rightarrow Y_0$. By replacing Y_0 by an open subset if necessary, we may further assume that there exists a locally free sheaf \mathcal{G}_0 on $X_0 \times Y_0$, with a connection relative to Y_0 , whose restriction to the generic fiber over Y_0 is \mathcal{F}_K , and with the connection ∇_K (to verify that the connection extends to an open set, one may think of it as a splitting of the jet sequence (*)). The given embedding $K \subset \mathbf{C}$ determines a closed point $y \in Y$, such that y maps to the generic point of Y_0 . If (\mathcal{G}, ∇) is the locally free sheaf with a connection relative to Y obtained on $X \times Y$, then $(\mathcal{G}, \nabla) \otimes \mathbf{C}(y) \simeq (\mathcal{F}_K, \nabla_K) \otimes_K \mathbf{C}$, which by choice is the sheaf \mathcal{F} with its given connection. Hence the proof is complete.

References

- [1] M. F. Atiyah, *Complex analytic connections in fibre bundles*, Trans. Amer. Math. Soc. **85** (1957) 181–207.
- [2] S. Bloch, *Applications of the dilogarithm function in algebraic K-theory and algebraic geometry*, Proc. Internat. Sympos. Algebraic Geometry (Kyoto, 1977), Kinokuniya, Tokyo, 1978.
- [3] H. Clemens & P. A. Griffiths, *The intermediate Jacobian of the cubic threefold* Ann. of Math. **95** (1972) 281–356.
- [4] H. Esnault & E. Viehweg, *Deligne-Beilinson cohomology*, Beilinson's conjectures on special values of L-functions, (M. Rapoport, P. Schneider & N. Schappacher, eds.), Academic Press, New York, 1988.
- [5] W. Fulton, *Intersection theory*, Ergebnisse Math. Grenzgeb. (3), B. 2, Springer, Berlin, 1984.

UNIVERSITÄT-GH-ESSEN

TATA INSTITUTE OF FUNDAMENTAL RESEARCH