JHEP HYPER VERSION

Received: *July 4, 2000*, Accepted: *September 5, 2000*

Measuring sparticle masses in non-universal string inspired models at the LHC

Ben C. Allanach

DAMTP, University of Cambridge
Wilberforce Road, Cambridge, CB3 0WA, UK
E-mail: b.c.allanach@damtp.cam.ac.uk

Christopher Gorham Lester, M.Andy Parker and Bryan R. Webber*

Cavendish Laboratory, University of Cambridge
Madingley Road, Cambridge, CB3 0HE, UK
E-mail: lester@hep.phy.cam.ac.uk, parker@hep.phy.cam.ac.uk, webber@hep.phy.cam.ac.uk

Abstract: We demonstrate that some of the suggested five supergravity points for study at the LHC could be approximately derived from perturbative string theories or M-theory, but that charge and colour breaking minima would result. As a pilot study, we then analyse a perturbative string model with non-universal soft masses that are optimised in order to avoid global charge and colour breaking minima. By combining measurements of up to six kinematic edges from squark decay chains with data from a new kinematic variable, designed to improve slepton mass measurements, we demonstrate that a typical LHC experiment will be able to determine squark, slepton and neutralino masses with an accuracy sufficient to permit an optimised model to be distinguished from a similar standard SUGRA point. The technique thus generalizes SUSY searches at the LHC.

Keywords: Supersymmetry Breaking, Beyond Standard Model, Supersymmetric Standard Model, Hadronic Colliders.

*Theory Division, CERN, 1211 Geneva 23, Switzerland.

JHEP09(2000)004

Contents

1. Introduction **1**

2. Theory **2**
- 2.1 Introduction 2
- 2.2 SUGRA point compatibility with string models 3
- 2.3 Optimized string model 5

3. Experimental observability **8**
- 3.1 Method 8
- 3.2 Other measurements 11
  - 3.2.1 Near, far, high and low $l^{\pm}q$ edges 11
  - 3.2.2 Zq edge 12
  - 3.2.3 M_{T2} 12
- 3.3 Event generation and detector simulation 13
- 3.4 Event selection cuts 14
  - 3.4.1 Cuts listed by observable 15
- 3.5 Edge resolutions 17
- 3.6 Reconstructing sparticle masses 23

4. Conclusions **25**

JHEP09(2000)004

1. Introduction

The purpose of this work is to extend the discussion of LHC supersymmetry (SUSY) searches to include string models. We begin by discussing whether string models can be used to motivate previous work on LHC SUSY searches, and then suggest a well-motivated non-universal string model for a new pilot study. We go on to examine how the SUSY particles can be detected and how the model can be distinguished from a similar well studied supergravity (SUGRA) model. We reconstruct sparticle masses by looking for kinematic edges in $\tilde{q}_L \to \tilde{\chi}_2^0 q \to \tilde{l}_R^{\pm} l^{\mp} q \to \tilde{\chi}_1^0 l^{\pm} l^{\mp} q$ and $\tilde{q}_L \to \tilde{\chi}_2^0 q \to \tilde{\chi}_1^0 X q \to \tilde{\chi}_1^0 l^{\pm} l^{\mp} q$ decay chains, and in doing so generalize the method of [1] by unifying the treatment of light and heavy sleptons. Additionally, with a novel method based on [2], we further constrain the $\tilde{\chi}_1^0$ and $\tilde{l}_R$ masses by studying the kinematics of events containing pair produced sleptons: $(gg/q\bar{q}) \to \tilde{l}_R^+ \tilde{l}_R^- \to l^+ \tilde{\chi}_1^0 l^- \tilde{\chi}_1^0$.

In particular, this allows the mass difference between the $\tilde{\chi}_1^0$ and $\tilde{l}_R$ to be determined with sufficient accuracy to permit discrimination between the string model and the most similar standard SUGRA model. We suggest that our analysis is likely to be applicable, not just to string motivated non-universal models, but to other non-universal models as well.

2. Theory

2.1 Introduction

Throughout this work we assume that the effective theory describing TeV scale physics is the R-parity conserving minimal supersymmetric standard model (MSSM). Within this framework, the collider phenomenology is strongly affected by the SUSY breaking terms

$$\mathcal{L} = \frac{1}{2}\sum_{a=1}^{3} M_a \lambda_a \bar{\lambda}_a - \sum_i m_i^2 |\phi_i|^2 - (A_{ijk} W_{ijk} + B\mu H_1 H_2 + \text{h.c..}) \,, \tag{2.1}$$

where $i, j, k = Q_L$, u_R^c, d_R^c, L_L, H_1, H_2 and ϕ_a, λ_a are the scalar and gaugino fields of the MSSM (see e.g. ref. [3]). W_{ijk} are the trilinear pieces of the MSSM superpotential, which written in terms of superfields is

$$W = \mathbf{Y}_E L H_1 \bar{E} + \mathbf{Y}_D Q H_1^b \bar{D} + \mathbf{Y}_U Q H_2 \bar{U} + \mu H_1 H_2 \,, \tag{2.2}$$

where we have suppressed all gauge and family indices. $\mathbf{Y}_D, \mathbf{Y}_E, \mathbf{Y}_U$ denote the down quark, charged lepton and up quark Yukawa matrices, respectively.

In eq. (2.1), a general parameterisation of possible SUSY breaking effects has been employed.[1] Usually, the parameters are constrained by the condition of universality,

$$m_i = m_0 \,, \qquad A_{ijk} = A_0 \,, \qquad M_a = M_{1/2} \tag{2.3}$$

deriving from simple SUGRA models. Equation (2.3) is subject to radiative corrections and should be imposed at the string scale M_S. In the usual formulation of perturbative string theory, this corresponds to $M_S \sim 5 \times 10^{17}\,\text{GeV}$, but eq. (2.3) is usually applied at the grand unified scale $M_{\text{GUT}} \sim 2 \times 10^{16}\,\text{GeV}$ as an approximation. Once B and μ are constrained by radiative electroweak symmetry breaking [5], the SUSY breaking sector is then characterised by one sign: $\text{sgn}\mu$, and four scalar parameters: $m_0, A_0, M_{1/2}$, and $\tan\beta$, the ratio v_2/v_1 of the two MSSM Higgs vacuum expectation values (VEVs). Once these are specified and current data are used to predict supersymmetric couplings such as the top Yukawa coupling and gauge couplings, the sparticle spectrum and decay chains are specified. Five points (denoted

[1] It has been assumed that non-standard terms such as those discussed in ref. [4] are disallowed because they cause a naturalness problem in the presence of gauge singlets.

JHEP09(2000)004

Model	m_0/GeV	$M_{1/2}$/GeV	A_0/GeV	$\tan\beta$	sgnμ	Weak	M-theory
S1	400	400	0	2	+	×	×
S2	400	400	0	10	+	×	×
S3	200	100	0	2	−	×	√
S4	800	200	0	10	+	×	√
S5	100	300	300	2.1	+	(√)	×

Table 1: Compatibility of LHC SUGRA points S1-S5 with strongly/weakly coupled string models. A tick in the "Weak"/"M-theory" column indicates that the sparticle spectrum could approximately be derived from the weakly/strongly coupled string models, respectively.

S1-S5) in the SUGRA parameter space $m_0, A_0, M_{1/2}$, sgnμ, $\tan\beta$ have been suggested for study of SUSY production at the LHC [6] and are catalogued in table 1. These models have been well studied in the context of the LHC [7]–[10], and we shall use them as a reference to compare and contrast with new models, which do not necessarily obey eq. (2.3).

2.2 SUGRA point compatibility with string models

In refs. [11, 12], the authors study the phenomenological viability of string and M-theory scenarios coming from the desirable absence of dangerous charge and colour breaking (CCB) minima or unbounded from below (UFB) directions in the effective potential. One of the models considered in [12] is weakly coupled string theory with orbifold compactifications. In this case, the soft masses evaluated at the string scale are dependent upon the modular weights n_i of the ϕ_i fields and do not necessarily conform with eq. (2.3) [13]. Other string scenarios exist [14, 15, 16] which erase the UFB/CCB global minima which we do not explicitly investigate. For all modular weights equal to -1 however, the tree-level pattern of soft-masses conforms with eq. (2.3), with the additional constraint

$$M_{1/2} = -A_0 = \sqrt{3} m_0 \,. \tag{2.4}$$

In table 1, we display under the "Weak" column whether each standard SUGRA point is approximately compatible with this sub-class of universal string models. None of S1-S5 fit exactly with this scenario, but S5 is the closest and would have a similar sparticle spectrum if m_0 were 173 GeV instead of 100 GeV. In regions of consistent radiative electroweak symmetry breaking (REWSB), the string model version of S5 has dangerous UFB minima [12]. We reject this class of model for further study, partly because S5 gives a similar spectrum, and partly because it is already well studied, but mainly because of the UFB problem in the potential mentioned above.

The "M-theory" column of table 1 refers to compatibility with the strong-coupling limit of $E_8 \times E_8$ Heterotic string theory [17]. Given simple assumptions

JHEP09(2000)004



Figure 1: Spectrum of M-theory valid at $M_{\rm GUT}$. (a) is valid for S1-S4, whereas (b) is valid for S5. Dotted lines show the relevant parameters for models that reproduce the LHC SUGRA point spectrum (S3-S5).

(that SUSY is spontaneously broken by the auxiliary components of the bulk moduli superfields), the soft terms of M-theory valid at $M_{\rm GUT}$ are [18]

$$M_{1/2} = \frac{\sqrt{3}M_{3/2}}{1+\epsilon}\left(\sin\theta + \frac{\epsilon}{\sqrt{3}}\cos\theta\right), \tag{2.5}$$

$$m_0^2 = M_{3/2}^2\left[1 - \frac{3}{(3+\epsilon)^2}\Big(\epsilon(6+\epsilon)\sin^2\theta + (3+2\epsilon)\cos^2\theta - 2\sqrt{3}\epsilon\sin\theta\cos\theta\Big)\right], \tag{2.6}$$

$$A_0 = -\frac{\sqrt{3}M_{3/2}}{3+\epsilon}\left[(3-2\epsilon)\sin\theta + \sqrt{3}\epsilon\cos\theta\right]. \tag{2.7}$$

So, the SUGRA parameters $M_{1/2}$, m_0, A_0 become replaced by the goldstino angle $\sin\theta$, the gravitino mass $M_{3/2}$ and a ratio of moduli VEVs $0 < \epsilon \leq 1$. $\epsilon = 0$ corresponds to the weakly-coupled perturbative string. For S1-S4, we notice from table 1 that $A_0 = 0$, allowing us to solve eq. (2.7):

$$\tan\theta = \frac{\sqrt{3}\epsilon}{2\epsilon - 3}. \tag{2.8}$$

Specifying ϵ then determines the ratio $|m_0/M_{1/2}|^2$, which is displayed in figure 1*a*. $|m_0/M_{1/2}|^2$ is always greater than one except as $\epsilon \to \infty$, ruling out M-theory derivations of S1 and S2. S3 and S4 are compatible with $\epsilon = 0.21$ and 0.53, respectively, as indicated in the figure.

Equation (2.8) is not relevant for S5 because $A_0 \neq 0$, but the condition $M_{1/2} = A_0$ may be solved yielding

$$\tan\theta = \frac{-\epsilon(3+2\epsilon)}{\sqrt{3}(3+\epsilon-\epsilon^2)}. \tag{2.9}$$

JHEP09(2000)004

$|m_0/M_{1/2}|^2$ is then plotted using this relation against ϵ in figure 1*b*. The figure indicates that M-theory does not reproduce S5 for any realistic value of ϵ.

To summarise, table 1 shows that S3-S4 are compatible with eqs. (2.5)–(2.7), and therefore M-theory. However, refs. [11, 12] show that each of the points corresponding to S3-S4 is in conflict either with UFB, CCB or REWSB constraints. In fact, all of the M-theory parameter space examined in ref. [12] was shown to be in conflict with one of these constraints.

We have therefore shown that while the LHC SUGRA points include models which may be derived from weakly or strongly coupled strings, they possess potentially catastrophic global CCB and/or UFB minima. The existence of a global CCB or UFB minimum does not necessarily rule out a model. Some models have metastable minima with lifetimes longer than the current age of the universe [19, 20, 21]. The question of which minimum the VEV of scalar fields rest in is one of cosmology, and beyond the scope of this paper. We therefore take the view that if the bounds from CCB/UFB global minima are not valid, examples of weakly/strongly coupled string models are approximated by or included within S1-S5 so there is no need to construct another similar model to examine the string-derived SUSY phenomenology at the LHC. If one should take the global CCB/UFB minimum bounds strictly however, ref. [12] indicates that no variation of parameters in the class of string models considered above will result in a model without CCB/UFB problems. We thus conclude that it is useful to examine a new class of string model which does not have difficulty evading CCB/UFB constraints. We note that it is also possible to evade the CCB/UFB constraints by lowering the string scale in type-I string models [15, 16].

2.3 Optimized string model

We now turn to the analysis of a model specifically designed to provide a large region of parameter space without CCB/UFB problems [12]. It is a weak coupling model, where the modular weights have been chosen so that

$$n_{Q_L} = n_{d_R^c} = n_{u_R^c} = -2\,, \qquad n_{L_L} = n_{e_R^c} = n_{H_{1,2}} = -1\,. \tag{2.10}$$

The model is non universal, but still in a family independent way, and so avoids serious problems associated with flavour changing neutral currents. As a consequence of the assignments in eq. (2.10), the string scale soft masses are

$$\begin{aligned} m^2_{H_{1,2}} = m^2_{L_L} = m^2_{e_R^c} = M^2_{3/2}\sin^2\theta\,, \qquad A_{L_L H_1 e_R^c} = -M_{1/2} = \sqrt{3}M_{3/2}\sin\theta\,, \\ m_{Q_L} = m_{d_R^c} = m_{u_R^c} = 0\,, \qquad A_t = M_{3/2}\left(\sqrt{2} - \frac{\sqrt{3}}{2}\right), \end{aligned} \tag{2.11}$$

so that the squarks are light and the sleptons heavy at the unification scale. This has the effect of ameliorating the CCB/UFB problem. It should be noted [12] that the

JHEP09(2000)004

Program	S1	S2	S3	S4	S5	O1
Slepton production						
ISAJET7.40	1.0×10^{-2}	1.1×10^{-2}	2.8×10^{-1}	8.7×10^{-4}	2.3×10^{-1}	1.2×10^{-1}
HERWIG6.0	1.0×10^{-2}	1.1×10^{-2}	2.7×10^{-1}	8.4×10^{-4}	2.2×10^{-1}	1.1×10^{-1}
SPYTHIA	1.1×10^{-2}	1.1×10^{-2}	3.1×10^{-1}	9.0×10^{-4}	2.5×10^{-1}	1.2×10^{-1}
Squark/gluino production						
ISAJET7.40	3.4×10^{0}	3.2×10^{0}	1.5×10^{3}	2.4×10^{1}	2.1×10^{1}	2.0×10^{1}
HERWIG6.0	3.2×10^{0}	3.2×10^{0}	1.4×10^{3}	2.3×10^{1}	2.0×10^{1}	1.7×10^{1}
SPYTHIA	3.7×10^{0}	2.8×10^{0}	1.3×10^{3}	2.0×10^{1}	1.7×10^{1}	1.6×10^{1}
Chargino/neutralino production						
ISAJET7.40	1.8×10^{-1}	2.1×10^{-1}	1.6×10^{1}	3.9×10^{0}	6.1×10^{-1}	6.7×10^{-1}
HERWIG6.0	2.1×10^{-1}	2.3×10^{-1}	1.8×10^{1}	3.9×10^{0}	7.1×10^{-1}	7.3×10^{-1}
SPYTHIA	2.1×10^{-1}	2.1×10^{-1}	1.6×10^{1}	4.0×10^{0}	6.5×10^{-1}	7.0×10^{-1}
Associated production						
ISAJET7.40	1.9×10^{-1}	1.8×10^{-1}	2.7×10^{1}	5.2×10^{-1}	9.4×10^{-1}	9.4×10^{-1}
HERWIG6.0	1.9×10^{-1}	1.9×10^{-1}	2.5×10^{1}	4.9×10^{-1}	9.6×10^{-1}	8.9×10^{-1}
SPYTHIA	2.1×10^{-1}	1.8×10^{-1}	2.7×10^{1}	4.7×10^{-1}	9.6×10^{-1}	8.9×10^{-1}

Table 2: Comparison of LHC SUSY production hard subprocess total cross sections (in picobarns) for the LHC SUGRA points S1-S5 and the optimized string model O1. By "associated production" we mean the production of a chargino or neutralino in association with a gluino or squark. CTEQ3L parton distributions were used and the statistical fractional error on each result is 2%.

A-terms of squarks other than the stop are not calculable for small $\tan\beta$. We shall approximate them here to be equal to A_t, but in fact they have a negligible effect upon the phenomenology/spectrum unless $\tan\beta$ is large, in which case $A_b = A_t$.

To be definite, we choose model parameters $\tan\beta = 10$, $M_{3/2} = 250\,\mathrm{GeV}$ and $\theta = \pi/4$. We call this optimized model O1 and numerically its parameters are

$$\begin{aligned} m_{H_{1,2}} = m_{L_L} = m_{e^c_R} = 177\,\mathrm{GeV}\,, \qquad -A_{L_L H_1 e^c_R} = M_{1/2} = 306\,\mathrm{GeV}\,, \\ m_{Q_L} = m_{d^c_R} = m_{u^c_R} = 0\,, \qquad A_{Q_L H_2 u^c_R} = A_{Q_L H_1 d^c_R} = 137\,\mathrm{GeV}\,. \end{aligned} \tag{2.12}$$

We make the approximation that these relations hold at $M_{\mathrm{GUT}} \sim 2\times10^{16}\,\mathrm{GeV}$, but it should be borne in mind that logarithmic corrections from renormalisation between $M_{\mathrm{string}} \sim 5\times10^{17}$ and M_{GUT} are expected. The spectrum and decay chains of the sparticles are calculated using ISASUGRA and ISASUSY [22] using eq. (2.12) as input.

Using three different Monte Carlo programs and the CTEQ3L parton distribution functions [23], we calculate the total cross sections of SUSY particles at the LHC for models S1-S5 and O1. Table 2 shows the comparison of HERWIG6.0 [2] total cross-section with those calculated by ISAJET7.40 [22] and SPYTHIA [25]. In each case, we

[2]This version of HERWIG, not officially released, was a developmental version of HERWIG6.1 [24].

JHEP09(2000)004



Figure 2: LHC SUSY production hard subprocess total cross sections for the LHC SUGRA points S1-S5 and the optimized string model O1. `HERWIG6.0` was used with the `CTEQ3L` parton distributions. Statistical fractional error on each point is 2%. Comparison with `SPYTHIA`, `ISAJET7.40` is shown in table 2.

M_g	m_{u_L}	$m_{u_R^c}$	m_{d_L}	$m_{d_R^c}$	m_{b_1}	m_{b_2}	m_{t_1}	m_{t_2}	m_{ν_e}	m_{e_L}	$m_{e_R^c}$	m_{ν_τ}
747	660	632	664	630	608	636	494	670	273	284	217	271
733	**654**	**631**	**657**	**628**	**600**	**629**	**460**	**671**	**230**	**239**	**157**	**230**
m_{τ_1}	m_{τ_2}	$m_{\chi_1^0}$	$m_{\chi_2^0}$	$m_{\chi_3^0}$	$m_{\chi_4^0}$	$m_{\chi_1^+}$	$m_{\chi_2^+}$	m_{h^0}	m_{H^0}	m_{A^0}	m_{H^+}	
209	285	−125	−234	371	−398	−233	−398	114	450	449	456	
157	**239**	−122	−233	**499**	−523	−232	−520	**94**	**612**	**607**	**612**	

Table 3: Comparison of sparticle and Higgs spectrum of O1 and **S5** (in bold type). These spectra were calculated using `ISASUGRA` and all masses are quoted in GeV. Sign conventions are as in `ISAJET`. The masses of the second family of sparticles are approximately equal to those of the first.

have used the `CTEQ3L` parton distribution functions and the spectrum is calculated by `ISASUGRA` with $m_t = 175\,\text{GeV}$. As can be seen from the table, the three Monte-Carlo programs agree to about 10%.

Figure 2 displays the result of the calculation using `HERWIG6.0`. It is noticeable from the figure that S5 and O1 have broadly similar SUSY production cross-sections except for the sleptons which are noticeably lower. This can be understood by comparing the spectra of the two models, displayed in table 3. The spectra are approximately similar, except for the sleptons which are heavier in O1 relative to

JHEP09(2000)004



Figure 3: "Sequential" decay.

the squarks. This is a consequence of the different choice of modular weights for the sleptons and the squarks. We therefore propose to analyse LHC SUSY production in O1 using S5 as a benchmark or comparison. We will show that the two models can be distinguished experimentally.

3. Experimental observability

3.1 Method

The primary experimental aim is to take previously developed model-independent methods for measuring SUSY particle masses (which were developed at standard SUGRA points) and by testing them in the context of a new optimised model, identify where these methods need to be generalised to perform well in both models. Secondarily it is to be shown that, after modifying these methods' treatments of the slepton sector, their performance is sufficient to distinguish between the optimised and standard SUGRA scenarios.

To accomplish these aims, some model-dependent assumptions have to be made. In this analysis, R-parity is taken to be conserved and certain sparticle decay chains are assumed to exist. In R-parity conserving (RPC) models, SUSY particles are only produced in pairs, and the lightest SUSY particle (LSP) is stable. SUSY events contain two of these LSPs which, being only weakly interacting, escape detection and lead to SUSY events with large amounts of missing transverse energy — the standard signature for RPC models. The fact that these massive particles go missing from all RPC SUSY events means that in these models it is not usually possible to measure particle masses by reconstructing entire decay chains. Thus, as in [1], endpoints in kinematic variables constructed from SUSY decay chains must instead be examined. Specifically, the "sequential" decay mode $\tilde{q}_L \to \tilde{\chi}^0_2 q \to \tilde{l}^{\mp}_R l^{\pm}_{\text{near}} q \to \tilde{\chi}^0_1 l^{\mp}_{\text{far}} l^{\pm}_{\text{near}} q$ (see figure 3) and the "branched" decay modes $\tilde{q}_L \to \tilde{\chi}^0_2 q \to \tilde{\chi}^0_1 X q \to \tilde{\chi}^0_1 l^{\pm} l^{\mp} q$ (see figure 4) form the starting point for this investigation.

The kinematic edges used in [1] to identify particle masses at S5 contain:

JHEP09(2000)004



Figure 4: "Branched" decays through the Z and Higgs bosons.

- l^+l^- edge: this picks out, from the "sequential" decays, the position of the very sharp edge in the dilepton invariant mass spectrum caused by $\tilde{\chi}_2^0 \to l\tilde{l}$ followed by $\tilde{l} \to l\tilde{\chi}_1^0$.

- l^+l^-q edge: in "sequential" decays, the llq invariant mass spectrum contains a linearly vanishing upper edge due to successive two-body decays. Our theoretical model of the l^+l^-q edge is more model independent than that used in earlier studies as it does do not assume any relation between the sparticle masses, other than the hierarchy $0 < m_{\tilde{\chi}_1^0} < m_{\tilde{l}_R} < m_{\tilde{\chi}_2^0} < m_{\tilde{q}}$ necessary for the "seqential" decay chain to exist at all.

- l^+l^-q threshold: this is the non-zero minimum in the "sequential" llq invariant mass spectrum, for the subset of events in which the angle between the two leptons (in the centre of mass frame of the slepton) is greater than $\pi/2$. This translates into the direct cut on m_{ll} described in section 3.4.1.

- hq edge: this is one of the two instances of the Xq edge. In general, the Xq edge is the upper edge of the distribution of the invariant mass of three visible particles in the "branched" decays of figure 4. The position of this edge is again determined by two-body kinematics. Depending on the Higgs mass and the mass difference between the $\tilde{\chi}_2^0$ and the $\tilde{\chi}_1^0$, one of the two "branched" decay chains will be strongly suppressed with respect to the other, so typically only one edge will be visible. At S5 the Higgs mode dominates, while the Z mode will dominate at O1.

As detailed later in this analysis, new edges are added to the list above. Two of them are more general versions of other edges treated in [1], while another is entirely new. They are the $l^{\pm}q$ high-, $l^{\pm}q$ low- and M_{T2} edges respectively. To summarise, the overall method we apply in this paper consists of:

As described later in this analysis, new edges are added to this list. Two of them are more general versions of other edges treated in [1], while another is entirely new. They are the $l^{\pm}q$ high-, $l^{\pm}q$ low- and M_{T2} edges, respectively. To summarise, the overall method we apply in this paper consists of:

- finding a model-independent set of cuts which can be used to select events from which the endpoints of all the kinematic edges may be measured,

JHEP09(2009)004

Related edge	Kinematic endpoint
l^+l^- edge	$(m_{ll}^{\max})^2 = (\tilde{\xi}-\tilde{l})(\tilde{l}-\tilde{\chi})/\tilde{l}$
l^+l^-q edge	$(m_{llq}^{\max})^2 = \begin{cases} \max\left[\frac{(\tilde{q}-\tilde{\xi})(\tilde{\xi}-\tilde{\chi})}{\tilde{\xi}}, \frac{(\tilde{q}-\tilde{l})(\tilde{l}-\tilde{\chi})}{\tilde{l}}, \frac{(\tilde{q}\tilde{l}-\tilde{\xi}\tilde{\chi})(\tilde{\xi}-\tilde{l})}{\tilde{\xi}\tilde{l}}\right] \\ \text{except for the special case in which } \tilde{l}^2 < \tilde{q}\tilde{\chi} < \tilde{\xi}^2 \\ \text{and } \tilde{\xi}^2\tilde{\chi} < \tilde{q}\tilde{l}^2 \text{ where one must use } (m_{\tilde{q}} - m_{\tilde{\chi}_1^0})^2. \end{cases}$
Xq edge	$(m_{Xq}^{\max})^2 = X + (\tilde{q}-\tilde{\xi})\left[\tilde{\xi} + X - \tilde{\chi} + \sqrt{(\tilde{\xi}-X-\tilde{\chi})^2 - 4X\tilde{\chi}}\right]/(2\tilde{\xi})$
l^+l^-q threshold	$(m_{llq}^{\min})^2 = \begin{cases} [2\tilde{l}(\tilde{q}-\tilde{\xi})(\tilde{\xi}-\tilde{\chi}) + (\tilde{q}+\tilde{\xi})(\tilde{\xi}-\tilde{l})(\tilde{l}-\tilde{\chi}) \\ \quad -(\tilde{q}-\tilde{\xi})\sqrt{(\tilde{\xi}+\tilde{l})^2(\tilde{l}+\tilde{\chi})^2 - 16\tilde{\xi}\tilde{l}^2\tilde{\chi}}]/(4\tilde{l}\tilde{\xi}) \end{cases}$
$l^{\pm}_{\text{near}}\ q$ edge	$(m_{l_{\text{near}}\ q}^{\max})^2 = (\tilde{q}-\tilde{\xi})(\tilde{\xi}-\tilde{l})/\tilde{\xi}$
$l^{\pm}_{\text{far}}\ q$ edge	$(m_{l_{\text{far}}\ q}^{\max})^2 = (\tilde{q}-\tilde{\xi})(\tilde{l}-\tilde{\chi})/\tilde{l}$
$l^{\pm}q$ high-edge	$(m_{lq(\text{high})}^{\max})^2 = \max\left[(m_{l_{\text{near}}\ q}^{\max})^2, (m_{l_{\text{far}}\ q}^{\max})^2\right]$
$l^{\pm}q$ low-edge	$(m_{lq(\text{low})}^{\max})^2 = \min\left[(m_{l_{\text{near}}\ q}^{\max})^2, (\tilde{q}-\tilde{\xi})(\tilde{l}-\tilde{\chi})/(2\tilde{l}-\tilde{\chi})\right]$
M_{T2} edge	$\Delta M = m_{\tilde{l}} - m_{\tilde{\chi}_1^0}$

Table 4: The absolute kinematic endpoints of invariant mass quantities formed from decay chains of the types mentioned in the text for known particle masses. The following shorthand notation has been used: $\tilde{\chi} = m^2_{\tilde{\chi}_1^0}, \tilde{l} = m^2_{\tilde{l}_R}, \tilde{\xi} = m^2_{\tilde{\chi}_2^0}, \tilde{q} = m^2_{\tilde{q}}$ and X is m_h^2 or m_Z^2 depending on which particle participates in the "branched" decay.

- obtaining an estimate of the accuracy with which the edge positions might be determined by an LHC experiment,
- performing chi-squared fits of the expected positions of these edges (as functions of the sparticle masses) to a set of "simulated edge measurements" one might expect from an ensemble of such experiments, and
- interpreting the results as the statistics contribution to model independent sparticle mass measurements of a typical LHC experiment.

Were all squarks to have the same mass, then (with perfect detector resolution and with infinite statistics) the endpoints of the edges listed above would be found at the positions given in table 4. Note that these positions depend on four unknown parameters, namely the masses of the squark, the slepton and the two neutralinos participating in each decay. The mass of the lightest Higgs boson also appears, but we assume this will already be known or will be obtained by other methods (e.g. [26]) to within 2%.

In a more realistic "non-degenerate squark masses" scenario, every distribution with a squark related edge will actually be a superposition of many under-

JHEP09(2000)004

lying distributions (one for each squark mass), each having its edge at a slightly different location. This results in some smearing of the edges, even before detector effects (resolutions, acceptances, jet energy calibrations etc.) are taken into account.

It is not possible to measure each of the squark masses separately. Consequently the quantity $m_{\tilde{q}}$ in table 4 will, after being obtained from the "smeared" edges, represent a squark mass *scale* rather than a specific squark mass. In all squark related edges, except the l^+l^-q threshold, the contribution to the outermost part of each edge is provided by the heaviest squarks. In the case of the l^+l^-q threshold, the "true" endpoint is set by the lightest squark. However, if (as at O1 and S5) the other eleven squarks are much heavier (see table 3), it is easier to measure the contribution coming from them. A strong correlation between $m_{\tilde{q}}$ and the mass of the heaviest squark would therefore be expected.

More work would be required to fully understand the "theoretical" systematic errors including the use of a full simulation of the expected edge positions. Better edge models than those used later in this analysis will definitely be needed to permit measurements of edge positions to be better correlated with functions of the particle masses. Other sources of systematic errors which require further analysis are the detector effects mentioned above, combinatorial backgrounds near the edges and possible cut biases. Such systematic errors can only meaningfully be studied when real data are available.

Although they are beyond the scope of this paper, it is assumed that the above investigations could be performed so as to leave the eventual edge resolutions determined only by statistics and detector resolution. In this analysis, then, all edges are fitted with simple shapes (see section 3.5) with the intention of extracting only an estimate of the *uncertainty* on the edge position, and not to obtain the edge position itself.

The first four edges listed in table 4 thus constitute a minimal constraint on the four unknown sparticle mass parameters, which may then be further constrained by the other measurements which follow.

3.2 Other measurements

3.2.1 Near, far, high and low $l^{\pm}q$ edges

In "sequential" decays there are three observable outgoing particles. There are only four different ways of grouping these particles together, and so only four different invariant masses may be formed from them: m_{ll}, m_{llq}, $m_{l_{\rm near}\ q}$ and $m_{l_{\rm far}\ q}$. The first two of these, m_{ll} and m_{llq}, may be formed from the observed momenta without knowledge of which lepton was $l_{\rm near}$ and which was $l_{\rm far}$; only the total lepton four-momentum is required. Edges in these invariant mass distributions have already been described. If it were possible to tag the near- and far-leptons separately, the

JHEP09(2000)004

third and fourth invariant mass combinations could also be formed unambiguously, allowing the positions of two more edges, $m^{\max}_{l_{\text{near}}\,q}$ and $m^{\max}_{l_{\text{far}}\,q}$, to play a part in the final fit. However, such tagging is impossible. If further information is to be gathered from these decays *in a model independent way*, it is then necessary to look for edges in variables (functions of $m_{l_{\text{near}}\,q}$ and $m_{l_{\text{far}}\,q}$) which *are* observable. On an event-by-event basis, then, we define

$$m_{lq(\text{high})} = \max\left(m_{l^+q}, m_{l^-q}\right) \equiv \max\left(m_{l_{\text{near}}\,q}, m_{l_{\text{far}}\,q}\right) \tag{3.1}$$

and

$$m_{lq(\text{low})} = \min\left(m_{l^+q}, m_{l^-q}\right) \equiv \min\left(m_{l_{\text{near}}\,q}, m_{l_{\text{far}}\,q}\right). \tag{3.2}$$

The simplest theoretical predictions for the positions of the corresponding edges, $m^{\max}_{lq(\text{high})}$ and $m^{\max}_{lq(\text{low})}$, are listed in table 4, along with the positions of the near- and far-edges. Note that although the position of the high-edge is just the higher of $m^{\max}_{l_{\text{near}}\,q}$ and $m^{\max}_{l_{\text{far}}\,q}$, the position of the low-edge has a more interesting form. This asymmetry arises because it is not always kinematically possible for the invariant mass of the lepton/quark pair coming from the lower near/far distribution to approach $\min[m^{\max}_{l_{\text{near}}\,q}, m^{\max}_{l_{\text{far}}\,q}]$ arbitrarily closely, while simultaneously requiring that this invariant mass is less than that of the other lepton/quark pair.

3.2.2 Zq edge

Since the neutralino mass difference at O1 (109 GeV) is too small to permit $\tilde{\chi}^0_2 \to h\tilde{\chi}^0_1$, "branched" decays are mediated by the Z. (Particle masses may be seen in table 3.) The cuts developed in [1] for picking out this special case of the Xq edge at S2 are found to perform equally well at O1, so they are adopted unchanged. Although it might be possible to benefit from adapting these cuts to O1 slightly, this temptation is resisted in order to retain model independence.

3.2.3 M_{T2}

In order to constrain the slepton and neutralino masses better, we construct another variable, $M_{T2}(\chi)$, whose distribution relates just these two masses. We base this on the variable, proposed in [2], which looks at events containing two identical two body decays: $xy \to X \to Ya_1a_2 \to Yb_1c_1b_2c_2$, where the particles of types a and c are of unknown mass, where particles of type c are undetectable, where the longitudinal momentum of the incoming particles is also unknown, and where it is assumed that Y does not contain any unobservable particles such as neutrinos. In such cases, the variable provides a kinematic constraint on the masses of a and c. We seek to apply this primarily to LHC dislepton events of the form $q\bar{q} \to \tilde{l}^+_R\tilde{l}^-_R \to l^+\tilde{\chi}^0_1 l^-\tilde{\chi}^0_1$ and so define our $M_{T2}(\chi)$ by:

$$M^2_{T2}(\chi) \equiv \min_{\not{\mathbf{p}}_1+\not{\mathbf{p}}_2=\not{\mathbf{p}}_T} \left[\max\left\{m^2_T\left(\mathbf{p}^{l_1}_T, \not{\mathbf{p}}_1, \chi\right), m^2_T\left(\mathbf{p}^{l_2}_T, \not{\mathbf{p}}_2, \chi\right)\right\}\right], \tag{3.3}$$

JHEP09(2000)004

where

$$m_T^2\left(\mathbf{p}_T^l, \mathbf{q}_T, \chi\right) \equiv m_l^2 + \chi^2 + 2\left(E_T^l E_T^\chi - \mathbf{p}_T^l \cdot \mathbf{q}_T\right), \tag{3.4}$$

$$E_T^l = \sqrt{{\mathbf{p}_T^l}^2 + m_l^2} \qquad \text{and} \qquad E_T^\chi = \sqrt{\mathbf{q}_T^2 + \chi^2}\,. \tag{3.5}$$

This definition includes the lepton masses for completeness, although these are neglected in all computations.

The value $M_{T2}(\chi)$ takes for a given candidate dislepton event is a function of: the transverse missing-momentum vector, $\not{\mathbf{p}}_T$; the transverse momentum vectors of the two leptons, $\mathbf{p}_T^{l_1}$ and $\mathbf{p}_T^{l_2}$; and one other parameter — an estimate of the neutralino mass, χ (not to be confused with the actual mass of the neutralino, $m_{\tilde{\chi}_1^0}$). Unlike the other parameters, the value of χ is not measured in each event — events may be reinterpreted for different values of χ. $M_{T2}(\chi)$ has the property that, for signal events in a perfect detector,

$$\max_{\text{events}}\left[M_{T2}\left(m_{\tilde{\chi}_1^0}\right)\right] = m_{\tilde{l}}\,. \tag{3.6}$$

Thus when χ is indeed $m_{\tilde{\chi}_1^0}$, then the distribution of M_{T2} has an end point at the slepton mass. Since the other observables allow $m_{\tilde{\chi}_1^0}$ to be measured in a model independent way, M_{T2} can then be included in the analysis to provide an additional constraint on the slepton/neutralino mass difference.

In practice the M_{T2} edge can be distorted by the finite resolution of the detector or missing energy from soft underlying events. Additionally, standard model (SM) backgrounds provide constraints on minimum detectable slepton/neutralino mass differences (see section 3.5). However the most important factor to control is the ability to correctly identify which particle species is contributing to an observed M_{T2} edge — particularly since it is possible to have multiple edges in the *non*-standard model contributions to M_{T2} distributions. At S5, for example, where both the right- and left-sleptons are lighter than at O1 (see table 3), both $\tilde{l}_R^+\tilde{l}_R^-$ and $\tilde{l}_L^+\tilde{l}_L^-$ events pass the cuts, and two edges are generated. The edge coming from the (lighter) right-slepton falls well under the SM background and so is not measurable, while the (heavier) left-slepton still has a cross section for pair-production high enough to let it form a good edge of its own at $M_{T2}(m_{\tilde{\chi}_1^0}) = m_{\tilde{l}_L}$. It is important that this edge is not mistaken for the $\tilde{l}_R$, so methods are needed to dismiss it. A detailed prescription of how to go about such a dismissal is beyond the scope of this paper, but it would clearly be accomplished by looking for inconsistency between a given edge-particle hypothesis and all the other sparticle masses, the branching ratios and (in particular) the strongly mass dependent pair-production cross sections, which could all be measured by other means.

3.3 Event generation and detector simulation

All events, except those from $q\bar{q} \to W^+W^-$ background processes, are simulated by `HERWIG6.0`. The W-pair events are generated by `ISAJET7.42` [22]. The detector

JHEP09(2009)004

chosen for simulation is the ATLAS detector [27, 28, 29], one of the two general purpose experiments scheduled for the LHC. The LHC is expected to start running at a luminosity of about $10^{33}\,\mathrm{cm}^{-2}\,\mathrm{s}^{-1}$ and this is expected to be increased over a period of about three years to the design luminosity of $10^{34}\,\mathrm{cm}^{-2}\,\mathrm{s}^{-1}$. These two periods are referred to, respectively, as the periods of low and high luminosity running.

The performance of the detector is simulated by `ATLFAST2.16` [30] which is primarily a fast calorimeter simulation which parametrises detector resolution and energy smearing and identifies jets and isolated leptons, in both the low and high luminosity environments. Throughout this analysis, the parameters controlling `ATLFAST`'s jet and lepton isolation criteria are left with the default values appropriate to the apparatus: i.e. jets must satisfy $p_T^j \geq 15\,\mathrm{GeV}$ and must lie in the pseudo-rapidity range $-5 \leq \eta^j \leq 5$, while electrons must have $p_T^l \geq 5\,\mathrm{GeV}$, muons $p_T^l \geq 6\,\mathrm{GeV}$ and both must lie in $-2.5 \leq \eta^l \leq 2.5$. For lepton isolation a maximum energy of 10 GeV may be deposited in a cone about the lepton of radius 0.2 in (η, ϕ)-space (ϕ being the azimuthal angle) while its separation from other jets must be at least 0.4 in the same units.

At high luminosity approximately 20 minimum bias events ("pile-up" events) are expected to occur in each bunch crossing. Pile-up events are not simulated by `ATLFAST2.16`, although it does alter its reconstruction resolutions to reflect the two different luminosity environments. It must be checked that any cuts applied at high luminosity will not be affected substantially by pile-up events. Within this article, events corresponding to $100\,\mathrm{fb}^{-1}$ are generated and are reconstructed in the high luminosity environment. This approximately corresponds to one year of high luminosity running.

3.4 Event selection cuts

Section 3.4.1 summarises all the cuts used to obtain the edges. The l^+l^- edge, l^+l^-q edge and l^+l^-q threshold cuts do not differ significantly from those used in [1]. The $l^\pm q$ edge cuts, however, do. The most significant change is the relaxation of the splitting requirement. The original splitting requirement tries to guarantee that the jet which comes from the quark produced in association with the observed dilepton pair is correctly identified. It achieves this by insisting that:

- both the dilepton pair and one of the two hardest jets j_1 and j_2 (ranked by p_T) are **consistent** with being the decay products of a squark (i.e. $m_{llj_i} < m_{\mathrm{cutoff}}$), and

- the invariant mass m_{llj_j} of the two leptons and the other of the two highest p_T jets is **inconsistent** with these being the decay products of a squark (i.e. $m_{llj_j} > m_{\mathrm{cutoff}}$).

JHEP09(2009)004

Although the above demand for inconsistency increases the purity of the signal, it has a significant detrimental effect on the efficiency. In our analysis the consistency requirement is retained but the inconsistency requirement dropped. Instead we require that m_{ll} is inside the expected region, given the l^+l^- edge measurement. We also perform background subtraction, modelling the opposite-sign same-lepton-family (OSSF) backgrounds by the distributions produced by those opposite-sign different-lepton-family (OSDF) events which pass the same cuts.

The production cross section for dislepton events suitable for M_{T2} analysis is, in all models, by far the smallest (see figure 2). So to show that M_{T2} events passing cuts are not in danger of being swamped by small variations in backgrounds or the cuts themselves, two set of cuts referred to as "hard" and "soft" are developed. The "hard" cuts attempt to maximise the signal to background ratio in the vicinity of the dislepton edge in the M_{T2} spectrum, while the much looser "soft" cuts try to maximise statistics at the edge at the expense of allowing in additional SUSY backgrounds.

3.4.1 Cuts listed by observable

This section summarises the cuts used in the analysis. For notational purposes, reconstructed leptons and jets are sorted by p_T. For example j_2, with corresponding transverse momentum $\mathbf{p}_T^{j_2}$, is the reconstructed jet with the second highest p_T. All cuts are requirements unless stated otherwise.

l^+l^- edge.

$n_{\text{leptons}} = 2$, both leptons OSSF and $p_T^{l_1} \geq p_T^{l_2} \geq 10\,\text{GeV}$.

$n_{\text{jets}} \geq 2$ and $p_T^{j_1} \geq p_T^{j_2} \geq 150\,\text{GeV}$. $\not{p}_T \geq 300\,\text{GeV}$.

The kinematics of OSSF leptons coming from background processes which produce uncorrelated leptons (for example tau decays) are modelled well by OSDF lepton combinations. Consequently, edge resolution is improved by "flavour subtracting" OSDF event distributions from OSSF event distributions.

l^+l^-q edge.

$n_{\text{leptons}} = 2$, both leptons OSSF and $p_T^{l_1} \geq p_T^{l_2} \geq 10\,\text{GeV}$.

$n_{\text{jets}} \geq 4$, $p_T^{j_1} \geq 100\,\text{GeV}$, and $p_T^{j_1} \geq p_T^{j_2} \geq p_T^{j_3} \geq p_T^{j_4} \geq 50\,\text{GeV}$.

$\not{p}_T \geq \max(100\,\text{GeV}, 0.2 M_{\text{effective}})$ and $M_{\text{effective}} \geq 400\,\text{GeV}$, where $M_{\text{effective}}$ is defined by the scalar sum:

$$M_{\text{effective}} = \not{E}_T + p_T^{j_1} + p_T^{j_2} + p_T^{j_3} + p_T^{j_4}\,. \tag{3.7}$$

Since the desired edge is a maximum, only the smaller of the two m_{llq} combinations which can be formed using j_1 or j_2 is used.

Flavour subtraction is employed here as at the l^+l^- edge.

JHEP09(2009)004

l^+l^-q threshold.

All the cuts for the l^+l^- edge as above.

In addition $m_{ll}^{\max}/\sqrt{2} \leq m_{ll} \leq m_{ll}^{\max}$, where the value of $m_{ll}^{\max}$ would be obtained from the l^+l^- edge in practice, but was determined theoretically in this analysis.

Since the desired edge is a minimum, only the larger of the two m_{llq} combinations which can be formed using j_1 or j_2 is used.

hq edge.

$n_{\text{leptons}} = 0$ and $\not{p}_T > 300\,\text{GeV}$.

Exactly two b jets with $p_T^{j_{b_1}} \geq p_T^{j_{b_2}} \geq 50\,\text{GeV}$. No other b jets, regardless of p_T.

At least two non-b jets with $p_T^{j_{q_1}} \geq p_T^{j_{q_2}} \geq 100\,\text{GeV}$, with at least one inside $-2.0 \leq \eta^j \leq 2.0$.

m_{bb} within 17 GeV of Higgs peak in m_{bb} spectrum.

Since the desired edge is a maximum, the non-b jet (i.e. j_{q_1} or j_{q_2}) chosen to form the m_{hq} invariant mass is that which minimises m_{hq}.

Zq edge.

$n_{\text{leptons}} = 2$, both leptons OSSF and $p_T^{l_1} \geq p_T^{l_2} \geq 10\,\text{GeV}$.

m_{ll} within 2.5 GeV of the centre of the Z-mass peak in the m_{ll} spectrum.

At least two non-b jets exist with $p_T^j \geq 100\,\text{GeV}$ and $\not{p}_T \geq 300\,\text{GeV}$.

Since the desired edge is a "maximum", the jet chosen to form the m_{Zq} invariant mass is the one (from those with $p_T^j \geq 100\,\text{GeV}$) which minimises m_{Zq}.

Again, flavour subtraction is used.

$l^\pm q$ high and $l^\pm q$ low edges.

All the cuts for the l^+l^-q edge above are required.

Additionally, events consistent with the l^+l^- edge measurement are selected by asking for $m_{ll} \leq m_{ll}^{\max} + 1\,\text{GeV}$.

To choose the jet from which to form $m_{l^\pm q}$ we select $j_i = j_1$ or j_2, whichever gives the smaller value of m_{llj}, and require $m_{llj_i} < m_{\text{cutoff}}$, where m_{cutoff} is chosen to be above the l^+l^-q edge, but is otherwise arbitrary. For easy comparison with [1], $m_{\text{cutoff}} = 600\,\text{GeV}$ was used in this analysis.

Finally, the two invariant mass combinations, $m_{l^\pm q}$, are assigned to $m_{lq(\text{high})}$ and $m_{lq(\text{low})}$ as defined earlier by eqs. (3.1) and (3.2).

JHEP09(2009)004

M_{T2} edge (hard cuts).

Events are required to have exactly one OSSF pair of isolated leptons satisfying $p_T^{l_1} > 50\,\text{GeV}$ and $p_T^{l_2} > 30\,\text{GeV}$.

$\delta_T < 20\,\text{GeV}$ is required, where $\delta_T = |\mathbf{p}_T^{l_1} + \mathbf{p}_T^{l_2} + \not{\mathbf{p}}_T|$. Large δ_T in signal events is indicative of an unidentifiable transverse boost to the centre of mass frame, perhaps due to initial state radiation. Large δ_T in other (not necessarily signal) events simply suggests an inconsistency with the desired event topology.

Events containing one or more jets with $p_T^j > 40$ to $50\,\text{GeV}$ are vetoed. This cut also helps to reduce standard model backgrounds, notably $t\bar{t}$. The lower this cut is placed, the better for the background rejection. However the cut cannot be placed too low (especially at high luminosity) due to the significant number of jets coming from the underlying event and other minimum bias events in the same bunch crossing. For order 25 minimum bias events per bunch crossing, [31] estimates that about 10% (1%) of bunch crossings will include a jet from the underlying event with a p_T greater than $40\,\text{GeV}$ ($50\,\text{GeV}$). Our results are not sensitive to variation of the jet veto cut between 40 and $50\,\text{GeV}$, where at least 90% signal efficiency is expected.

Events with $|m_{l_1 l_2} - m_Z| < 5\,\text{GeV}$ are vetoed to exclude lepton pairs from Z bosons.

$m_{l_1 l_2} > 80\,\text{GeV}$ and $\not{p}_T > 80\,\text{GeV}$ are also required.

M_{T2} edge (soft cuts).

These are as above, but

- the $\not{p}_T$ requirement is lowered from 80 to $50\,\text{GeV}$,
- the upper limit for δ_T is extended from 20 to $90\,\text{GeV}$, and
- the dilepton invariant mass cut is removed altogether.

3.5 Edge resolutions

By applying at S5 the cuts listed in section 3.4.1 we reproduce the m_{ll} and m_{llq} distributions whose edges are analyzed in [1] (see figure 5: *a*, *b* and *d*). We also confirm that the same cuts may also be used to generate edges of similar quality at O1 (see figure 6: *a*, *b* and *d*). In addition, plots c1 and c2 in figures 5 and 6 display the $m_{lq(\text{high})}$ and $m_{lq(\text{low})}$ distributions generated from the modified $l^{\pm}q$ edge cuts also listed above. Numerical results are summarised in table 5.

Similar pictures are seen at S5 and O1. The most obvious difference between the two sets of data is the peak at the Z mass present in figure 6*a* but absent in figure 5*a*. This peak comes from direct $\tilde{\chi}_2^0 \to \tilde{\chi}_1^0 Z \to \tilde{\chi}_1^0 l^+ l^-$ decays, which have a branching fraction of 39% at O1 compared with 0.65% at S5. The most

JHEP09(2009)004



Figure 5: The S5 distributions whose endpoints are described in table 4: (a) the l^+l^- edge, (b) the l^+l^-q edge, (c1) the $l^{\pm}q$ high-edge (c2) the $l^{\pm}q$ low-edge, (d) the l^+l^-q threshold and (e) the hq edge. Plots were produced with the cuts described in table 3.4.1. The number of events corresponds to $100\,\mathrm{fb}^{-1}$ of high luminosity running.

common direct neutralino decay mode at S5 is through the h_0 (65%) of which only a tiny proportion (0.02%) decay to the two light leptons — the partial decay width $\Gamma(h_0 \to ff)$ going approximately as m_f^2. Scenarios in which the l^+l^- edge happens to coincide with the Z peak require special treatment and are not considered here.

In order to make useful statements about the degree to which model parameters may be extracted from the endpoints and edges of these distributions, it is necessary

JHEP09(2000)004



Figure 6: The O1 distributions whose endpoints are described in table 4: (a) the l^+l^- edge, (b) the l^+l^-q edge, (c1) the $l^{\pm}q$ high-edge (c2) the $l^{\pm}q$ low-edge, (d) the l^+l^-q threshold and (e) the Zq edge. Plots were produced with the cuts described in table 3.4.1. The number of events corresponds to 100 fb^{-1} of high luminosity running.

to obtain an estimate of the accuracy with which these observables may be measured. It is expected that the errors on all of the observables considered here will eventually be statistics dominated, so simple fits have been made to the data to obtain estimates of the statistical errors on the edge or end point locations. The shapes fitted to the data (see figure 7) and the algorithms for determining the boundaries of the fitted regions have been kept as simple and generic as possible, with the intention of making

JHEP09(2000)004



Figure 7: Line-shapes used for fits to the mass distributions. Shape (a) (or its reflection in the vertical axis) is the standard "straight line" used to extract the l^+l^-q edge, $l^{\pm}q$ high-edge, $l^{\pm}q$ low-edge and l^+l^-q threshold resolutions. Shape (b) models the expected shape of the sharp l^+l^- edge in the absence of detector effects. The l^+l^- edge is actually fitted with shape (c), which is identical to (b) except that it is smeared with a gaussian resolution whose width is a free parameter of the fit.

	S5		O1		Table 4 values	
Endpoint	Fit	Fit error	Fit	Fit error	S5	O1
l^+l^- edge	109.10	0.13	70.47	0.15	109.12	70.50
l^+l^-q edge	532.1	3.2	544.1	4.0	536.7	530.5
$l^{\pm}q$ high-edge	483.5	1.8	515.8	7.0	464.2	513.6
$l^{\pm}q$ low-edge	321.5	2.3	249.8	1.5	337.0	231.3
l^+l^-q threshold	266.0	6.4	182.2	13.5	264.9	168.1
Xq edge	514.1	6.6	525.5	4.8	509.2	503.4
ΔM (M_{T2} edge)	——	——	——	10%	35.7	92.5

Table 5: Endpoints and associated fit uncertainties. The results of the naive fits shown in figures 5 and 6 may be seen in the "fit" and (one-sigma) "fit error" columns. For the reasons outlined in section 3.1 there is presently a lack of good theoretical predictions for these edge positions. (In this context a "good prediction" is one capable of taking into account, up to a level compatible with the statistical/fit errors, either the naivety of the fits, or the effects introduced by edge distortions and the presence of backgrounds.) The best guides presently available are the quantities listed in table 4, evaluated at $m_{\tilde{q}}$ equal to the largest squark mass. These values are also shown for comparison purposes. All values are GeV except where stated otherwise.

the fit results both conservative and simple to interpret. Given real data, it would be worth putting substantial effort into understanding how detector and physics effects affect the shape of each distribution in turn. This could significantly improve particle mass resolutions. The resolutions obtained from the fits to the distributions in figure 5 and figure 6 are listed in table 5.

The M_{T2} distributions obtained after the cuts listed in section 3.4.1 are shown in figure 8. The signal dislepton region having the desired edge is the unhatched region on each plot. Conveniently, most of the other SUSY events passing the cuts (mainly events containing gauginos) are also distributed with an edge located at a similar

JHEP09(2000)004



Figure 8: These M_{T2} distributions for O1 are generated from 100 events fb^{-1} at high luminosity, and from the M_{T2} cuts described in section 3.4. (a) and (b) use the "hard" cuts, while (c) and (d) use the "soft" cuts. In (a) and (c), events containing a jet with $p_T > 40\,\text{GeV}$ were vetoed. In (b) and (d), this cut was relaxed to 50 GeV. In all plots, $M_{T2}(\chi)$ is evaluated at $\chi = m_{\tilde{\chi}_1^0}$. The results are presented in the form $M_{T2}(m_{\tilde{\chi}_1^0}) - m_{\tilde{\chi}_1^0}$ in order to show the edge of the signal region located at the difference between the slepton and neutralino masses (92.5 GeV).

position to the disleptons'. This is due to the fact that such events often include pairs of slepton decays, and these may sometimes occur in combination with low jet activity and without additional leptons being produced inside detector acceptance. Consequently, for much of the SUSY background which passes cuts, the cuts are accepting M_{T2} values which do not degrade edge performance significantly. Note that $\Delta M(\chi) = M_{T2}(\chi) - \chi \geq 0$ is plotted evaluated at $\chi = m_{\tilde{\chi}_1^0}$, so this should have an edge at $\Delta M_{\text{max}} = m_{\tilde{l}} - m_{\tilde{\chi}_1^0}$. Of course, $m_{\tilde{\chi}_1^0}$ is not actually known *a priori*, so generating this graph in a real experiment requires an estimate of $m_{\tilde{\chi}_1^0}$ to be obtained by other means. For our purposes it will be sufficient if the position of

JHEP09(2009)004



Figure 9: These two plots are identical to figure 8*d* except that the true neutralino mass (125 GeV) is not presupposed. Instead, (a) uses $\chi = 70\,\text{GeV}$ and (b) uses $\chi = 170\,\text{GeV}$.

the edge of the distribution can be measured to about 10%, and this determines the accuracy required of the neutralino mass estimate χ. The width $\Delta M(\chi) = M_{T2}(\chi) - \chi$ remains approximately stable at the 10% level for $m_{\tilde{\chi}_1^0}/2 < \chi < 2m_{\tilde{\chi}_1^0}$. To illustrate the relative insensitivity of $\Delta M_{\max}(\chi)$ to χ near $m_{\tilde{\chi}_1^0}$, plots generated from the same data as before but with $\chi \approx m_{\tilde{\chi}_1^0} \pm 50\,\text{GeV}$ are shown in figure 9. Satisfying the 10% requirement above by obtaining a suitable value of χ within such a $\pm 40\%$ range is not difficult and may be accomplished by first performing a "cut down" version of the analysis described later in the text, but omitting the M_{T2} data, and then choosing the estimate, χ, to be the resulting reconstructed neutralino mass.

Determining the likely resolution for the M_{T2} edge requires a slightly different approach to that used for the other edges. Whereas the cuts used to obtain the invariant mass distributions have high SM rejections (primarily due to the presence of at least one high mass $\not{p}_T$ or $M_{\text{effective}}$ cut), the M_{T2} cuts cannot be so hard, primarily because the desired dislepton events have have very little jet activity. With the dislepton production cross sections typically two orders of magnitude smaller than the squark/gluino production cross sections (table 2) a low efficiency is not affordable. There are also irreducible SM backgrounds (primarily $W^+W^- \to l^+l^-\nu\bar{\nu}$ but also $t\bar{t} \to b\bar{b}W^+W^- \to jjl^+l^-\nu\bar{\nu}$ in cases where jets are below the p_T cut or outside detector acceptance) which have signatures identical to dislepton events. These backgrounds are clearly visible in figure 8 and would cause problems for naive straight line fitting techniques.

The standard model backgrounds in figure 8, although large, are easy to control. The SM edge is very clean, since the events which pass the cuts are well reconstructed and there are no noticeable tails. The SM edge will fall at m_W for $\chi = 0$, corresponding to the mass of the missing neutrino in SM events. As χ increases, the SM edge

JHEP09(2000)004



Figure 10: Variation with χ of the SM contributions to $\max(M_{T2}(\chi) - \chi)$ (for a number of events corresponding to $100\,\mathrm{fb}^{-1}$) which provides a measure of the minimum $\tilde{l}$-$\tilde{\chi}_1^0$ mass difference which is needed for signal M_{T2} events to be able to extend beyond the SM backgrounds. Events contributing to this plot were selected using the "hard" M_{T2} cuts.

recedes to lower masses as shown in figure 10, falling to 60 GeV for $\chi = 100\,\mathrm{GeV}$. A significant excess of events above this threshold would be a clear signal for non-SM processes. Such an excess will appear when the $\tilde{l}$-$\tilde{\chi}_1^0$ mass difference exceeds 80 GeV for low $m_{\tilde{\chi}_1^0}$ and 50 GeV for high $m_{\tilde{\chi}_1^0}$.

We estimate that, using M_{T2} with either set of cuts, it is possible to measure $\Delta M_{\max} = m_{\tilde{l}} - m_{\tilde{\chi}_1^0}$ to 10% or better. The "hard" M_{T2} cuts successfully remove almost all SUSY background above the SM threshold at the expense of only retaining half of the events passing the "soft" cuts. For both sets of cuts the SM threshold, at about 60 GeV, would be approximately three sigma away from $\Delta M_{\max}$ at this accuracy.

3.6 Reconstructing sparticle masses

Sparticle masses are reconstructed by performing a chi-squared fit with between four and six free parameters p. The main parameters of the fit are $m_{\tilde{\chi}_1^0}$, $m_{\tilde{l}}$, $m_{\tilde{\chi}_2^0}$ and $m_{\tilde{q}}$. Where appropriate, m_h and m_Z also appear as fit parameters, although if they do, they are strongly constrained (particularly in the case of the Z) by present LEP measurements or expected LHC errors (0.0077% and 2.0%, respectively).

The chi-squared, as a function of the free parameters p, is then formulated as:

$$\chi^2(p) = \sum_1^n \frac{(O_i^{\mathrm{sm}} - O_i(p))^2}{{\sigma_i}^2}\,, \tag{3.8}$$

where $O_i^{\mathrm{sm}} = O_i(p_{\mathrm{model}}) + \sigma_i X_i$ are n "smeared observables", $X_i \sim N(0,1)$ are the n random variables from the Normal distribution (with mean 0 and standard

JHEP09(2000)004

	Fractional RMS $\sqrt{\overline{\varepsilon_p^2}}$		Fractional mean $\overline{\varepsilon_p}/10^{-3}$		Reconstruction width $\sqrt{\overline{(m_p - \overline{m_p})^2}}$	
p	S5	O1	S5	O1	S5	O1
$\tilde{\chi}_1^0$	0.140	0.175	11.4	8.0	17 GeV	22 GeV
$\tilde{l}_R$	0.112	0.091	8.8	5.7	17 GeV	20 GeV
$\tilde{\chi}_2^0$	0.074	0.084	6.0	5.3	17 GeV	20 GeV
$\tilde{q}$	0.034	0.047	2.7	2.4	22 GeV	29 GeV

Table 6: RMS and mean values, for each of the particles p, of the fractional mass-error distributions (ε_p) in figures 13 and 14. Also shown are the widths of the distributions of the reconstructed particle masses (from figures 11 and 12).

deviation 1) which accomplish the smearing, σ_i is the anticipated statistical error for the i-th observable (approximated by the fit error listed in table 5), and $O_i(p)$ is the value one would expect for the i-th observable given the parameters p (for examples see table 4). p_{model} indicates the actual masses of the sparticles in the particular model being considered. The results of the fit, p_{fit}, are then those which minimise the chi-squared. p_{fit} may be interpreted as the sparticle masses which might be reconstructed by an LHC experiment after obtaining 100 fb^{-1} at high luminosity, with the X_i parametrising the experimental errors. In order to determine the accuracy with which this reconstruction can be performed, the above fitting process is repeated many times for different values of the X_i, producing the distributions of reconstructed particle masses which follow.

Figures 11 and 12 show the probability distributions expected for the reconstructed $\tilde{\chi}_1^0$, $\tilde{l}$, $\tilde{\chi}_2^0$ and $\tilde{q}$ masses, while figures 13 and 14 show the corresponding fractional errors, ε_p, for the same quantities. All are approximately gaussian and have reassuringly small tails. Statistics summarising these plots (widths, means and RMS values) are listed in table 6.

It may be seen that in both O1 and S5 the widths of the mass distributions for all four particles are very similar. This is because, in localized regions of parameter space, the edges tend to constrain mass differences far better than absolute masses. Evidence of this may be seen in figure 15 which shows the scatter of reconstructions in the $m_{\tilde{\chi}_1^0}$-$m_{\tilde{l}_R}$ plane. It is interesting to note that *without* the M_{T2} constraint at O1, the fit's chi-squared commonly has two distinct and competing comparable minima — one at high and one at low values of $m_{\tilde{l}_R}$. The simultaneous existence of these minima is a direct consequence of putting the near and far $l^{\pm}q$ edges on an equal footing in this analysis, allowing more than one interpretation for each of the high and low edges. The need to resolve this kind of ambiguity in model-independent investigations of this type illustrates the importance of establishing reliable model-independent ways of measuring the absolute scale of $m_{\tilde{l}_R} - m_{\tilde{\chi}_1^0}$ (or a related quantity) even if only to an accuracy of 20–30%.

JHEP09(2009)004



Figure 11: Reconstructed $\tilde{\chi}_1^0$, $\tilde{l}$, $\tilde{\chi}_2^0$ and $\tilde{q}$ masses at S5. The small arrows indicate the masses used as the input parameters, $p_{\rm model}$.

The clear gap between the reconstruction regions for O1 and S5 in figure 15 supports the original claim that, systematic errors permitting, it will be possible to distinguish between the two scenarios.

4. Conclusions

Some of the five standard LHC SUGRA points are compatible with universal perturbative string and M-theory, but dangerous CCB/UFB breaking minima are present in each example. We therefore studied a perturbative string model which is optimized to ameliorate the CCB/UFB problems present in the other models. The optimized model is non universal because the squarks and sleptons are split in mass at the string scale. We identify the SUGRA point with the most similar spectrum and hard

JHEP09(2000)004



Figure 12: Reconstructed $\tilde{\chi}_1^0$, $\tilde{l}$, $\tilde{\chi}_2^0$ and $\tilde{q}$ masses at O1. The small arrows indicate the masses used as the input parameters, p_{model}.

SUSY production cross sections (S5) to compare the optimized model with. The main difference is that the sleptons are heavier and therefore have lower production cross-sections. We have demonstrated the existence of a method by which an LHC experiment will be able to measure the masses of the (lighter) sleptons and the two neutralinos at O1 in a largely model independent way. In a specific comparison of S5 and O1 we have shown that this method will be able to distinguish a SUGRA model from an optimised string model with very similar properties.

More importantly, we expect that the techniques developed here are general enough to be used to discriminate between other pairings of optimised and non-optimised models with similar characteristics. The optimized model analysis applies to a more general class of models than the string model itself. We could apply it to models with non-universal SUSY breaking terms at M_{GUT} in which the squarks and sleptons are explicitly split in mass. These constitute a superset of the particular string model considered here.

JHEP09(2000)004



Figure 13: Fractional errors in reconstructed $\tilde{\chi}_1^0$, $\tilde{l}$, $\tilde{\chi}_2^0$ and $\tilde{q}$ masses at S5.

Acknowledgments

This work was partially supported by the U.K. Particle Physics and Astronomy Research Council. We thank D.J. Summers for helpful discussions. CGL also wishes to thank A.J. Barr, L.M. Drage, J.P.J. Hetherington and C. Jones for their help on numerous occasions.

References

[1] ATLAS collaboration, *Supergravity models*, in [32], May, 1999. sections 20.2.3-20.2.6.

[2] C.G. Lester and D.J. Summers, *Measuring masses of semi-invisibly decaying particles pair produced at hadron colliders*, *Phys. Lett.* **B 463** (1999) 99 [hep-ph/9906349].

JHEP09(2000)004



Figure 14: Fractional errors in reconstructed $\tilde{\chi}_1^0$, $\tilde{l}$, $\tilde{\chi}_2^0$ and $\tilde{q}$ masses at O1.

[3] H.E. Haber and G.L. Kane, *The search for supersymmetry: probing physics beyond the standard model*, *Phys. Rep.* **117** (1985) 75.

[4] I. Jack and D.R. T. Jones, *Non-standard soft supersymmetry breaking*, *Phys. Lett.* **B 457** (1999) 101 [hep-ph/9903365].

[5] A. Dedes, A.B. Lahanas and K. Tamvakis, *Radiative electroweak symmetry breaking in the MSSM and low-energy threshold*, *Phys. Rev.* **D 53** (1996) 3793 [hep-ph/9504239].

[6] LHCC, *supersymmetry workshop*, October 1996.

[7] E. Richter-Was, D. Froidevaux and J. Söderqvist, *Precision SUSY measurements with ATLAS for SUGRA points 1 and 2*, ATLAS Internal Note, 1997, PHYS-No-108.

JHEP09(2000)004



Figure 15: Reconstructed $m_{\tilde{l}}$ versus $m_{\tilde{\chi}_1^0}$ for O1 and S5. The stars show the true masses for each model.

[8] I. Hinchliffe et al., *Precision SUSY measurements at LHC: point 3*, ATLAS Internal Note, 1997, ATLAS-NOTE-Phys-109.

[9] F. Gianotti, *Precision SUSY measurements with ATLAS for SUGRA: point 4*, ATLAS Internal Note, 1997, PHYS-No-110.

[10] G. Polesello, L. Poggioli, E. Richter-Was and J. Söderqvist, *Precision SUSY measurements with ATLAS for SUGRA: point 5*, ATLAS Internal Note, 1997, PHYS-No-111.

[11] S.A. Abel and C.A. Savoy, *Charge and colour breaking constraints in the MSSM with non-universal SUSY breaking*, *Phys. Lett.* **B 444** (1998) 119 [hep-ph/9809498].

[12] J.A. Casas, A. Ibarra and C. Munoz, *Phenomenological viability of string and M-theory scenarios*, *Nucl. Phys.* **B 554** (1999) 67 [hep-ph/9810266].

[13] A. Dedes and A.E. Faraggi, *D-term spectroscopy in realistic heterotic-string models*, *Phys. Rev.* **D 62** (2000) 016010 [hep-ph/9907331].

[14] H. Baer, M.A. Diaz, P. Quintana and X. Tata, *Impact of physical principles at very high energy scales on the superparticle mass spectrum*, *J. High Energy Phys.* **04** (2000) 016 [hep-ph/0002245].

[15] S.A. Abel, B.C. Allanach, F. Quevedo, L. Ibáñez and M. Klein, *Soft SUSY breaking, dilaton domination and intermediate scale string models*, hep-ph/0005260.

[16] S.A. Abel and B.C. Allanach, *Quasi-fixed points and charge and colour breaking in low scale models*, *J. High Energy Phys.* **07** (2000) 037 [hep-ph/9909448].

JHEP09(2000)004

[17] P. Hořava and E. Witten, *Heterotic and type-I string dynamics from eleven dimensions*, *Nucl. Phys.* **B 460** (1996) 506 [hep-th/9510209].

[18] K. Choi, H.B. Kim and C. Munoz, *Four-dimensional effective supergravity and soft terms in M-theory*, *Phys. Rev.* **D 57** (1998) 7521 [hep-th/9711158].

[19] S.A. Abel and C.A. Savoy, *On metastability in supersymmetric models*, *Nucl. Phys.* **B 532** (1998) 3 [hep-ph/9803218].

[20] A. Kusenko, P. Langacker and G. Segre, *Phase transitions and vacuum tunneling into charge and color breaking minima in the MSSM*, *Phys. Rev.* **D 54** (1996) 5824 [hep-ph/9602414].

[21] A. Kusenko, *Color and charge breaking minima in the MSSM*, *Nucl. Phys.* **52A** *(Proc. Suppl.)* (1997) 67 [hep-ph/9607287].

[22] F.E. Paige, S.D. P. pescu, H. Baer and X. Tata, *Isajet 7.40: a Monte Carlo event generator for pp, $\bar{p}p$ and e^+e^- reactions*, hep-ph/9810440.

[23] H.L. Lai et al., *Global QCD analysis and the CTEQ parton distributions*, *Phys. Rev.* **D 51** (1995) 4763 [hep-ph/9410404].

[24] G. Corcella et al., *Herwig 6.1 release note*, hep-ph/9912396.

[25] S. Mrenna, *Spythia, a supersymmetric extension of Pythia 5.7*, *Comput. Phys. Commun.* **101** (1997) 232 [hep-ph/9609360].

[26] ATLAS collaboration, *Higgs bosons*, ch. 19, in [32], May, 1999.

[27] ATLAS collaboration, *ATLAS inner detector technical design report*, no. CERN/LHCC/97-16. April, 1997.

[28] ATLAS collaboration, *ATLAS calorimeter performance technical design report*, no. CERN/LHCC/96-40, December, 1996.

[29] ATLAS collaboration, *ATLAS muon spectrometer: Technical design report*, no. CERN-LHCC-97-22. 1997.

[30] E. Richter-Was, D. Froidevaux and L. Poggioli, *ATLFAST 2.0: a fast simulation package for ATLAS*, Tech. Rep. ATL-PHYS-98-131, 1998.

[31] G. Ciapetti and A. Di Ciaccio, *Monte carlo simulation of minimum bias events at the LHC energy*, in *Aachen LHC Workshop*, G. Jarlskog and D. Rein eds., vol. 2, pp. 155–162, Dec, 1990.

[32] ATLAS collaboration, *ATLAS detector and physics performance* Technical Design Report 2, no. CERN-LHCC-99-015 ATLAS-TDR-15. May, 1999.

JHEP09(2000)004