

# Constraints on Regge models from perturbation theory

---

**Jorge Mondejar**

*Dept. d'Estructura i Constituents de la Matèria, U. Barcelona,*

*Diagonal 647, E-08028 Barcelona, Spain*

*E-mail: mondejar@ecm.ub.es*

**Antonio Pineda**

*Grup de Física Teòrica and IFAE, Universitat Autònoma de Barcelona,*

*E-08193 Bellaterra, Barcelona, Spain*

*E-mail: pineda@ifae.es*

**ABSTRACT:** We study the constraints that the operator product expansion imposes on large  $N_c$  inspired QCD models for current-current correlators. We focus on the constraints obtained by going beyond the leading-order parton computation. We explicitly show that, assumed a given mass spectrum: linear Regge behavior in  $n$  (the principal quantum number) plus corrections in  $1/n$ , we can obtain the logarithmic (and constant) behavior in  $n$  of the decay constants within a systematic expansion in  $1/n$ . Our example shows that it is possible to have different large  $n$  behavior for the vector and pseudo-vector mass spectrum and yet comply with all the constraints from the operator product expansion.

**KEYWORDS:** Phenomenological Models, Asymptotic Freedom, QCD, Sum Rules.

---

## Contents

<b>1. Introduction</b>	<b>1</b>
<b>2. Correlators</b>	<b>2</b>
<b>3. Matching</b>	<b>3</b>
3.1 LO matching	4
3.2 NLO matching	6
3.3 NNLO matching	7
<b>4. Axial versus vector correlators</b>	<b>8</b>
<b>5. Numerical analysis</b>	<b>9</b>
<b>6. Conclusions</b>	<b>12</b>

---

## 1. Introduction

The Operator Product Expansion (OPE) has been used since long in order to gain information on the non-perturbative dynamics of the hadronic spectrum and decays [1–9]. In this article we revisit this problem. We want to obtain the constraints that the knowledge of the perturbative expansion in  $\alpha_s(Q^2)$  of the current-current correlators in the Euclidean regime poses on the relation between the decay constants and the mass spectrum for excitations with a large quantum number  $n$  (where  $n$  is the the quantum number of the bound state). We put special emphasis in going beyond the leading-order parton computation. We will work with a specific model for the hadronic spectrum. This is compulsory, since different spectral functions<sup>1</sup> may yield the same OPE expression, yet we believe some aspects of our discussion may hold beyond the assumptions of our model.

In order to have a well defined bound state it is crucial to consider the large  $N_c$  approximation [10]. This ensures infinitely narrow resonances at arbitrarily large energies. We will consider to be in the large  $N_c$  limit in what follows, as well as in the exact chiral (massless) limit. We will then set a specific model for the hadronic spectrum, valid for large values of  $n$  (we only need the behavior of the spectrum and decays for large  $n$ , we do not aim to get any information from perturbation theory for low values of  $n$ ). This model will be based on the Regge behavior plus corrections in  $1/n$  that will be included in a systematic way. The model is based on the assumption that the Regge behavior is a good description

---

<sup>1</sup>In particular the one derived directly from perturbation theory, which we do not consider, since we will work in the large  $N_c$  limit with infinitely narrow resonances.

of the spectrum for large  $n$  (this can be explicitly seen in the 't Hooft model [11] and it is also consistent with phenomenology). Given the  $1/n$  corrections to the mass spectrum, the expression of the correlator can also be written as a systematic expansion in  $1/n$ , where higher powers in  $1/n$  are equivalent to higher orders in  $1/Q^2$  in its OPE. By matching the OPE and hadronic expressions order by order in  $1/Q^2$ , we will be able to predict the logarithmic dependence on  $n$  of the decay constants (actually also the constant terms). This result can also be systematically organized within an expansion in  $1/n$  together with an expansion in  $1/\ln n$ . We will give explicit expressions up to order  $1/n^2$  and  $1/\ln^3 n$ . We will also make some numerical estimates of the impact of these corrections. Finally, we would like to stress that we are able to introduce power corrections in  $1/n$  to the Regge behavior and yet comply with the OPE. This is in contrast with ref. [5], where, besides the Regge behavior, only exponentially suppressed terms are introduced (parametrically smaller than any finite power of  $1/n$  for large  $n$ ). This parameterization is however fine if considered as a fit not emanated from the large  $n$  limit.

## 2. Correlators

For definiteness, we will consider the vector-vector correlator but most of the discussion applies to any other current-current correlator (axial-vector, scalar, ...).

$$\Pi_V^{\mu\nu}(q) \equiv (q^\mu q^\nu - g^{\mu\nu} q^2) \Pi_V(q) \equiv i \int d^4x e^{iqx} \langle vac | T \{ J_V^\mu(x) J_V^\nu(0) \} | vac \rangle, \quad (2.1)$$

where  $J_V^\mu = \sum_f Q_f \bar{\psi}_f \gamma^\mu \psi_f$ . In order to avoid divergences, we will consider the Adler function

$$\mathcal{A}(Q^2) \equiv -Q^2 \frac{d}{dQ^2} \Pi(Q^2) = Q^2 \int_0^\infty \frac{1}{(t+Q^2)^2} \frac{1}{\pi} \text{Im} \Pi_V(t) dt, \quad (2.2)$$

where  $Q^2 = -q^2$  is the Euclidean momentum.

Since we are working in the large  $N_c$  limit, the spectrum consists of infinitely narrow resonances, and the Adler function can be written in the following way

$$\mathcal{A}(Q^2) = Q^2 \sum_{n=0}^\infty \frac{F_V^2(n)}{(Q^2 + M_V^2(n))^2}. \quad (2.3)$$

On the other hand, for large positive  $Q^2$ , one may try to approximate the Adler function by its OPE, which reads

$$\begin{aligned} \mathcal{A}_{\text{OPE}}(Q^2) = \sum_f Q_f^2 \left[ \frac{4}{3} \frac{N_c}{16\pi^2} \left( 1 + \frac{3}{8} N_c \frac{\alpha_{\mathcal{A}}(Q^2)}{\pi} \right) \right. \\ \left. + \frac{C(\alpha_s(Q^2))}{Q^4} \beta(\alpha_s(\nu)) \langle vac | G^2(\nu) | vac \rangle + \mathcal{O} \left( \frac{1}{Q^6} \right) \right], \end{aligned} \quad (2.4)$$

where  $\alpha_{\mathcal{A}}(Q^2)$  admits an analytic expansion in terms of  $\alpha_s(Q^2)$  (computed in the  $\overline{\text{MS}}$  scheme),

$$\beta(\alpha_s) = -\beta_0 \frac{\alpha_s(Q^2)}{4\pi} - \beta_1 \left( \frac{\alpha_s(Q^2)}{4\pi} \right)^2 + \dots, \quad (2.5)$$

with  $\beta_0 = 11/3N_c$ ,  $\beta_1 = 34/3N_c^2$ ,  $\beta_2 = 2857/54N_c^3$ , and [12]

$$C(\alpha_s(Q^2)) = -\frac{2}{11N_c} \left( 1 - \frac{35}{22} N_c \frac{\alpha_s(Q^2)}{4\pi} + \dots \right). \quad (2.6)$$

### 3. Matching

High excitations of the QCD spectrum are believed to satisfy linear Regge trajectories:

$$\lim_{n \rightarrow \infty} \frac{M_{V,n}^2}{n} = \text{constant}.$$

For generic current-current correlators, such behavior is consistent with perturbation theory in the Euclidean region at leading order in  $\alpha_s$  if the decay constants are taken to be “constants”, ie. independent of the principal quantum number  $n$ .

The inclusion of subleading effects in  $\alpha_s$  can be incorporated into this model by changing the  $n$  dependence of the decay constants without changing the ansatz for the spectrum. The inclusion of these effects has consequences on subleading sum-rules and the relation with the non-perturbative condensates.

Here we would like to go beyond the analysis at leading order in  $\alpha_s$ , as well as to consider power-like corrections in  $1/n$ . We will consider that the large  $n$  expression for the mass spectrum can be organized within a  $1/n$  expansion in a systematic way starting from the asymptotic linear Regge behavior. In order to fix (and simplify) the problem we will assume that no  $\ln n$  term appears in the mass spectrum.<sup>2</sup> Therefore, we write the mass spectrum in the following way (for large  $n$ )

$$M_V^2(n) = \sum_{s=-1}^{\infty} B_V^{(-s)} n^{(-s)} = B_V^{(1)} n + B_V^{(0)} + \frac{B_V^{(-1)}}{n} + \dots, \quad (3.1)$$

where  $B_V^{(-s)}$  are constants. We will usually denote  $M_{V,LO}^2(n) = B_V^{(1)} n$ ,  $M_{V,NLO}^2(n) = B_V^{(1)} n + B_V^{(0)}$  and so on. To shorten the notation, we will denote  $B_V^{(1)} = B_V$ ,  $B_V^{(0)} = A_V$  and  $B_V^{(-1)} = C_V$ .

For the decay constants, we will have a double expansion in  $1/n$  and  $1/\ln n$ .

$$F_V^2(n) = \sum_{s=0}^{\infty} F_{V,s}^2(n) \frac{1}{n^s} = F_{V,0}^2(n) + \frac{F_{V,1}^2(n)}{n} + \frac{F_{V,2}^2(n)}{n^2} + \dots, \quad (3.2)$$

where the coefficients  $F_{V,s}^2(n)$  have a logarithmic dependence on  $n$ :

$$F_{V,s}^2(n) = \sum_{r=0}^{\infty} C_{V,s}^{(r)}(n) \frac{1}{\ln^r n}. \quad (3.3)$$

As we did with the masses, we will define  $F_{V,LO}^2(n) = F_{V,0}^2(n)$ ,  $F_{V,NLO}^2(n) = F_{V,0}^2(n) + \frac{F_{V,1}^2(n)}{n}$ , and so on. Note that in this case we also have an expansion in  $1/\ln n$ .

---

<sup>2</sup>This is a simplification. If one considers, for instance, the 't Hooft model [11],  $\ln n$  terms do indeed appear.

We are now in position to start the computation. Our aim is to compare the hadronic and OPE expressions of the Adler function within an expansion in  $1/Q^2$ , but keeping the logarithms of  $Q$ . In order to do so we have to arrange the hadronic expression appropriately. Our strategy is to split the sum over hadronic resonances into two pieces, for  $n$  above or below some arbitrary but formally large  $n^*$ , such that  $\Lambda_{\text{QCD}} n^* \ll Q$ . The sum up to  $n^*$  can be analytically expanded in  $1/Q^2$  and will not generate  $\ln Q^2$  terms (neither a constant term at leading order in  $1/Q^2$ ). For the sum from  $n^*$  up to  $\infty$ , we can use eqs. (3.1) and (3.2) and the Euler-MacLaurin formula to transform the sum in an integral plus corrections in  $1/Q^2$ . Whereas the latter do not produce logarithms, the integral does. These logarithms of  $Q$  are generated by the large  $n$  behavior of the bound states and the introduction of powers of  $1/n$  is equivalent (once introduced in the integral representation, and for large  $n$ ) to the introduction of (logarithmically modulated)  $1/Q^2$  corrections in the OPE expression.

Therefore, by using the Euler-MacLaurin formula, we write the Adler function in the following way ( $B_2 = 1/6$ ,  $B_4 = -1/30$ , ...)

$$\begin{aligned}
 A(Q^2) = & Q^2 \int_0^\infty dn \frac{F_V^2(n)}{(Q^2 + M_V^2(n))^2} + Q^2 \left[ \sum_{n=0}^{n^*-1} \frac{F_V^2(n)}{(Q^2 + M_V^2(n))^2} - \int_0^{n^*} \frac{F_V^2(n)}{(Q^2 + M_V^2(n))^2} \right] \\
 & + \frac{Q^2}{2} \frac{F_V^2(n^*)}{(Q^2 + M_V^2(n^*))^2} + Q^2 \sum_{k=1}^\infty (-1)^k \frac{|B_{2k}|}{(2k)!} \frac{d^{(2k-1)}}{dn^{(2k-1)}} \frac{F_V^2(n)}{(Q^2 + M_V^2(n))^2} \Bigg|_{n=n^*}, \quad (3.4)
 \end{aligned}$$

where  $n^*$  stands for the subtraction point we mentioned above, such that for  $n$  larger than  $n^*$  one can use the asymptotic expressions (3.1) and (3.2). This allows us to eliminate terms that vanish when  $n \rightarrow \infty$ . Note that the last sum in eq. (3.4) is an asymptotic series, and in this sense the equality should be understood.

Note also that for  $n$  below  $n^*$ , we will not distinguish between LO, NLO, etc... in masses or decay constants, since for those states we will not assume that one can do an expansion in  $1/n$  and use eqs. (3.1) and (3.2).

Finally, note that the expressions we have for the masses and decay constants become more and more infrared singular as we go to higher and higher orders in the  $1/n$  expansion. This is not a problem, since we always cut off the integral for  $n$  smaller than  $n^*$ . Either way, the major problems would come from the decay constants, since, in the case of the mass,  $Q^2$  effectively acts as an infrared regulator.

### 3.1 LO matching

We want to match the hadronic, eq. (3.4), and OPE, eq. (2.4), expressions for the Adler function at the lowest order in  $1/Q^2$ . This means that we have to consider the lowest order expressions in  $1/n$  for the masses and decay constants, i.e.  $F_{V,LO}^2(n)$  and  $M_{V,LO}^2(n)$ , since the corrections in  $1/n$  give contributions suppressed by powers of  $1/Q^2$ .

Only the first term in eq. (3.4) can generate logarithms or terms that are not suppressed by powers of  $1/Q^2$ . Therefore we obtain the following equality,

$$A^{pt.}(Q^2) \equiv Q^2 \int_0^\infty dn \frac{F_{V,LO}^2(n)}{(Q^2 + M_{V,LO}^2(n))^2} = \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2 \left( 1 + \frac{3}{8} N_c \frac{\alpha_A(Q^2)}{\pi} \right). \quad (3.5)$$

This equation can be fulfilled by demanding that

$$\frac{F_{V,LO}^2(n)}{|dM_{V,LO}^2(n)/dn|} = \frac{1}{\pi} \text{Im} \Pi_V^{\text{pert.}}(M_{V,LO}^2(n)). \quad (3.6)$$

By using the perturbative expression for  $\text{Im} \Pi_V^{\text{pert.}}$  (see [13]), we obtain

$$F_{V,LO}^2(n) = B_V \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2 \left\{ 1 + \frac{3}{8\pi} N_c \alpha_s(nB_V) + \frac{243 - 176 \zeta(3)}{128\pi^2} N_c^2 \alpha_s^2(nB_V) \right. \\ \left. + \frac{346201 - 2904\pi^2 - 324528 \zeta(3) + 63360 \zeta(5)}{27648\pi^3} N_c^3 \alpha_s^3(nB_V) + \mathcal{O}(\alpha_s^4(nB_V)) \right\}, \quad (3.7)$$

where  $\alpha_s(nB_V)$  should actually be understood as a function of  $\alpha_s(B_V)$  and  $\ln n$ . Therefore, it is obvious that the above expression is resumming powers of  $\alpha_s(B_V) \ln n$ :

$$F_{V,LO}^2(n) = B_V \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2 \left\{ 1 + \frac{3}{2} \frac{1}{1 + \frac{11}{3} N_c \frac{\alpha_s(B_V)}{4\pi} \ln(n)} N_c \frac{\alpha_s(B_V)}{4\pi} \right. \\ + \frac{(2673 - 1936 \zeta(3) - 408 \ln(1 + \frac{11}{12\pi} N_c \alpha_s(B_V) \ln(n)))}{88(1 + \frac{11}{3} N_c \frac{\alpha_s(B_V)}{4\pi} \ln(n))^2} \frac{N_c^2 \alpha_s^2(B_V)}{(4\pi)^2} \\ + \frac{N_c^3 \alpha_s^3(B_V)}{(4\pi)^3} \frac{1}{52272\pi(1 + \frac{11}{3} N_c \frac{\alpha_s(B_V)}{4\pi} \ln(n))^3} [-350427 N_c \alpha_s(B_V) \ln(n) \\ + 121\pi(346201 - 2904\pi^2 - 324528 \zeta(3) + 63360 \zeta(5)) \\ - 3672\pi(2877 - 1936 \zeta(3)) \ln(1 + \frac{11}{12\pi} \alpha_s(B_V) \ln(n)) \\ \left. + 749088\pi \ln^2(1 + \frac{11}{12\pi} \alpha_s(B_V) \ln(n)) \right] + \mathcal{O}(\alpha_s^4(B_V)) \left. \right\}. \quad (3.8)$$

Doing so we see that we are able to obtain the dependence of the decay constant in  $\ln n$  (somewhat we are assuming that  $\alpha_s(B_V)$  is a small parameter,  $B_V \sim 1 \text{ GeV}$ ).

We can also rewrite the decay constant as an expansion in  $1/\ln n$  by using the equality

$$\ln \tilde{n} = \frac{1}{\beta_0} \left( \frac{4\pi}{\alpha_s(nB_V)} + \frac{\beta_1}{\beta_0} \ln \left( \beta_0 \frac{\alpha_s(nB_V)}{4\pi} \right) + \left( \frac{\beta_2}{\beta_0} - \left( \frac{\beta_1}{\beta_0} \right)^2 \right) \frac{\alpha_s(nB_V)}{4\pi} \right), \quad (3.9)$$

where  $\tilde{n} = nB_V/\Lambda_{\overline{\text{MS}}}$ . We then obtain

$$F_{V,LO}^2(n) = B_V \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2 \left\{ 1 + \frac{9}{22} \frac{1}{\ln \tilde{n}} + \frac{1}{\ln^2 \tilde{n}} \left[ -\frac{459}{1331} \ln \ln \tilde{n} + \frac{144}{121} \left( \frac{243}{128} - \frac{11}{8} \zeta(3) \right) \right] \right. \\ + \frac{1}{\ln^3 \tilde{n}} \left[ \frac{46818}{161051} \ln^2 \ln \tilde{n} + \frac{459}{322102} (-2877 + 1936 \zeta(3)) \ln \ln \tilde{n} + \frac{42272605}{2576816} \right. \\ \left. \left. - \frac{3\pi^2}{22} - \frac{20283 \zeta(3)}{1331} + \frac{360 \zeta(5)}{121} \right] + \mathcal{O} \left( \frac{1}{\ln^4 n} \right) \right\}. \quad (3.10)$$

Note that the lowest contribution in  $1/\ln n$  for the decay constant,  $B_V \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2$ , which, usually, is the only one considered, reproduces the leading-order partonic prediction for the Adler function.

Note also that there is no problem with the Landau pole, even if the result is written in the form of eq. (3.10), since it holds only for  $n$  larger than an  $n^*$  such that  $\Lambda_{\overline{\text{MS}}} \ll n^* B_V$  (the integral has an infrared cutoff at  $n^*$ ).

Finally, we remind that, strictly speaking, we can only fix the ratio between the decay constant and the derivative of the mass. We have fixed this ambiguity by arbitrarily imposing the  $n$  dependence of the mass spectrum.

### 3.2 NLO matching

We now want to obtain extra information on the decay constant by demanding the validity of the OPE at  $\mathcal{O}(1/Q^2)$ , in particular the absence of condensates at this order. We then have to use the NLO expressions for  $M_V^2(n)$  and  $F_V^2(n)$ . With the ansatz we are using for the mass at NLO, it is compulsory to introduce (logarithmically modulated)  $1/n$  corrections to the decay constant if we want this constraint to hold. Note that it is possible to shift all the perturbative corrections to the decay constant.

Imposing that the  $1/Q^2$  condensate vanishes produces the following sum rule:

$$\begin{aligned}
 & A \frac{d}{dQ^2} \mathcal{A}^{pt.} - \frac{A}{Q^2} \mathcal{A}^{pt.} + \frac{1}{Q^2} \left[ \sum_{n=0}^{n^*-1} F_V^2(n) - \int_0^{n^*} dn F_{V,LO}^2(n) \right] + \frac{F_V^2(n^*)}{2Q^2} \\
 & + \frac{1}{Q^2} \sum_{k=1}^{\infty} (-1)^k \frac{|B_{2k}|}{(2k)!} \frac{d^{(2k-1)}}{dn^{(2k-1)}} F_V^2(n) \Big|_{n=n^*} - Q^2 \int_0^{n^*} dn \frac{F_{V,1}^2(n)/n}{(Q^2 + M_{V,LO}^2(n))^2} \\
 & + Q^2 \int_0^{\infty} dn \frac{F_{V,1}^2(n)/n}{(Q^2 + M_{V,LO}^2(n))^2} = 0.
 \end{aligned} \tag{3.11}$$

This equality should hold independently of the value of  $n^*$ , which formally should be taken large enough so that  $\alpha_s(n^* B_V) \ll 1$ , i.e. the limit  $\Lambda_{\overline{\text{MS}}} \ll n^* B_V \ll Q^2$ . Again, the meaning of the asymptotic series appearing in eq. (3.11) should be taken with care. If we forget about this potential problem, only a few terms in eq. (3.11) can generate  $\ln Q^2$  terms, which should cancel at any order. Those are the first two and the last two terms. Actually, the next to last term does not generate logarithms, but it allows to regulate possible infrared divergences appearing in the calculation. Therefore, asking for the cancellation of the  $1/Q^2$  suppressed logarithmic terms produced by the first two and the last term in eq. (3.11) fixes  $F_{V,1}^2$ . The non-logarithmic terms should also be cancelled but they cannot be fixed from perturbation theory.

One can actually find an explicit solution to the above constraint for  $F_{V,1}^2$  by performing some integration by parts. We obtain

$$\begin{aligned}
 \frac{F_{V,1}^2(n)}{n} &= \frac{A_V}{B_V} \frac{d}{dn} F_{V,0}^2(n) \\
 &= A_V \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2 \frac{1}{n} \left\{ -\frac{9}{22} \frac{1}{\ln^2 \tilde{n}} - \left[ \frac{459}{1331} (1 - 2 \ln(\ln \tilde{n})) + \frac{2187}{484} - \frac{36 \zeta(3)}{11} \right] \frac{1}{\ln^3 \tilde{n}} \right. \\
 &+ \frac{3}{2576816} [-45794053 + 351384\pi^2 + 41637552 \zeta(3) - 7666560 \zeta(5) \\
 &\left. - 3672 \ln(\ln \tilde{n}) (-3013 + 1936 \zeta(3) + 204 \ln(\ln \tilde{n})) \right] \frac{1}{\ln^4 \tilde{n}} + \mathcal{O}\left(\frac{1}{\ln^5 \tilde{n}}\right) \Big\},
 \end{aligned} \tag{3.12}$$

or in terms of  $\alpha_s(nB_V)$  or  $\alpha_s(B_V)$ ,

$$\frac{F_{V,1}^2(n)}{n} = A_V \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2 \frac{1}{n} \left\{ -\frac{11}{32\pi^2} N_c^2 \alpha_s^2(nB_V) - \frac{2877 - 1936\zeta(3)}{768\pi^3} N_c^3 \alpha_s^3(nB_V) \right. \quad (3.13)$$

$$\left. - \frac{11(376357 - 2904\pi^2 - 344112\zeta(3) + 63360\zeta(5))}{110592\pi^4} N_c^4 \alpha_s^4(nB_V) + \mathcal{O}(\alpha_s^5(nB_V)) \right\},$$

$$\frac{F_{V,1}^2(n)}{n} = A_V \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2 \frac{1}{n} \left\{ -\frac{11}{2} \frac{1}{\left(1 + \frac{11}{3} N_c \frac{\alpha_s(B_V)}{4\pi} \ln(n)\right)^2} N_c^2 \frac{\alpha_s^2(B_V)}{(4\pi)^2} \quad (3.14)$$

$$- \frac{(2877 - 1936\zeta(3) - 408 \ln\left(1 + \frac{11}{12\pi} N_c \alpha_s(B_V) \ln(n)\right))}{12\left(1 + \frac{11}{3} N_c \frac{\alpha_s(B_V)}{4\pi} \ln(n)\right)^3} N_c^3 \frac{\alpha_s^3(B_V)}{(4\pi)^3}$$

$$- \frac{1}{4752\pi\left(1 + \frac{11}{3} N_c \frac{\alpha_s(B_V)}{4\pi} \ln(n)\right)^4} N_c^4 \frac{\alpha_s^4(B_V)}{(4\pi)^4} [-233618 N_c \alpha_s(B_V) \ln(n)$$

$$+ 121\pi(376357 - 2904\pi^2 - 344112\zeta(3) + 63360\zeta(5))$$

$$- 3672\pi(3013 - 1936\zeta(3)) \ln\left(1 + \frac{11}{12\pi} \alpha_s(B_V) \ln(n)\right)$$

$$+ 749088\pi \ln^2\left(1 + \frac{11}{12\pi} \alpha_s(B_V) \ln(n)\right)] + \mathcal{O}(\alpha_s^5(B_V)) \right\}.$$

Note that besides the  $1/n$  suppression, we also have an extra  $\alpha_s^2(nB_V)$  suppression.

In principle one could think of the existence of  $1/n \times \text{constant}$  terms in the decay constant, i.e. without any associated logarithm. However, such terms produce  $\ln(Q^2)$  contributions in the Euclidean regime that do not appear in the perturbative computation, so they can be ruled out. This appears to be a general statement since  $1/n^m \times \text{constant}$  for any  $m$  integer also produces logarithms. Note that in order to give meaning to these integrals it is implicit that the integral over  $n$  has an infrared cutoff at  $n^*$ . Nevertheless, the logarithm does not appear to multiply powers of the infrared cutoff (as expected).

Finally, we would like to mention that, besides the constraints coming from the logarithmic related behavior of the OPE, there is also the constraint from its constant terms, which should sum up to zero. Nevertheless, for this constraint we cannot give a closed expression. This is due to the fact that the  $\ln Q^2$  independent terms may receive contributions from any subleading order in the  $1/n$  expansion of the masses and decay constants. The reason is that the decay constant at a given order in  $1/n$  is obtained after performing some integration by parts, which generate new ( $\ln Q^2$ -independent) terms that can be  $Q^2$  enhanced. This statement is general and also applies to any subleading power in the  $1/Q^2$  matching computation.

### 3.3 NNLO matching

We now consider expressions for the mass and decay constants at NNLO. For the first time we have to consider condensates. Simplifying terms that do not produce logs, we obtain



the following equation,

$$\frac{35}{121} \frac{\alpha_s(Q^2)}{4\pi} \frac{\beta(\alpha_s(\nu)) \langle vac | G^2(\nu) | vac \rangle}{Q^4} \tag{3.15}$$

$$\doteq Q^2 \int_{n^*}^{\infty} \frac{dn}{(Q^2 + B_V n)^2} \left[ \frac{F_{V,2}^2(n)}{n^2} - \frac{1}{B_V} \frac{d}{dn} \left( \frac{C_V F_{V,0}^2(n)}{n} + \frac{A_V F_{V,1}^2(n)}{2n} \right) \right],$$

where  $\doteq$  stands for the fact that we can only predict the  $\ln Q^2$  dependence. Constant terms are not fixed by this relation.

In order to get a more closed expression is convenient to use the following equality,

$$\frac{1}{Q^4} \alpha_s(Q^2) \doteq Q^2 \int_{n^*}^{\infty} \frac{dn}{(Q^2 + B_V n)^2} \frac{1}{B_V n^2} \frac{\beta_0}{8\pi} \alpha_s^2(n B_V), \tag{3.16}$$

valid up to terms that do not produce logarithms or those that are subleading.

We get then

$$F_{V,2}^2(n) = -C_V \frac{4}{3} \frac{N_c}{16\pi^2} \sum_f Q_f^2 \left\{ 1 + \frac{3}{8\pi} N_c \alpha_s(n B_V) \right. \tag{3.17}$$

$$+ \left[ \frac{287 - 176 \zeta(3)}{128\pi^2} - \frac{11 A_V^2}{64\pi^2 B_V C_V} - \frac{35}{88} \frac{\beta(\alpha_s(\nu)) \langle vac | G^2(\nu) | vac \rangle}{B_V C_V N_c^2} \right] N_c^2 \alpha_s^2(n B_V)$$

$$\left. + \mathcal{O}(\alpha_s^3(n B_V)) \right\}.$$

Note that in this case we only consider up to  $\mathcal{O}(\alpha_s^2(n B_V))$  corrections, since higher order loops are unknown. The accuracy is set by our knowledge of the matching coefficient of the gluon condensate. Note also that  $F_{V,2}^2(n)$  does not have  $\alpha_s$  suppression. Therefore, for low  $n$ , this contribution could be practically of the same size than, formally, more important terms.

#### 4. Axial versus vector correlators

The above discussion has been performed for the vector-vector correlator Adler function. It goes without saying that we could perform a similar analysis with axial-vector currents, since the perturbative expansions for both correlators are equal. Here it comes an important observation. We could change the coefficients for the mass spectrum  $B_A \neq B_V$ ,  $A_A \neq A_V$ ,  $\dots$ , yet we would obtain the same expression for the OPE (at the order we are working, the first chiral breaking related effects are  $\mathcal{O}(1/Q^6)$ ). Therefore, we conclude that the OPE does not fix  $B_A = B_V$  as it is sometimes claimed in the literature [1, 3].<sup>3</sup> Our computation gives a specific counter example. Moreover, it is nice to see what the role played by  $B_A$  and  $B_V$  is in our case. When one goes to the Euclidean regime,  $B_A$  and  $B_V$  become renormalization factorization scales and, obviously, the physical result does not depend on them (for large  $Q^2$  in the Euclidean). On the other hand, it is evident that having different constants:  $B_A, B_V, \dots$  produces different physical predictions for the

---

<sup>3</sup>Another issue, on which we do not enter, is whether some other kind of arguments (relying on the specific model used), like semiclassical arguments, may fix those parameters to be equal.

masses and decay widths for vector or axial-vector channels. The point to be emphasized is that  $B_A = B_V$  cannot come from an OPE analysis alone. This point has already been stressed in refs. [4, 8], what we think is novel in our analysis is that we have seen that the inclusion of corrections in  $\alpha_s$  does not affect that conclusion, and that  $B_A$  and  $B_V$  play the role of the renormalization scale in the analogous perturbative analysis in the Euclidean regime, and are therefore unobservable. Finally, we cannot avoid mentioning the analysis of ref. [14] where, using AdS/CFT, they explicitly find Regge behavior with different slopes for vector and axial-vector channels.

In any case, even though the constants that characterize the spectrum can be different for the vector and axial-vector channel, they have to yield the same expressions for the OPE when combined with the decay constants. This produces some relations that we list in what follows. We first define  $t \equiv B_V n = B_A n'$  and take  $n$  and  $n'$  as continuous variables. We then obtain the following equalities,

$$\frac{F_{V,LO}^2(n)}{B_V} = \frac{F_{A,LO}^2(n')}{B_A} = \frac{1}{\pi} \text{Im} \Pi_V^{pert.}(t) \equiv f_0(t), \tag{4.1}$$

$$\frac{1}{A_V B_V} \frac{F_{V,1}^2(n)}{n} = \frac{1}{A_A B_A} \frac{F_{A,1}^2(n')}{n'} = \frac{d}{dt} f_0(t), \tag{4.2}$$

$$\begin{aligned} \frac{1}{B_V} \left[ \frac{F_{V,2}^2(n)}{n^2} - \frac{1}{B_V} \frac{d}{dn} \left( \frac{C_V F_{V,0}^2(n)}{n} + \frac{A_V F_{V,1}^2(n)}{2n} \right) \right] \\ = \frac{1}{B_A} \left[ \frac{F_{A,2}^2(n')}{n'^2} - \frac{1}{B_A} \frac{d}{dn'} \left( \frac{C_A F_{A,0}^2(n')}{n'} + \frac{A_A F_{A,1}^2(n')}{2n'} \right) \right] \\ = \frac{1}{t^2} \beta(\alpha_s(\nu)) \langle vac | G^2(\nu) | vac \rangle f_1(t), \end{aligned} \tag{4.3}$$

where


$$f_1(t) = \frac{35}{121} \frac{\beta_0}{2} \frac{\alpha_s^2(t)}{(4\pi)^2} + \dots \tag{4.4}$$

## 5. Numerical analysis

We restrict ourselves to the SU(2) case (non-strange sector) and study the vector and axial-vector channels. We would like here to assess the importance of including perturbative corrections to a standard analysis based on the OPE. We do not aim to perform a full fledged analysis, but only to see the importance of the corrections. In table 1, we give the values of the masses and decay constants. In figure 1 we show the changes in both  $F_{V,LO}(I)$  and  $F_{V,LO}(II)$  as we include higher orders in the expansion in  $\alpha_s$ , and in figure 2 the changes in the full  $F_V(I)$  and  $F_V(II)$  as we include higher orders in  $1/n$ . In figure 3 we show the same plots for the axial-vector case. We take the experimental values from ref. [15]. In principle there are more states in the particle data book, in particular in the vector channel. Nevertheless, it is not clear whether they belong to the same Regge trajectory or whether they belong to some daughter one, see, for instance, the discussion in ref. [5]. For the time being we will disregard the study of other possible (vector) Regge trajectories and restrict the analysis to a single trajectory. We will consider the two possibilities listed in table 1.

	$n = 1$	$n = 2$	$n = 3$	$n = 4$
$M_\rho(\text{I})$	781.3(775.5 $\pm$ 0.4)	1440.2(1459 $\pm$ 11)	1891.8(1870 $\pm$ 20)	2257(2265 $\pm$ 40)
$M_\rho(\text{II})$	771.5(775.5 $\pm$ 0.4)	1471.7(1459 $\pm$ 11)	1855(1870 $\pm$ 20)	2154.8(2149 $\pm$ 17)
$M_{a_1}$	1235.6(1230 $\pm$ 40)	1621.7(1647 $\pm$ 22)	1962(1930 $^{+30}_{-70}$ )	2257.8(2270 $^{+55}_{-40}$ )
$F_V(\text{I})$	156(156 $\pm$ 1)	155	154	153
$F_V(\text{II})$	185(156 $\pm$ 1)	147	139	135
$F_A$	123(122 $\pm$ 24)	137	139	139

**Table 1:** We give the experimental values of the masses (in MeV) and electromagnetic decay constants (when available) for vector and axial vector particles (within parenthesis), compared with the values obtained from the fit. For the vector states we consider two possible Regge trajectories that we label I and II respectively. We take  $\alpha_s(1 \text{ GeV}) = 0.5$  and  $\beta\langle G^2 \rangle = -(352 \text{ MeV})^4$ .



**Figure 1:** In this plot we show  $F_{V,LO}(\text{I})$  and  $F_{V,LO}(\text{II})$  at different orders in  $\alpha_s$ .


Our choice of states for the set (I) is motivated by the discussion of ref. [16] on the possible formation of multiplets in the case of chiral symmetry restoration. The set (II) is based on the assignment of states made in ref. [5] (based on the existence of  $S$  and  $D$ -wave daughter trajectories) and in particular on the analysis of ref. [17], where the state 2265 is argued to belong to the  $D$ -wave Regge trajectory.<sup>4</sup>

In order to fix the parameters of the mass spectrum we use the experimental values of the masses we list in the table. We obtain the values:


$$\begin{aligned}
 B_V(\text{I}) &= 1.525 \times 10^6 \text{ MeV}^2, & A_V(\text{I}) &= -1.038 \times 10^6 \text{ MeV}^2, & C_V(\text{I}) &= 0.123 \times 10^6 \text{ MeV}^2, \\
 B_V(\text{II}) &= 1.128 \times 10^6 \text{ MeV}^2, & A_V(\text{II}) &= 0.353 \times 10^6 \text{ MeV}^2, & C_V(\text{II}) &= -0.885 \times 10^6 \text{ MeV}^2, \\
 B_A &= 1.278 \times 10^6 \text{ MeV}^2, & A_A &= -0.100 \times 10^6 \text{ MeV}^2, & C_A &= 0.349 \times 10^6 \text{ MeV}^2.
 \end{aligned}
 \tag{5.1}$$

We should mention that the values obtained for these parameters are not very stable under the change of number of data points, except for  $B_V$  and  $B_A$ , which are roughly stable, although with quite sizeable uncertainties. For the subleading terms  $A$  and  $C$ , their values are basically random with the fit. We roughly find  $B_V \simeq B_A$  within the uncertainties.

<sup>4</sup>We also thank S. Afonin for discussions on this point.



**Figure 2:** In this plot we show  $F_V(I)$  and  $F_V(II)$  at different orders in the  $1/n$  expansion.



**Figure 3:** In this plot we show  $F_{A,LO}$  and  $F_A$  at different orders in  $\alpha_s$  and in the  $1/n$  expansion, respectively.

The  $n$  dependence of the axial and vector (model II) decay constants is small but sizeable (and it goes in the right direction for low  $n$ ). The  $1/n$  corrections are always corrections compared with the leading order terms. Nevertheless, the  $1/n^2$  correction is much larger than the  $1/n$  one for the range of values of  $n$  that we explore. This appears to be due to the  $\alpha_s^2/(4\pi)^2$  suppression of the  $1/n$  term, as well as to the difference in size between the constants  $A$  and  $C$ . This is so for the axial and vector (model II) decay constants. Nevertheless, for the vector (model I) decay constants the  $n$  dependence appears to be quite small also at NNLO. This appears to be due to the small value of the coefficient  $C_V(I)$ . The gluon condensate contribution is a small correction to the total NNLO term. Either way, our predictions compare favorably with experiment when this comparison is possible.

We should keep in mind that these results have been obtained for a specific model, so we are testing the impact of the perturbative corrections for this specific model. On the other hand, if one believes that the large  $n$  behavior of the spectrum is dictated by the Regge behavior and that the corrections can be obtained as an expansion in  $1/n$ , the set up is general. The only ambiguity comes from where the logarithms should be introduced (masses or decays). At this respect it is worth mentioning that, as a matter of principle, this ambiguity could be fixed if enough experimental information were available for the masses and decays.

## 6. Conclusions

We have studied the constraints that the OPE imposes on large  $N_c$  inspired QCD models for current-current correlators. We have focused on the constraints obtained by going beyond the leading-order parton computation. We have explicitly showed that, assumed a given mass spectrum (Regge plus corrections in  $1/n$ ), we can obtain the logarithmic (and constant) behavior in  $n$  of the decay constants within a systematic expansion in  $1/n$ . More than that, power-like  $1/n$  corrections can only be incorporated in the analysis if full consideration to the perturbative corrections in the Euclidean regime is made. This is due to the fact that this type of contributions produce logarithms of  $Q$  in the Euclidean (this is one of the reasons why this sort of corrections are not usually considered in quark-hadron duality analysis). On the other hand, the existence of  $\ln n$  in the decay constants may point to the existence of two scales in the problem,  $\Lambda_{\text{QCD}}$  and  $n\Lambda_{\text{QCD}}$ , in the Minkowski regime.

We have also performed some numerical estimates of the importance of these corrections. The  $n$  dependence of the decay constants is small but sizeable for the axial and vector (model II) channel, for the vector (model I) one this dependence is small. On the other hand the uncertainties of the calculation are large. Either way, our predictions compare favorably with experiment when this comparison is possible.

Our example shows that it is possible to have different large  $n$  behavior for the vector and pseudo-vector mass spectrum and yet comply with all the constraints from the OPE.

An important caveat of our analysis is that we have not considered what the effect of renormalons could be. We have focused on the effect of low orders in perturbation theory to the decay constants. It would be interesting to see whether the knowledge of the higher order behavior of perturbation theory may give some extra constraints on the values of these constants and the mass spectrum. At this respect we have to say that we have obtained approximated expressions for the decay constants as an expansion in  $\alpha_s(nB_V)$ , with just the low order contributions in  $\alpha_s$ . It is quite likely that this expansion is asymptotic and that different orders in  $1/n$  are related in a similar way to the one found in the renormalon analysis for the OPE expansion for different orders in  $1/Q^2$ . Therefore, the results obtained for the  $1/n$  corrections could be affected as well by the asymptotic behavior of the  $1/\ln n$  expansion in the leading-order term. This is obviously related with renormalons. We expect to come back to this issue in the future.

## Acknowledgments

We thank S. Afonin, A. Andrianov, D. Espriu, and S. Peris for discussions and L. Glozman for correspondence. This work is partially supported by the network Flavianet MRTN-CT-2006-035482, by the spanish grant FPA2004-04582-C02-01, by the catalan grant SGR2005-00564 and by a *Distinció* from the *Generalitat de Catalunya*.

## References

- [1] S.R. Beane, *Constraining quark hadron duality at large- $N_c$* , *Phys. Rev. D* **64** (2001) 116010 [[hep-ph/0106022](#)].
- [2] M. Golterman, S. Peris, B. Phily and E. De Rafael, *Testing an approximation to large- $N_c$  QCD with a toy model*, *JHEP* **01** (2002) 024 [[hep-ph/0112042](#)].
- [3] T.D. Cohen and L.Y. Glozman, *Does one observe chiral symmetry restoration in baryon spectrum?*, *Int. J. Mod. Phys. A* **17** (2002) 1327 [[hep-ph/0201242](#)].
- [4] M. Golterman and S. Peris, *On the use of the operator product expansion to constrain the hadron spectrum*, *Phys. Rev. D* **67** (2003) 096001 [[hep-ph/0207060](#)].
- [5] S.S. Afonin, A.A. Andrianov, V.A. Andrianov and D. Espriu, *Matching Regge theory to the OPE*, *JHEP* **04** (2004) 039 [[hep-ph/0403268](#)].
- [6] J.J. Sanz-Cillero, *Spin-1 correlators at large- $N_c$ : matching OPE and resonance theory up to  $O(\alpha_s)$* , *Nucl. Phys. B* **732** (2006) 136 [[hep-ph/0507186](#)].
- [7] M. Shifman, *Highly excited hadrons in QCD and beyond*, [hep-ph/0507246](#).
- [8] O. Catà, M. Golterman and S. Peris, *The operator product expansion does not imply parity doubling of hadrons*, *Phys. Rev. D* **74** (2006) 016001 [[hep-ph/0602194](#)].
- [9] S.S. Afonin and D. Espriu, *Qualitative solution of QCD sum rules*, *JHEP* **09** (2006) 047 [[hep-ph/0602219](#)].
- [10] G. 't Hooft, *A planar diagram theory for strong interactions*, *Nucl. Phys. B* **72** (1974) 461.
- [11] G. 't Hooft, *A two-dimensional model for mesons*, *Nucl. Phys. B* **75** (1974) 461.
- [12] L.R. Surguladze and F.V. Tkachov, *Two loop effects in QCD sum rules for light mesons*, *Nucl. Phys. B* **331** (1990) 35.
- [13] K.G. Chetyrkin, *Corrections of order  $\alpha_s^3$  to  $R_{\text{had}}$  in pQCD with light gluinos*, *Phys. Lett. B* **391** (1997) 402 [[hep-ph/9608480](#)].
- [14] R. Casero, E. Kiritsis and A. Paredes, *Chiral symmetry breaking as open string tachyon condensation*, [hep-th/0702155](#).
- [15] PARTICLE DATA GROUP collaboration, W.M. Yao et al., *Review of particle physics*, *J. Phys. G* **33** (2006) 1.
- [16] L.Y. Glozman, *Chiral multiplets of excited mesons*, *Phys. Lett. B* **587** (2004) 69 [[hep-ph/0312354](#)].
- [17] D.V. Bugg, *Four sorts of mesons*, *Phys. Rept.* **397** (2004) 257 [[hep-ex/0412045](#)].