

Bile alcohols function as the ligands of membrane-type bile acid-activated G protein-coupled receptor

Yusuke Iguchi,^{1,*†} Masafumi Yamaguchi,* Hiroyuki Sato,[§] Kenji Kihira,[†]
Tomoko Nishimaki-Mogami,** and Mizuho Une*

Laboratory of Organic and Bio-molecular Chemistry,* Faculty of Pharmaceutical Science, Hiroshima International University, Kure, Japan; Division of Clinical Pharmaceutical Science,[†] Programs for Pharmaceutical Sciences, Graduate School of Biomedical Sciences, Hiroshima University, Hiroshima, Japan; Department of Epidemiology,[§] National Cardiovascular Center Research Institute, Osaka, Japan; and Division of Functional Biochemistry and Genomics,** National Institute of Health Sciences, Tokyo, Japan

Abstract TGR5 is a G protein-coupled receptor that is activated by bile acids, resulting in an increase in cAMP levels and the subsequent modulation of energy expenditure in brown adipose tissue and muscle. Therefore, the development of a TGR5-specific agonist could lead to the prevention and treatment of various metabolic disorders related to obesity. In the present study, we evaluated the ability of bile alcohols, which are structurally and physiologically similar to bile acids and are produced as the end products of cholesterol catabolism in evolutionarily primitive vertebrates, to act as TGR5 agonists. In a cell-based reporter assay and a cAMP production assay performed in vitro, most bile alcohols with a side chain containing hydroxyl group(s) were highly efficacious agonists for TGR5 comparable to its most potent ligand in the naturally occurring bile acid, lithocholic acid. However, the abilities of the bile alcohols to activate TGR5 varied with the position and number of the hydroxyl substituent in the side chain. Additionally, the conformation of the steroidal nucleus of bile alcohols is also important for its activity as a TGR5 agonist. Thus, we have provided new insights into the structure-activity relationships of bile alcohols as TGR5 agonists.—Iguchi, Y., M. Yamaguchi, H. Sato, K. Kihira, T. Nishimaki-Mogami, and M. Une. Bile alcohols function as the ligands of membrane-type bile acid-activated G protein-coupled receptor. *J. Lipid Res.* 2010. 51: 1432–1441.

Supplementary key words TGR5 • structure-activity relationship • FXR • drug development • metabolic syndrome

Bile acids are synthesized from cholesterol in mammalian liver and are excreted into the bile. Bile acids facilitate the formation of micelles, which promote the processing of dietary fat. Recently, bile acids have been shown to act as physiological ligands for farnesoid X receptor (FXR) (1–3). Through the activation of FXR, bile acids regulate

the transcription of their own biosynthetic enzymes and of transporter proteins for bile acids. In humans, more than 99% of all bile acids are localized in tissues participating in enterohepatic circulation. However, some bile acids are also present in the systemic circulation at concentrations of 15 μ M after a meal or at concentrations of 5 μ M even between meals (4), whereas their concentrations are several ten- to several hundred-fold higher in the gallbladder, liver, and intestine (5). In addition, bile acids are known to exhibit immunosuppressive effects on cell-mediated immunity and macrophage functions (6–8). In view of these findings, bile acids might be expected to function not only in the limited tissues involved in enterohepatic circulation, but also in the whole body as signaling molecules.

Recently, a G-protein-coupled plasma membrane receptor responsive to bile acids was discovered using high-throughput screening (9, 10). This receptor, named TGR5, has been shown to stimulate adenylate cyclase, causing a subsequent increase in the intracellular production of cAMP. This signal transduction occurs independently of FXR. Also, the stimulation of TGR5 by bile acids further enhances energy expenditure in adipocytes and myocytes via these cell-specific signal transductions (11). Furthermore, TGR5 expression has been demonstrated in enteroendocrine cells, where bile acids stimulate the secretion of glucagon-like peptide-1 via TGR5 (12). These observations suggest that TGR5 might be an attractive target for the treatment of obesity, diabetes, and metabolic syndrome.

During the course of the development of new reagents with the ability to activate TGR5, a diverse variety of bile acids and chemically modified bile acid analogs were evaluated and their structure-activity relationships were studied in regard to TGR5 activation (13). In a previous study,

Abbreviations: CDCA, chenodeoxycholic acid; CRE, cAMP response element; LCA, lithocholic acid; FXR, farnesoid X receptor; THC, 5 β -cholestane-3 α ,7 α ,12 α -triol.

[†]To whom correspondence should be addressed.
e-mail: y-iguchi@ps.hirokoku-u.ac.jp

This work was supported in part by a grant from the Japan Health Sciences Foundation.

Manuscript received 17 August 2009 and in revised form 16 December 2009.

Published, JLR Papers in Press, December 16, 2009

DOI 10.1194/jlr.M004051

Copyright © 2010 by the American Society for Biochemistry and Molecular Biology, Inc.

This article is available online at <http://www.jlr.org>




Fig. 1. Structures of bile alcohols. A: [1], 5β-cholestane-3α,7α,12α-triol (THC); [2], 5β-cholestane-3α,7α,12α,26-tetrol; [3], 5β-cholestane-3α,7α,12α,25-tetrol; [4], (24R)-5β-cholestane-3α,7α,12α,24-tetrol; [5], (24S)-5β-cholestane-3α,7α,12α,24-tetrol; [6], (23R)-5β-cholestane-3α,7α,12α,23-tetrol; [7], (23S)-5β-cholestane-3α,7α,12α,23-tetrol; [8], (22R)-5β-cholestane-3α,7α,12α,22-tetrol; [9], (22S)-5β-cholestane-3α,7α,12α,22-tetrol; [10], 24-nor-5β-cholane-3α,7α,12α,23-tetrol; [11], 5β-cholane-3α,7α,12α,24-tetrol; [12], 26,27-dinor-5β-cholestane-3α,7α,12α,25-tetrol; [13], 27-nor-5β-cholestane-3α,7α,12α,26-tetrol; [14], 23,24-dinor-5β-cholane-3α,7α,22-triol; [15], 24-nor-5β-cholane-3α,7α,23-triol; [16], 5β-cholane-3α,7α,24-triol; [17], 26,27-dinor-5β-cholestane-3α,7α,25-triol; [18], 27-nor-5β-cholestane-3α,7α,26-triol; [19], 5β-cholestane-3α,7α,12α,26,27-pentol; [20], 5β-cholestane-3α,7α-diol; [21], 5β-cholestane-3α,7α,26-triol; [22], 5β-cholestane-3α,7α,26,27-tetrol. B: [23], 5α-cholestane-3α,7α,12α-triol; [24], 5α-cholestane-3α,7α,12α,26-tetrol; [25], 5α-cholestane-3α,7α,12α,26,27-pentol; [26], 5α-cholestane-3α,7α,26-triol; [27], 5α-cholestane-3α,7α,26,27-tetrol. The compounds in the figure are indicated by the bracketed numbers.

TGR5 agonist activity and affinity were shown to be increased by neutral compounds obtained from the reduction of the side chain of bile acids. Therefore, we investigated the TGR5 agonist activity and affinity of bile alcohols, which are neutral polyhydroxylated sterols with the same steroid nucleus as bile acids and are abundantly present in nature.

In the present study, we demonstrated that TGR5 activation and cAMP production in response to the bile alcohols varied with the position and number of the hydroxyl groups in the steroid side chain. In addition, we showed that the conformation of the steroidal nucleus of bile alcohols is also important for their activities as TGR5 agonists. Furthermore, we report that the selectivity of bile alcohols toward TGR5 increased as the number of hydroxyl groups in the side chain increased. Thus, we have provided new insights into the structure-activity relationships of bile alcohols as TGR5 agonists.

MATERIALS AND METHODS

Bile acids and bile alcohols

We obtained our general chemicals from Sigma-Aldrich (St. Louis, MO). Naturally occurring bile acids were purchased from Sigma-Aldrich or Steraloids (Newport, RI). 5 β -Cholestane-

3 α ,7 α ,12 α -triol (THC) [1] and 5 β -cholestane-3 α ,7 α -diol [20] were synthesized as described previously (14). 5 β -Cholestane-3 α ,7 α ,12 α ,26-tetrol [2] (15), 5 β -cholestane-3 α ,7 α ,12 α ,25-tetrol [3] (16), (24R)- and (24S)- 5 β -cholestane-3 α ,7 α ,12 α ,24-tetrols [4 and 5] (17, 18), (23R)- and (23S)- 5 β -cholestane-3 α ,7 α ,12 α ,23-tetrols [6 and 7] (19), (22R)- and (22S)-5 β -cholestane-3 α ,7 α ,12 α ,22-tetrols [8 and 9] (19), 24-nor-5 β -cholestane-3 α ,7 α ,12 α ,23-tetrol [10] (20), 5 β -cholestane-3 α ,7 α ,12 α ,24-tetrol [11] (20), 26,27-dinor-5 β -cholestane-3 α ,7 α ,12 α ,25-tetrol [12], and 27-nor-5 β -cholestane-3 α ,7 α ,12 α ,26-tetrol [13] were synthesized from cholic acid, as described previously (21). 23,24-Dinor-5 β -cholestane-3 α ,7 α ,22-triol [14], 26,27-dinor-5 β -cholestane-3 α ,7 α ,25-triol [17], and 27-nor-5 β -cholestane-3 α ,7 α ,26-triol [18] were obtained by the reduction of 3 α ,7 α -dihydroxy-23,24-dinor-5 β -cholestane-22-al, 3 α ,7 α -dihydroxy-26,27-dinor-5 β -cholestane-25-oic acid, and 3 α ,7 α -dihydroxy-27-nor-5 β -cholestane-26-oic acid using lithium aluminum hydride, respectively (22–24). 24-Nor-5 β -cholestane-3 α ,7 α ,23-triol [15] (20), 5 β -cholestane-3 α ,7 α ,24-triol [16] (25), 5 β -cholestane-3 α ,7 α ,26-triol [21], and 5 β -cholestane-3 α ,7 α ,26,27-tetrol [22] were synthesized from chenodeoxycholic acid (CDCA), as described previously (26, 27), and 5 α - and 5 β -cholestane-3 α ,7 α ,12 α ,26,27-pentols [25 and 19] and 5 α -cholestane-3 α ,7 α ,12 α ,26-tetrol [24] were isolated from the bile of carp (28–30). 5 α -Cholestane-3 α ,7 α ,12 α -triol [23] and 5 α -cholestane-3 α ,7 α ,26-triol [26] were synthesized from 5 α -cholestane-3 α ,7 α ,12 α ,26-tetrol [24], as described previously (31–33). 5 α -Cholestane-3 α ,7 α ,26,27-tetrol [27] was synthesized from 5 α -cholestane-3 α ,7 α ,12 α ,26,27-pentol [25], as described previously (34).




Fig. 2. Effect of the presence of a hydroxyl group at different positions in the side chain of THC on TGR5 activation in a cellular transactivation assay. Hek293T cells were transfected with CRE-luciferase reporter constructs and a human TGR5 expression plasmid. The cells were exposed to the vehicle alone or 0.1–10 μ M of the indicated bile alcohols. Firefly luciferase activity in the cell extract was normalized to *Renilla* luciferase activity and expressed as the fold induction relative to vehicle-exposed cells. The values are the means \pm SD of three experiments. The compounds in the figure are indicated by the bracketed numbers. *: The parameter was not used for the calculation of EC₅₀ value.

Plasmids

The NIH Mammalian Gene Collection clone MGC: 40597 (also named pCMVSPORT6/hTGR5) was obtained from Invitrogen (Carlsbad, CA). cAMP response element (CRE)-driven luciferase reporter construct was obtained from SA Bioscience (Frederick, MD).

Transient transfections and reporter gene assays

The human embryonic kidney cell line 293T (HEK293T cells) was seeded at 50–60% confluence in 96-well plates and was maintained in DMEM containing 10% FBS for 24 h prior to transfection. For transient transfections, the cells were rinsed with PBS and were transfected with 12.5 ng/well of CRE-driven luciferase reporter constructs and 18.75 ng/well of human TGR5 expression plasmids (pCMVSPORT6/hTGR5) using Atractene Transfection Reagent (Qiagen, Chatsworth, CA) in OPTI-MEM I reduced-serum medium (Invitrogen). Six hours after transfection, a volume of DMEM containing 20% fetal calf serum and equivalent to the amount of transfection media that was used was added to the cells. After a further 18 h of incubation, the cells were washed once with PBS and were treated for 5 h with different concentrations of each compound in fresh DMEM containing 0.5% delipidated serum. After treatment, the cells were lysed and luciferase activity was determined using the Dual-Luciferase® Reporter Assay System (Promega, Madison, WI) according to the manufacturer's instructions. The firefly luciferase activity was normalized to the *Renilla* luciferase activity for each well.

Huh-7 cells were maintained in DMEM containing 10% FBS and were seeded onto 24-well plates at 24 h prior to transfection. The cells were transfected with 188 ng of pFXRE-tk-Luc (35), 63 ng each of the pcDNA3.1 expression vectors for human FXR α and RXR α , and 313 ng of the pSV- β -galactosidase vector (Promega, Madison, WI) using Atractene Transfection Reagent (Qiagen, Chatsworth, CA). Three hours after transfection, the cells were exposed to each compound for 20 h in a medium containing 0.5% delipidated FBS. The cells were then lysed and the luciferase activity level was determined. The firefly luciferase activity was normalized to the β -galactosidase activity for each well.

cAMP production analysis

HEK293T cells, in which the human TGR5 expression plasmid was transiently transfected using the above method, were washed with PBS and the medium was exchanged for cAMP assay medium [DMEM containing 0.1% (w/v) BSA and 0.5 mM 3-isobutyl-1-methylxanthine]. After incubation for 30 min at 37°C, the cells were treated with each compound in fresh cAMP assay medium for 30 min. After treatment, the cAMP amounts were determined using a cAMP screening kit (Applied Biosystems, Foster City, CA) according to the manufacturer's instructions. The Lowry protein assay was used to determine the total protein level (36).

50% Effective concentrations and efficacy determination

Assays were performed in triplicate for each condition. The EC₅₀ values of each compound were calculated using a probit




Fig. 3. Effect of length of steroid side chain of bile alcohol on TGR5 activation. TGR5 activation was evaluated using the luciferase reporter assay as described in the legend to Fig. 2. The data represent the mean \pm SD of three determinations. The compounds in the figure are indicated by the bracketed numbers. *: The parameter was not used for the calculation of EC₅₀ value.

analysis. Efficacy was determined by calculating the percentages of the TGR5 agonist activity of 10 μM of lithocholic acid (LCA) and the FXR agonist activity of 10 μM of CDCA.

Statistical analysis

The statistical analysis was performed using the Student *t*-test; a value of $P < 0.05$ was considered statistically significant.

RESULTS

As depicted in **Fig. 1**, 27 compounds of naturally occurring and chemically modified bile alcohols were evaluated using a transient transfection assay for TGR5 activity. The HEK293T cells were cotransfected with CRE-driven luciferase reporter constructs and expression plasmids for human TGR5. LCA, which has the highest TGR5 agonist activity among the endogenous bile acids produced in humans, was used as a positive control. Because of their toxicity, the maximum concentration of bile alcohols examined in this study was 20 μM . The efficacy of each bile alcohol was expressed as the relative percentage of the TGR5 agonist activity of 10 μM of LCA. The affinity of bile alcohols to TGR5 was represented as the EC_{50} value.

As shown in **Fig. 2**, THC [1], the side chain of which is the same as that of cholesterol, had a lower TGR5 activity (efficacy = 36.1%) and affinity (EC_{50} = 3.39 μM) than LCA (efficacy = 100%, EC_{50} = 0.18 μM). Next, we evaluated whether a series of bile alcohols [2-9] possessing one hydroxyl group at C-22 - C-26 in the side chain of THC were potent TGR5 agonists. The exposure of the cells to 5 β -cholestane-3 α ,7 α ,12 α ,26-tetrol [2] or 5 β -cholestane-3 α ,7 α ,12 α ,25-tetrol [3] led to the induction of luciferase

activity at levels comparable to that of LCA (**Fig. 2**). Bile alcohols possessing a hydroxyl group at C-22, C-23, or C-24 each have two stereoisomers, and the activity of each isomer differed significantly (**Fig. 2**). In particular, the 23(R)-isomer [6] and the 22(S)-isomer [9] exhibited minimal TGR5 agonist activity.

The effects of the length of the side chain in the bile alcohols on TGR5 agonist activity are shown in **Fig. 3**. Among the bile alcohols with 23 to 26 carbon atoms and a CA-nucleus ([10]-[13]) and with 22 to 26 carbon atoms and a CDCA-nucleus ([14]-[18]), no significant differences in the TGR5-activation ability and affinity were observed (**Fig. 3**).

We also evaluated whether the A/B ring juncture conformation in the steroid nucleus of bile acids and bile alcohols affected TGR5 activation. As shown in **Fig. 4**, 5 β -bile acids containing a *cis*-oriented A/B ring had a higher TGR5 agonist activity and affinity than their 5 α -counterparts. **Figure 5** shows the effects on TGR5 activation of a series of bile alcohols differing in their conformations at the C-5 position. In a group of bile alcohols with three hydroxyl groups at the 3 α , 7 α , and 12 α positions in the steroid nucleus, the 5 β -bile alcohols ([1], [2], [19]) exhibited higher TGR5 agonist activities and affinities than the corresponding 5 α -bile alcohols ([23], [24], [25]), as expected (**Fig. 5A**). Similar results were also obtained in a group of bile alcohols with two hydroxyl groups at the 3 α and 7 α positions in the nucleus (**Fig. 5B**). In addition, the abilities of these bile alcohols, especially those of the 5 β -bile alcohols, to activate TGR5 increased as the number of hydroxyl groups in their side chain increased (**Fig. 5**).




Fig. 4. Effect of A/B ring juncture conformation of bile acids on TGR5 activation. The TGR5 activation ability in a series of both 5 α - (allo-) and 5 β -bile acids was evaluated using the luciferase reporter assay, as described in the legend to **Fig. 2**. The data represent the mean \pm SD of three determinations. *: The parameter was not used for the calculation of EC_{50} value.




Fig. 5. Effect of the number of hydroxyl groups in the side chain of bile alcohols and their A/B ring juncture conformation on TGR5 activation. The TGR5 activation ability of a series of bile alcohols with the hydroxyl group in the 3 α -, 7 α -, and 12 α -positions (A) and in the 3 α - and 7 α -positions (B) of their steroid nucleus was evaluated using the luciferase reporter assay, as described in the legend to Fig. 2. The data represent the mean \pm SD of three determinations. The compounds in the figure are indicated by the bracketed numbers. *: The parameter was not used for the calculation of EC₅₀ value.

To further confirm the ability of these bile alcohols to modulate intracellular signaling, the intracellular cAMP concentration was measured as a predictive parameter of biological activity. The cAMP-production potencies of the bile alcohols that were tested were strongly correlated with the TGR5 agonist activities evaluated for this set of compounds using the luciferase assay (Fig. 6). In a series of bile alcohols with or without a hydroxyl group in

their side chain ([1][9]), a remarkable difference in the cAMP-production levels was noted between each isomer of 5 β -cholestane-3 α ,7 α ,12 α ,23-tetrols ([6] and [7]) or 5 β -cholestane-3 α ,7 α ,12 α ,22-tetrols ([8] and [9]). Additionally, the cAMP production ability of 5 β -cholestane-3 α ,7 α ,12 α ,26,27-pentol [19] was significantly higher than that of its 5 α -isomer [25]. Further, we observed a stepwise increase in the cAMP concentration for bile alcohols with




Fig. 6. hTGR5-transfected HEK293T cells produced cAMP after stimulation with LCA or bile alcohols. HEK293T cells were transfected with an hTGR5 expression plasmid. Then, the cells were stimulated with vehicle (DMSO) only or LCA or various bile alcohols at 37°C for 30 min. cAMP production and the protein concentration were then determined. The data represent the mean \pm SD of three determinations. The compounds in the figure are indicated by the bracketed numbers. *: $P < 0.05$; **: $P < 0.01$ versus vehicle (DMSO)-treated cells.

an increasing number of hydroxyl groups in their side chains ([20][22]).

Because bile alcohols were also reported to activate FXR (37–39), we assessed the TGR5 selectivity of bile alcohols using the TGR5/FXR EC_{50} ratio calculated from the EC_{50} values for TGR5 and FXR. The EC_{50} values computed using a probit analysis were both statistically significant ($P < 0.01$). A selective agonist for TGR5 was not found among the bile alcohols tested in the present study. However, an increasing number of hydroxyl groups in the side chain led to a decrease in the TGR5/FXR EC_{50} ratio, indicating an increase in TGR5 selectivity (Table 1). Among the bile alcohols that were tested in the present study, 5 β -cholestane-3 α ,7 α ,26,27-tetrol [22], which possess two hydroxyl groups in its side chain and has a CDCA-nucleus, exhibited the highest level of TGR5 selectivity.

DISCUSSION

TGR5 was recently identified as a metabolic receptor that mediates the nongenomic potency of bile acids (9, 10), whereas FXR is a nuclear receptor that modulates genomic effects on them (40). TGR5 that has been activated by bile acids is thought to be involved in immunomodulation and hepatoprotection via the regulation of nitric oxide production and an increase in energy expenditure followed by the activation of thyroid hormone in brown adipose tissue, subsequently leading to the prevention of obesity (10, 11, 41). Furthermore, TGR5 could play a critical in vivo role in the maintenance of glucose homeostasis (42). Thus, TGR5 is an attractive target for the treatment of metabolic disorders.

Bile alcohols have long been known as the final metabolites in cholesterol elimination and are major constituents of bile in evolutionarily primitive animals, such as fishes and frogs (43, 44). In addition, some bile alcohols are formed as intermediates in the bile acid biosynthesis pathway in mammals, including humans (44). Bile alcohols are present as minor components in bile, feces and urine obtained from healthy controls (44–46). We previously demonstrated that bile alcohols are potent FXR ligands as well as bile acids (37–39). Therefore, investigating whether bile alcohols that are structurally and physiologically re-


lated to bile acids are potent TGR5 agonists is of interest. In this study, we comprehensively elucidated the ability of naturally occurring bile alcohols, including intermediates in bile acid biosynthesis as well as chemically modified bile alcohol derivatives, to act as TGR5 agonists.

First, we found that THC, which is an intermediate in the bile acid biosynthesis pathway, exhibited minimal TGR5 agonist activity in a cell-based luciferase assay, whereas 5 β -cholestane-3 α ,7 α ,12 α ,26-tetrol, a hydroxylation product at the terminal methyl group of THC, was a potential ligand for TGR5 (Fig. 2). These findings suggested that the presence of a hydroxyl group (or oxygen atom) in the side chain is essential for TGR5 agonist activity. In addition, bile alcohols having a hydroxyl group at the C-22 to C-25 position in the side chain also activated TGR5, though the activity varied with the position and the configuration of the hydroxyl group (Fig. 2). Significant differences were also produced by the configuration of the hydroxyl group at the C-22 and C-23 positions. Four isomers at C-22 and C-23 of 3 α ,7 α -dihydroxy-22,23-methylene-5 β -cholan-24-oic acid and two isomers at C-23 of 3 α ,7 α -(12 α)-di(tri)hydroxy-23-methyl-5 β -cholan-24-oic acid have reportedly exhibited remarkable differences in TGR5 agonist activity (13, 47). These previous results together with the present findings strongly suggest that the stereochemistry of the substituent at the C-22 or C-23 position, irrespective of its hydrophilic (hydroxyl) or hydrophobic (alkyl) properties, substantially affects the ability of bile acids and bile alcohols to activate TGR5.


Second, the shortening of the side chain in bile alcohols did not significantly affect the TGR5 agonist activity (Fig. 3). Even a C₂₂-bile alcohol [14] exhibited some activity (EC_{50} : 1.65 μ M, efficacy: 83.3%). This result is consistent with the fact that C₂₁- and C₂₂-steroid hormones are able to activate TGR5 (13). In contrast, progressive shortening of the side chain of bile acids reportedly attenuates TGR5 agonist activity and affinity (13). These contradictory findings between the neutral compounds and bile acids remain to be elucidated.

Third, we examined the effect of the A/B ring junction in bile alcohols on the ability to activate TGR5. A series of 5 α -bile alcohols ([23], [24], [25], [26], [27]) exhibited little TGR5 agonist activity compared with their corresponding 5 β -isomers ([1], [2], [19], [21], [22])

TABLE 1. TGR5 selectivity of bile alcohols

A

Compound No.	R ₁	R ₂	TGR5		FXR		TGR5 selectivity index, EC ₅₀ ratio (TGR5/FXR)
			EC ₅₀	Efficacy	EC ₅₀	Efficacy	
[1]	-H	-H	3.39	36.1	4.29	130.8	0.79
[2]	-OH	-H	1.17	85.9	4.18	86.0	0.28
[19]	-OH	-OH	0.58	94.3	3.99	63.7	0.15

B

Compound No.	R ₁	R ₂	TGR5		FXR		TGR5 selectivity index, EC ₅₀ ratio (TGR5/FXR)
			EC ₅₀	Efficacy	EC ₅₀	Efficacy	
[20]	-H	-H	5.07	43.7	2.31	53.3	2.20
[21]	-OH	-H	1.02	84.9	3.18	126.6	0.32
[22]	-OH	-OH	0.27	94.0	3.88	50.3	0.07


TGR5, FXR: Units are μM for EC₅₀ and % of 10 μM LCA or CDCA value for efficacy, respectively.

(Fig. 5). These results were supported by a cAMP production analysis, shown in Fig. 6. These findings suggest that the A/B ring juncture conformation of bile alcohols is an important factor for the expression of TGR5 agonist activity.

We recently reported that bile alcohols are potent ligands for FXR (39). Therefore, the TGR5/FXR selectivity of bile alcohols was examined by calculating the ratio of the EC₅₀ for TGR5 to that for FXR as the agonist (Table 1). The selectivity seemed to increase as the number of hydroxyl groups in the side chain increased. This result may be attributed to the tertiary protein structure of the ligand binding region in both receptors. Therefore, a site that interacts with the side chain of bile alcohols in the ligand-binding region of TGR5 might contain a larger number of

charged amino acids residues compared with that of FXR. Additionally, the site of TGR5 that interacts with the side chain of bile alcohols is likely to be spatially broader than that of FXR. To elucidate these assumptions, a further study, such as crystal structure analysis of TGR5 protein with the ligand, would be required.

We previously demonstrated using 7 ξ -methyl-LCA and 7 β -methyl-CDCA that when a methyl group is inserted into the C-7 position of LCA and CDCA, respectively, the resulting compound has a higher affinity and selectivity for TGR5 than its parent compound (13). While this finding might indicate that the introduction of the hydrophobic methyl group into the steroid nucleus of bile acids elicits TGR5-selectivity, we could not examine such derivatives in the present study.

We focused on bile alcohols that exist in abundance in nature and discussed the structure-activity relationship between TGR5 and bile alcohols. Consequently, we have shown that some bile alcohols exert a TGR5 agonist activity equal to that of endogenous bile acids, although none of the agonist activities were more robust than that of 7 ξ -LCA. Nevertheless, we are firmly convinced that the new insights into the structure-activity relationship obtained from the present study may lead to the development of a new agent for the treatment of metabolic syndrome. 

REFERENCES

- Wang, H., J. Chen, K. Hollister, L. C. Sowers, and B. M. Forman. 1999. Endogenous bile acids are ligands for the nuclear receptor FXR/BAR. *Mol. Cell.* **3**: 543–553.
- Parks, D. J., S. G. Blanchard, R. K. Bledsoe, G. Chandra, T. G. Consler, S. A. Kliewer, J. B. Stimmel, T. M. Willson, A. M. Zavacki, D. D. Moore, et al. 1999. Bile acids: natural ligands for an orphan nuclear receptor. *Science*. **284**: 1365–1368.
- Makishima, M., A. Y. Okamoto, J. J. Repa, H. Tu, R. M. Learned, A. Luk, M. V. Hull, K. D. Lustig, D. J. Mangelsdorf, and B. Shan. 1999. Identification of a nuclear receptor for bile acids. *Science*. **284**: 1362–1365.
- Everson, G. T. 1987. Steady-state kinetics of serum bile acids in healthy human subjects: single and dual isotope techniques using stable isotopes and mass spectrometry. *J. Lipid Res.* **28**: 238–252.
- Ho, K. J. 1976. Circadian distribution of bile acid in the enterohepatic circulatory system in hamsters. *J. Lipid Res.* **17**: 600–604.
- Keane, R. M., T. R. Gadacz, A. M. Munster, W. Birmingham, and R. A. Winchurh. 1984. Impairment of human lymphocyte function by bile salts. *Surgery*. **95**: 439–443.
- Kimmings, A. N., S. J. van Deventer, H. Obertop, E. A. Rauws, and D. J. Gouma. 1995. Inflammatory and immunologic effects of obstructive jaundice: pathogenesis and treatment. *J. Am. Coll. Surg.* **181**: 567–581.
- Drivas, G., O. James, and N. Wardle. 1976. Study of reticuloendothelial phagocytic capacity in patients with cholestasis. *BMJ*. **1**: 1568–1569.
- Maruyama, T., Y. Miyamoto, T. Nakamura, Y. Tamai, H. Okada, E. Sugiyama, H. Itadani, and K. Tanaka. 2002. Identification of membrane-type receptor for bile acids (M-BAR). *Biochem. Biophys. Res. Commun.* **298**: 714–719.
- Kawamata, Y., R. Fujii, M. Hosoya, M. Harada, H. Yoshida, M. Miwa, S. Fukusumi, Y. Habata, T. Itoh, Y. Shintani, et al. 2003. A G protein-coupled receptor responsive to bile acids. *J. Biol. Chem.* **278**: 9435–9440.
- Watanabe, M., S. M. Houten, C. Matak, M. A. Christoffoleto, B. W. Kim, H. Sato, N. Messaddeq, J. W. Harney, O. Ezaki, T. Kodama, et al. 2006. Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature*. **439**: 484–489.
- Katsuma, S., A. Hirasawa, and G. Tsujimoto. 2005. Bile acids promote glucagon-like peptide-1 secretion through TGR5 in a murine enteroendocrine cell line STC-1. *Biochem. Biophys. Res. Commun.* **329**: 386–390.
- Sato, H., A. Macchiarulo, C. Thomas, A. Gioiello, M. Une, A. F. Hofmann, R. Saladin, K. Schoonjans, R. Pellicciari, and J. Auwerx. 2008. Novel potent and selective bile acid derivatives as TGR5 agonists: biological screening, structure-activity relationships, and molecular modeling studies. *J. Med. Chem.* **51**: 1831–1841.
- Bergstrom, S., and L. Krabich. 1957. Preparation of some hydroxycoprostanes: 3 α ,7 α - and 3 α ,12 α -dihydroxycoprostanes. *Acta Chem. Scand. A*. **11**: 1067.
- Dayal, B., G. S. Tint, A. K. Batta, S. Shefer, and G. Salen. 1981. Synthesis of biological precursors of cholic acid II. *Steroids*. **37**: 205–211.
- Dayal, B., S. Shefer, G. S. Tint, G. Salen, and E. H. Mosbach. 1976. Synthesis of 5 β -cholestane-3 α ,7 α ,12 α ,25-tetrol and 5 β -cholestane-3 α ,7 α ,12 α ,24 ξ ,25-pentol. *J. Lipid Res.* **17**: 74–77.
- Masui, T., and E. Staple. 1967. The separation of the stereo-isomers of bile steroids, 5-beta-cholestane-3-alpha, 7-alpha, 12-alpha, 24-alpha-tetrol and 5-beta-cholestane-3-alpha, 7-alpha, 12-alpha, 24-beta-tetrol, by thin layer chromatography. *Steroids*. **9**: 443–450.
- Dayal, B., A. K. Batta, S. Shefer, G. S. Tint, and G. Salen. 1978. Synthesis of biological precursors of cholic acid. *Steroids*. **32**: 337–344.
- Kuramoto, T., N. Matsumoto, and T. Hoshita. 1978. Syntheses of 22- and 23-hydroxylated bile alcohols. *Chem. Pharm. Bull. (Tokyo)*. **26**: 2788–2792.
- Kihira, K., T. Mikami, S. Ikawa, A. Okamoto, M. Yoshii, S. Miki, E. H. Mosbach, and T. Hoshita. 1992. Synthesis of sulfonate analogs of bile acids. *Steroids*. **57**: 193–198.
- Kazuno, T., T. Masui, and T. Hoshita. 1961. Stero-bile acids and bile sterols. 36. Isolation of a new bile sterol, 3 alpha, 7 alpha, 12 alpha, 26-tetrahydroxy-delta23-bishomocholene, from bull frog bile. *J. Biochem.* **50**: 12–19.
- Kihira, K., Y. Morioka, and T. Hoshita. 1981. Synthesis of (22R and 22S)-3 alpha, 7 alpha, 22-trihydroxy-5 beta-cholan-24-oic acids and structure of haemulcholic acid, a unique bile acid isolated from fish bile. *J. Lipid Res.* **22**: 1181–1187.
- Cohen, B. I., G. S. Tint, T. Kuramoto, and E. H. Mosbach. 1975. New bile alcohols—synthesis of 5beta-cholestane-3alpha, 7alpha, 25-triol and 5beta-cholestane-3alpha, 7alpha, 25–24 (14C)-triol. *Steroids*. **25**: 365–378.
- Mikami, T., E. H. Mosbach, B. I. Cohen, N. Ayyad, M. Yoshii, K. Kihira, and T. Hoshita. 1995. Synthesis and metabolism of sodium 3 alpha, 7 alpha-dihydroxy-25,26-bishomo-5 beta-cholane-26-sulfonate in the hamster. *Lipids*. **30**: 71–78.
- Kihira, K., M. Yoshii, A. Okamoto, S. Ikawa, H. Ishii, and T. Hoshita. 1990. Synthesis of new bile salt analogues, sodium 3 alpha, 7 alpha-dihydroxy-5 beta-cholane-24-sulfonate and sodium 3 alpha, 7 beta-dihydroxy-5 beta-cholane-24-sulfonate. *J. Lipid Res.* **31**: 1323–1326.
- Dayal, B., A. K. Batta, S. Shefer, G. S. Tint, G. Salen, and E. H. Mosbach. 1978. Preparation of 24(R)- and 24(S)-5beta-cholestane-3alpha,7alpha,24-triols and 25(R)- and 25(S)-5beta-cholestane-3alpha,7alpha,26-triols by a hydroboration procedure. *J. Lipid Res.* **19**: 191–196.
- Une, M., Y. Shinonaga, N. Matoba, S. Kuroki, K. Kihira, and T. Hoshita. 1986. Identification of new bile alcohols, 5 beta-cholestane-3 alpha,7 alpha,24,26-tetrol, 5 beta-cholestane-3 alpha,7 alpha,25,26-tetrol, and 5 beta-cholestane-3 alpha,7 alpha,26,27-tetrol in human gallbladder bile. *J. Lipid Res.* **27**: 1318–1323.
- Hoshita, T., T. Sasaki, Y. Tanaka, S. Betsuki, and T. Kazuno. 1965. Stero-bile acids and bile sterols. LXXIV. Biosynthesis of bile acids and bile alcohols in toad. *J. Biochem.* **57**: 751–757.
- Une, M., N. Matsumoto, K. Kihira, M. Yasuhara, T. Kuramoto, and T. Hoshita. 1980. Bile salts of frogs: a new higher bile acid, 3 alpha, 7 alpha, 12 alpha, 26-tetrahydroxy-5 beta-cholestanoic acid from the bile Rana plancyi. *J. Lipid Res.* **21**: 269–276.
- Hoshita, T., T. Sasaki, and T. Kazuno. 1965. Isolation of a bile alcohol, 5alpha-cholestane-3alpha,7alpha,12alpha,26-tetrol from carp bile. *Steroids*. **5**: 241–247.
- Hoshita, T., S. Hirofujii, T. Nakagawa, and T. Kazuno. 1967. Studies on the bile salts of the newt and synthesis of 5alpha-cholestane-3alpha,7alpha,12alpha,25,26-pentol (5alpha-bufol). *J. Biochem.* **62**: 62–66.
- Hoshita, T., S. Shefer, and E. H. Mosbach. 1968. Conversion of 7-alpha,12-alpha-dihydroxycholest-4-en-3-one to 5-alpha-cholestane-3-alpha, 7-alpha,12-alpha-triol by iguana liver microsomes. *J. Lipid Res.* **9**: 237–243.
- Une, M., K. Kihira, T. Kuramoto, and T. Hoshita. 1978. Two new bile alcohols, 3-epi-16-deoxymyxinol from the hagfish, *Heptatretus burgeri*. *Tetrahedron Lett.* : 2527–2530.
- Kuramoto, T., H. Kikuchi, H. Sanemori, and T. Hoshita. 1973. Bile salts of anura. *Chem. Pharm. Bull. (Tokyo)*. **21**: 952–959.
- Fujino, T., M. Une, T. Imanaka, K. Inoue, and T. Nishimaki-Mogami. 2004. Structure-activity relationship of bile acids and bile acid analogs in regard to FXR activation. *J. Lipid Res.* **45**: 132–138.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**: 265–275.
- Nishimaki-Mogami, T., M. Une, T. Fujino, Y. Sato, N. Tamehiro, Y. Kawahara, K. Shudo, and K. Inoue. 2004. Identification of intermediates in the bile acid synthetic pathway as ligands for the farnesoid X receptor. *J. Lipid Res.* **45**: 1538–1545.
- Nishimaki-Mogami, T., Y. Kawahara, N. Tamehiro, T. Yoshida, K. Inoue, Y. Ohno, T. Nagao, and M. Une. 2006. 5Alpha-bile alcohols function as farnesoid X receptor antagonists. *Biochem. Biophys. Res. Commun.* **339**: 386–391.

39. Iguchi, Y., K. Kihira, T. Nishimaki-Mogami, and M. Une. 2010. Structure-activity relationship of bile alcohols as human farnesoid X receptor agonist. *Steroids*. **75**: 95–100.
40. Makishima, M. 2005. Nuclear receptors as targets for drug development: regulation of cholesterol and bile acid metabolism by nuclear receptors. *J. Pharmacol. Sci.* **97**: 177–183.
41. Keitel, V., R. Reinehr, P. Gatsios, C. Rupprecht, B. Gorg, O. Selbach, D. Haussinger, and R. Kubitz. 2007. The G-protein coupled bile salt receptor TGR5 is expressed in liver sinusoidal endothelial cells. *Hepatology*. **45**: 695–704.
42. Thomas, C., A. Gioiello, L. Noriega, A. Strehle, J. Oury, G. Rizzo, A. Macchiarulo, H. Yamamoto, C. Matak, M. Pruzanski, et al. 2009. TGR5-mediated bile acid sensing controls glucose homeostasis. *Cell Metab.* **10**: 167–177.
43. Hoshita, T., and T. Kazuno. 1968. Chemistry and metabolism of bile alcohols and higher bile acids. *Adv. Lipid Res.* **6**: 207–254.
44. Karlaganis, G., and J. Sjøvall. 1984. Formation and metabolism of bile alcohols in man. *Hepatology*. **4**: 966–973.
45. Kuroki, S., K. Shimazu, M. Kuwabara, M. Une, K. Kihira, T. Kuramoto, and T. Hoshita. 1985. Identification of bile alcohols in human bile. *J. Lipid Res.* **26**: 230–240.
46. Ludwig-Kohn, H., H. V. Henning, A. Sziedat, D. Matthaei, G. Spiteller, J. Reiner, and H. J. Egger. 1983. The identification of urinary bile alcohols by gas chromatography-mass spectrometry in patients with liver disease and in healthy individuals. *Eur. J. Clin. Invest.* **13**: 91–98.
47. Pellicciari, R., H. Sato, A. Gioiello, G. Costantino, A. Macchiarulo, B. M. Sadeghpour, G. Giorgi, K. Schoonjans, and J. Auwerx. 2007. Nongenomic actions of bile acids. Synthesis and preliminary characterization of 23- and 6,23-alkyl-substituted bile acid derivatives as selective modulators for the G-protein coupled receptor TGR5. *J. Med. Chem.* **50**: 4265–4268.