

On the mechanism of accumulation of cholestanol in the brain of mice with a disruption of sterol 27-hydroxylase

Ann Båvner,^{1,*} Marjan Shafaati,^{1,*} Magnus Hansson,^{*} Maria Olin,^{*} Shoshi Shpitzen,[†] Vardiella Weiner,[§] Eran Leitersdorf,[†] and Ingemar Björkhem^{2,*}

Department of Laboratory Medicine,^{*} Division of Clinical Chemistry, Karolinska Institutet, Karolinska University Hospital, Huddinge, Sweden; and Department of Medicine B,[†] and Department of Human Genetics,[§] Hadassah-Hebrew University Hospital, Israel

Abstract The rare disease cerebrotendinous xanthomatosis (CTX) is due to a lack of sterol 27-hydroxylase (CYP27A1) and is characterized by cholestanol-containing xanthomas in brain and tendons. Mice with the same defect do not develop xanthomas. The driving force in the development of the xanthomas is likely to be conversion of a bile acid precursor into cholestanol. The mechanism behind the xanthomas in the brain has not been clarified. We demonstrate here that female *cyp27a1*^{-/-} mice have an increase of cholestanol of about 2.5-fold in plasma, 6-fold in tendons, and 12-fold in brain. Treatment of *cyp27a1*^{-/-} mice with 0.05% cholic acid normalized the cholestanol levels in tendons and plasma and reduced the content in the brain. The above changes occurred in parallel with changes in plasma levels of 7 α -hydroxy-4-cholesten-3-one, a precursor both to bile acids and cholestanol. Injection of a *cyp27a1*^{-/-} mouse with ²H₇-labeled 7 α -hydroxy-4-cholesten-3-one resulted in a significant incorporation of ²H₇-cholestanol in the brain. The results are consistent with a concentration-dependent flux of 7 α -hydroxy-4-cholesten-3-one across the blood-brain barrier in *cyp27a1*^{-/-} mice and subsequent formation of cholestanol. It is suggested that the same mechanism is responsible for accumulation of cholestanol in the brain of patients with CTX.—Båvner, A., M. Shafaati, M. Hansson, M. Olin, S. Shpitzen, V. Weiner, E. Leitersdorf, and I. Björkhem. On the mechanism of accumulation of cholestanol in the brain of mice with a disruption of sterol 27-hydroxylase. *J. Lipid Res.* 2010. 51: 2722–2730.

Supplementary key words CYP27A1 • 7 α -hydroxy-4-cholesten-3-one • blood-brain barrier • tendon xanthomas

Sterol 27-hydroxylase (CYP27A1) is a mitochondrial cytochrome P-450 present in most cells and tissues (1). In the liver, the enzyme catalyzes the initial step in the degradation of the steroid side-chain in bile acid biosynthesis. Part of the latter synthesis may start extrahepatically with

a 27-hydroxylation of cholesterol followed by a flux of 27-hydroxycholesterol or cholestenic acid to the liver. This mechanism may be regarded as an alternative to reversed cholesterol transport, and CYP27A1 can thus be regarded to be an antiatherogenic enzyme.

CYP27A1 deficiency [cerebrotendinous xanthomatosis (CTX)] is a rare familial lipid storage disease characterized by accumulation of cholesterol and cholestanol in most tissues, in particular in tendon and brain xanthomas [for a review, see (2)]. The most serious symptoms: dementia, cerebellar ataxia, and spinal cord paresis, are thought to be caused by the brain xanthomas. It is noteworthy that the xanthomas develop in spite of normal circulating levels of cholesterol. As a consequence of the reduced capacity to form normal bile acids and the reduced negative feedback inhibition of the cholesterol 7 α -hydroxylase, CTX patients may excrete gram amounts of 7 α -hydroxylated bile alcohols in feces.

In contrast to humans, disruption of the CYP27A1 gene in mice does not lead to formation of xanthomas in tendons or in brain (3–6). In similarity with CTX patients, *Cyp27a1*^{-/-} mice have a reduced formation of bile acid, but there is a low excretion only of 7 α -hydroxylated bile alcohols. As a consequence of the lack of bile acids and reduced cholesterol absorption, CYP7A1 is modestly up-regulated and cholesterol synthesis is increased, at least in the liver.

It is well documented that treatment of CTX patients with bile acids, in particular chenodeoxycholic acid, reverses the cholestanol accumulation and the symptoms (2, 7). Even the size of the xanthomas in the brain may be reduced as a result of such treatment (8). In *cyp27a1*^{-/-} mice, treatment with cholic acid normalizes CYP7A1 activity and cholesterol synthesis (9).

This study was funded by the Swedish Research Council, Swedish Brain Power (to I.B.), and The Sarah and Moshe Mayer Foundation for Research (to E.L.).

Manuscript received 10 May 2010 and in revised form 28 May 2010.

Published, JLR Papers in Press, May 28, 2010

DOI 10.1194/jlr.M008326

²A. Båvner and M. Shafaati contributed equally to this work.

[†]To whom correspondence should be addressed.

e-mail: Ingemar.Bjorkhem@karolinska.se

Copyright © 2010 by the American Society for Biochemistry and Molecular Biology, Inc.

This article is available online at <http://www.jlr.org>

To understand the pathogenetic mechanism behind formation of xanthomas in patients with CTX, it is of interest to compare the effect of CYP27A1 deficiency on some key enzyme systems and some key metabolic products in mice and humans. It has been reported that the bile acid deficiency leads to a 22-fold upregulation of cholesterol 7 α -hydroxylase activity in CTX patients but only 2-7-fold in Cyp27a1^{-/-} mice (4). In accordance with this, the circulating levels of 7 α -hydroxycholesterol are increased about 4-fold in Cyp27a1^{-/-} mice (6) but often more than 50-fold in CTX-patients (unpublished observation).

Another interesting difference between CTX patients and Cyp27a1^{-/-} mice is the degree of accumulation of cholestanol in the circulation. In patients with CTX, the levels of plasma cholestanol are generally increased about 7-fold or more, whereas the levels in Cyp27a1^{-/-} mice have been reported to be increased by a factor of about 2-fold (5, 7).

There is a clear link between the increased cholesterol 7 α -hydroxylase activities in patients with CTX and the accumulation of cholestanol. By injecting 7 α -tritium-labeled cholesterol and measuring incorporation of the label in cholestanol, we could demonstrate that the major part of the cholestanol formed in CTX involves 7 α -hydroxylated intermediates (10). We have also shown that the bile acid precursor 7 α -hydroxy-4-cholesten-3-one is metabolized into cholestanol under in vitro conditions, with cholesta-4,6-dien-3-one and 4-cholesten-3-one as intermediates (10). The latter intermediates are markedly accumulated in the circulation of patients with CTX (7).

The origin of the cholestanol present in the brain of patients is not obvious. The simplest mechanism, direct blood-to-brain passage of cholestanol formed extracerebrally, is dependent upon the ability of cholestanol to cross the blood-brain barrier. Buchmann and Claussen (11) reported that rabbits fed a diet enriched with cholestanol for 8 weeks had brain cholestanol about twice that of animals fed a control diet. Additionally, Buyn et al. (12) showed that feeding mice with 1% cholestanol for 8 months led to a significant enrichment of this sterol in the cerebellum. Because of the possibility of a contamination from blood vessels, it is difficult to draw firm conclusions from these studies. If a passage of cholestanol does occur from the circulation into the brain, the efficiency of such transfer must be very low. There is a close structural similarity between cholesterol and cholestanol, and it is well established that there is no significant transfer of cholesterol over the blood-brain barrier [for a review, see ref (13)].

We have suggested an alternative mechanism in which there is a passage of a circulating precursor such as 7 α -hydroxycholesterol or 7 α -hydroxy-4-cholesten-3-one into the brain, with subsequent conversion into cholestanol. In accordance with this hypothesis, we recently demonstrated a very efficient transfer of 7 α -hydroxy-4-cholesten-3-one across cultured porcine brain endothelial cells (a model for the blood-brain barrier) that was about 100 times more efficient than the transfer of cholestanol (14). Furthermore, there was an efficient conversion of 7 α -hydroxy-4-cholesten-3-one into cholestanol in cultured neuronal and

glial cells as well as in monocyte-derived macrophages of human origin.

In the present work, we measured the content of cholestanol in the circulation, liver, brain, and tendons of cyp27a1^{-/-} mice. In accordance with previous reports, no xanthomas were formed, but there was a marked and hitherto unreported accumulation of cholestanol in brain and tendons of the cyp27a1^{-/-} mice. The accumulation was less marked in liver and plasma. In addition, we measured the concentration of the cholestanol precursor, 7 α -hydroxy-4-cholesten-3-one, in plasma under different conditions. The latter precursor was also injected in cyp27a1^{-/-} mice. The results are consistent with the hypothesis that 7 α -hydroxy-4-cholesten-3-one is an important precursor of cholestanol in the brain of the cyp27a1^{-/-} mice.

MATERIALS AND METHODS

Materials

7 α -Hydroxy-4-cholesten-3-one was obtained from Steraloids. ²H₇-labeled 7 α -hydroxycholesterol, with a purity of >99%, was obtained from Avanti. ²H₇-labeled 7 α -hydroxy-4-cholesten-3-one was synthesized from ²H₇-labeled 7 α -hydroxycholesterol by enzymatic oxidation as described (15). The ²H₇-labeled 7 α -hydroxy-4-cholesten-3-one obtained was purified by preparative thin-layer chromatography using toluene/ethyl acetate 1/1 (v/v) as moving phase.

CYP27A1 knockout mice

Generation of these mice in Israel has been described previously (6). The breeding of these cyp27a1^{-/-} mice in our Swedish animal facility became complicated by low mothering nurture instincts and aggressive behavior toward pups. Because of this, we generated the cyp27a1^{-/-} mice from heterozygotes on a C57BL/6J background. This breeding was uncomplicated and also generated the cyp27a1^{+/+} mice that were used as controls. The mice had free access to normal chow and water. In one set of experiments, mice were fed with normal chow containing 5% cholestyramine or 0.05% cholic acid. In the case of treatment with cholestyramine, the mice were 8–9 weeks old at the start of the experiment. In the case of treatment with cholic acid, the mice were 3 weeks old at the start of the experiment. The treatment continued for 8–9 weeks, following which they were euthanized by carbon dioxide inhalation. Cerebrum, cerebellum, liver, and tendons were collected immediately after collection of blood by cardiac puncture. There were 4–5 mice in each group, with the exception of the group of cyp27a1^{-/-} females treated with cholestyramine that consisted of three animals and the group with male cyp27a1^{-/-} mice on chow diet, which consisted of eight animals.

In one experiment, a cyp27a1^{-/-} mouse was subjected to daily injections of unlabeled 7 α -hydroxy-4-cholesten-3-one, 200 μ g dissolved in 500 μ l saline containing BSA 1% (w/v) and 10% ethanol (v/v) for 3 months. In another experiment, a cyp27a1^{-/-} mouse was injected daily with ²H₇-labeled 7 α -hydroxy-4-cholesten-3-one, 100 μ g dissolved in 500 μ l saline containing BSA 1% (w/v) and 10% ethanol (v/v) for 9 days. In accordance with previous work (6), there were no signs of xanthomas in the tendons, liver, or brain of the cyp27a1^{-/-} mice.

All experimental protocols were approved by the local ethics committee for animal experiments.

Lipid extractions

Lipids were extracted from the brain as described elsewhere with some modifications (16, 17). Approximately 50–100 mg of brain tissue was added to 1 ml of homogenization buffer (5 mM EDTA, 50 μ g/ml butylated hydroxytoluene in phosphate-buffered saline, pH 7.4) in a clean glass tube, and the tissue was disrupted using a polytron homogenizer. Five milliliters of chloroform:methanol (2:1, v/v) were added to the homogenate, and the vials were mixed by vortexing and shaken at room temperature overnight. Mouse plasma (25 μ l) was added to 1 ml of NaCl (0.9% solution) and 4 ml of chloroform:methanol (2:1, v/v) and incubated on a shaking platform for 1 h at room temperature. Mouse livers were extracted in 3 ml of chloroform:methanol (2:1, v/v) for 24 h at room temperature. Liver pieces were removed and 1 ml of NaCl (0.9% solution) was added. Samples were centrifuged at 10,000 g for 10 min. The organic phase was transferred to a new vial. The aqueous phase was reextracted one more time. Mouse tendons were extracted in 3 ml of chloroform:methanol (2:1, v/v) for 5 days at +4°C. The organic phases from different tissues were dried under a stream of nitrogen gas, and chloroform:methanol (2:1, v/v) was added to the dried samples to achieve suitable concentrations.

Sterol analysis

Sterols in the lipid extract were assayed by combined gas chromatography-MS of the trimethylsilyl ether derivative similar to previously described methods from our laboratory (14). The assay of cholestanol was different from the previously used method, however, by using $^2\text{H}_4$ -lathosterol (18) as internal standard and the ion at m/z 460 and m/z 462 in the selected ion monitoring of cholestanol and $^2\text{H}_4$ lathosterol, respectively. The latter compound was used as internal standard also in the assay of lathosterol. Cholesterol was assayed by isotope dilution-MS as described previously using $^2\text{H}_7$ -cholesterol as internal standard (14). Cholesterol precursors were analyzed by isotope dilution MS as previously described (19). Content of deuterium in cholestanol isolated from the mouse injected with $^2\text{H}_7$ -7 α -hydroxy-4-cholesten-3-one was calculated by use of combined gas chromatography-MS and selected monitoring of the molecular ions at m/z 460 and m/z 467. 7 α -Hydroxy-4-cholesten-3-one was analyzed by LC-MS-MS as described previously (15).

Isolation and measurement of mRNA levels

RNA was isolated from mouse brains with TRIzol Reagent according to the manufacturer's instructions (Invitrogen). The mRNA levels of the different genes relative to the housekeeping gene, hprt, were measured as described previously (20).

Analysis of data

Data are reported as the mean \pm SEM in the specified number of individual animals. Differences between mean values were tested for statistical significance (* $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$) by the two-tailed Student's *t*-test assuming equal variance.

RESULTS

Cholestanol levels in plasma, liver, tendons, and brain of wild-type and cyp27a1-deficient mice

In accordance with previous work (5), the levels of cholestanol in the circulation of both male and female cyp27a1^{-/-} mice were increased about two-fold in relation to wild type (Fig. 1). Also in accordance with previous investigations (3), the levels of cholesterol were significantly

reduced by about 50% in female mice. The levels of cholestanol were increased 2- to 3-fold in the liver of the female cyp27a1^{-/-} mice and about 2-fold in the liver of the males (Fig. 1). The cholesterol content in the liver was significantly increased in the female cyp27a1^{-/-} mice and decreased in the liver of the male cyp27a1^{-/-} mice (Fig. 1). Cholesterol synthesis was markedly increased in the liver of both male and female cyp27a1^{-/-} mice as shown by the lathosterol levels (Fig. 1).

In view of the fact that tendons are preferential sites for formation of cholestanol-containing xanthomas in patients with CTX, we also measured the cholestanol content in the Achilles tendons of the cyp27a1^{-/-} mice and wild-type mice. In this case, we got more reproducible results when expressing the levels of cholestanol in relation to cholesterol than to weight of tissue. The levels of cholesterol in the tendons were not significantly affected by the knockout of cyp27a1, but the individual variations were great (results not shown). As shown in Fig. 2, the content of cholestanol in relation to cholesterol was about 6-fold higher in female cyp27a1^{-/-} mice than in wild-type mice. The corresponding figure for male mice was about 4-fold. The lathosterol levels were increased under all conditions, suggesting increased cholesterol synthesis (Fig. 2). In the case of female mice on a control diet, this level did not reach statistical significance due to the great inter-individual variations ($P > 0.05$).

The levels of cholestanol in the brain of the cyp27a1-deficient female mice were increased about 12-fold compared with their wild-type littermates (Fig. 3). In cyp27a1-deficient males, the corresponding increase was about 6-fold. The lack of cyp27a1 did not affect the cholesterol levels in the brain (Fig. 3). The levels of lathosterol were, however, increased about 2-fold, indicating an increased rate of cholesterol synthesis (Fig. 3).

In view of the fact that the cerebellum is a preferential site for accumulation of cholestanol-containing xanthomas in the brain of CTX patients, we also measured the cholestanol content in the cerebellum of cyp27a1-deficient female mice. These levels were similar to those in the whole brain, and there was no effect on the cholesterol content (results not shown).

It is evident from above that the accumulation of cholestanol is higher in plasma and tissues of female than in male cyp27a1-deficient mice. Because of this, a more detailed investigation was performed with female mice only.

Levels of 7 α -hydroxy-4-cholesten-3-one in the circulation of cyp27a1^{-/-} mice and wild-type mice

As shown in Table 1, the levels of 7 α -hydroxy-4-cholesten-3-one were about 35 times higher in the circulation of cyp27a1^{-/-} mice than in the circulation of the corresponding control mice. Treatment with cholestyramine would be expected to increase the activity of the cholesterol 7 α -hydroxylase due to a reduced negative feedback by bile acids. This would be expected to increase the levels of 7 α -hydroxy-4-cholesten-3-one. In accordance with this, the levels of 7 α -hydroxy-4-cholesten-3-one increased about




Fig. 1. Levels of sterols (A) in plasma of female wild-type (WT) and CYP27KO mice on different treatments (WT and KO; ctrl n3 and n3, cholestyramine n3 and n3, cholic acid n5 and n5) and male WT (n5) and CYP27KO (n3) mice on control diet (B) in liver of female WT (n4) and CYP27KO (n5) and male WT (n5) and CYP27KO (n8) on control diet.

3-fold in the wild-type mice and about twice in the *cyp27a1*^{-/-} mice treated with cholestyramine. Treatment with cholic acid, the major suppressor of cholesterol 7 α -hydroxylase activity in mice (21), would be expected to reduce cholesterol 7 α -hydroxylase activity as well as the levels of 7 α -hydroxy-4-cholesten-3-one. Under the conditions employed, with a dose of cholic acid corresponding to the normal production of cholic acid (21), there was a very small reduction only of the levels of 7 α -hydroxy-4-cholesten-3-one in the wild-type mice. In *cyp27a1*^{-/-} mice, however, with a very low endogenous production of cholic acid (6), the treatment caused a marked reduction in the levels of 7 α -hydroxy-4-cholesten-3-one by about 75% (Table 1).

Effect of cholestyramine and cholic acid on cholestanol levels in tissue and plasma of female wild-type and *cyp27a1*^{-/-} mice

As shown in Figs. 1 and 3, neither treatment with cholestyramine nor treatment with cholic acid had any significant effect on cholestanol levels in plasma or brain of the wild-type mice. In tendons, treatment with cholic acid reduced the levels of cholestanol in the wild-type mice (Fig. 2). In the *cyp27a1*^{-/-} mice, treatment with cholestyramine increased the levels of cholestanol in plasma but not in tendons or brain (Figs. 1–3). Treatment with

cholic acid markedly reduced the levels of cholestanol in plasma as well as in tendons and brain of the *cyp27a1*^{-/-} mice (Figs. 1, 3, and 4).

As shown in Fig. 4, there was no relation between levels of cholestanol and 7 α -hydroxy-4-cholesten-3-one in plasma or brain of wild-type mice under the different conditions. In plasma of the *cyp27a1*^{-/-} mice, however, the changes in levels of cholestanol closely followed the changes in levels of cholestanol under the different conditions. In brain of the *cyp27a1*^{-/-} mice, the changes in levels of cholestanol induced by treatment with cholic acid were similar to those of 7 α -hydroxy-4-cholesten-3-one in plasma. The increased levels of 7 α -hydroxy-4-cholesten-3-one in plasma induced by treatment of the *cyp27a1*^{-/-} mice with cholestyramine were, however, not followed by a similar increase in cholestanol.

Direct evidence for a transfer of 7 α -hydroxy-4-cholesten-3-one from the circulation into the brain and its conversion into cholestanol

The above results are consistent with a transfer of 7 α -hydroxy-4-cholesten-3-one from the circulation into the brain under conditions when the level of this oxysterol is sufficiently high (at least higher than about 200 ng/ml plasma). To demonstrate such a transfer, we treated a female *cyp27a1*^{-/-} mouse with daily injections with 200 μ g




Fig. 2. Levels of sterols in tendons of female wild-type (WT) and CYP27KO mice on different treatments (WT and KO; ctrl n4 and n5, cholestyramine n4 and n3, cholic acid n5 and n5) and male WT (n5) and CYP27KO (n8) mice on control diet.

of 7 α -hydroxy-4-cholesten-3-one during 3 months. As a result of this treatment, the levels of cholestanol in the brain of this mouse increased to 1.47 μ g/mg, almost 4-fold higher than the corresponding level in untreated *cyp27a1*^{-/-} mice. In spite of this high level, corresponding to about 10% of the sterol fraction, no xanthomas could be found in the brain. Further direct evidence for a formation of cholestanol from 7 α -hydroxy-4-cholesten-3-one in the circulation was obtained by treating a *cyp27a1*^{-/-} mouse with daily injections of 100 μ g of ³H₇-labeled 7 α -hydroxy-4-cholesten-3-one for 10 days. After this treatment, the content of ³H₇ in plasma cholestanol was 2.6% and in brain cholestanol was 0.19%.

Effect of the *cyp27a1* deficiency on cholesterol homeostasis in the brain

There was no significant difference between *cyp27a1*^{-/-} mice and wild-type mice with respect to content of cholesterol in the brain (Fig. 3). The levels of lathosterol were, however, increased in the brain of the *cyp27a1*^{-/-} mice, suggesting an increased cholesterol synthesis. Such increase was observed under all conditions, including treatment with cholic acid. It was shown that not only lathosterol but also some other precursors to cholesterol (lanosterol, 24,25-dihydroxycholesterol, and 7-dehydrocholesterol) were increased in the *cyp27a1*^{-/-} mice (results not




Fig. 3. Levels of sterols in cerebrum of female wild-type (WT) and CYP27KO mice on different treatments (WT and KO; ctrl n4 and n5, cholestyramine n4 and n3, cholic acid n5 and n5) and male WT (n5) and CYP27KO (n8) mice on control diet.

shown). The apparent increase in cholesterol synthesis was not, however, associated with increased mRNA levels of 3-hydroxy-3-methyl-glutaryl-CoA reductase or synthase (results not shown). The mRNA levels of a gene involved in the metabolism of cholesterol in the brain, *cyp46a1*, was also not affected by *cyp27a1* deficiency (results not shown).

DISCUSSION

Mice with a disruption of the CYP27A1 are not ideal model systems for CTX because of the fact that no xanthomas are formed in tendons and brain. We show here, however, that both the brain and tendons of *cyp27*^{-/-} mice contain a relatively high accumulation of cholestanol, which is the most specific feature of the xanthomas in CTX.

TABLE 1. Plasma 7 α -hydroxy-4-cholesten-3-one levels in wild-type and cyp27^{-/-} female mice on different treatments

Mice	Diet	7 α -Hydroxy-4-cholesten-3-one $\mu\text{g/ml}$
WT	Control	0.053
WT	Cholestyramine	0.19
WT	Cholic acid	0.036 \pm 0.014
CYP27 ^{-/-}	Control	1.9 \pm 0.15
CYP27 ^{-/-}	Cholestyramine	3.8 \pm 0.38
CYP27 ^{-/-}	Cholic acid	0.49 \pm 0.093

Mice were treated as indicated in the table. Data from wild-type control and cholestyramine-treated mice were received from pooled plasma from four animals per group. Individual animals analyzed per group: wild-type and CYP27^{-/-} cholic acid fed, n = 5, CYP27^{-/-} control and cholestyramine fed, n = 3.

Female mice were found to have higher accumulation of cholestanol than male mice, and a detailed investigation was therefore made on female mice only. The reason for the gender difference is unknown. Although there is no obvious direct link to atherosclerosis, it is noteworthy that female mice are more sensitive than male mice to develop atherosclerosis under different experimental conditions (22, 23). Evidently, this is due to a reduced capacity for reverse cholesterol transfer, and female mice tend to have lower levels of HDL. In view of our finding that cholestanol can be eliminated from cultured cells by the same lipoprotein-mediated mechanism as cholesterol, this could be the explanation for the gender difference.

The two pathways involved in the formation of cholestanol from cholesterol are shown in Fig. 5. Under normal conditions, cholestanol is formed from cholesterol by a pathway involving the action of a 3 β -hydroxy- Δ -5-dehydrogenase as a first step (24) (pathway I in the figure). This reaction

leads to formation of 4-cholesten-3-one, which is further converted into cholestanol in two reductive steps (2). In the alternative pathway (pathway II in the figure), 4-cholesten-3-one is not formed directly from cholesterol but from 7 α -hydroxy-4-cholesten-3-one by dehydration followed by saturation of the Δ 6 double bond (2, 10). The enzymes involved in the latter two reactions have been characterized with respect to reaction mechanisms (25) and are analogous to reactions involved in the bacterial 7 α -dehydroxylation of cholic acid into deoxycholic acid (26).

It is evident that cholestanol is mainly formed by the classical pathway in the wild-type mouse and not by the alternative pathway. The small changes in plasma cholestanol levels induced by cholestyramine and cholic acid were similar to those of cholesterol. The slight reducing effect of treatment with cholestyramine is likely to be due to increased activity of the cholesterol 7 α -hydroxylase with increased consumption of substrate. It is well established that cholestanol is a substrate for this enzyme (27). When treating the cyp27a1^{-/-} mice with cholestyramine, however, opposite effects were obtained on plasma levels of cholestanol. Under these conditions, the plasma levels of 7 α -hydroxy-4-cholesten-3-one increased about 3-fold with a parallel increase in the levels of cholestanol. This is consistent with 7 α -hydroxy-4-cholesten-3-one as a precursor of cholestanol under conditions when its level in plasma is considerably higher than about 200 ng/ml (corresponding to the levels obtained in a wild-type mouse treated with cholestyramine).

Increased brain levels of cholestanol were found in the cyp27a1-deficient mice when the plasma levels of 7 α -hydroxy-4-cholesten-3-one were between 500 and 4,500 ng/ml. Reducing the plasma levels of 7 α -hydroxy-4-




Fig. 4. Levels of cholestanol and 7 α -hydroxy-4-cholesten-3-one (C4) in female wild-type (WT) and CYP27KO mice (A) in plasma. B: Cerebrum cholestanol levels and plasma 7 α -hydroxy-4-cholesten-3-one levels shown in the same diagrams. Number of mice, used to measure 7 α -hydroxy-4-cholesten-3-one levels in plasma, per group were: WT and KO; ctrl n4 (pool) and n3, cholestyramine n4 (pool) and n3, cholic acid n5 and n5). Number of mice, used to measure cholestanol levels in plasma and cerebrum, per group were: WT and KO; ctrl n4 and n5, cholestyramine n4 and n3, cholic acid n5 and n5.




Fig. 5. Sequence of reactions in connection with formation of cholestanol by the “classical” pathway (I) and the “alternative” pathway (II) involving intermediates in bile acid synthesis.

cholesten-3-one in the *cyp27a1*-deficient mice by treatment with cholic acid led to expected reduction in levels of cholestanol in the brain.

Surprisingly, an increase of the levels of 7α-hydroxy-4-cholesten-3-one, by a factor of two, by treating the *cyp27a1*^{−/−} mice with cholestyramine did not lead to the expected increase of cholestanol concentration in the brain. The reason for this may have been the relatively short duration of the treatment. Another possibility could be presence of a metabolism of cholestanol in the brain that may be activated when the accumulation reaches a certain level.

The major mechanism by which cholesterol is removed from the brain involves a 24S-hydroxylation catalyzed by the brain-specific cholesterol 24S-hydroxylase. This enzyme is also active toward cholestanol (28), but the role of this enzyme in connection with the accumulation of cholestanol is uncertain. A treatment of a *cyp27a1*-deficient mouse with daily injections of very large amounts of 7α-hydroxy-4-cholesten-3-one during 3 months led, however, to a marked increase in brain levels of cholestanol. If a mechanism is present in the brain of the *cyp27a1*^{−/−} mice that counteracts the accumulation, it is evident that this mechanism can be overcome by a sufficiently high flux of the cholestanol precursor into the brain. Further evidence for a direct flux of 7α-hydroxy-4-cholesten-3-one into the brain with subsequent formation of cholestanol was obtained by the demonstration of incorporation of deuterium in brain cholestanol in a mouse injected with ²H₇-labeled 7α-hydroxy-4-cholesten-3-one.

The mechanism for accumulation of cholestanol in the brain of *cyp27a1*^{−/−} mice demonstrated here is likely to also be valid for patients with CTX. The suppressive effect of cholic acid on the accumulation of cholestanol in the brain

of the *cyp27a1*^{−/−} mice is analogous to the suppressive effect of chenodeoxycholic acid on brain xanthomas reported for patients with CTX (8). It should be noted that cholic acid is the major suppressor of cholesterol 7α-hydroxylase in mice (21), whereas chenodeoxycholic acid is the major suppressor in humans (1). Unexpectedly, the treatment with cholic acid appeared to increase, rather than decrease, lathosterol levels in the *cyp27a1*^{−/−} mice. The reason for this is unknown, and further experiments have been initiated to elucidate the mechanism behind this effect.

The accumulation of cholestanol in relation to cholesterol was considerably higher in tendons and brain of the *cyp27a1*^{−/−} mice than in circulation or liver. Tendons and brain are also the preferential sites for formation of xanthomas in patients with CTX. The CYP27A1-mediated conversion of cholesterol into 27-hydroxycholesterol and cholestenic acid can be regarded as an antiatherogenic mechanism by which cholesterol can be eliminated from macrophages (29) and possibly also from glial cells. Not only cholesterol but also cholestanol can be eliminated by this mechanism (30). The classical HDL-mediated mechanism for reversed cholesterol transfer may be less effective in tendons and brain than in other tissues and organs, increasing the importance of CYP27A1 for removal of excess cholesterol and cholestanol. We have shown that human monocyte-derived macrophages eliminate cholestanol less efficiently than cholesterol, both by the classical HDL-mediated mechanism and by CYP27A1 (30). A reduced capacity for removal of sterols is thus likely to result in a preferential accumulation of cholestanol.

The reason for development of cholesterol-containing xanthomas in the brain and tendons of patients with CTX but not in *cyp27a1*-deficient mice is still not known. The

accumulation of cholestanol in brain and tendons was thus not accompanied by a parallel accumulation of cholesterol in our mouse model. The lathosterol levels were slightly increased, suggesting increased cholesterol synthesis. Whether this increase is a consequence of the accumulation of cholestanol or the lack of 27-hydroxycholesterol is not possible to evaluate from the present study. In CTX, the upregulation of the cholesterol 7 α -hydroxylase due to the low production of chenodeoxycholic acid is considerably higher than the corresponding upregulation in the mouse model. As a consequence, both production and accumulation of cholestanol are considerably higher in CTX than in the mouse model. The production of cholestanol in patients with CTX has been estimated to be about 40–50 mg/24 h (31, 32), which is about 10% of the normal rate of formation of bile acids in humans. The accumulation of cholestanol in the brain of CTX patients has been reported to be 20–40% of the sterol fraction (31). In our untreated cyp27a1-deficient mice, cholestanol corresponded only to about 3% of the sterol fraction. In the cyp27a1-deficient mouse treated with 7 α -hydroxy-4-cholesten-3-one, the corresponding enrichment was about 10%.

In contrast to the situation in human deficiency of the CYP27A1, cyp27^{-/-} mice have a marked upregulation of the nuclear xenobiotic receptor pregnane X receptor and its target genes (33). 7 α -Hydroxy-4-cholesten-3-one is one of the activators of this receptor. One of the metabolic consequences of this activation is an increased capacity to convert bile acid intermediates into more polar 25-hydroxylated alcohols. The excretion of such bile alcohols is, however, very modest in cyp27^{-/-} mice compared to the situation in CTX (4, 6). The reason for this is probably that 7 α -hydroxylation of cholesterol is rate limiting not only for the synthesis of cholestanol by the alternative pathway but also for the production of 25-hydroxylated bile alcohols. The cholesterol 7 α -hydroxylase is not upregulated in cyp27^{-/-} mice to the same extent as in patients with CTX. In any case, it is difficult to link differences in the degree of activation of the pregnane X receptor to the differences in the accumulation of cholestanol and cholesterol at the present state of knowledge.

To summarize, we have an explanation for the accumulation of cholestanol in the brain of CYP27A1-deficient mice and humans and for the higher accumulation in the human situation. We still do not know, however, why the accumulation of cholestanol in the human situation is associated with accumulation of cholesterol in the form of xanthomas. ■

REFERENCES

- Russell, D. W. 2003. The enzymes, regulation, and genetics of bile acid synthesis. *Annu. Rev. Biochem.* **72**: 137–174.
- Björkhem, I., Muri-Boberg, K., Leitersdorf, E. 2001. Inborn errors in bile acid biosynthesis: an storage of sterols other than cholesterol. In *Metabolic and Molecular Bases of Inherited Diseases*. C. R. Scriver, A. L. Beaudet, W. S. Sly, and D. Valle, editors. McGraw-Hill, 2961–2988.
- Dubrac, S., S. R. Lear, M. Ananthanarayanan, N. Balasubramanian, J. Bollineni, S. Shefer, H. Hyogo, D. E. Cohen, P. J. Blanche, R. M. Krauss, et al. 2005. Role of CYP27A in cholesterol and bile acid metabolism. *J. Lipid Res.* **46**: 76–85.
- Honda, A., G. Salen, Y. Matsuzaki, A. K. Batta, G. Xu, E. Leitersdorf, G. S. Tint, S. K. Erickson, N. Tanaka, and S. Shefer. 2001. Side chain hydroxylations in bile acid biosynthesis catalyzed by CYP3A are markedly up-regulated in Cyp27^{-/-} mice but not in cerebrotendinous xanthomatosis. *J. Biol. Chem.* **276**: 34579–34585.
- Honda, A., G. Salen, Y. Matsuzaki, A. K. Batta, G. Xu, E. Leitersdorf, G. S. Tint, S. K. Erickson, N. Tanaka, and S. Shefer. 2001. Differences in hepatic levels of intermediates in bile acid biosynthesis between Cyp27^{-/-} mice and CTX. *J. Lipid Res.* **42**: 291–300.
- Rosen, H., A. Reshef, N. Maeda, A. Lippoldt, S. Shpizen, L. Triger, G. Eggertsen, I. Björkhem, and E. Leitersdorf. 1998. Markedly reduced bile acid synthesis but maintained levels of cholesterol and vitamin D metabolites in mice with disrupted sterol 27-hydroxylase gene. *J. Biol. Chem.* **273**: 14805–14812.
- Björkhem, I., S. Skrede, M. S. Buchmann, C. East, and S. Grundy. 1987. Accumulation of 7 α -hydroxy-4-cholesten-3-one and cholesta-4,6-dien-3-one in patients with cerebrotendinous xanthomatosis: effect of treatment with chenodeoxycholic acid. *Hepatology*. **7**: 266–271.
- Berginer, V. M., G. Salen, and S. Shefer. 1989. Cerebrotendinous xanthomatosis. *Neurol. Clin.* **7**: 55–74.
- Repa, J. J., E. G. Lund, J. D. Horton, E. Leitersdorf, D. W. Russell, J. M. Dietsch, and S. D. Turley. 2000. Disruption of the sterol 27-hydroxylase gene in mice results in hepatomegaly and hypertriglyceridemia. Reversal by cholic acid feeding. *J. Biol. Chem.* **275**: 39685–39692.
- Skrede, S., I. Björkhem, M. S. Buchmann, G. Hopen, and O. Fausa. 1985. A novel pathway for biosynthesis of cholestanol with 7 α -hydroxylated C27-steroids as intermediates, and its importance for the accumulation of cholestanol in cerebrotendinous xanthomatosis. *J. Clin. Invest.* **75**: 448–455.
- Buchmann, M. S., and O. P. Clausen. 1986. Effects of cholestanol feeding and cholestyramine treatment on the tissue sterols in the rabbit. *Lipids*. **21**: 738–743.
- Byun, D. S., T. Kasama, T. Shimizu, H. Yorifuji, and Y. Seyama. 1988. Effect of cholestanol feeding on sterol concentrations in the serum, liver, and cerebellum of mice. *J. Biochem.* **103**: 375–379.
- Björkhem, I., and S. Meaney. 2004. Brain cholesterol: long secret life behind a barrier. *Arterioscler. Thromb. Vasc. Biol.* **24**: 806–815.
- Panzenboeck, U., U. Andersson, M. Hansson, W. Sattler, S. Meaney, and I. Björkhem. 2007. On the mechanism of cerebral accumulation of cholestanol in patients with cerebrotendinous xanthomatosis. *J. Lipid Res.* **48**: 1167–1174.
- Lovgren-Sandblom, A., M. Heverin, H. Larsson, E. Lundström, J. Wahren, U. Diczfalusy, and I. Björkhem. 2007. Novel LC-MS/MS method for assay of 7 α -hydroxy-4-cholesten-3-one in human plasma. Evidence for a significant extrahepatic metabolism. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **856**: 15–19.
- Folch, J., I. Ascoli, M. Lees, J. A. Meath, and B. N. Le. 1951. Preparation of lipid extracts from brain tissue. *J. Biol. Chem.* **191**: 833–841.
- Folch, J., M. Lees, and G. H. Sloane Stanley. 1957. A simple method for the isolation and purification of total lipides from animal tissues. *J. Biol. Chem.* **226**: 497–509.
- Lund, E., L. Sisfontes, E. Reihner, and I. Björkhem. 1989. Determination of serum levels of unesterified lathosterol by isotope dilution-mass spectrometry. *Scand. J. Clin. Lab. Invest.* **49**: 165–171.
- Acimovic, J., A. Lovgren-Sandblom, K. Monostory, D. Rozman, M. Golcnik, D. Lutjohann, and I. Björkhem. 2009. Combined gas chromatographic/mass spectrometric analysis of cholesterol precursors and plant sterols in cultured cells. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **877**: 2081–2086.
- Heverin, M., S. Meaney, A. Brafman, M. Shafir, M. Olin, M. Shafaati, S. von Bahr, L. Larsson, A. Lovgren-Sandblom, U. Diczfalusy, et al. 2007. Studies on the cholesterol-free mouse: strong activation of LXR-regulated hepatic genes when replacing cholesterol with desmosterol. *Arterioscler. Thromb. Vasc. Biol.* **27**: 2191–2197.
- Li-Hawkins, J., M. Gafvels, M. Olin, E. G. Lund, U. Andersson, G. Schuster, I. Björkhem, D. W. Russell, and G. Eggertsen. 2002. Cholic acid mediates negative feedback regulation of bile acid synthesis in mice. *J. Clin. Invest.* **110**: 1191–1200.
- Caligiuri, G., A. Nicoletti, X. Zhou, I. Tornberg, and G. K. Hansson. 1999. Effects of sex and age on atherosclerosis and autoimmunity in apoE-deficient mice. *Atherosclerosis*. **145**: 301–308.
- Paigen, B., A. Morrow, P. A. Holmes, D. Mitchell, and R. A. Williams. 1987. Quantitative assessment of atherosclerotic lesions in mice. *Atherosclerosis*. **68**: 231–240.

24. Bjorkhem, I., and K. E. Karlmar. 1974. Biosynthesis of cholestanol: conversion of cholesterol into 4-cholesten-3-one by rat liver microsomes. *Biochim. Biophys. Acta*. **337**: 129–131.
25. Bjorkhem, I., M. Buchmann, and S. Bystrom. 1992. Mechanism and stereochemistry in the sequential enzymatic saturation of the two double bonds in cholesta-4,6-dien-3-one. *J. Biol. Chem.* **267**: 19872–19875.
26. Hylemon, P. B., and J. Harder. 1998. Biotransformation of monoterpenes, bile acids, and other isoprenoids in anaerobic ecosystems. *FEMS Microbiol. Rev.* **22**: 475–488.
27. Shefer, S., S. Hauser, and E. H. Mosbach. 1968. 7- α -hydroxylation of cholestanol by rat liver microsomes. *J. Lipid Res.* **9**: 328–333.
28. Mast, N., R. Norcross, U. Andersson, M. Shou, K. Nakayama, I. Bjorkhem, and I. A. Pikuleva. 2003. Broad substrate specificity of human cytochrome P450 46A1 which initiates cholesterol degradation in the brain. *Biochemistry*. **42**: 14284–14292.
29. Babiker, A., O. Andersson, E. Lund, R. J. Xiu, S. Deeb, A. Reshef, E. Leitersdorf, U. Diczfalusy, and I. Bjorkhem. 1997. Elimination of cholesterol in macrophages and endothelial cells by the sterol 27-hydroxylase mechanism. Comparison with high density lipoprotein-mediated reverse cholesterol transport. *J. Biol. Chem.* **272**: 26253–26261.
30. von Bahr, S., T. Movin, N. Papadogiannakis, I. Pikuleva, P. Ronnow, U. Diczfalusy, and I. Bjorkhem. 2002. Mechanism of accumulation of cholesterol and cholestanol in tendons and the role of sterol 27-hydroxylase (CYP27A1). *Arterioscler. Thromb. Vasc. Biol.* **22**: 1129–1135.
31. Bhattacharyya, A. K., D. S. Lin, and W. E. Connor. 2007. Cholestanol metabolism in patients with cerebrotendinous xanthomatosis: absorption, turnover, and tissue deposition. *J. Lipid Res.* **48**: 185–192.
32. Salen, G., and S. M. Grundy. 1973. The metabolism of cholestanol, cholesterol, and bile acids in cerebrotendinous xanthomatosis. *J. Clin. Invest.* **52**: 2822–2835.
33. Goodwin, B., K. C. Gauthier, M. Umetani, M. A. Watson, M. I. Lochansky, J. L. Collins, E. Leitersdorf, D. J. Mangelsdorf, S. A. Kliewer, and J. J. Repa. 2003. Identification of bile acid precursors as endogenous ligands for the nuclear xenobiotic pregnane X receptor. *Proc. Natl. Acad. Sci. USA*. **100**: 223–228.