

1 Chemically amplified ^{19}F - ^1H nuclear Overhauser effects
I. Kuprov and P.J. Hore

8 Towards quantitative measurements in solid-state CPMAS NMR: A Lee–Goldburg frequency modulated cross-polarization scheme
Riqiang Fu, Jun Hu, and Timothy A. Cross

18 The effects of cholesterol on magnetically aligned phospholipid bilayers: a solid-state NMR and EPR spectroscopy study
Jun-Xia Lu, Marc A. Caporini, and Gary A. Lorigan

31 Self-diffusion measurements with chemical shift resolution in inhomogeneous magnetic fields
Daniel Topgaard and Alexander Pines

36 CORCEMA refinement of the bound ligand conformation within the protein binding pocket in reversibly forming weak complexes using STD-NMR intensities
V. Jayalakshmi and N. Rama Krishna

46 Reduced apparent longitudinal relaxation times in slice-selective experiments in strong magnetic field gradients
Nikolaus Nestle, Bernadeta Walaszek, and Markus Nolte

53 Subspace-based MRS data quantitation of multiplets using prior knowledge
T. Laudadio, Y. Selén, L. Vanhamme, P. Stoica, P. Van Hecke, and S. Van Huffel

66 Low temperature solid-state NMR experiments of half-integer quadrupolar nuclides: caveats and data analysis
Andrew S. Lipton, Robert W. Heck, Jesse A. Sears, and Paul D. Ellis

75 2D TRIPLE in orientationally disordered samples—a means to resolve and determine relative orientation of hyperfine tensors
D. Goldfarb, B. Epel, H. Zimmermann, and G. Jeschke

88 Hyperfine decoupling in electron paramagnetic resonance as a powerful tool for unraveling complicated ESEEM spectra of $S = 1/2$, $I \geq 1/2$ systems
George Mitrikas and Arthur Schweiger

97 Frequency shifts in parametrically enhanced low-field MR detection
Daniel W. Sinkovits and Mark S. Conradi

103 Design of a constant adiabaticity pulse for selective population inversion
Lorenz Mitschang, Bernd Ittermann, Florian Schubert, and Herbert Rinneberg

110 Tailored sinc pulses for uniform excitation and artifact-free radio frequency time-domain EPR imaging
N. Devasahayam, R. Murugesan, K. Matsumoto, J.B. Mitchell, J.A. Cook, S. Subramanian, and M.C. Krishna

Continued

118 The application of the ERETIC method to 2D-NMR
Nathalie Michel and Serge Akoka

124 Resolution of ^{13}C - ^{19}F interactions in the ^{13}C NMR of spinning solids and liquid crystals
Giancarlo Antonioli and Paul Hodgkinson

132 Electron spin-relaxation via vibronic level of nickel (I) and nickel (III) cyanide complexes in NaCl single crystals
N.V. Vugman, M.B. de Araújo, N.M. Pinhal, C.J. Magon, and A.J. da Costa Filho

137 Rotational resonance in uniformly ^{13}C -labeled solids: effects on high-resolution magic-angle spinning NMR spectra and applications in structural studies of biomolecular systems
Aneta T. Petkova and Robert Tycko

147 Cross-polarization schemes for peptide samples oriented in hydrated phospholipid bilayers
Hyeongnam Kim, Timothy A. Cross, and Riqiang Fu

153 Orientation of the antimicrobial peptide PGLa in lipid membranes determined from ^{19}F -NMR dipolar couplings of 4-CF₃-phenylglycine labels
Ralf W. Glaser, Carsten Sachse, Ulrich H.N. Dürr, Parvesh Wadhwan, and Anne S. Ulrich

164 Distortion-free single point imaging of multi-layered composite sandwich panel structures
Andrew E. Marble, Igor V. Mastikhin, Rod P. MacGregor, Mohamad Akl, Gabriel LaPlante, Bruce G. Colpitts, Pearl Lee-Sullivan, and Bruce J. Balcom

175 NMR acceleration mapping in percolation model objects
Bogdan Buhai, Aidar Hakimov, Ioan Ardelean, and Rainer Kimmich