

4-Anilino-6,7-dialkoxyquinoline-3-carbonitrile Inhibitors of Epidermal Growth Factor Receptor Kinase and Their Bioisosteric Relationship to the 4-Anilino-6,7-dialkoxyquinazoline Inhibitors

Allan Wissner,* Dan M. Berger, Diane H. Boschelli, M. Brawner Floyd, Jr., Lee M. Greenberger, Brian C. Gruber, Bernard D. Johnson, Nellie Mamuya, Ramaswamy Nilakantan, Marvin F. Reich, Ru Shen, Hwei-Ru Tsou, Erik Upeslakis, Yu Fen Wang, Biqi Wu, Fei Ye, and Nan Zhang

Wyeth-Ayerst Research, A Division of American Home Products, 401 North Middletown Road, Pearl River, New York 10965-1215

Received May 15, 2000

The synthesis and SAR of a series of 4-anilino-6,7-dialkoxyquinoline-3-carbonitrile inhibitors of epidermal growth factor receptor (EGF-R) kinase are described. Condensation of 3,4-dialkoxyanilines with ethyl (ethoxymethylene)cianoacetate followed by thermal cyclization gave, regiospecifically, 6,7-dialkoxy-4-oxo-1,4-dihydroquinoline-3-carbonitriles. Chlorination (POCl_3) followed by the reaction with substituted anilines furnished the 4-anilino-6,7-dialkoxyquinoline-3-carbonitrile inhibitors of EGF-R kinase. An alternate synthesis of these compounds starts with a methyl 3,4-dialkoxybenzoate. Nitration followed by reduction (Fe , NH_4Cl , $\text{MeOH-H}_2\text{O}$) gave a methyl 2-amino-4,5-dialkoxybenzoate. Amidine formation using DMF-acetal followed by cyclization using LiCH_2CN furnished a 6,7-dialkoxy-4-oxo-1,4-dihydroquinoline-3-carbonitrile, which was transformed as before. Compounds containing acid, ester, amide, carbinol, and aldehyde groups at the 3-position of the quinoline ring were also prepared for comparison, as were several 1-anilino-6,7-dimethoxyisoquinoline-4-carbonitriles. The compounds were evaluated for their ability to inhibit the autophosphorylation of the catalytic domain of EGF-R. The SAR of these inhibitors with respect to the nature of the 6,7-alkoxy groups, the aniline substituents, and the substituent at the 3-position was studied. The compounds were further evaluated for their ability to inhibit the growth of cell lines that overexpress EGF-R or HER-2. It was found that 4-anilinoquinoline-3-carbonitriles are effective inhibitors of EGF-R kinase with activity comparable to the 4-anilinoquinazoline-based inhibitors. A new homology model of EGF-R kinase was constructed based on the X-ray structures of Hck and FGF receptor-1 kinase. The model suggests that with the quinazoline-based inhibitors, the N3 atom is hydrogen-bonded to a water molecule which, in turn, interacts with Thr 830. It is proposed that the quinoline-3-carbonitriles bind in a similar manner where the water molecule is displaced by the cyano group which interacts with the same Thr residue.

Introduction

Protein tyrosine kinases play a role in normal cell growth. Many of the growth factor receptor proteins have intracellular domains that function as tyrosine kinases, and it is with this that they effect signaling. The interaction of growth factors with these receptors is a necessary event in normal regulation of cell growth. However, under certain conditions, as a result of overexpression, mutation, or coexpression of the ligand and the receptor, these receptors can become hyperactivated; the result of this is uncontrolled cell proliferation.¹ Among the growth factor receptor kinases that have been identified as being important in cancer is epidermal growth factor receptor (EGF-R) kinase (also known as erb-B1 or HER-1) and the related human epidermal growth factor receptor HER-2 (also known as erbB-2 or *neu*). EGF-R is overexpressed in numerous tumors² including those derived from the brain, lung, bladder, head, and neck. Such overexpression has been correlated with poor prognosis in some of these diseases. The gene product is frequently mutated in the extra-

cytoplasmic domain of the receptor in gliomas,³ prostate cancer,⁴ breast cancer,⁵ and non-small-cell lung cancer;⁶ this mutation makes the receptor constitutively active.⁷ In addition, the receptor need not be overexpressed or mutated to be activated, since the ligands for the receptor (EGF or transforming growth factor- α (TGF α)) can be produced within the same cancerous tissue or cell that expresses EGF-R. This suggests that paracrine or autocrine loops stimulate hyperproliferation as in prostate cancer,⁸ head and neck cancer,⁹ ovarian cancer,¹⁰ non-small-cell lung cancer,¹¹ and bladder cancer.¹² EGF-R hyperactivation has also been implicated in other diseases including polycystic kidney disease,¹³ psoriasis,¹⁴ and asthma.¹⁵

Since in many types of proliferative diseases, the phosphorylation event mediated by EGF-R or HER-2 kinases is a necessary signal for cell division to occur and since hyperactivation of these kinases has been associated with these diseases, an inhibitor of this event, an EGF-R or HER-2 kinase inhibitor, may have potential therapeutic value. Since these two receptor kinases have a high sequence homology in their catalytic

* To whom correspondence should be addressed. Tel: 914-732-3580. Fax: 914-732-5561. E-mail: wissnea@war.wyeth.com.


domains,¹⁶ one might expect that an inhibitor of one would have similar effects on the other.

The middle of the past decade was marked by the discovery of the 4-anilinoquinazolines, a series of potent and selective ATP-competitive inhibitors of EGF-R kinase.¹⁷ Several pharmaceutical firms,¹⁸ including ourselves,¹⁹ established programs based on these inhibitors, and now at least three such compounds have entered or will be entering clinical trials.²⁰ In extending our work with this series of inhibitors, we realized that the nitrogen atom located at the 3-position of these quinazoline inhibitors is an important feature needed for good activity. Replacing this atom with a carbon leads to a significant loss in the ability of the compound to inhibit the enzyme.²¹ We reasoned that given the importance of this nitrogen atom for activity, it is likely that it has a specific interaction with the enzyme. Presumably, it functions as a hydrogen bond acceptor. In designing our inhibitors, we made use of a homology model of the enzyme that is similar to the model previously constructed.²² More recently we have developed a refined model as discussed below. With either model, docking experiments using our inhibitors did not reveal any residues close by to which the N3 atom could directly interact. We therefore made the hypothesis that this nitrogen atom could be interacting with a water molecule and that this water molecule could then serve as a bridge between the drug and enzyme. Such water bridging is a well-known phenomenon.²³ Since the N3 atom of the quinazoline inhibitor, in solution, surely forms a hydrogen bond with a water molecule, retaining this interaction after binding to the enzyme should have a minimal effect on the entropy of binding. In addition, it occurred to us that if this nitrogen atom were removed and replaced with a carbon atom that had an attached electron-withdrawing group (such as a cyano group), such a molecule should retain a similar overall shape and charge distribution compared to the quinazoline hydrogen-bonded to a water molecule. In this communication we would like to report that applying the above concept has resulted in a new series of 4-anilinoquinoline-3-carbonitriles that are potent inhibitors of EGF-R kinase.

Chemistry

Since earlier work by the Parke-Davis group with the quinazoline-based inhibitors of EGF-R established that 6,7-dialkoxy substitution is compatible with good activity,²⁴ we decided to retain this feature in our initial compounds. One of the methods we used to prepare the intermediate quinolones **3a–c** follows the method of Bredereck²⁵ as shown in Scheme 1. Condensation of anilines **1a,b** with diethyl ethoxymethylenemalonate or ethyl (ethoxymethylene)cyanoacetate followed by thermal cyclization in refluxing Dowtherm gave, regiospecifically, **3a–c**. These quinolones were converted, in good yield, to the corresponding chloroquinolines by refluxing in an excess of POCl₃. The preparations of **3c**^{25b} and **4a**²⁶ have been described previously. In the final step of the sequence, refluxing a solution of a chloroquinoline and a substituted aniline derivative in ethoxyethanol gave the 4-anilino-6,7-dialkoxyquinoline-3-carbonitriles **5–38** (see Table 1). In most cases, the


Scheme 1^a



^a (a) For **2a**: C₂H₅OCH=C(CO₂C₂H₅)₂, neat, 100 °C; for **2b,c**: C₂H₅OCH=C(CO₂C₂H₅)CN, toluene, reflux; (b) Dowtherm, 258 °C; (c) POCl₃, reflux; (d) C₂H₅O(CH₂)₂OH, reflux; for **35**: NaH, DMF, reflux; (e) (i) NaNO₂, HOAc, H₂O, 0 °C, (ii) NaN₃; (f) (CH₃CO)₂O, CH₃CO₂H.

products can be isolated as the hydrochloride salt by filtering directly from the reaction mixture, or alternatively, the free base can be isolated after neutralization of the salt with base. Due to the hindered nature of 2-bromoaniline, it was necessary to prepare compound **35** using sodium hydride in refluxing DMF. Compound **40** was prepared by acetylation of **16** with acetic anhydride in acetic acid. Compound **39** was prepared by a two-step process involving formation of the diazonium salt of **16** using sodium nitrite and acetic acid followed by the reaction with sodium azide.

An alternate synthesis of some of these compounds is shown in Scheme 2. Compounds containing unsymmetrical 6,7-dialkoxy substitution were prepared from the isomeric methylbenzoates **41a,b** by alkylation with ethyl iodide and potassium carbonate in refluxing DMF. Compound **42c** was prepared as already described.²⁷ Nitration of **42a–c** in acetic acid at 50 °C gave the nitro derivatives **43a–c**. Reduction using iron and NH₄Cl in a water–methanol mixture at reflux followed by amidine formation using *N,N*-dimethylformamide dimethylacetal in DMF at reflux gave **45a–c** which were used without additional purification. The lithium anion of acetonitrile was prepared in THF by the addition of *n*-butyllithium at –78 °C. Addition of the amidines to


Scheme 2^a

^a (a) C₂H₅I, K₂CO₃, DMF, 100 °C; (b) HNO₃, HOAc, 50 °C; (c) Fe, NH₄Cl, CH₃OH, H₂O, reflux; (d) DMA-DMF, DMF, reflux; (e) (i) *n*-BuLi, CH₃CN, THF, -78 °C, (ii) HOAc, -78 to 25 °C; (f) for **47a,c**: POCl₃, reflux; for **47b**: (COCl)₂, CH₂Cl₂, reflux; (g) C₂H₅O(CH₂)₂OH or C₂H₅OH, reflux.

this solution containing 2 equiv of the anion followed by the addition of acetic acid and warming to 25 °C furnished the quinolones **46a–c**. These were then converted to the 4-anilino-6,7-dialkoxyquinoline-3-carbonitriles **48–50** as described above.

Compounds containing varying substitution at the 3-position were prepared by simple functional group manipulations as shown in Scheme 3. Reduction of the ester **5** with DIBAL gave the carbinol derivative **51** which, in turn, was oxidized to the aldehyde **52** using pyridinium chlorochromate buffered with sodium acetate in methylene chloride. Base hydrolysis of **5** furnished the carboxylic acid **53** which was then converted to the amide **54** using a two-step procedure consisting of first activating the carboxylate group as the acyl imidazole derivative followed by the addition of ammonia.

Additional variation of the 6,7-alkoxy groups was accomplished as shown in Scheme 3 by first demethylating **8** by heating in neat pyridine hydrochloride at 210 °C to give the 6,7-dihydroxy derivative **55**. Bis-alkylation of **55** with ethyl iodide, chloromethyl methyl ether, or bromoethyl methyl ether using K₂CO₃ in refluxing DMF gave compounds **6**, **56**, and **57**, respectively. Compounds **58** and **59** which have, respectively, a fused five- or seven-membered diether ring were prepared by alkylation using bromochloromethane and

Scheme 3^a

^a (a) DIBAL, toluene; (b) PCC, NaOAc, CH₂Cl₂; (c) NaOH, H₂O, C₂H₅OH, reflux; (d) (i) CDI, DMF, (ii) NH₃; (e) pyridine·HCl, 210 °C; (f) C₂H₅I, K₂CO₃, DMF; (g) CH₃OCH₂Cl, K₂CO₃, DMF; (h) CH₃O(CH₂)₂Br, K₂CO₃, DMF; (i) BrCH₂Cl, Cs₂CO₃, DMF; (j) Br(CH₂)₃Br, K₂CO₃, DMF; (k) substituted aniline, isopropanol, reflux.

Cs₂CO₃ in DMF or 1,3-dibromopropane and potassium carbonate in DMF, respectively.

Finally, we were interested in seeing if replacement of the N1 atom of a quinazoline inhibitor with a carbon bearing a cyano group would also lead to an active EGF-R inhibitor. Accordingly, we prepared the 1-anilinoquinoline-4-carbonitrile derivatives **61** and **62** from the corresponding known²⁸ chloro derivative **60** by treating it with the substituted aniline in refluxing 2-propanol.

Molecular Modeling

In a prior publication,¹⁹ we discussed modeling work on EGF-R kinase using a homology model similar to that described by Knighton et al.²² This model was based on the first crystal structure of a protein kinase, viz. cAMP-dependent protein kinase, a serine/threonine kinase. While the sequence homology between cAMP-dependent protein kinase and EGF-R kinase is 41%, the sequence identity is only 27%. Since then, a number of additional kinase structures have been solved, including several tyrosine kinases. We therefore decided to build a homology model of EGF-R kinase using more closely related crystal structures as templates. We used the BLAST server at NCBI to search the Protein Data Bank. The top-ranked structure was fibroblast growth factor receptor-1 kinase,²⁹ FGF receptor-1 (PDB code: 1fgk), with a sequence identity of 33%. However, several residues in the activation loop are missing from this




Figure 1. Proposed binding models for (a) quinazoline **63** and (b) 3-cyanoquinoline **8** at the ATP site in EGF-R based on the homology model.

structure. Another high-ranked hit in the BLAST search, **hematopoietic cell kinase**,³⁰ Hck (PDB code: 1qcf) was also explored as a possible template. However, its alignment with EGF-R involved a major insertion in the glycine loop which is close to the ATP binding site. The resulting model had major distortions in the ATP binding site. We therefore decided to use a combination of these two templates: FGF receptor-1 for the N-terminal lobe and Hck for the C-terminal lobe. We used the homology modeling facility within the Molecular Operating Environment (MOE) software to do this. The resulting model was minimized using Quanta/CHARMm and used for subsequent studies.

A recent paper from a group at Glaxo Wellcome³¹ revealed the crystal structures of the 4-anilinoquinazoline class of compounds complexed with two kinases: CDK2 and p38-MAP kinase. The binding mode of the quinazoline ring in both structures is similar, the N1 atom of the quinazoline being hydrogen-bonded to the backbone nitrogen of a residue on the domain connector strand (Leu 83 in CDK2 and Met 109 in p38). In the p38 structure, the N3 atom forms a hydrogen bond with a water molecule, which, in turn, makes an interaction with the hydroxyl group of a Thr residue. These two crystal structures strongly suggest that there is a single binding orientation of the quinazoline ring in kinases. We therefore revised our earlier binding model to reflect these new data from the crystal structures. The water molecules found in the Hck structure were added to the homology model. We also used a pentapeptide model substrate AEYLR by positioning it so that the tyrosine hydroxyl is close to the catalytic aspartate (Asp 813) and close to the γ -phosphate of ATP. The enzyme-ATP-substrate model was built and optimized separately. We then removed the ATP molecule and replaced it with quinazoline inhibitor **63** (PD 153035),²⁴ using the figures and descriptions of the published X-ray structures as a guide to orient the molecule. The entire complex was

optimized by several cycles of molecular dynamics and minimization using Quanta/CHARMm.

In the final model with the quinazoline **63** (see Figure 1a), the N1 atom of the quinazoline forms a hydrogen bond with the backbone NH of Met 769 and the N3 atom forms a hydrogen bond to a water molecule. This water molecule, in turn, forms a hydrogen bond with the hydroxyl group of Thr 830. The C2 atom of the quinazoline is 3.28 Å from the backbone carbonyl oxygen of Gln 767 and the C8 atom is 4.20 Å from the backbone carbonyl oxygen of Met 769.

The corresponding model with the quinoline-3-carbonitrile **8** (see Figure 1b) was generated in a similar manner. The orientation of the molecule and its interactions with the protein are similar to that observed in the quinazoline model. The hydrogen bond of the N1 atom to the backbone NH of Met 769 is retained. In this model the 3-cyano group displaces the water molecule that was previously hydrogen-bonded to the N3 atom of the quinazoline and instead makes its own interaction with the hydroxyl group of Thr 830.

As indicated by these models, the similar manner in which **63** and **8** bind suggests that the carbon with its attached cyano group (C-CN) of **8** is bioisosteric with the N3 atom of **63** hydrogen-bonded to a water molecule. To lend more credence to this idea, we performed single-point AM1 calculations on **63** and **8** each in their binding conformations as extracted from the models. The results of these calculations are shown in Figure 2. It is evident from these calculations that both systems are similar in terms of their overall shape and the location of charge.

After this work was completed, a report appeared that described a series of quinazoline and benztriazine inhibitors of Scytalone dehydratase along with the 3-cyanoquinoline and 3-cyanocinnoline inhibitors derived from them, respectively.³² This report compared the X-ray structures of a benztriazine and its corre-




Figure 2. (a) Electrostatic potential surfaces (0.005 contour level) calculated for **63** hydrogen-bonded to the water molecule and (b) the corresponding surface for **8**. Single-point calculations of the structures extracted from the binding models were accomplished using the semiempirical AM1 method as implemented in MacSpartan.

sponding 3-cyanocinnoline bound at the active site of this enzyme. Although this protein is not a kinase, the manner in which the molecules bind has relevance to the discussion at hand. In this structure, the N3 atom of the benztriazine forms a hydrogen bond with a crystallographic water molecule. The water molecule, in turn, forms a bridge to the hydroxyl groups of two Tyr residues. In addition, the authors prepared a corresponding 3-cyanocinnoline, and in the resulting X-ray structure, the cyano group occupies a position corresponding to the position of the water molecule in the benztriazine structure where the nitrogen atom of the cyano groups bridges the same two Tyr residues. This is analogous to the situation we are now proposing for the binding of the quinazoline and quinoline-3-carbonitrile inhibitors at the ATP binding pocket of EGF-R.

Our homology model of EGF-R differs somewhat from the model proposed by the Parke-Davis group³³ in that while the orientation of the bound quinazoline is similar and has the same interaction between N1 and Met 769, the latter model has a direct hydrogen bond between N3 and the hydroxyl group of Thr 766; it would seem that the EGF-R inhibition activity exhibited by the quinoline-3-carbonitriles, as discussed below, is not easily explained by this latter model.

Results and Discussion

The compounds shown in Table 1 were evaluated for their ability to inhibit EGF-R kinase. To compare these data with similar data already reported for the quinazoline-based inhibitors, the inhibitory activity of the standard compound, **63**, under our assay conditions, is incorporated into the data set. First, it is immediately evident that the IC_{50} value we obtain with **63** for inhibition of EGF-R is significantly higher than was found by previous workers.²⁴ Additionally, for our compounds, we do not see the extremely large range in potency found by these workers for the quinazoline inhibitors. We attribute these observations to differ-


ences in the nature of the enzyme and substrate as well as to the overall assay conditions. We use a solid-phase ELISA-based assay. Our enzyme consists of the purified cytoplasmic domain of EGF-R, and we measure the inhibition of autophosphorylation of this protein. Other workers have measured inhibition using, in a soluble format, the entire enzyme, purified from A431 cells, along with an exogenous peptide substrate. To further address this issue, two compounds, **63** and its corresponding quinoline-3-carbonitrile analogue **8**, were additionally evaluated in EGF-R kinase assays using a soluble format, purified cytoplasmic domain, and exogenous substrate as described previously.¹⁹

In the ELISA-based EGF-R assay **63** is only about 3 times more potent than **8**. The difference in potency between the two compounds is similar when using the alternate protocol with the same enzyme preparation but in a soluble format and with an exogenous peptide substrate where the IC_{50} values for **63** and **8** are 0.0022 and 0.0075 μ M, respectively. As expected, the IC_{50} values are lower in the latter assay. These results establish that the quinoline-3-carbonitriles are potentially comparable to the quinazolines in their ability to inhibit EGF-R kinase.

We were interested in determining the effect on activity of replacing the 3-cyano group with other functionalities. With respect to EGF-R inhibition, replacement with an ester (**5**), carboxylic acid (**53**), carbinol (**51**), or amide (**54**) group leads to significant (79-fold to greater than 530-fold) loss in potency. While the aldehyde derivative **52** was closest in activity to **8**, it is still about 18-fold less potent. A possible rationalization of these data is that one might expect for **5**, **53**, **54**, and **52** that the low-energy conformations would be characterized by an intramolecular hydrogen bond between the carbonyl oxygen of the 3-substituent and the NH group of the aniline moiety. Our binding model would suggest that this hydrogen bond would have to be sacrificed for these compounds to adopt an appropriate binding conformation; this will result in a commensurate loss in binding energy. In addition, the sp³-hybridization of the cyano substituent could result in a more favorably directed interaction with Thr 830 than when the substituents are sp²-hybridized.

Since replacement of the N3 atom of the quinazoline with a carbon atom bearing a cyano group is compatible with good activity, it was obviously of interest to see what would happen when the same replacement was made with the N1 atom. Accordingly, we found that the isoquinoline-4-carbonitriles **61** and **62** are, respectively, 368- and 496-fold less potent than their corresponding quinoline-3-carbonitrile isomers **8** and **9**. This is an expected result in view of the binding model, which indicates a direct interaction of N1 of the quinoline, and quinazoline-based inhibitors with the backbone NH of Met 769.

We surveyed a variety of substituents at different positions on the aniline ring of these quinoline-3-carbonitriles. Our molecular model suggests that the aniline ring resides in a large, mostly lipophilic, pocket buried deep within the active site. Good activity is seen when the aniline ring is monosubstituted with a bromine (**8**) or chlorine (**9**) atom at the *meta*-position. A similar observation was reported for the quinazoline

Table 1. Inhibition of EGF-R Kinase and Inhibition of Cell Proliferation


compd ^a	X	Y	R ¹	R ²	R''	IC ₅₀ (μM)			
						EGF-R ^b	A431 ^c	SKBR3 ^c	SW620 ^c
5	C-CO ₂ Et	N	CH ₃	CH ₃	3-Br	>81.3	3.01	6.21	6.93
6	C-CN	N	C ₂ H ₅	C ₂ H ₅	3-Br	0.42	0.11	0.09	0.16
7	C-CN	N	C ₂ H ₅	C ₂ H ₅	3-Cl, 4-F	0.98	0.27	0.49	0.49
8	C-CN	N	CH ₃	CH ₃	3-Br	0.19 ^e	0.78	0.48	3.72
9	C-CN	N	CH ₃	CH ₃	3-Cl	0.18	0.59	1.76	7.06
10	C-CN	N	CH ₃	CH ₃	3-CF ₃	1.45	2.53	13.42	35.63
11	C-CN	N	CH ₃	CH ₃	3,4-di-OCH ₃	50.49	5.28	13.16	8.76
12	C-CN	N	CH ₃	CH ₃	3-F	0.74	3.07	8.29	13.52
13	C-CN	N	CH ₃	CH ₃	3-CN	14.18	3.51	25.97	43.29
14	C-CN	N	CH ₃	CH ₃	4-F	47.94	1.43	0.74	2.87
15	C-CN	N	CH ₃	CH ₃	3-COCH ₃	7.38	7.37	5.12	6.30
16	C-CN	N	CH ₃	CH ₃	3-NH ₂	0.82	6.62	2.45	14.33
17 ^d	C-CN	N	CH ₃	CH ₃	3-N(CH ₃) ₂	0.85	0.65	0.36	0.99
18	C-CN	N	CH ₃	CH ₃	3-NO ₂	0.87	2.14	2.00	17.27
19	C-CN	N	CH ₃	CH ₃	3-Cl, 4-F	0.54	0.70	0.47	2.3
20	C-CN	N	CH ₃	CH ₃	4-Cl, 2-F	13.85	0.818	0.54	5.34
21	C-CN	N	CH ₃	CH ₃	3-OH	6.36	1.52	0.95	7.8
22	C-CN	N	CH ₃	CH ₃	4-CH ₃	2.73	0.96	0.64	60.43
23	C-CN	N	CH ₃	CH ₃	3-CONH ₂	7.69	5.91	8.44	6.29
24	C-CN	N	CH ₃	CH ₃	3-Br, 4-CH ₃	1.79	0.75	0.55	2.89
25	C-CN	N	CH ₃	CH ₃	3-Cl, 4-OH	9.04	3.01	1.29	2.75
26	C-CN	N	CH ₃	CH ₃	3,5-di-Cl, 4-OH	3.15	3.136	2.017	3.43
27 ^d	C-CN	N	CH ₃	CH ₃	5-Cl, 2-OH	17.02	0.94	1.19	3.01
28 ^d	C-CN	N	CH ₃	CH ₃	3-SCH ₃	0.76	1.451	1.54	4.98
29 ^d	C-CN	N	CH ₃	CH ₃	3-Cl, 4-CH ₃	1.48	0.93	0.94	24.96
30 ^d	C-CN	N	CH ₃	CH ₃	2-F, 4-Br	>91.2	0.89	1.69	19.17
31	C-CN	N	CH ₃	CH ₃	4-CH(CH ₃) ₂	0.23	3.02	2.91	3.51
32	C-CN	N	CH ₃	CH ₃	2-CH(CH ₃) ₂	0.36	13.2	16.12	4.98
33	C-CN	N	CH ₃	CH ₃	3-CH(CH ₃) ₂	62.32	4.35	6.74	3.80
34	C-CN	N	CH ₃	CH ₃	4-Br	6.65	0.76	1.47	3.76
35	C-CN	N	CH ₃	CH ₃	2-Br	4.18	21.03	0.80	17.02
36 ^d	C-CN	N	CH ₃	CH ₃	3-CF ₃ , 4-F	0.16	7.72	0.52	29.53
37	C-CN	N	CH ₃	CH ₃	2-CH ₃ , 3-Br	0.84	3.64	0.39	11.83
38	C-CN	N	CH ₃	CH ₃	3-CH ₃ , 4-Br	0.62	0.73	1.38	3.09
39	C-CN	N	CH ₃	CH ₃	3-N ₃	2.21	1.48	1.16	4.74
40	C-CN	N	CH ₃	CH ₃	3-NHCOCH ₃	69.40	2.62	1.55	2.74
48	C-CN	N	C ₂ H ₅	CH ₃	3-Br	21.19	0.57	0.64	4.24
49	C-CN	N	CH ₃	C ₂ H ₅	3-Br	33.02	0.14	0.06	0.06
50	C-CN	N	CH ₃	-CH ₂ CH ₂ -	3-Br	>104.7	0.42	0.27	0.43
51	C-CH ₂ OH	N	CH ₃	CH ₃	3-Br	>102.2	7.91	15.36	23.15
52	C-CHO	N	CH ₃	CH ₃	3-Br	3.37	1.95	2.97	5.24
53	C-CO ₂ H	N	CH ₃	CH ₃	3-Br	34.22	18.23	35.71	53.82
54	C-CONH ₂	N	CH ₃	CH ₃	3-Br	15.17	4.65	7.78	9.87
55	C-CN	N	H	H	3-Br	0.35	2.23	2.74	4.52
56	C-CN	N	CH ₃ OCH ₂	CH ₃ OCH ₂	3-Br	6.75	14.50	14.20	9.79
57	C-CN	N	CH ₃ O(CH ₂) ₂	CH ₃ O(CH ₂) ₂	3-Br	1.10	0.71	0.56	4.32
58	C-CN	N	CH ₃	-CH ₂ -	3-Br	6.79	2.62	1.94	4.18
59	C-CN	N	CH ₃	-(CH ₂) ₃ -	3-Br	5.15	2.08	5.07	2.90
61 ^d	N	C-CN	CH ₃	CH ₃	3-Br	70.84	18.76	24.96	48.49
62	N	C-CN	CH ₃	CH ₃	3-Cl	88.29	26.13	27.67	64.45
63	N	N	CH ₃	CH ₃	3-Br	0.07 ^f	1.52	0.82	13.94

^a Unless indicated compounds were tested as the free base. ^b Concentration needed to inhibit the autophosphorylation of the cytoplasmic domain of EGF-R by 50% as determined from the dose-response curve. Determinations were done in duplicate and repeat values agreed, on average, within 40%. ^c Dose-response curves were determined at five concentrations. The IC₅₀ values are the concentrations needed to inhibit cell growth by 50% as determined from these curves. ^d Compound tested as the HCl salt. ^e Using the assay protocol described in ref 19 and purified cytoplasmic domain, the IC₅₀ for inhibition is 0.0075 μM. ^f Using the assay protocol described in ref 19 and purified cytoplasmic domain, the IC₅₀ for inhibition is 0.0022 μM.

series.²⁴ Moving the bromine atom to the *ortho*- or *para*-position as in **35** and **34** results in a 22- and 35-fold decrease, respectively, in the ability of the compound to inhibit EGF-R. Surprisingly, with the weakly electron-donating isopropyl group, this trend is reversed; the *meta*-substituted derivative **33** is by far less potent than either the *para*- or *ortho*-isomers **31** and **32**, respectively. Modeling offers no immediate explanation for this trend. For the *meta*-trifluoromethyl-substituted compounds **10** and **36**, adding an additional *para*-fluoro

substituent leads to a 9-fold increase in potency, while for the *meta*-chloro derivatives **9** and **19** this same modification leads to a modest (3-fold) decrease. Compounds **21** and **25–27**, having a hydroxyl group in various positions on the aniline ring, each showed poor activity, while compounds **16** and **17**, which bear amino groups at the *meta*-position, had only a moderate loss in potency.

We made a number of modifications to the 6,7-dialkoxy substituents. For both compounds **8** and **19**,

replacing 6,7-dimethoxy groups with ethoxy groups (**6** and **7**, respectively) leads, in each case, to a small (2-fold) decrease in EGF-R inhibitory activity. This is in contrast to what was reported in the quinazoline series, where replacing the 6,7-dimethoxy groups with 6,7-diethoxy groups led to a gain in potency.²⁴ Surprisingly, the 7-ethoxy-6-methoxy and 6-ethoxy-7-methoxy derivatives **48** and **49**, respectively, are both very much less potent than **8**; we can offer no explanation for this result. The 6,7-dihydroxy derivative **55** has comparable activity to **8**. Incorporation of the 6,7-alkoxy substituents into a fused ring as in **50**, **58**, and **59** results in significant loss of activity. Likewise, placing polyether chains at the 6,7-positions as in **56** and **57** decreases potency compared to **8**.

The compounds were also evaluated for their ability to inhibit the growth of certain cell lines. Three human carcinoma cell lines were used: A431 (epidermoid) which highly overexpresses EGF-R, SKBR3 (breast) which highly overexpresses HER-2 and, to a lesser extent, EGF-R, and SW620 (colon) which serves as a control line expressing low levels of EGF-R and HER-2. It is immediately obvious from the data shown in Table 1 that inhibition of cell proliferation exhibited by these compounds is likely the result of multiple mechanisms; cell lines such as SW620 that have little dependence on EGF-R or HER-2 are inhibited. It is possible that this result reflects a lack of selectivity and that these compounds inhibit other kinases. In fact, we know that the selectivity of these compounds for a particular kinase can be modulated by changing the substitution pattern on the aniline ring; this will be the subject of future communications. While multiple kinase activities would make a straightforward interpretation of these results difficult, some trends are evident. There is an approximate correlation between IC₅₀ values observed for growth inhibition of the A431 and SKBR3 lines. In addition, for the majority of the compounds, we find that they are better inhibitors of proliferation of the A431 and SKBR3 lines than the control SW620 line suggesting that, at least in part, inhibition of proliferation of these lines is the result of EGF-R or HER-2 kinase inhibition. In comparing the quinoline-3-carbonitrile **8** with its quinazoline counterpart **63**, we find that while **8** is about twice as potent as **63** in inhibiting the growth of both A431 and SKBR3 lines, it is also a more potent inhibitor of the SW620 control line. Some of the most potent inhibitors of cell growth are compounds that contain an ethoxy group at the 7-position such as **6**, **7**, and **49**, but these are also some of the least selective inhibitors since they inhibit the control line at comparable levels suggesting that with these compounds, a major component of the inhibition is unrelated to the EGF-R or HER-2 kinase activity.

Conclusions

The above results demonstrate that 4-anilino-6,7-dialkoxyquinoline-3-carbonitriles are effective inhibitors of EGF-R kinase. The SAR profile exhibited by these compounds is, in a number of respects, similar to that shown by the quinazoline series of EGF-R inhibitors. In addition, we constructed a homology model of EGF-R with which we proposed a binding model for the quinazoline- and quinoline-3-carbonitrile-based inhibi-

tors that is consistent with much of the observed activities. In future reports, we will attempt to show the versatility of this new series of tyrosine kinase inhibitors.

Of additional significance is our proposal that a carbon atom bearing a cyano group can sometimes be bioisosteric with an azomethine group that is hydrogen-bonded to a water molecule. Potentially, this concept may have utility in other areas of drug design.

Experimental Section

Biology. Preparation and purification of EGF-R DNA

constructs: A 1.6-kb cDNA for the EGF-R cytoplasmic domain (amino acids 45–1186) was cloned in baculoviral expression vectors pBlueBacHis2B (Invitrogen, Carlsbad, CA) and pFASTBacHTc (GIBCO, Rockville, MD). A sequence that encodes (His)₆ is located 5' upstream to the EGF-R sequence. Sf-9 cells were infected at moi = 10 for 3 days for protein expression. Sf-9 cell pellets were solubilized at 0 °C in a buffer at pH 7.4 containing 50 mM HEPES, 10 mM NaCl, 1% Triton, 10 μ M ammonium molybdate, 100 μ M sodium vanadate, 10 μ g/mL aprotinin, 10 μ g/mL leupeptin, 10 μ g/mL pepstatin, and 16 μ g/mL benzamidinium HCl for 20 min followed by 20-min centrifugation. Crude extract supernatant was passed through an equilibrated Ni-NTA superflow packed column (Qiagen, Valencia, CA) and washed with 10 and 100 mM imidazole to remove nonspecifically bound material. Histidine-tagged proteins were eluted with 250 and 500 mM imidazole and dialyzed against 50 mM NaCl, 20 mM HEPES, 10% glycerol and 1 μ g/mL each of aprotinin, leupeptin and pepstatin for 2 h. The entire purification procedure was performed at 4 °C or on ice. Purified materials were subjected to electrophoresis followed by Coomassie blue stain to determine the purity of the preparation and Western blot for the identity of protein. Enzyme preparations were stored at –80 °C.

EGF-R kinase autophosphorylation assay by DELFIA/

time-resolved fluorometry: The EGF-R kinase assay was set up to assess the level of autophosphorylation based on DELFIA/time-resolved fluorometry. Compounds were dissolved in 100% DMSO and diluted to the appropriate concentrations with 25 mM HEPES at pH 7.4. In each well, 10 μ L of compound was incubated with 10 μ L of recombinant enzyme (1:80 dilution in 100 mM HEPES) for 10 min at room temperature (>90% EGF-R binds to maxisorp plate within 1 h of incubation). Then, 10 μ L of 5x buffer (containing 20 mM HEPES, 2 mM MnCl₂, 100 μ M Na₃VO₄ and 1mM DTT) and 20 μ L of 0.1 mM ATP-50 mM MgCl₂ was added and the mixture was incubated for 1 h. Positive and negative controls were included in each plate by incubation of enzyme with or without ATP-MgCl₂. At the end of incubation, liquid was aspirated and plates were washed three times with wash buffer. A 75- μ L (400 ng) sample of europium-labeled anti-phosphotyrosine antibody was added to each well for another 1 h of incubation. After washing, enhancement solution was added and the signal was detected by Victor (Wallac Inc.) with excitation at 340 nm and emission at 615 nm. All reagents were supplied by Wallac Inc. (Wallac/Perkin-Elmer, Gaithersburg, MD). The percentage of autophosphorylation inhibition by the compound was calculated using the equation: $100 - [(test - negative\ control)/(positive\ control - negative\ control)]$. The IC₅₀ was obtained from curves of percentage inhibition with 8 concentrations of compound. As the purity of EGF-R enzyme preparation is greater than 80%, the majority of the signal detected by the anti-phosphotyrosine antibody is from EGF-R. The IC₅₀ values reported in Table 1 are averages of duplicate determinations.

Cell proliferation assay: The cell proliferation assays were done with three human carcinoma cell lines: A431 (epidermoid carcinoma), SKBR3 (breast carcinoma), and SW620 (colon carcinoma), as previously described.¹⁹ All cell lines were obtained from the American Type Culture Collection. Cells were plated in 96-well plates at densities of 5.0×10^4 /mL in RPMI-1640 medium supplemented with 5% fetal bovine

serum. On the next day, compounds were dosed at 0.5, 5, 50, 500, and 5000 ng/mL concentrations and the cells were cultured for 2 days. At the end of incubation, cell survival was determined by the sulforhodamine B assay as previously described.³⁴ The IC₅₀ values were obtained from the growth curves.

Molecular Modeling. Semiempirical calculations (AM1 basis) were carried out on the quinazoline and quinoline-3-carbonitrile compounds. In the quinazoline case, the bound molecule along with the water molecule hydrogen-bonded to the N3 atom were extracted from the modeled complex, and a single-point calculation was carried out. In the quinoline-3-carbonitrile case, only the bound compound was extracted and used for the single-point calculation. The electrostatic potential was mapped onto the surface (electron density contoured at the 0.005 level) and color-coded with red representing negative potential and blue representing positive potential. These calculations were done using MacSpartan Plus (Wavefunction Inc., 18401 Von Karman, Suite 370, Irvine, CA 92612).

Homology modeling was done using the MOE (Molecular Operating Environment) software (Chemical Computing Group Inc., 1255 University St., Suite 1600, Montreal, Quebec, Canada H3B 3X). Initial crude minimization was done within the homology modeling function of MOE. However, the resulting model needed further energy minimization. This was done using Quanta/CHARMM (Molecular Simulations Inc., 9685 Scranton Rd., San Diego, CA 92121). The crude model from MOE was minimized with a few thousand cycles of minimization using the ABNR (adopted-basis Newton–Raphson) method.

Ligands were modeled by positioning them in the active site in accordance with the published crystal structures of quinazoline derivatives bound to CDK2 and MAP kinase (p38).³¹ The entire complex was then subjected to alternate cycles of minimization and dynamics. Each dynamics run was short, about 3 ps. The intent was to get a satisfactory structure for the complex that was consistent with the published crystal structure.

Chemistry. ¹H NMR spectra were determined with a NT-300 WB spectrometer at 300 MHz. Chemical shifts (δ) are expressed in parts per million relative to the internal standard tetramethylsilane. Electrospray mass spectra were recorded in positive mode on a Micromass Platform spectrometer. Electron impact and high-resolution mass spectra were obtained on a Finnigan MAT-90 spectrometer. Chromatographic purifications were by flash chromatography using Baker 40- μ m silica gel. Melting points were determined in open capillary tubes on a Meltemp melting point apparatus and are uncorrected.

2-Cyano-3-(3,4-dimethoxyphenylamino)acrylic Acid Ethyl Ester (2c). A mixture of 3,4-dimethoxyaniline (30.6 g, 200 mmol), ethyl (ethoxymethylene)cyanoacetate (33.8 g, 200 mmol), and 80 mL of toluene was stirred at 100 °C for 1 h and at 125 °C for 15 min. After evaporation of the toluene, the residue was recrystallized from EtOAc to give 40.0 g (72%) of a tan solid: mp 166–170 °C; MS (ES+) m/z 277.2 (M + H)⁺. Anal. (C₁₄H₁₆N₂O₄) C, H, N.

6,7-Dimethoxy-4-oxo-1,4-dihydroquinoline-3-carbonitrile (3c). A stirred mixture of **2c** (40 g, 145 mmol) and 1.2 L of Dowtherm A was heated at reflux under N₂ for 10 h. After cooling to 50 °C the mixture was diluted with hexane. The product was filtered off, washed with hexane followed by methylene chloride, and dried to give 21.1 g (63%) of a brown solid: mp 330–350 °C dec; ¹H NMR (DMSO-*d*₆) δ 12.57 (s, 1H), 8.59 (s, 1H), 7.44 (s, 1H), 7.03 (s, 1H), 3.89 (s, 3H), 3.87 (s, 3H); MS (ES+) m/z 231.0 (M + H)⁺. Anal. (C₁₂H₁₀N₂O₃) C, H, N.

4-Chloro-6,7-dimethoxyquinoline-3-carbonitrile (4c). A stirred mixture of **3c** (20 g, 87 mmol) and 87 mL of POCl₃ was heated at reflux for 2 h. Volatile materials were removed under vacuum at about 70 °C. The residue was stirred at 0 °C with methylene chloride and H₂O as solid K₂CO₃ was carefully added until the pH was 8–9. After stirring for 30 min at 25 °C the organic layer was separated, washed with H₂O, dried, filtered through Celite, and concentrated to give 19.8 g (92%)

of an off-white solid. A sample recrystallized from CH₂Cl₂ gave an off-white solid: mp 220–223 °C; ¹H NMR (DMSO-*d*₆) δ 8.98 (s, 1H), 7.54 (s, 1H), 7.42 (s, 1H), 4.02 (s, 3H), 4.01 (s, 3H).

4-(3-Bromophenylamino)-6,7-dimethoxyquinoline-3-carboxylic Acid Ethyl Ester (5). A stirred mixture of **4a**²⁶ (14.8 g, 50 mmol), 3-bromoaniline (9.46 g, 55 mmol), pyridine (4.05 mL, 50 mmol), and 150 mL of EtOH was heated at reflux for 30 min. The EtOH was evaporated, and the residue was partitioned between CH₂Cl₂ and aqueous NaHCO₃. The organic layer was washed with water, dried over MgSO₄, and concentrated. The product was recrystallized from EtOH to give 15.4 g (71%) of a white solid: mp 155–158 °C; ¹H NMR (DMSO-*d*₆) δ 9.44 (s, 1H), 8.82 (s, 1H), 7.37 (s, 1H), 7.30 (s, 1H), 7.25–6.90 (m, 5H), 4.03 (q, J = 7.0 Hz, 2H), 3.95 (s, 3H), 3.71 (s, 3H), 1.16 (t, J = 7.0 Hz, 2H); MS (ES+) m/z 431.2, 433.2 (M + H)⁺. Anal. (C₂₀H₁₉BrN₂O₄) C, H, N.

4-(3-Bromophenylamino)-6,7-diethoxyquinoline-3-carbonitrile (6). To a stirred mixture of **5** (5.34 g, 15 mmol), K₂CO₃ (8.29 g, 60 mmol), and 60 mL of DMF at 0 °C was added 4.8 mL (60 mmol) of iodoethane. The resulting mixture was warmed to 25 °C and stirred for 4.5 h. The reaction mixture was partitioned between EtOAc and H₂O as 4N HCl was added slowly with cooling until the pH reached about 8. The organic layer was separated, washed well with H₂O, dried and concentrated. The residue was subjected to flash chromatography on silica gel with 60:30:1 CH₂Cl₂–EtOAc–TEA to give the product (3.27 g, 53%); homogeneous on TLC with the above solvent system. Recrystallization from EtOAc gave a white solid: mp 173–175 °C; ¹H NMR (DMSO-*d*₆) δ 9.50 (s, 1H), 8.54 (s, 1H), 7.67 (s, 1H), 7.35 (s, 1H), 7.55–7.15 (m, 4H), 4.35–4.10 (m, 4H), 1.45–1.35 (m, 6H); MS (ES+) m/z 412.3, 414.2 (M + H)⁺. Anal. (C₂₀H₁₈BrN₃O₂) C, H, N.

4-(3-Bromophenylamino)-6,7-diethoxyquinoline-3-carbonitrile (8). A stirred mixture of **4c** (0.40 g, 1.61 mmol), 3-bromoaniline (0.28 g, 1.61 mmol), and 15 mL of 2-ethoxyethanol was heated at reflux for 3 h and cooled to 25 °C. The crude HCl salt was filtered off and washed with 2-propanol and Et₂O: yield 0.62 g (92%). The free base was obtained by partition between EtOAc and aqueous Na₂CO₃. The organic layer was dried (MgSO₄) and solvent was removed to give 0.58 g of crude product. Recrystallization (EtOAc–hexane) gave a white solid: mp 224–228 °C; ¹H NMR (DMSO-*d*₆) δ 9.55 (s, 1H), 8.55 (s, 1H), 7.69 (s, 1H), 7.41 (s, 1H), 7.45–7.15 (m, 4H), 3.96 (s, 3H), 3.92 (s, 3H); MS (ES+) m/z 384.1, 386.1 (M + H)⁺. Anal. (C₁₈H₁₄BrN₃O₂) C, H, N, Br.

4-(3-Chloro-4-fluorophenylamino)-6,7-diethoxyquinoline-3-carbonitrile (7). The compound was prepared in 88% yield according to the procedure for **8** from 4-chloro-6,7-diethoxyquinoline-3-carbonitrile (**4b**) and 3-chloro-4-fluoroaniline: mp 194–198 °C (EtOAc); ¹H NMR (DMSO-*d*₆) δ 9.50 (s, 1H), 8.47 (s, 1H), 7.70 (s, 1H), 7.54–7.25 (m, 3H), 7.32 (s, 1H), 4.26–4.15 (m, 4H), 1.42 (t, 6H, J = 6.0 Hz); MS (ES+) m/z 386.2 (M + H)⁺. Anal. (C₂₀H₁₇ClF₂N₃O₂) C, H, N.

4-(3-Chlorophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (9). The compound was prepared in 55% yield according to the procedure for **8** using 3-chloroaniline: mp 214–217 °C (EtOAc–hexane); ¹H NMR (DMSO-*d*₆) δ 9.66 (s, 1H), 8.58 (s, 1H), 7.58 (s, 1H), 7.39 (s, 1H), 7.52–7.20 (m, 4H), 3.97 (s, 3H), 3.92 (s, 3H); MS (ES+) m/z 340.1, 342.2 (M + H)⁺. Anal. (C₁₈H₁₄ClN₃O₂) C, H, N.

6,7-Dimethoxy-4-(3-trifluoromethylphenylamino)-quinoline-3-carbonitrile (10). The compound was prepared in 72% yield according to the procedure for **8** using 3-trifluoromethylaniline: mp 190–193 °C (EtOAc–hexane); ¹H NMR (DMSO-*d*₆) δ 9.67 (s, 1H), 8.58 (s, 1H), 7.71 (s, 1H), 7.65–7.45 (m, 4H), 7.39 (s, 1H), 3.97 (s, 3H), 3.92 (s, 3H); MS (ES+) m/z 374.2 (M + H)⁺. Anal. (C₁₉H₁₄F₃N₃O₂) C, H, N.

4-(3,4-Dimethoxyphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (11). The compound was prepared in 66% yield according to the procedure for **8** using 3,4-dimethoxyaniline: mp 230–240 °C (EtOAc–hexane); ¹H NMR (DMSO-*d*₆) δ 9.45 (s, 1H), 8.36 (s, 1H), 7.98 (s, 1H), 7.30 (s, 1H), 7.00–6.83 (m, 3H), 3.94 (s, 3H), 3.92 (s, 3H), 3.78 (s, 3H), 3.76 (s, 3H); MS (ES+) m/z 366.3 (M + H)⁺. Anal. (C₂₀H₁₉N₃O₄) C, H, N.

4-(3-Fluorophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (12). The compound was prepared in 74% yield according to the procedure for **8** using 3-fluoroaniline: mp 226–230 °C (EtOAc); ¹H NMR (DMSO-*d*₆) δ 9.57 (s, 1H), 8.54 (s, 1H), 7.70 (s, 1H), 7.37 (s, 1H), 7.45–6.95 (m, 4H), 3.96 (s, 3H), 3.92 (s, 3H); MS (ES+) *m/z* 324.3 (M + H)⁺. Anal. (C₁₈H₁₄FN₃O₂) C, H, N.

4-(3-Cyanophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (13). The compound was prepared in 89% yield according to the procedure for **8** using 3-cyanoaniline: mp 285–288 °C; ¹H NMR (DMSO-*d*₆) δ 9.66 (s, 1H), 8.58 (s, 1H), 7.58 (s, 1H), 7.75–7.50 (m, 4H), 7.39 (s, 1H), 3.97 (s, 3H), 3.92 (s, 3H); MS (ES+) *m/z* 331.3 (M + H)⁺. Anal. (C₁₉H₁₄N₄O₂·0.5H₂O) C, H, N.

4-(4-Fluorophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (14). The compound was prepared in 91% yield according to the procedure for **8** using 4-fluoroaniline: mp 282–285 °C (EtOAc); ¹H NMR (DMSO-*d*₆) δ 9.49 (s, 1H), 8.42 (s, 1H), 7.76 (s, 1H), 7.40–7.23 (m, 4H), 7.32 (s, 1H), 3.95 (s, 3H), 3.92 (s, 3H); MS (ES+) *m/z* 324.3 (M + H)⁺. Anal. (C₁₈H₁₄FN₃O₂) C, H, N.

4-(3-Acetylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (15). The compound was prepared in 69% yield according to the procedure for **8** using 3-acetylaniline: mp 204–206 °C (acetone–hexane); ¹H NMR (DMSO-*d*₆) δ 9.62 (s, 1H), 8.53 (s, 1H), 7.77 (d, *J* = 1.7 Hz, 1H), 7.74 (s, 1H), 7.65–7.45 (m, 4H), 3.96 (s, 3H), 3.92 (s, 3H), 2.61 (s, 3H); MS (ES+) *m/z* 347.9 (M + H)⁺. Anal. (C₂₀H₁₇N₃O₃) C, H, N.

4-(3-Aminophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (16). A stirred mixture of **4c** (3.73 g, 15.0 mmol), 1,3-phenylenediamine (4.86 g, 45 mmol), pyridine (1.21 mL, 15 mmol), and 45 mL of 2-ethoxyethanol was heated at reflux for 30 min, cooled to 25 °C, and stirred with aqueous NaHCO₃. The resulting solid was filtered, washed with H₂O, and dried. Recrystallization from EtOH gave 4.32 g (90%) of a brown solid: mp 222–228 °C; ¹H NMR (DMSO-*d*₆) δ 9.28 (s, 1H), 8.43 (s, 1H), 7.74 (s, 1H), 7.31 (s, 1H), 6.99 (m, 1H), 6.43 (m, 3H), 5.21 (bs, 2H), 3.94 (s, 3H), 3.90 (s, 3H); MS (ES+) *m/z* 321.0 (M + H)⁺. Anal. (C₁₈H₁₆N₄O₂·0.5H₂O) C, H, N.

4-(3-Dimethylaminophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (17). The compound was prepared according to the procedure for **8** using *N,N*-dimethyl-1,3-phenylenediamine dihydrochloride. The product was recrystallized from water to give 0.4 g (18%) of a yellow solid: mp 246–249 °C; ¹H NMR (DMSO-*d*₆) δ 8.88 (s, 1H), 8.12 (s, 1H), 7.48 (s, 1H), 7.28 (m, 1H), 6.78 (m, 2H), 6.70 (d, 1H, *J* = 11.49), 3.99 (s, 3H), 3.98 (s, 3H), 2.94 (s, 6H). MS (ES+) *m/z* 349.2 (M + H)⁺, 174.9 (M + 2H)²⁺. Anal. (C₂₀H₂₀N₄O₂·HCl·0.75H₂O) C, H, N.

4-(3-Nitrophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (18). The compound was prepared according to the procedure for **8** using 3-nitroaniline. Recrystallization from EtOH gave a 69% yield of **18** as yellow crystals: mp 221–222 °C; ¹H NMR (DMSO-*d*₆) δ 3.92 (s, 3H), 3.98 (s, 3H), 7.42 (s, 1H), 7.65 (m, 3H), 7.96 (m, 2H), 8.64 (s, 1H), 9.79 (s, 1H); MS (ES+) *m/z* 351.0 (M + H)⁺. Anal. (C₁₈H₁₄N₄O₄) C, H, N.

4-(3-Chloro-4-fluorophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (19). The compound was prepared in 87% yield according to the procedure for **8** using 3-chloro-4-fluoroaniline: mp 223–225 °C; ¹H NMR (DMSO-*d*₆) δ 9.65 (s, 1H), 8.52 (s, 1H), 7.75 (s, 1H), 7.62–7.28 (m, 3H), 7.35 (s, 1H), 3.96 (s, 3H), 3.93 (s, 3H); MS (ES+) *m/z* 358.2 (M + H)⁺. Anal. (C₁₈H₁₃ClFN₃O₂) C, H, N.

4-(4-Chloro-2-fluorophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (20). Prepared in 52% yield according to the procedure for **8** using 4-chloro-2-fluoroaniline: mp 184–186 °C; ¹H NMR (DMSO-*d*₆) δ 9.49 (br s, 1H), 8.47 (s, 1H), 7.79 (s, 1H), 7.60 (dd, *J* = 10.7, 2.1 Hz, 1H), 7.48 (t, *J* = 8.7 Hz, 1H), 7.35 (m, 2H), 3.96 (s, 3H), 3.94 (s, 3H); MS (ES+) *m/z* 358 (M + H)⁺. Anal. (C₁₈H₁₃ClFN₃O₂) C, H, N.

4-(3-Hydroxyphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (21). Prepared (91% yield) according to the procedure for **8** using 3-hydroxyaniline: mp > 250 °C; ¹H NMR (DMSO-*d*₆) δ 3.90 (s, 3H), 3.95 (s, 3H), 6.58–6.66 (m, 3H), 7.17

(t, *J* = 8.25 Hz, 1H), 7.33 (s, 1H), 7.72 (s, 1H), 8.46 (s, 1H), 9.37 (s, 1H), 9.52 (s, 1H). Anal. (C₁₈H₁₅N₃O₃·0.2H₂O) C, H, N.

4-(4-Methylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (22). Prepared in 88% yield according to the procedure for **8** using 4-methylaniline: mp 128–130 °C; ¹H NMR (DMSO-*d*₆) δ 2.33 (s, 3H), 3.91 (s, 3H), 3.94 (s, 3H), 7.15 (d, *J* = 6.1, 2H Hz), 7.22 (d, *J* = 6.1 Hz, 2H), 7.32 (s, 1H), 7.76 (s, 1H), 8.41 (s, 1H), 9.40 (s, 1H). Anal. (C₁₉H₁₇N₃O₂·0.5H₂O) C, H, N.

3-(3-Cyano-6,7-dimethoxyquinolin-4-ylamino)benzamide (23). Prepared in 89% yield according to the procedure for **8** using 3-aminobenzamide: mp 253–255 °C; ¹H NMR (DMSO-*d*₆) δ 9.56 (br s, 1H), 8.51 (s, 1H), 8.02 (s, 1H), 7.76 (s, 2H), 7.71 (d, *J* = 7.8 Hz, 1H), 7.47 (t, *J* = 7.8 Hz, 1H), 7.41 (s, 1H), 7.38 (d, *J* = 7.8 Hz, 1H), 7.36 (s, 1H), 3.96 (s, 3H), 3.92 (s, 3H); MS (ES+) *m/z* 349 (M + H)⁺. Anal. (C₁₉H₁₆N₄O₃·0.3HCl) C, H, N.

4-(3-Bromo-4-methylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (24). Prepared in 72% yield according to the procedure for **8** using 3-bromo-4-methylaniline: mp 292–294 °C; ¹H NMR (DMSO-*d*₆) δ 11.05 (br s, 1H), 8.10 (s, 1H), 7.71 (d, *J* = 2.1 Hz, 1H), 7.48 (d, *J* = 8.3 Hz, 1H), 7.44 (s, 1H), 7.38 (dd, *J* = 8.3, 2.1 Hz, 1H), 4.00 (s, 3H), 3.98 (s, 3H), 2.41 (s, 3H); MS (ES+) *m/z* 398 (M + H)⁺. Anal. (C₁₉H₁₆BrN₃O₂·0.3H₂O) C, H, N.

4-(3-Chloro-4-hydroxyphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (25). Prepared in 65% yield according to the procedure for **8** using 4-amino-2-chlorophenol: mp 230–232 °C; ¹H NMR (DMSO-*d*₆) δ 10.29 (br s, 1H), 9.39 (br s, 1H), 8.37 (s, 1H), 7.76 (s, 1H), 7.31 (d, *J* = 2.4 Hz, 1H), 7.30 (s, 1H), 7.12 (dd, *J* = 8.4, 2.8 Hz, 1H), 6.99 (d, *J* = 8.4 Hz, 1H), 3.94 (s, 3H), 3.92 (s, 3H); MS (ES+) *m/z* 356 (M + H)⁺. Anal. (C₁₉H₁₆BrN₃O₂·0.4EtOAc) C, H, N.

4-(3,5-Dichloro-4-hydroxyphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (26). By the procedure as described above for **8** using 3,5-dichloro-4-hydroxyaniline, the title compound was obtained in 89% yield: mp > 250 °C; ¹H NMR (DMSO-*d*₆) δ 3.96 (s, 3H), 3.98 (s, 3H), 7.37 (s, 1H), 7.49 (s, 2H), 7.94 (s, 1H), 8.75 (s, 1H), 10.34 (bs, 1H), 10.40 (bs, 1H); MS (ES+) *m/z* 389.8, 391.8 (M + H)⁺. Anal. (C₁₈H₁₃Cl₂N₃O₃·1.0H₂O) C, H, N.

4-(5-Chloro-2-hydroxyphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile Hydrochloride (27). By the procedure as described above for **8**, using 5-chloro-2-hydroxyaniline, the title compound was obtained in 83% yield as the hydrochloride salt directly from the reaction mixture: mp > 250 °C; ¹H NMR (DMSO-*d*₆) δ 3.98 (s, 3H), 3.99 (s, 3H), 7.02 (d, *J* = 9 Hz, 1H), 7.33 (dd, *J* = 2.7, 8.7 Hz, 1H), 7.41 (s, 1H), 7.44 (d, *J* = 2.7 Hz, 1H), 8.07 (s, 1H), 8.90 (s, 1H), 10.51 (bs, 1H), 10.63 (bs, 1H); MS (ES+) *m/z* 355.8, 357.8 (M + H)⁺. Anal. (C₁₈H₁₄ClN₃O₃·0.9HCl) C, H, N.

6,7-Dimethoxy-4-(3-methylthiophenylamino)quinoline-3-carbonitrile Hydrochloride (28). By the procedure as described above for **8**, using 3-methylthioaniline, the title compound was obtained in 35% yield as the hydrochloride salt directly from the reaction mixture: mp > 250 °C; ¹H NMR (DMSO-*d*₆) δ 2.51 (s, 3H), 3.98 (s, 3H), 4.00 (s, 3H), 7.21 (d, *J* = 8.3 Hz, 1H), 7.29 (d, *J* = 8.1 Hz, 1H), 7.34 (s, 1H), 7.41 (dd, *J* = 7.0, 7.8 Hz, 1H), 7.45 (s, 1H), 8.10 (s, 1H), 8.96 (s, 1H), 11.10 (bs, 1H); MS (ES+) *m/z* 351.9 (M + H)⁺. Anal. (C₁₉H₁₇N₃O₂S·1HCl) C, H, N.

4-(3-Chloro-4-methylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile Hydrochloride (29). By the procedure as described above for **8**, using 3-chloro-4-methylaniline, the title compound was obtained in 90% yield as the hydrochloride salt directly from the reaction mixture: mp > 250 °C; ¹H NMR (DMSO-*d*₆) δ 2.39 (s, 3H), 4.00 (s, 3H), 4.02 (s, 3H), 7.34 (dd, *J* = 2.13, 8.1 Hz, 1H), 7.43 (s, 1H), 7.48 (d, *J* = 8.3 Hz, 1H), 7.56 (d, *J* = 2.0 Hz, 1H), 8.09 (s, 1H), 8.96 (s, 1H), 10.98 (bs, 1H); MS (ES+) *m/z* 353.9, 355.8 (M + H)⁺. Anal. (C₁₉H₁₆ClN₃O₂·1HCl) C, H, N.

4-(4-Bromo-2-fluorophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile Hydrochloride (30). By the procedure as described above for **8**, using 4-bromo-2-fluoroaniline, the

title compound was obtained in 52% yield as the hydrochloride salt directly from the reaction mixture: mp > 250 °C; ^1H NMR (DMSO- d_6) δ 4.00 (s, 6H), 7.46 (s, 1H), 7.58 (m, 2H), 7.83 (d, J = 9.21 Hz, 1H), 8.16 (s, 1H), 9.00 (s, 1H), 11.00 (bs, 1H); MS (ES+) m/z 410.8, 403.8 ($M + H$) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{13}\text{BrFN}_3\text{O}_2 \cdot 1\text{HCl} \cdot 0.2\text{H}_2\text{O}$) C, H, N.

4-(4-Isopropylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (31). Prepared in 79% yield according to the procedure for **8** using 4-isopropylaniline giving a pale yellow solid: mp 198–200 °C; ^1H NMR (DMSO- d_6) δ 9.41 (s, 1H), 8.42 (s, 1H), 7.72 (s, 1H), 7.32 (s, 1H), 7.27 (d, 2H, J = 8.4 Hz, 2H), 7.17 (d, 2H, J = 8.4 Hz, 2H), 3.94 (s, 3H), 3.89 (s, 3H), 2.50 (m, 1H), 1.21 (d, J = 6.9 Hz, 6H); HRMS m/z 348.1705 ($M + H$) $^{+1}$. Anal. ($\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_2 \cdot 0.2\text{H}_2\text{O}$) C, H, N.

4-(2-Isopropylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (32). Prepared in 60% yield according to the procedure for **8** using 2-isopropylaniline giving a gray solid: mp 90–91 °C; ^1H NMR (DMSO- d_6) δ 9.35 (s, 1H), 8.30 (s, 1H), 7.87 (s, 1H), 7.42 (m, 2H), 7.30 (s, 1H), 7.26 (m, 2H), 3.95 (s, 3H), 3.91 (s, 3H), 2.50 (m, 1H), 1.21 (d, J = 6.9 Hz, 6H); HRMS m/z 348.1705 ($M + H$) $^{+1}$. Anal. ($\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_2 \cdot 0.6\text{H}_2\text{O}$) C, H, N.

4-(3-Isopropylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (33). Prepared in 70% yield according to the procedure for **8** using 3-isopropylaniline giving a yellow solid: mp 160–162 °C; ^1H NMR (DMSO- d_6) δ 9.47 (s, 1H), 8.45 (s, 1H), 7.75 (s, 1H), 7.33 (s, 1H), 7.32 (t, J = 7.9 Hz, 1H), 7.11 (s, 1H), 7.08 (m, 2H), 3.93 (d, J = 13.3 Hz, 6H), 2.50 (m, 1H); HRMS m/z 348.1705 ($M + H$) $^{+1}$. Anal. ($\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_2 \cdot 0.2\text{H}_2\text{O}$) C, H, N.

4-(4-Bromophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (34). The compound was prepared according to the procedure for **8** using 4-bromoaniline in 47% yield, light yellow solid: mp 199–200 °C; ^1H NMR (DMSO- d_6) δ 9.49 (s, 1H), 8.51 (s, 1H), 7.71 (s, 1H), 7.56 (d, J = 9 Hz, 2H), 7.36 (s, 1H), 7.19 (d, J = 9 Hz, 2H), 3.95 (s, 3H), 3.91 (s, 3H); MS (ES+) m/z 383.8, 385.8 ($M + H$) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{14}\text{BrN}_3\text{O}_2 \cdot 0.5\text{H}_2\text{O}$) C, H, N.

4-(2-Bromophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (35). To a suspension of NaH (60% in mineral oil, 90 mg, 2.25 mmol) in 5 mL of DMF was added 2-bromoaniline (387 mg, 2.25 mmol). The mixture was heated to reflux then cooled and **4c** (250 mg, 1.05 mmol) was added. The mixture was heated at reflux for 40 min then cooled to room temperature and partitioned between water and ethyl acetate. The organic layer was washed with dilute aqueous ammonium hydroxide, dried over MgSO_4 , filtered and concentrated in vacuo. EtOAc and diethyl ether were added to the residue and the solid was collected to provide 168 mg (43%) of **35** as a tan solid: mp 217–219 °C; ^1H NMR (DMSO- d_6) δ 9.58 (s, 1H), 8.38 (s, 1H), 7.85 (s, 1H), 7.76 (d, J = 6 Hz, 1H), 7.58–7.42 (m, 2H), 7.36–7.32 (m, 2H), 3.95 (s, 3H), 3.93 (s, 3H); MS (ES+) m/z 383.8, 385.8 ($M + H$) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{14}\text{BrN}_3\text{O}_2$) C, H, N.

4-(4-Fluoro-3-trifluoromethylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile Hydrochloride (36). The compound was prepared as a hydrochloride salt in 72% yield according to the procedure for **8** using 4-fluoro-3-trifluoromethylaniline: mp 240–243 °C; ^1H NMR (DMSO- d_6) δ 10.21 (broad s, 1H), 8.69 (s, 1H), 7.88 (s, 1H), 7.79–7.69 (m, 2H), 7.64–7.57 (m, 1H), 7.39 (s, 1H), 3.98 (s, 3H), 3.96 (s, 3H); MS (ES+) m/z 391.9 ($M + H$) $^{+1}$. Anal. ($\text{C}_{19}\text{H}_{13}\text{F}_4\text{N}_3\text{O}_2 \cdot 0.6\text{HCl}$) C, H, N.

4-(3-Bromo-2-methylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (37). By the procedure as described above for **8** using 3-bromo-2-methylaniline, the title compound was obtained in 42% yield: mp 183–185 °C; ^1H NMR (DMSO- d_6) δ 2.28 (s, 3H), 3.93 (s, 3H), 3.94 (s, 3H), 7.21 (dd, J = 7.92, 7.89 Hz, 1H), 7.31 (d, J = 7.70, 1H), 7.33 (s, 1H), 7.63 (d, J = 7.62 Hz, 1H), 7.82 (s, 1H), 8.37 (s, 1H), 9.49 (bs, 1H); MS (ES+) m/z 397.8, 399.8 ($M + H$) $^{+1}$. Anal. ($\text{C}_{19}\text{H}_{16}\text{BrN}_3\text{O}_2 \cdot 0.21\text{C}_6\text{H}_{14}$) C, H, N.

4-(4-Bromo-3-methylphenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (38). By the procedure as described above for **8** using 4-bromo-3-methylaniline, the title compound was obtained in 61% yield: mp 168–169 °C; ^1H NMR (DMSO-

d_6) δ 2.35 (s, 3H), 3.91 (s, 3H), 3.95 (s, 3H), 7.01 (dd, J = 2.67, 8.52 Hz, 1H), 7.24 (d, J = 2.55, 1H), 7.36 (s, 1H), 7.55 (d, J = 8.49 Hz, 1H), 7.72 (s, 1H), 8.51 (s, 1H), 9.46 (bs, 1H); MS (ES+) m/z 397.8, 399.8 ($M + H$) $^{+1}$. Anal. ($\text{C}_{19}\text{H}_{16}\text{BrN}_3\text{O}_2 \cdot 0.26\text{C}_6\text{H}_{14} \cdot 0.1\text{H}_2\text{O}$) C, H, N.

4-(3-Azidophenylamino)-6,7-dimethoxyquinoline-3-carbonitrile (39). Compound **16** (0.643 g, 2.00 mmol) was dissolved in 25 mL of 80% aqueous acetic acid and chilled to 0 °C. A solution of 0.152 g (2.21 mmol) of NaNO_2 in 2.2 mL of H_2O was added, followed 10 min later by a solution of 0.144 g (2.21 mmol) of NaN_3 in 2.2 mL of H_2O . Volatile material was removed after the reaction had stirred at 25 °C for 1.5 h. The crude product was dissolved in EtOAc and filtered and the filtrate was treated with saturated NaHCO_3 . The organic layer was separated, washed with brine, dried (Na_2SO_4) and evaporated. The residue was filtered through silica gel (40% CH_2Cl_2 in EtOAc). The eluent was evaporated and the product was dried in vacuo (50 °C) to give 0.526 g (76%) of product as brown crystals: ^1H NMR (DMSO- d_6) δ 3.91 (s, 3H), 3.96 (s, 3H), 6.98 (m, 3H), 7.43 (m, 2H), 7.69 (s, 1H), 8.54 (s, 1H), 9.55 (s, 1H); HRMS (CI) m/z ($M + H$) $^{+1}$ calcd for $\text{C}_{18}\text{H}_{15}\text{N}_6\text{O}_2$ 347.1256, found 347.1255. Anal. ($\text{C}_{18}\text{H}_{14}\text{N}_6\text{O}_2$) C, H, N.

N-[3-(3-Cyano-6,7-dimethoxyquinolin-4-ylamino)phenyl]acetamide (40). A solution of **16** (0.96 g, 3.0 mmol) in 9 mL of HOAc was treated with 0.85 mL of Ac_2O at 25 °C. After 2 h the solution was evaporated to dryness at 40 °C, stirred in MeOH, and evaporated to dryness. The residue was recrystallized from EtOH to give 0.50 g (46%) of a brown solid: mp 147–150 °C; ^1H NMR (DMSO- d_6) δ 10.00 (s, 1H), 9.48 (s, 1H), 8.48 (s, 1H), 7.75 (s, 1H), 7.66 (s, 1H), 7.34 (s, 1H), 7.29 (m, 2H), 6.89 (m, 1H), 3.96 (s, 3H), 3.91 (s, 3H), 2.05 (s, 3H); MS (ES+) m/z 363.2 ($M + H$) $^{+1}$. Anal. ($\text{C}_{20}\text{H}_{18}\text{N}_4\text{O}_3 \cdot 1.5\text{H}_2\text{O}$) C, H, N.

3-Ethoxy-4-methoxybenzoic Acid Methyl Ester (42a). A mixture of 24.3 g (134 mmol) of 3-hydroxy-4-methoxybenzoic acid methyl ester, 36.8 g (267 mmol) of K_2CO_3 and 16 mL (31.4 g, 201 mmol) of $\text{C}_2\text{H}_5\text{I}$ in 500 mL of DMF was stirred at 100 °C for 6 h. An additional 16 mL (31.4 g, 201 mmol) of $\text{C}_2\text{H}_5\text{I}$ and 18.4 g (133 mmol) of K_2CO_3 was added and the reaction was heated for 3 h more. The reaction was then filtered, the insoluble material was washed with DMF and the combined filtrate and wash were evaporated. The residue was mixed with H_2O and filtered and the insoluble material was washed with H_2O and dried. Recrystallization from heptane gave 22.5 g (80%) of product as white crystals: mp 81–83 °C; ^1H NMR (DMSO- d_6) δ 1.34 (t, J = 6.9 Hz, 3H), 3.81 (s, 3H), 3.84 (s, 3H), 4.04 (q, J = 6.9 Hz, 2H), 7.07 (d, J = 8.5 Hz, 1H), 7.42 (d, J = 2.0, 1H), 7.58 (dd, J = 8.5, J = 2.0 Hz, 1H); MS (ES+) m/z 210.9 ($M + H$) $^{+1}$. Anal. ($\text{C}_{11}\text{H}_{14}\text{O}_4$) C, H.

4-Ethoxy-3-methoxybenzoic Acid Methyl Ester (42b). Using the procedure described above for **42a** and a total reaction time of 2 h gave, after recrystallization from hexane, 82% of product as white crystals: ^1H NMR (DMSO- d_6) δ 1.35 (t, 3H, J = 7.0), 3.81 (s, 3H), 3.82 (s, 3H), 4.09 (q, J = 7.0 Hz, 2H), 7.05 (d, J = 8.5 Hz, 1H), 7.44 (d, J = 1.95 Hz, 1H), 7.56 (dd, J = 8.5, 1.95 Hz, 1H); MS (EI) m/z 210 (M) $^{+1}$.

5-Ethoxy-4-methoxy-2-nitrobenzoic Acid Methyl Ester (43a). Concentrated HNO_3 (15 mL) was slowly added to a slurry of 15.0 g (71.4 mmol) of **42a** in 45 mL of glacial acetic acid. After an exotherm had subsided, the reaction was reheated to 50–60 °C for 45 min. It was then poured into ice H_2O and extracted with CH_2Cl_2 . The extracts were washed with H_2O and 0.5 M NaOH, dried (MgSO_4) and evaporated to give 17.8 g (98% crude yield) of product as yellow crystals. An analytical sample was obtained from CH_2Cl_2 –hexane: mp 85–88 °C; ^1H NMR (DMSO- d_6) δ 1.36 (t, J = 7.0 Hz, 3H), 3.83 (s, 3H), 3.91 (s, 3H), 4.18 (q, 2H, J = 7.0 Hz, 2H), 7.30 (s, 1H), 7.63 (s, 1H); MS (ES+) m/z 256.0 ($M + H$) $^{+1}$. Anal. ($\text{C}_{11}\text{H}_{13}\text{NO}_6$) C, H, N.

4-Ethoxy-5-methoxy-2-nitrobenzoic Acid Methyl Ester (43b). The compound was prepared from **42b** using the procedure described above for **43a** and a reaction time of 1 h. A crude yield of 87% of the product was obtained. A purified sample (light yellow crystals) was obtained by recrystallization

from heptane: ^1H NMR (DMSO- d_6) δ 1.36 (t, J = 7.02 Hz, 3H), 3.83 (s, 3H), 3.92 (s, 3H), 4.18 (q, J = 7.02 Hz, 2H), 7.32 (s, 1H), 7.61 (s, 1H); MS (ES+) m/z 255.8 (M + H) $^{+1}$.

7-Nitro-2,3-dihydrobenzo[1,4]dioxine-6-carboxylic Acid Methyl Ester (43c). This compound was prepared from **42c**²⁷ using the procedure described for **43a** and a reaction time of 3 h at 70 °C. Recrystallization from heptane–toluene gave 91% of product as yellow crystals: ^1H NMR (DMSO- d_6) δ 3.80 (s, 3H), 4.39 (s, 4H), 7.32 (s, 1H), 7.66 (s, 1H); MS (EI) m/z 239 (M) $^{+1}$.

2-Amino-5-ethoxy-4-methoxybenzoic Acid Methyl Ester (44a). A mixture of 17.0 g (66.7 mmol) of **43a**, 13.1 g (233 mmol) of powdered Fe and 17.7 g (334 mmol) of NH_4Cl in 95 mL of H_2O and 245 mL of MeOH was heated at reflux for 4.5 h. More Fe (13.1 g, 233 mmol) was added and the reaction was heated for 2.5 h. Additional amounts of Fe (13.1 g, 233 mmol) and NH_4Cl (17.7 g, 334 mmol) were then added and heating was continued for 12 h more. The reaction was filtered and the insoluble material was washed with H_2O and MeOH. The combined filtrate and wash was evaporated and the residue was partitioned between CHCl_3 and H_2O . The organic material was treated with activated carbon, dried (MgSO_4), evaporated and dried in vacuo (50 °C) to give 11.0 g (73% crude yield) of product as tan crystals. An analytical sample was prepared by recrystallization from EtOH: mp 120–122 °C; ^1H NMR (DMSO- d_6) δ 1.26 (t, J = 5.1 Hz, 3H), 3.74 (s, 6H), 3.85 (q, 2H, J = 5.1 Hz), 6.36 (s, 1H), 6.44 (s, 2H), 7.14 (s, 1H); MS (ES+) m/z 225.9 (M + H) $^{+1}$. Anal. ($\text{C}_{11}\text{H}_{15}\text{NO}_4$) C, H, N.

2-Amino-4-ethoxy-5-methoxybenzoic Acid Methyl Ester (44b). This compound was prepared from **43b** in a manner similar to **44a**. Additional powdered Fe and NH_4Cl were added as necessary to drive the reaction to completion. The crude yield was 83% as pink crystals. A purified sample was prepared by recrystallization from Et $_2$ O: ^1H NMR (DMSO- d_6) δ 1.34 (t, J = 6.99 Hz, 3H), 3.64 (s, 3H), 3.74 (s, 3H), 3.98 (q, 2H, J = 6.99 Hz), 6.35 (s, 1H), 6.42 (s, 2H), 7.12 (s, 1H); MS (ES+) m/z 226.2 (M + H) $^{+1}$.

7-Amino-2,3-dihydrobenzo[1,4]dioxine-6-carboxylic Acid Methyl Ester (44c). This compound was prepared from **43c** in a manner similar to **44a**. Additional powdered Fe and NH_4Cl were added to complete the reaction. Filtration of the crude product through silica (CHCl_3) helped to remove colored impurities. Solids washed with MeOH to give a 90% yield of product as tan crystals: ^1H NMR (DMSO- d_6) δ 3.75 (s, 3H), 4.13 (m, 2H), 4.23 (m, 2H), 6.24 (s, 3H), 7.14 (s, 1H); MS (ES+) m/z 209.9 (M + H) $^{+1}$.

6-Ethoxy-7-methoxy-4-oxo-1,4-dihydroquinoline-3-carbonitrile (46a). A solution of 10.2 g (45.3 mmol) of **44a** and 12.6 mL (10.8 g, 90.7 mmol) of dimethylformamide dimethylacetal in 50 mL of DMF was heated at reflux under N_2 for 2.3 h. Volatile material was removed and the residue was azeotroped twice with toluene and dried in vacuo (25 °C). The formamidine product **45a** was a purple syrup that crystallized on standing.

A solution of 5.32 mL (4.18 g, 102 mmol) of CH_3CN in 80 mL of THF was added over 15 min to a solution of 40 mL (100 mmol) of 2.5 M *n*-BuLi in hexane and 60 mL of THF at –78 °C under N_2 . After 20 min, a solution of the crude **45a** in 80 mL of THF was added over 0.5 h. The reaction was stirred at –78 °C for 1 h and then quenched with 13 mL (13.7 g, 227 mmol) of glacial acetic acid. The mixture was warmed to 25 °C and volatile material was removed. The residue was slurried with H_2O , collected and dried. This material was washed twice with CHCl_3 to give 7.95 g of product as yellow crystals. An analytical sample (cream crystals) was obtained by boiling a portion of the crude product in MeOH and drying in vacuo (50 °C): mp > 310 °C; ^1H NMR (DMSO- d_6) δ 1.38 (t, J = 6.96 Hz, 3H), 3.89 (s, 3H), 4.11 (q, J = 6.96 Hz, 2H), 7.04 (s, 1H), 7.43 (s, 1H), 8.59 (s, 1H); MS (ES+) m/z 243.2 (M + H) $^{+1}$. Anal. ($\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_3$) C, H, N.

7-Ethoxy-6-methoxy-4-oxo-1,4-dihydroquinoline-3-carbonitrile (46b). Starting with **44b** and using the procedure described for **46a** a crude yield of 83% (yellow solid) of the title compound was obtained: ^1H NMR (DMSO- d_6) δ 1.40 (t,

J = 6.96 Hz, 3H), 3.86 (s, 3H), 4.12 (q, J = 6.96 Hz, 2H), 7.03 (s, 1H), 7.45 (s, 1H), 8.59 (s, 1H); MS (ES+) m/z 245.2 (M + H) $^{+1}$.

9-Oxo-2,3,6,9-tetrahydro[1,4]dioxino[2,3-*g*]quinoline-8-carbonitrile (46c). Starting with **44c** and using the procedure described for **46a** a crude yield of 72% (tan powder) of the title compound was obtained: ^1H NMR (DMSO- d_6) δ 4.34 (m, 2H), 4.37 (m, 2H), 7.05 (s, 1H), 7.46 (s, 1H), 8.58 (s, 1H); MS (ES+) m/z 228.8 (M + H) $^{+1}$.

4-Chloro-6-ethoxy-7-methoxyquinoline-3-carbonitrile (47a). A slurry of 12.9 g (52.7 mmol) of **46b** in 200 mL of CH_2Cl_2 under N_2 was treated with 23.1 mL (33.5 g, 263 mmol) of oxalyl chloride and 1 mL of DMF for 2 days. The reaction was then diluted with CHCl_3 and mixed with saturated aqueous NaHCO_3 . The organic layer was separated, dried (MgSO_4), evaporated and dried in vacuo. The crude yield was 86% of a gray solid: ^1H NMR (DMSO- d_6) δ 1.43 (t, 3H, J = 6.99 Hz), 4.02 (s, 3H), 4.27 (q, 2H, J = 6.99 Hz), 7.43 (s, 1H), 7.55 (s, 1H), 8.99 (s, 1H); MS (ES+) m/z 262.8, 264.8 (M + H) $^{+1}$.

4-Chloro-7-ethoxy-6-methoxyquinoline-3-carbonitrile (47b). This compound was prepared from **46b** and oxalyl chloride as previously described. The crude yield was 86% of a gray solid: ^1H NMR (DMSO- d_6) δ 1.44 (t, 3H, J = 6.96 Hz), 4.02 (s, 3H), 4.30 (q, 2H, J = 6.96 Hz), 7.46 (s, 1H), 7.54 (s, 1H), 8.99 (s, 1H); MS (ES+) m/z 262.8, 264.8 (M + H) $^{+1}$.

9-Chloro-2,3-dihydro[1,4]dioxino[2,3-*g*]quinoline-8-carbonitrile (47c). This compound was prepared from **46c** and POCl_3 as previously described for **4c**. The crude yield was 97% of a tan solid: ^1H NMR (DMSO- d_6) δ 4.48 (m, 4H), 7.55 (s, 1H), 7.57 (s, 1H), 8.94 (s, 1H); MS (ES+) m/z 246.8, 248.8 (M + H) $^{+1}$.

4-(3-Bromophenylamino)-6-ethoxy-7-methoxyquinoline-3-carbonitrile (48). This compound was prepared from **47a** and 3-bromoaniline as previously described for **8**. Recrystallization from EtOH gave an 86% yield of tan crystals: mp 190–191.5 °C; ^1H NMR (DMSO- d_6) δ 1.40 (t, 3H, J = 6.99 Hz), 3.96 (s, 3H), 4.16 (q, 2H, J = 6.99 Hz), 7.22 (m, 1H), 7.34 (m, 3H), 7.40 (s, 1H), 7.67 (s, 1H), 8.54 (s, 1H), 9.50 (s, 1H); HRMS (EI) m/z calcd for $\text{C}_{19}\text{H}_{16}\text{BrN}_3\text{O}_2$ 397.0426, found 397.0429. Anal. ($\text{C}_{19}\text{H}_{16}\text{BrN}_3\text{O}_2 \cdot \text{EtOH}$) C, H, N.

4-(3-Bromophenylamino)-7-ethoxy-6-methoxyquinoline-3-carbonitrile (49). This compound was prepared from **47b** and 3-bromoaniline as previously described for **8**. The crude yield was 73% of a tan solid: ^1H NMR (DMSO- d_6) δ 1.42 (t, J = 6.93 Hz, 3H), 3.91 (s, 3H), 4.25 (q, J = 6.93 Hz, 2H), 7.22 (m, 1H), 7.33 (m, 3H), 7.40 (s, 1H), 7.68 (s, 1H), 8.54 (s, 1H), 9.54 (s, 1H); HRMS (EI) m/z calcd for $\text{C}_{19}\text{H}_{16}\text{BrN}_3\text{O}_2$ 397.0426, found 397.0436. Anal. ($\text{C}_{19}\text{H}_{16}\text{BrN}_3\text{O}_2 \cdot 0.5\text{H}_2\text{O}$) C, H, N.

9-(3-Bromophenylamino)-2,3-dihydro[1,4]dioxino[2,3-*g*]quinoline-8-carbonitrile (50). This compound was prepared from **47c** and 3-bromoaniline as previously described for **8**. Recrystallization from EtOAc gave a 65% yield of tan crystals: ^1H NMR (DMSO- d_6) δ 4.42 (m, 4H), 7.21 (m, 1H), 7.33 (m, 3H), 7.41 (s, 1H), 7.86 (s, 1H), 8.50 (s, 1H), 9.57 (s, 1H); HRMS (FAB+) m/z (M + H) $^{+1}$ calcd for $\text{C}_{18}\text{H}_{13}\text{BrN}_3\text{O}_2$ 382.0191, found 382.0215. Anal. ($\text{C}_{18}\text{H}_{12}\text{BrN}_3\text{O}_2 \cdot 0.3\text{H}_2\text{O}$) C, H, N.

[4-(3-Bromophenylamino)-6,7-dimethoxyquinolin-3-yl]-methanol (51). To a stirred mixture of **5** (1.45 g, 3.36 mmol) and 67 mL of toluene at 0 °C was added 1.53 M diisobutylaluminum hydride in toluene (11 mL, 16.8 mmol) during 10 min. After an additional 10 min, the solution was treated with 7 mL of MeOH at 0 °C then stirred at 25 °C for 10 min. The resulting gel was diluted with THF and stirred with $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ for 30 min. Solids were removed by filtration, and the filtrate was concentrated. The residue was recrystallized from CH_2Cl_2 –hexane to give 0.56 g (43%) of a white solid: mp 176–180 °C; ^1H NMR (DMSO- d_6) δ 8.76 (s, 1H), 8.44 (s, 1H), 7.38 (s, 1H), 7.17 (s, 1H), 7.15–6.60 (m, 4H), 5.31 (t, J = 5.3 Hz, 1H), 4.49 (d, J = 5.3 Hz, 2H), 3.93 (s, 3H), 3.73 (s, 3H); MS (ES+) m/z 389.1, 391.1 (M + H) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{17}\text{BrN}_2\text{O}_3$) C, H, N.

4-(3-Bromophenylamino)-6,7-dimethoxyquinoline-3-

carbaldehyde (52). To a stirred solution of **51** (3.20 g, 8.22 mmol) in 82 mL of CH_2Cl_2 were added NaOAc (0.20 g, 2.5 mmol) and pyridinium chlorochromate (2.65 g, 12.3 mmol). After 2 h at 25 °C the mixture was stirred at 0 °C with 1 N NaOH (50 mL). After filtration through Celite the CH_2Cl_2 layer was washed with water, dried, and concentrated. The residue was chromatographed on silica gel with CH_2Cl_2 -EtOAc to give 1.92 g (60%) of a white solid after recrystallization from EtOAc: mp 175–177 °C; ^1H NMR (DMSO- d_6) δ 10.07 (s, 1H), 10.05 (s, 1H), 8.83 (s, 1H), 7.38 (s, 1H), 7.27 (s, 1H), 7.14 (s, 1H), 7.30–7.08 (s, 3H), 3.96 (s, 3H), 3.60 (s, 3H); MS (ES+) m/z 387.2, 389.2 (M + H) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{15}\text{BrN}_2\text{O}_3$) C, H, N, Br.

4-(3-Bromophenylamino)-6,7-dimethoxyquinoline-3-carboxylic Acid (53). A mixture of **5** (0.86 g, 2.0 mmol), 10 N NaOH (1.0 mL, 10 mmol), and 20 mL of EtOH was stirred at 25 °C for 2 h. The resulting solution was concentrated to remove EtOH, and the residue was dissolved in H_2O and acidified with NaH_2PO_4 . The resulting white solid was filtered off, washed with water, and dried to give 0.81 g (100%) of product: mp 282–285 °C dec; ^1H NMR (DMSO- d_6) δ 14.0 (bs, 1H), 9.12 (s, 1H), 7.64 (s, 1H), 8.30–7.60 (m, 3H), 7.47 (s, 1H), 7.45 (s, 1H), 7.26 (s, 1H), 4.01 (s, 6H); MS (ES+) m/z 403.1, 405.1 (M + H) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{15}\text{BrN}_2\text{O}_4$) C, H, N.

4-(3-Bromophenylamino)-6,7-dimethoxyquinoline-3-carboxylic Acid Amide (54). A mixture of **53** (4.03 g, 10 mmol), 1,1'-carbonyldiimidazole (3.24 g, 20 mmol), and 100 mL of DMF was stirred at 55 °C for 30 min. The resulting solution was cooled to 0 °C, saturated with NH_3 gas, and stirred at 25 °C for 45 min. The residue obtained after evaporation of solvents was stirred in H_2O . The solid was filtered, washed with H_2O , dried, and recrystallized from acetone to give a gray solid (4.0 g, 100%): mp 239–242 °C; ^1H NMR (DMSO- d_6) δ 10.1 (s, 1H), 8.84 (s, 1H), 8.15 (s, 1H), 7.61 (s, 1H), 7.36 (s, 1H), 7.25–6.80 (m, 5H), 3.94 (s, 6H); MS (ES+) m/z 402.1, 404.2 (M + H) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{16}\text{BrN}_3\text{O}_3$) C, H, N, Br.

4-(3-Bromophenylamino)-6,7-dihydroxyquinoline-3-carbonitrile (55). A 5.11 g (13.3 mmol) quantity of **8** and 30.74 g (266 mmol) of pyridine hydrochloride were mixed and then heated in an oil bath at 207 °C giving a melt. This was heated for 1 h, then cooled and treated with 100 mL of water. The resulting solid was filtered and digested with 2-methoxyethanol. This gave 3.0 g (63%) of a gray solid: mp 270–275 °C; ^1H NMR (DMSO- d_6) δ 12.09 (s, 1H, broad), 10.86 (s, 1H), 10.49 (s, 1H, broad), 8.94 (s, 1H), 7.94 (s, 1H), 7.68 (s, 1H), 7.57 (m, 2H), 7.48 (m, 2H); MS (ES+) m/z 356.1, 358.1 (M + H) $^{+1}$. Anal. ($\text{C}_{16}\text{H}_{10}\text{N}_3\text{O}_2$) $\text{Br} \cdot 1.0\text{H}_2\text{O}$) C, H, N.

4-(3-Bromophenylamino)-6,7-bis(methoxymethoxy)-quinoline-3-carbonitrile (56). This compound was prepared in 32% yield using chloromethyl methyl ether according to the procedure for **6**, except that the reaction was run entirely at 0 °C: mp 169–177 °C (EtOAc); ^1H NMR (DMSO- d_6) δ 9.92 (s, 1H), 8.37 (s, 1H), 7.80 (bs, 1H), 7.25 (s, 1H), 7.20–6.80 (m, 4H), 5.44 (s, 2H), 5.31 (s, 2H), 3.44 (s, 3H), 3.27 (s, 3H); MS (ES+) m/z 444.0, 446.0 (M + H) $^{+1}$. Anal. ($\text{C}_{20}\text{H}_{18}\text{BrN}_3\text{O}_4$) C, H, N.

4-(3-Bromophenylamino)-6,7-bis(2-methoxyethoxy)-quinoline-3-carbonitrile (57). This compound was prepared in 24% yield using bromoethyl methyl ether according to the procedure for **6**, except the reaction was run at 50 °C: mp 135–138 °C (MeOH); ^1H NMR (DMSO- d_6) δ 9.50 (s, 1H), 8.54 (s, 1H), 7.71 (s, 1H), 7.40 (s, 1H), 7.36–7.05 (m, 4H), 4.33–4.23 (m, 4H), 3.77–3.73 (m, 4H), 3.35–3.30 (m, 6H); MS (ES+) m/z 472.0, 473.9 (M + H) $^{+1}$. Anal. ($\text{C}_{22}\text{H}_{22}\text{BrN}_3\text{O}_4$) C, H, N.

8-(3-Bromophenylamino)[1,3]dioxolo[4,5-g]quinoline-7-carbonitrile (58). A solution of 2.17 g (6.09 mmol) of **55**, 0.59 mL (1.18 g, 9.14 mmol) of bromochloromethane, and 2.98 g (9.14 mmol) of Cs_2CO_3 in 20 mL of DMF was stirred and heated in an oil bath at 111 °C for 2 h. The reaction was poured into 75 mL of water and the resulting mixture was extracted with four 50-mL portions of CH_2Cl_2 . The combined extracts were dried over anhydrous MgSO_4 , then concentrated with the addition of hexanes until an oil precipitated. This oil was separated and taken up in ethyl acetate, and this solution was

washed with several portion of water, then with brine. After drying over anhydrous MgSO_4 , the solution was evaporated in vacuo to a solid. This solid was suspended in CH_2Cl_2 and poured into hexanes. The product was collected to give 0.95 g (42%) of a tan solid: mp 201–205 °C; ^1H NMR (DMSO- d_6) δ 9.44 (s, 1H), 8.57 (s, 1H), 7.74 (s, 1H), 7.33 (m, 4H), 7.16 (m, 1H), 6.28 (s, 2H); MS (ES+) m/z 368.1, 370.1 (M + H) $^{+1}$. Anal. ($\text{C}_{17}\text{H}_{10}\text{N}_3\text{O}_2\text{Br}$) C, H, N, Br.

4-(3-Bromophenylamino)-3,4-dihydro-2H-[1,4]dioxepino[2,3-g]quinoline-3-carbonitrile (59). To a stirred mixture of **55** (0.71 g, 2.0 mmol), K_2CO_3 (1.10 g, 8.0 mmol), and 8 mL of DMF at 65 °C was added a solution of 1,3-dibromopropane (0.20 mL, 2.0 mmol) during 30 min. After an additional 30 min at 65 °C the DMF was evaporated, and the residue was partitioned between CH_2Cl_2 and H_2O . After filtration, the CH_2Cl_2 layer was washed with H_2O , dried, and concentrated. The residue chromatographed on silica gel, eluting the product with 60:30:1 CH_2Cl_2 -EtOAc-HOAc to give 0.32 g (40%) of white solid: mp 120–130 °C; ^1H NMR (DMSO- d_6) δ 9.68 (s, 1H), 8.55 (s, 1H), 8.02 (s, 1H), 7.45–7.12 (m, 5H), 7.35 (s, 1H), 4.40–4.24 (m, 4H), 2.30 (s, 3H), 2.24–2.19 (m, 2H); MS (ES+) m/z 396.1, 398.1 (M + H) $^{+1}$. Anal. ($\text{C}_{19}\text{H}_{14}\text{BrN}_3\text{O} \cdot \text{HO}_2\text{CCH}_3$) C, H, N.

1-(3-Bromophenylamino)-6,7-dimethoxyisoquinoline-4-carbonitrile (61). A mixture of 6 mmol (0.65 mL, 1.03 g) of 3-bromoaniline and 5 mmol (1.25 g) of **60** 28 in 10 mL of 2-propanol was stirred and heated at reflux in an oil bath at 114 °C for 23 h. On cooling, the product was collected and washed with cold 2-propanol and ether giving 2.04 g (97%) of the product: mp 291–293 °C; ^1H NMR (DMSO- d_6) δ 10.11 (t, J = 0.5 Hz, 1H), 8.42 (s, 1H), 8.06 (d, J = 12.2 Hz, 1H), 8.03 (d, J = 1.6 Hz, 1H), 7.80 (m, 1H), 7.36 (m, 2H), 7.17 (s, 1H), 4.02 (s, 3H), 4.01 (s, 3H); MS (ES+) m/z 384.1, 386.2 (M + H) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{14}\text{N}_3\text{O}_2\text{Br} \cdot 1.0\text{HCl}$) C, H, N, Br.

1-(3-Chlorophenylamino)-6,7-dimethoxyisoquinoline-4-carbonitrile (62). A mixture of 1.24 g (5 mmol) of **60** 28 and 0.63 mL (0.76 g, 6 mmol) of 3-chloroaniline in 10 mL of 2-methoxyethanol was heated at reflux in an oil bath at 149 °C for 1 h. On cooling the solid was collected and washed with ether. The salt was stirred for about 2 h with 25 mL of saturated aqueous NaHCO_3 . The resulting solid was filtered, washed with water, and dried giving 1.62 g (95%) of the product: mp 278–280 °C; ^1H NMR (DMSO- d_6) δ 9.64 (s, 1H), 8.42 (s, 1H), 7.92 (m, 2H), 7.75 (m, 1H), 7.41 (t, J = 8.1 Hz, 1H), 7.16 (m, 2H), 4.00 (s, 3H), 3.99 (s, 3H); MS (ES+) m/z 340.2 (M + H) $^{+1}$. Anal. ($\text{C}_{18}\text{H}_{14}\text{N}_3\text{O}_2\text{Cl}$) C, H, N, Cl.

Acknowledgment. The authors thank Drs. Philip Frost, Tarek Mansour, Dennis Powell, and Janis Upeslakis for their support and encouragement. We also thank the members of the Wyeth-Ayerst Discovery Analytical Chemistry group for analytical and spectral determinations.

References

- (1) Plowman, G. D.; Ullrich, A.; Shawver, L. K. Receptor Tyrosine Kinases as Targets for Drug Intervention. *Drug News Perspect.* **1994**, 7, 334–337.
- (2) (a) Salomon, D. S.; Brandt, R.; Ciadiello, F.; Normanno, N. Epidermal Growth Factor-related Peptides and Their Receptors in Human Malignancies. *Crit. Rev. Oncol. Haematol.* **1995**, 19, 183–232. (b) Gullick, W. J. Prevalence of Aberrant Expression of the Epidermal Growth Factor Receptor in Human Cancers. *Br. Med. Bull.* **1991**, 47, 87–98. (c) Woodburn, J. R. The Epidermal Growth Factor Receptor and its Inhibition in Cancer Therapy. *Pharmacol. Ther.* **1999**, 82, 241–250.
- (3) (a) Ekstrand, A. J.; Sugawa, N.; James, C. D.; Collins, V. P. Amplified and Rearranged Epidermal Growth Factor Receptor Genes in Human Glioblastomas Reveal Deletions of Sequences Encoding Portions of the N- and/or C-Terminal Tails. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, 89, 4309–4313. (b) Wikstrand, C. J.; McLendon, R. E.; Friedman, A.; Bigner, D. D. Cell Surface Localization and Density of the Tumor-associated Variant of the Epidermal Growth Factor Receptor, EGFRvIII. *Cancer Res.* **1997**, 57, 4130–4140.

- (4) Olapade-Olaopa, E. O.; Moscatello, D. K.; MacKay, E. H.; Horsburgh, T.; Sandhu, D. P. S.; Terry, T. R.; Wong, A. J.; Habib, F. K. Evidence for the Differential Expression of a Variant EGF Receptor Protein in Human Prostate Cancer. *Br. J. Cancer* **2000**, *82*, 186–194.
- (5) Moscatello, D. K.; Holgado-Madruga, M.; Godwin, A. K.; Ramirez, G.; Gunn, G.; Zoltick, P. W.; Biegel, J. A.; Hayes, R. L.; Wong, A. J. Frequent Expression of a Mutant Epidermal Growth Factor Receptor in Multiple Human Tumors. *Cancer Res.* **1995**, *55*, 5536–5539.
- (6) Garcia de Palazzo, I. E.; Adams, G. P.; Sundaresan, P.; Wong, A. J.; Testa, J. R.; Bigner, D. D.; Weiner, L. M. Expression of Mutated Epidermal Growth Factor Receptor by Nonsmall Cell Lung Carcinomas. *Cancer Res.* **1993**, *53*, 3217–3220.
- (7) Ekstrand, A. J.; Longo, N.; Hamid, M. L.; Olson, J. J.; Liu, L.; Collins, V. P.; James, C. D. Functional Characterization of an EGF Receptor with a Truncated Extracellular Domain Expressed in Glioblastomas with EGFR Gene Amplification. *Oncogene* **1994**, *9*, 2313–2320.
- (8) Cohen, D. W.; Simak, R.; Rair, W. R.; Melamed, J.; Scher, H. I.; Cordon-Cardo, C. Expression of Transforming Growth Factor- α and the Epidermal Growth Factor Receptor in Human Prostate Tissues. *J. Urol.* **1994**, *152*, 2120–2124.
- (9) Grandis, J. R.; Melhem, M. F.; Gooding, W. E.; Day, R.; Holst, V. A.; Wagener, M. M.; Drenning, S. D.; Tweardy, D. J. Levels of TGF α and EGFR Protein in Head and Neck Squamous Cell Carcinoma and Patient Survival. *J. Natl. Cancer Inst.* **1998**, *90*, 824–832.
- (10) Morishige, K. I.; Kurachi, H.; Ameniya, K.; Fujita, Y.; Yamamoto, T.; Mikaye, A.; Tanizawa, O. Evidence for the Involvement of TGF α and EGFR Autocrine Mechanism in Primary Ovarian Cancers In Vitro. *Cancer Res.* **1991**, *51*, 5322–5328.
- (11) Rusch, V.; Klimstra, D.; Venkatraman, E.; Pisters, P. W. T.; Langenfeld, J.; Dmitrovsky, E. Overexpression of the Epidermal Growth Factor Receptor and its Ligand Transforming Growth Factor Alpha is Frequent in Resectable Nonsmall Cell Lung Cancer but does not Predict Tumor Progression. *Clin. Cancer Res.* **1997**, *3*, 515–522.
- (12) Thogersen, V. B.; Jorgensen, P. E.; Sorensen, B. S.; Bross, P.; Orntoft, T.; Wolf, H.; Nexø, E. Expression of Transforming Growth Factor Alpha and Epidermal Growth Factor Receptor in Human Bladder Cancer. *Scand. J. Clin. Lab Invest.* **1999**, *59*, 267–277.
- (13) Sweeney, W. E., Jr.; Chen, Y.; Nakanishi, K.; Frost, P.; Avner, E. Treatment of Polycystic Kidney Disease with a Novel Tyrosine Kinase Inhibitor. *Kidney Int.* **2000**, *57*, 33–40.
- (14) (a) Elder, J. T.; Fisher, G. J.; Lindquist, P. B.; Bennett, G. J.; Pittelkow, M. R.; Coffey, R. J.; Ellingsworth, L.; Derynck, R.; Voorhees, J. J. Overexpression of Transforming Growth Factor- α in Psoriatic Epidermis. *Science* **1989**, *243*, 811–814. (b) Cook, P. W.; Peipkorn, M.; Clegg, C. H.; Plowman, G. D.; DeMay, J. M.; Brown, J. R.; Pittelkow, M. R. Transgenic Expression of the Human Amphiregulin Gene Induces a Psoriasis-like Phenotype. *J. Clin. Invest.* **1997**, *100*, 2286–2294.
- (15) Davies, D. E.; Polosa, R.; Puddicombe, S. M.; Richter, A.; Holgate, S. T. The Epidermal Growth Factor Receptor and its Ligand Family: Their Potential Role in Repair and Remodelling in Asthma. *Allergy* **1999**, *54*, 771–783.
- (16) (a) Yamamoto, T.; Ikawa, S.; Akiyama, T.; Semba, K.; Nomura, N.; Miyajima, N.; Saito, T.; Toyoshima, K. Similarity of Protein Encoded by the Human *c-erb-B-2* Gene to the Epidermal Growth Factor Receptor. *Nature* **1986**, *319*, 230–234. (b) Coussens, L.; Yang-Feng, T. L.; Liao, Y.-C.; Chen, E.; Gray, A.; McGrath, J.; Seeburg, P. H.; Libermann, T. A.; Schlessinger, J.; Francke, U.; Levinson, A.; Ullrich, A. Tyrosine Kinase Receptor with Extensive Homology to EGF Receptor Shares Chromosomal Location with *neu* Oncogene. *Science* **1985**, *230*, 1132–1139.
- (17) (a) Fry, D. W.; Kraker, A. J.; McMichael, A.; Ambrosio, L. A.; Nelson, J. M.; Leopold, W. R.; Connors, R. W.; Bridges, A. J. A Specific Inhibitor of the Epidermal Growth Factor Receptor Tyrosine Kinase. *Science* **1994**, *265*, 1093–1095. (b) Ward, W. H. J.; Cook, P. N.; Slater, A. M.; Davies, D. H.; Holdgate, G. A.; Green, L. R. Epidermal Growth Factor Tyrosine Kinase. Investigation of Catalytic Mechanism, Structure-Based Searching and Discovery of a Potent Inhibitor. *Biochem. Pharmacol.* **1994**, *48*, 659–666.
- (18) For recent reviews, see: (a) Bridges, A. J. The Rational and Strategy Used to Develop a Series of Highly Potent, Irreversible Inhibitors of the Epidermal Growth Factor Receptor Family of Tyrosine Kinases. *Cur. Med. Chem.* **1999**, *6*, 825–843. (b) Traxler, P. M. Protein Tyrosine Kinase Inhibitors in Cancer Treatment. *Exp. Opin. Ther. Patents* **1997**, *7*, 571–588. (c) Traxler, P. M. Tyrosine Kinase Inhibitors in Cancer Treatment (Part II). *Exp. Opin. Ther. Patents* **1998**, *8*, 1599–1625.
- (19) Discifani, C. M.; Carroll, M.; Floyd, M. B.; Hollander, I. J.; Husain, Z.; Johnson, B. D.; Kitchen, D.; May, M. K.; Minnick, A. A.; Nilakantan, R.; Shen, R.; Wang, Y.; Wissner, A.; Greenberger, L. M. CL-387,785: An Irreversible Inhibitor of Epidermal Growth Factor Receptor Tyrosine Kinase with In Vivo Activity. *Biochem. Pharmacol.* **1999**, *57*, 917–925.
- (20) (a) Kris, M.; Ranson, M.; Ferry, D.; Hammond, L.; Averbuch, S.; Ochs, J.; Rowinsky, E. Phase I Study of Oral ZD1839 (Iressa), a Novel Inhibitor of Epidermal Growth Factor Receptor Tyrosine Kinase (EGFR-TK): Evidence of Good Tolerability and Activity. *Clin. Cancer Res.* **1999**, *5*, 3749s–3750s. (b) Moyer, J. D.; Barbacci, E. G.; Iwata, K. K.; Arnold, L.; Bowman, B.; Cunningham, A.; DiOrio, C.; Doty, J.; Morin, M. J.; Moyer, M. P.; Neuveu, M.; Pollack, V. A.; Pustilnik, L. R.; Reynolds, M. M.; Sloan, D.; Theleman, A.; Miller, P. Induction of Apoptosis and Cell Cycle Arrest by CP-358,774, an Inhibitor of Epidermal Growth Factor Receptor Tyrosine Kinase. *Cancer Res.* **1997**, *57*, 4838–4848. (c) Fry, D. W.; Bridges, A. J.; Denny, W.; Doherty, A.; Greis, K. D.; Hicks, J. L.; Hook, K. E.; Keller, P. R.; Leopold, W. R.; Loo, J. H.; McNamara, D. J.; Nelson, J. M.; Sherwood, V.; Smail, J. B.; Trumpp-Kallmeyer, S.; Dobrusin, E. M. Specific, Irreversible Inhibitors of the Epidermal Growth Factor Receptor and *erbB2*, by a New Class of Tyrosine Kinase Inhibitor. *Proc. Natl. Acad. Sci. U.S.A.* **1998**, *95*, 12022–12027.
- (21) Rewcastle, G. W.; Denny, W. A.; Bridges, A. J.; Zhou, H.; Cody, D. R.; McMichael, A.; Fry, D. W. Tyrosine Kinase Inhibitors. 5. Synthesis and Structure–Activity Relationships for 4-(Phenylmethylamino)- and 4-(Phenylamino)quinazolines as Potent Adenosine 5'-Triphosphate Binding Site Inhibitors of the Tyrosine Kinase Domain of the Epidermal Growth Factor Receptor. *J. Med. Chem.* **1995**, *38*, 3482–3487.
- (22) Knighton, D. R.; Zheng, J.; Ten Eyck, L. F.; Taylor, S. S.; Sowadski, J. M.; Gill, G. N. Structural Features that Specify Tyrosine Kinase Activity Deduced from Homology Modeling of Epidermal Growth Factor Receptor Kinase. *Proc. Natl. Acad. Sci. U.S.A.* **1993**, *90*, 5001–5005.
- (23) Wang, H.; Ben-Naim, A. A Possible Involvement of Solvent-Induced Interactions in Drug Design. *J. Med. Chem.* **1996**, *39*, 1531–1539.
- (24) Bridges, A. J.; Zhou, H.; Cody, D. R.; Rewcastle, G. W.; McMichael, A.; Showalter, H. D. H.; Fry, D. W.; Kraker, A. J.; Denny, W. A. Tyrosine Kinase Inhibitors. 8. Unusually Steep Structure–Activity Relationship for Analogues of 4-(3-Bromo-anilino)-6,7-dimethoxyquinazoline (PD 153035), a Potent Inhibitor of the Epidermal Growth Factor Receptor. *J. Med. Chem.* **1996**, *39*, 267–276.
- (25) (a) Bredereck, H.; Effenberger, F.; Botsch, H.; Rehn, H. Synthesen in der Heterocyclischen Reihe V. Umsetzungen von Vinylgen Carbonsäureamiden zu Heterocyclen. *Chem. Ber.* **1965**, *98*, 1081–1086. (b) Egri, J.; Halmos, J.; Rakoczy, J. Synthesis of Substituted 4-Hydroxyquinoline-3-carboxylic esters, IV. *Acta Chim. (Budapest)* **1973**, *78*, 217–225.
- (26) Burke, T. R., Jr.; Lim, B.; Marquez, V. E.; Li, Z.-H.; Bolen, J. B.; Stefanova, I.; Horak, I. D. Bicyclic Compounds as Ring-Constrained Inhibitors of Protein-Tyrosine Kinase p56. *J. Med. Chem.* **1993**, *36*, 425–432.
- (27) Lipp, M.; Dallacker, F.; Schaffranek, R. Darstellung von Derivaten des 1,2-Athylendioxy-benzols. *Chem. Ber.* **1958**, *91*, 2247–2250.
- (28) Ernest, W.; Haugwitz, R. D. 4-Substituted 6,7-Dimethoxyisoquinolines. *Can. J. Chem.* **1968**, *46*, 1160–1163.
- (29) Mohammadi, M.; Schlessinger, J.; Hubbard, S. R. Structure of the FGF Receptor Tyrosine Kinase Domain Reveals a Novel Autoinhibitory Mechanism. *Cell* **1996**, *86*, 577–587.
- (30) Schindler, T.; Sicheri, F.; Pico, A.; Gazit, A.; Levitzki, A.; Kuriyan, J. Crystal Structure of Hck in Complex with a Src Family-Selective Tyrosine Kinase Inhibitor. *Mol. Cell* **1999**, *3*, 639–648.
- (31) Shewchuk, L.; Hassell, A.; Wisely, B.; Rocque, W.; Holmes, W.; Veal, J.; Kuyper, L. F. Binding Mode of the 4-Anilinoquinazoline Class of Protein Kinase Inhibitor: X-ray Crystallographic Studies of 4-Anilinoquinazolines Bound to Cyclin-Dependent Kinase 2 and p38 Kinase. *J. Med. Chem.* **2000**, *43*, 133–138.
- (32) Chen, J. M.; Xu, S. L.; Wawrzah, Z.; Basarab, G. S.; Jordan, D. B. Structure-based Design of Potent Inhibitors of Scytalone Dehydratase: Displacement of a Water Molecule from the Active Site. *Biochemistry* **1998**, *37*, 17735–17744.
- (33) Palmer, B. D.; Trumpp-Kallmeyer, S.; Fry, D. W.; Nelson, J. M.; Showalter, H. D. H.; Denny, W. A. J. Tyrosine Kinase Inhibitors. 11. Soluble Analogues of Pyrrolo- and Pyrazoloquinazolines as Epidermal Growth Factor Receptor Inhibitors: Synthesis, Biological Evaluation, and Modeling of the Mode of Binding. *J. Med. Chem.* **1997**, *40*, 1519–1529.
- (34) Skehan, P.; Stornet, R.; Scudiero, D.; Monks, A.; McMahon, J.; Vistica, D.; Warren, J.; Bokkosh, H.; Kenny, S.; Boyd, M. New Colorimetric Cytotoxic Assay for Anticancer-Drug Screening. *J. Natl. Cancer Inst.* **1990**, *82*, 1107–1112.