

2000, Volume 43

David Bebbington,* Nathaniel J. T. Monck, Suneel Gaur, Alan M. Palmer, Karen Benwell, Victoria Harvey, Craig S. Malcolm, and Richard H. P. Porter: 3,5-Disubstituted-4-hydroxyphenyls Linked to 3-Hydroxy-2-methyl-4(1*H*)-pyridinone: Potent Inhibitors of Lipid Peroxidation and Cell Toxicity.

Page 2781. Figures 2 and 3 have incorrect scaling on the *y*-axis (0% to 12%). The correct *y*-axis scale is from 0% to 120%. The corrected figures are shown below.




Figure 2. Inhibition of IAA-induced cell toxicity. CGC were exposed to 30 μ M IAA for 30 min in a physiological salt solution. This was replaced with maintenance media containing 1 μ M test compound, and the cells were tested for viability 24 h later. Compound **3b** was tested as its mesylate salt. The error bars represent SE bars. Statistical analysis was performed using two-tailed pair *t*-test. Statistical significance is defined as $p < 0.05$; **significantly different from **7** and **9** ($p < 0.01$); #significantly different from IAA alone ($p < 0.05$).




Figure 3. Inhibition of H₂O₂-induced cell toxicity. CGC were exposed to 100 μ M H₂O₂ for 10 min prior to addition of test compounds at a concentration of 10 μ M. The cells were tested for viability 24 h later. Compound **3b** was tested as its mesylate salt. The error bars represent SE bars. Statistical analysis was performed using two-tailed paired *t*-test. Statistical significance is defined as $p < 0.05$; *significantly different from **7** and **9** ($p < 0.05$); #significantly different from H₂O₂ alone ($p < 0.05$).

JM0010872

10.1021/jm0010872

Published on Web 11/30/2000