

Identification of Shc Src Homology 2 Domain-Binding Peptoid–Peptide Hybrids

Won Jun Choi,[†] Sung-Eun Kim,[†] Andrew G. Stephen,[‡] Iwona Weidlich,^{†,§} Alessio Giubellino,^{||} Fa Liu,[†] Karen M. Worthy,[‡] Lakshman Bindu,[‡] Matthew J. Fivash,[⊥] Marc C. Nicklaus,[†] Donald P. Bottaro,^{||} Robert J. Fisher,[‡] and Terrence R. Burke, Jr.*[†]

Laboratory of Medicinal Chemistry, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Frederick, Maryland 21702, Protein Chemistry Laboratory, Advanced Technology Program, SAIC–Frederick, Frederick, Maryland 21702, Urologic Oncology Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, Maryland 20892, and Data Management Services, Inc., NCI–Frederick, Frederick, Maryland 21702

Received June 30, 2008

A fluorescence anisotropy (FA) competition-based Shc Src homology 2 (SH2) domain-binding was established using the high affinity fluorescein isothiocyanate (FITC) containing peptide, FITC-NH-(CH₂)₄-CO-pY-Q-G-L-S-amide (**8**; K_d = 0.35 μM). Examination of a series of open-chain bis-alkenylamide containing peptides, prepared as ring-closing metathesis precursors, showed that the highest affinities were obtained by replacement of the original Gly residue with N^α-substituted Gly (NSG) “peptoid” residues. This provided peptoid–peptide hybrids of the form “Ac-pY-Q-[NSG]-L-amide.” Depending on the NSG substituent, certain of these hybrids exhibited up to 40-fold higher Shc SH2 domain-binding affinity than the parent Gly-containing peptide (IC₅₀ = 248 μM) (for example, for *N*-homoallyl analogue **50**, IC₅₀ = 6 μM). To our knowledge, this work represents the first successful example of the application of peptoid–peptide hybrids in the design of SH2 domain-binding antagonists. These results could provide a foundation for further structural optimization of Shc SH2 domain-binding peptide mimetics.

Introduction

Disrupting protein–protein interactions is evolving as a challenging but potentially rewarding approach to therapeutic development.¹ Src homology 2 (SH2^{cc}) domains facilitate the formation of signaling complexes by mediating the recognition and binding to phosphotyrosyl (pTyr) containing sequences,^{2–4} and SH2 domain-binding antagonists may down-regulate protein–tyrosine kinase signaling in a fashion that is complementary to kinase-directed inhibitors.^{5,6} Although significant effort has focused on development of p56^{lck}, Src, and growth factor receptor bound protein 2 (Grb2) SH2 domain-binding antagonists,^{7–11} relatively little has been reported on agents directed against the Shc SH2 domain.^{12,13} Members of the Shc family of noncatalytic SH2 domain-containing adapter proteins serve as transforming elements involved with mitogenic signal transduction,^{14,15} and they have been shown to be essential for induction of the proangiogenic vascular endothelial growth factor (VEGF).¹⁶ Pathways downstream of Shc or Grb2 have also been shown to be critical for transformation and metastasis.^{17,18} The importance of Shc in Ras activation pathways has made antagonists of Shc signaling potentially attractive therapeutic targets. The current report details unexpected finding of potent

inhibitory motifs that grew out of unsuccessful attempts to develop macrocyclic Shc SH2 domain binding inhibitors.

Results and Discussion

Development of Shc SH2 Domain-Binding Fluorescence Anisotropy Assays: Identification of a High Affinity Fluorescein Isothiocyanate Containing Peptide. Previous analyses of Shc SH2 domain-binding constants have relied on relatively labor-intensive NMR or surface plasmon resonance techniques.¹³ Assays based on fluorescence anisotropy (FA) afford attractive alternatives for determining binding constants, but they can be limited by the need to append fluorescein isothiocyanate (FITC) to every peptide examined. However, this requirement can be circumvented by the availability of a high affinity reference FITC-containing peptide that can be used in FA assays to determine affinities of non-FITC labeled peptides by competition techniques. Therefore, our first objective of the current study was to identify a FITC-containing reference peptide having high Shc SH2 domain binding affinity for use in FA competition assays.

Modeling based on NMR solution studies of the ζ chain T-cell-receptor-derived 14-mer peptide “GHDGLpYQGL-STATK” (**1**) bound to the Shc SH2 domain^{19,20} indicated that significant interactions only occur for residues “pYQGL.” Accordingly, the slightly longer “LpYQGLS” sequence (**2**) was used as a starting point for structural exploration of FITC-containing peptides. Initial studies were conducted to determine the optimal introduction of an N-terminal FITC group using a variety of spacer lengths (**3–14**, Table 1). Direct N-terminal attachment of the FITC group (**3**) resulted in a complete loss of binding affinity. However, deletion of the N-terminal Leu residue from **2** prior to addition of the FITC tag gave high affinity (**4**, K_d ≈ 12 μM). Adding the FITC label to the latter “pY-Q-G-L-S-amide” sequence by means of a series of progressively longer amino-*n*-alkylcarboxylic acid amide linkers (**5–9**) gave the highest affinity using a pentanoyl chain (**8**, K_d = 0.35 μM). The best affinities from a similar series of FITC-containing

* To whom correspondence should be addressed. Phone: (301) 846-5906. Fax: (301) 846-6033. E-mail: tburke@helix.nih.gov.

[†] Laboratory of Medicinal Chemistry, National Cancer Institute.

[‡] SAIC–Frederick.

[§] On leave from the Poznan University of Medical Sciences, Faculty of Pharmacy, Poland.

^{||} Urologic Oncology Branch, National Cancer Institute.

[⊥] Data Management Services, Inc.

^{cc} Abbreviations: FA, fluorescence anisotropy; FITC, fluorescein isothiocyanate; SH2, Src homology 2; NSG, N^α-substituted glycine; pTyr, phosphotyrosyl; VEGF, vascular endothelial growth factor; Grb2, growth factor receptor bound protein 2; PEG, polyethylene glycol; HGF, hepatocyte growth factor; RCM, ring-closing metathesis; FEB, free energy of binding; NMP, *N*-methyl-2-pyrrolidone; DIPCDI, *N,N'*-diisopropylcarbodiimide; HOBT, 1-hydroxybenzotriazole; PyBOP, (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate; DIPEA, *N,N*-diisopropylethylamine; EDT, 1,2-ethanedithiol.














Table 1. Shc SH2 Domain-Binding Affinities Determined Using a Fluorescence Anisotropy Assay

compd	sequence	$K_d \pm SD$ (μM) ^a
3	FITC-L-pY-Q-G-L-S-amide	nb ^b
4	FITC-pY-Q-G-L-S-amide	12.4 ± 2.6
5	FITC-NH-(CH ₂) ₂ -CO-pY-Q-G-L-S-amide	nb ^b
6	FITC-NH-(CH ₂) ₂ -CO-pY-Q-G-L-S-amide	1.09 ± 0.04
7	FITC-NH-(CH ₂) ₃ -CO-pY-Q-G-L-S-amide	7.2 ± 0.2
8	FITC-NH-(CH ₂) ₄ -CO-pY-Q-G-L-S-amide	0.35 ± 0.13
9	FITC-NH-(CH ₂) ₅ -CO-pY-Q-G-L-S-amide	4.7 ± 0.1
10	FITC-NH-(CH ₂) ₂ -CO-L-pY-Q-G-L-S-amide	nb ^b
11	FITC-NH-(CH ₂) ₂ -CO-L-pY-Q-G-L-S-amide	7.6 ± 0.5
12	FITC-NH-(CH ₂) ₃ -CO-L-pY-Q-G-L-S-amide	8.6 ± 0.3
13	FITC-NH-(CH ₂) ₄ -CO-L-pY-Q-G-L-S-amide	8.9 ± 0.3
14	FITC-NH-(CH ₂) ₅ -CO-L-pY-Q-G-L-S-amide	nb ^b
15	Ac-NH-(CH ₂) ₄ -CO-pY-Q-G-L-S-amide	158 ± 10 ^c

^a Except as noted, determined by fluorescence anisotropy techniques as described in the Experiment Section. ^b No binding. ^c IC₅₀ values determined by FA competition experiments using peptide **8** as the reference peptide.

peptides prepared using the longer “L-pY-Q-G-L-S-amide” sequence (**10–14**) were approximately 20-fold less potent. Replacement of the FITC group in **8** with *N*-acetyl (**15**, IC₅₀ = 158 μM) resulted in a significant loss of binding affinity as determined using a FA competition assay against **8** as the reference peptide. This indicated that the FITC moiety in **8** contributes substantially to the binding affinity.

Table 2. Shc SH2 Domain-Binding Affinities for FITC-X-pY-Q-G-L-S-amide

No	Spacer (X)	$K_d \pm s.d$ (μM) ^a	No	Spacer (X)	$K_d \pm s.d$ (μM) ^a
16		8.6 ± 0.9	23		5.0 ± 0.9
17		9.0 ± 3.2	24		0.77 ± 0.04
18		5.3 ± 0.6	25		14.2 ± 2.1
19		9.5 ± 1.1	26		25.4 ± 1.6
20		18.0 ± 0.8	27		0.5 ± 0.3
21		23.9 ± 2.0	28		12.3 ± 2.3
22		poor			

^a Determined by a direct binding fluorescence anisotropy assay as described in the Experimental Section.

To examine the possible effects in binding affinity of the linker structure other than chain length, analogues were prepared based on the FITC-X-pY-Q-G-L-S-amide sequence (**16–28**, Table 2). These more structurally complex linkers could potentially affect binding affinity by interacting directly with the SH2 domain protein or by positioning the FITC ring system. Most of the peptides exhibited binding affinities in the range 5–25 μM . However, peptides **24** ($K_d \approx 0.8 \mu M$) and **27** ($K_d \approx 0.5 \mu M$) exhibited approximately 10-fold higher affinities than other members of the series. Of particular note was the 50-fold higher affinity of **27** relative to its trans isomer (**26**). Since none of the peptides in Table 2 exhibited higher affinity than the original FITC-containing **8**, this latter peptide was chosen for use as a reference in subsequent FA competition assays.

Blockade of Shc Binding to c-Met Tyrosine Kinase in Whole Cells. In order to confirm that the highest affinity FITC-labeled peptide (**8**) is capable of binding to the Shc SH2 domain in a physiologically relevant manner, studies were performed that examined the ability of **8** to inhibit the binding of intracellular Shc protein to the cytoplasmic domain of the c-Met protein–tyrosine kinase in intact human mammary B5/589 cells, which express high levels of c-Met. For this purpose, Universal PEG Novatag resin²¹ was used to prepare peptides **29** and **30** (Figure 1), which contained the membrane carrier peptide




Figure 1. Inhibition by peptides **29** (A) and **30** (B) of binding of Shc to hepatocyte growth factor receptor (c-Met) tyrosine kinase in intact B5/589 cells as described in Experimental Section. Cells were incubated for 1 h in the presence (10 μ M) or absence (0 μ M) of the indicated peptides, then untreated (control) or briefly stimulated with HGF (+HGF). Cells were lysed and the amount of Shc bound to c-Met were determined (upper panels) and the amount c-Met recovered (lower panels).

sequences “RRRRRRRR”²² and “RQIKIWFQNRRMKWKK”²³ attached to the C-terminus of **8** by means of a polyethylene glycol (PEG) spacer. The Shc SH2 domain-binding affinities of

29 and **30** as determined by extracellular FA competition assays were $K_d = 5$ and 1 μ M, respectively. Following incubation of cells with the peptides in the presence and absence of hepatocyte growth factor (HGF, a ligand for c-Met receptor), the cells were washed and lysed and binding of Shc to activated c-Met was measured as described in the Experimental Section. At 10 μ M, both **29** and **30** were able to achieve a consistent blockade of intracellular binding of Shc to c-Met (Figure 1).


Examination of Shc SH2 Domain-Binding Affinities of N-Alkenyl-Containing Peptides (31–44) Designed as Ring-Closing Metathesis Precursors. We had previously shown that induction of conformational constraint through macrocyclization using ring-closing metathesis (RCM)²⁴ could improve the binding affinity of Grb2 SH2 domain-binding peptides.²⁵ In the current study we performed molecular modeling experiments to explore the potential utility of RCM macrocyclization in the design of Shc SH2 domain-binding peptides. However, these modeling studies failed to identify obvious macrocyclic candidates. Nonetheless, we undertook work directed at the preparation of a library of macrocycles. This was intended as an empirical alternative to a “structure-based design” approach. For this purpose, two series of peptides were prepared, predicated on the sequences Ac-pY-Q-G-L-S-NHR and Ac-pY-Q-G-L-NHR (A and B, respectively, Table 3), where R = allyl and homoallyl. For each series of peptides, additional N-allyl groups were placed at various nitrogen positions within the peptide backbone to form bis-alkenyl open-chain RCM precursors. Shc SH2 domain binding affinities of these putative precursors were determined prior to macrocyclization studies.

In the A series, the highest affinities were obtained by N-allylation of the Gln-Gly amide nitrogen, with the N-terminal homoallyl peptide (**37**; $IC_{50} = 8 \mu$ M) exhibiting higher affinity than the corresponding N-terminal allyl congener (**33**; $IC_{50} = 48 \mu$ M). Placement of the backbone N-alkenyl group at other positions in the peptide chain (position 1 or 3, Table 3) or deletion entirely (peptides **31** and **35**) gave generally lower affinities. Similar results were found in the B series. The data presented in Table 3 were of note in that the highest affinities were derived by replacing the original Gly residues with N $^{\alpha}$ -substituted glycines (NSG). The NSG groups are the fundamental units of the “peptoid” class of peptide mimetics, and

Table 3. Shc SH2 Domain-Binding Affinities of Bis-N-Olefin Containing Peptides

peptide A analogues				peptide B analogues			
compd	position ^a	n	$IC_{50} \pm SD$ (μ M) ^b	compd	position ^a	n	$IC_{50} \pm SD$ (μ M) ^b
31	0	1	189 \pm 15	38	0	1	91 \pm 12
32	1	1	353 \pm 34	39	1	1	81 \pm 6
33	2	1	48 \pm 2	40	2	1	47 \pm 2
34	3	1	41 \pm 7	41	3	1	450 \pm 38
35	0	2	165 \pm 4	42	0	2	84 \pm 4
36	1	2	89 \pm 5	43	1	2	84 \pm 6
37	2	2	8 \pm 2	44	2	2	14 \pm 3

^a Site of N-allyl substitution. A designation of “0” indicates the lack of N-allyl substitution. ^b Determined by a fluorescence anisotropy based competition assay using peptide **8** as a reference as described in the Experimental Section.

Table 4. Shc SH2 Domain-Binding Affinities of *N*-Alkylglycine Containing Peptides

No	R	IC ₅₀ ± s.d (μM) ^a	No	R	IC ₅₀ ± s.d (μM) ^a	No.	R	IC ₅₀ ± s.d (μM) ^a
45		248 ± 5	52		83 ± 3	59		16 ± 4
46		19 ± 0.5	53		17 ± 3	60		17 ± 3
47		25.7 ± 0.7	54		40 ± 6	61		33 ± 1
48		26.6 ± 0.7	55		23 ± 3	62		34 ± 1
49		8.9 ± 0.1	56		56 ± 13	63		20 ± 3
50		6.0 ± 0.2	57		29 ± 7	64		14 ± 0
51		18.9 ± 0.3	58		10 ± 0.7	65		44 ± 2

^a Determined by a fluorescence anisotropy based competition assay using peptide **8** as the reference peptide as described in the Experimental Section.

peptides such as **37** and **44** represent peptoid–peptide hybrids. To our knowledge, this is the first identification of peptoid–peptide hybrids as high affinity SH2 domain-binding inhibitors.

As indicated above, the bis-alkenylamine-containing peptides were originally intended as precursors for the preparation of

macrocyclic peptide mimetics. However, RCM macrocyclization studies carried out using second generation Grubbs catalyst²⁶ were largely unsuccessful and only a small number of ring-closed products could be obtained. Consistent with preliminary molecular modeling studies that failed to predict binding enhancement through




Figure 2. Hydrophobic and hydrophilic surface maps for two conformations (A and B) of peptide ligand **50** binding to Shc SH2 domain formed after 10 ps of molecular dynamics simulations. The coloring of atoms is as follows: carbon, green; nitrogen, blue; oxygen, red. The *N*-homoallyl group interacts with hydrophobic pockets. Hydrophobic and hydrophilic surface maps are colored in orange and blue, respectively. Peptide **50** is displayed in ball and stick representation.

macrocyclization, none of the obtained final cyclic peptides showed significantly enhanced affinity (data not shown), and further work with peptide macrocyclization was not pursued.

Examination of *N*-Substituted Glycine Residues in Peptides Lacking C-Terminal Alkenylamide Substituents.

Because *N*-alkenylation of the Gln-Gly amide nitrogen appeared to be advantageous in the open-chain series bearing C-terminal alkenylamides (for example, **37** and **44**), studies were undertaken to examine an array of NSG residues in a simplified peptide platform lacking any C-terminal carboxamide substituents (Table 4). For the parent nonsubstituted Gly-containing peptides **38** and **42**, deletion of the C-terminal allyl and homoallyl amide functionalities, respectively, gave "Ac-pY-Q-G-L-amide", which resulted in an approximate 2- to 3-fold loss of binding affinity (**45**, Table 4; $IC_{50} = 248 \mu M$). Relative to **45**, addition of a C-terminal Ser residue ("Ac-pY-Q-G-L-S-amide") resulted in an approximate doubling of binding affinity ($IC_{50} = 112.5 \pm 0.7 \mu M$), while shortening **45** by removing the N-terminal Leu residue abrogated nearly all binding affinity ("Ac-pY-Q-G-amide" $IC_{50} > 500 \mu M$). Therefore, peptide **45** was taken as a starting point for peptoid-peptide synthesis because it represented a balance between size and binding affinity.

Starting from **45**, formation of NSG residues by alkylation of the Gln-Gly amide bond enhanced binding affinity by an order of magnitude or more (Table 4). The highest affinities were obtained with 4-carbon linear chains (*n*-butyl (**49**), $IC_{50} = 8.9 \mu M$ and homoallyl (**50**), $IC_{50} = 6.0 \mu M$). Increasing chain length beyond 4 units resulted in a loss of affinity (*n*-pentyl (**58**) $IC_{50} = 10 \mu M$ and *n*-hexyl (**63**), $IC_{50} = 20 \mu M$), as did insertion of heteroatoms (oxygen (**56**), $IC_{50} = 56 \mu M$; sulfur (**62**), $IC_{50} = 34 \mu M$ or nitrogen (**65**), $IC_{50} = 44 \mu M$). Although an approximate 2- to 3-fold loss of potency was incurred by the original deletion of C-terminal alkenylamide functionality, subsequent introduction of a homoallyl-NSG residue (**50**) more than compensated for the potency loss to yield a peptide that exhibited twice the affinity of the parent bis-homoallyl-containing **44**.

Molecular Modeling. Molecular dynamics (MD) simulations of *N*-homoallyl-containing **50** bound to the Shc SH2 domain protein were conducted. The two conformations with the two lowest energy complexes are depicted in Figure 2. Subjecting these complexes to FEB (free energy of binding) calculations using the program eMBrAcE developed by Schrödinger showed

that both complexes were energetically similar (conformation A, $-21\,957.656$ kJ/mol; conformation B, $-21\,957.133$ kJ/mol). NMR and crystallographic studies have previously indicated that interactions of peptide ligands with Shc SH2 domain protein can vary significantly.^{19,27} Therefore, it was not surprising that binding of peptide **50** differed from that reported for the 14-mer T-cell-derived parent peptide (**1**).¹⁹ For peptide **50**, the slightly more favorable conformation A (Figure 2) has the *N*-homoallyl group interacting with a hydrophobic pocket created by residues Leu54, Leu76, Ile87, and Ile88. In conformation B, the Ala90 residue has shifted position, allowing the *N*-homoallyl group to bind within a newly created hydrophobic pocket. In both conformations, the homoallyl group makes significant binding interactions that may explain the relatively high affinity of the peptoid-peptide hybrids.

Conclusions

By use of a FA-based competition assay based on the high affinity FITC-containing peptide **8**, evaluation of a number of bis-alkenylamide peptides for Shc SH2 domain-binding affinity resulted in the discovery of pTyr + 2 *N*-alkyl-Gly residues as favorable structural motifs. Deletion of the C-terminal alkenylamide could be achieved without overall loss of binding affinity. Molecular modeling studies identified specific pockets within the protein surface that appear to afford regions of hydrophobic interactions with the NSG side chain. These findings represent the first examples of peptoid-peptide hybrids as preferred SH2 domain-binding motifs. As such, this work could provide the foundation for further structure-based optimization of Shc SH2 domain-binding peptide mimetics.

Experimental Section

Preparation of Shc SH2 Domain Protein. Shc protein was prepared using previously published protocols.²⁸ A total of 1.2 mg of lyophilized Shc protein was resuspended in a 17 μL of 100% (v/v) dimethyl sulfoxide (DMSO), and then 159 μL of a buffer solution (BS) consisting of 10 mM HEPES, 150 mM NaCl, pH 7.5, was added for a stock of 500 μM protein.

Steady State Fluorescence Anisotropy Assays. Aliquots of a solution of 5 nM peptide labeled with fluorescein at the N-terminus in 1 \times BS, 10% (v/v) dimethyl sulfoxide were placed in 96-well Costar polypropylene plates (Corning, NY). Shc protein was serially diluted in the peptides with a starting concentration of 100 μM . Then 40 μL was removed and transferred into 384-well Costar

polypropylene plates (Corning, NY). Samples were excited at 485 nm, and the emission intensities at 535 nm from the parallel and perpendicular planes were measured using a Tecan Ultra plate reader (Durham, NC).

Curves were fit assuming a single binding site per peptide, using the relation

$$\text{An}(C) = ((R\text{An}_{\text{bound}} - \text{An}_{\text{free}})((K_d + C + \text{Lt}) - [(K_d + C + \text{Lt})^2 - 4\text{Lt}C]^{1/2}) / (2\text{Lt})) + \text{An}_{\text{free}}) / ((R - 1)((K_d + C + \text{Lt}) - [(K_d + C + \text{Lt})^2 - 4\text{Lt}C]^{1/2}) / (2\text{Lt})) + 1) \quad (1)$$

where $\text{An}(C)$ is the measured anisotropy as a function of concentration, Lt is the peptide concentration, An_{free} is the anisotropy of unbound peptide, An_{bound} is the anisotropy of the completely bound peptide solution, R is the ratio of the fluorescence intensity of saturated bound peptide relative to free peptide, C is the protein concentration, and K_d is the measured dissociation constant resulting from the fit.

Fluorescence Anisotropy Competition Assays. A complex of 2 μM Shc and 10 nM peptide **8** in $1 \times \text{BS}$, 10% (v/v) DMSO was placed in 96-well Costar polypropylene plates (Corning, NY). Unlabeled competitor peptides were serially diluted into the complex with starting concentrations of 500 μM . A total of 40 μL was removed and transferred into 384-well Costar polypropylene plates (Corning, NY). Samples were excited at 485 nm, and the emission intensities at 535 nm from the parallel and perpendicular planes were measured using a Tecan Ultra plate reader (Durham, NC). To estimate the IC_{50} values for these competition experiments, the fractional binding of the competitor ligand was determined from the binding part of the anisotropy signal using estimates of the maximum and minimum anisotropy signal.²⁹ The fractional binding signal is described using a least-squares spline approximate.³⁰ Three knots in a spline of order 3 were used, with the value at which the spline achieved one-half being taken as an estimate of the IC_{50} value.

Inhibition of Shc SH2 Domain Binding to Cytoplasmic c-Met Tyrosine Kinase Receptor in Intact Cells. Intact B5/589 human mammary epithelial cells were serum-deprived (48 h), then incubated (1 h) in presence or absence of peptides **29** and **30**. Cells were treated with hepatocyte growth factor (R&D Systems, Minneapolis, MN; 50 ng/mL, 20 min) and then lysed in cold buffer (pH 7.4) containing nonionic detergent and protease and phosphatase inhibitors. After incubation with a biotin-tagged, affinity-purified c-Met antibody (BAF358, R&D Systems, Minneapolis, MN; 1 h on ice), immunocomplexes were captured using a streptavidin resin (Pierce, Rockford, IL, 1 h at 4 °C with rotation). The beads were washed with cold lysis buffer (3 \times), and samples were eluted with SDS sample buffer and resolved by SDS–PAGE prior to electrophoretic transfer to PVDF membranes (Immobilon P; Millipore, Billerica, MA). Chemiluminescent detection of anti-Shc (Santa Cruz Biotechnology) or anti-c-Met (AF276, R&D Systems, Minneapolis, MN) antibodies was performed using ECL (GE Healthcare).

Molecular Modeling. Calculations were performed on Silicon Graphics computer running the IRIX operating system. All simulations were performed using the molecular modeling tools available within MacroModel 9.1 (Schrödinger Inc.). The peptide GHDG-LyQGLSTATK (**1**) extracted from the Shc SH2 domain NMR structure (PDB code 1TCE) was used as a starting geometry for the modeling study. The structure of the bound ligand was modified using the 3D-sketcher tools available within MacroModel 9.1 to yield initial structures of peptide **50** complexed with the protein. The resulting protein–ligand complex was then subjected to minimization using the OPLS2005 method until a gradient convergence threshold of 0.05 was reached. This was followed by 10 ps of equilibration and 50 ps of molecular dynamics simulation using the OPLS2005 force field. During minimization, the phosphotyrosine and all atoms of the protein were held fixed. Two bound conformations of **50** were obtained and subjected to molecular dynamics simulations, during which the phosphotyrosine and all protein residues were held fixed except for those within a 10 Å sphere around the ligand. A total of 100 structures were saved for

each simulation, and the two conformations with the two lowest energy complexes are depicted in Figure 2. These two complexes were subjected to FEB (free energy of binding) calculations using the program eMBrAcE developed by Schrödinger. Both complexes were energetically similar (conformation A, −21 957.656 kJ/mol; conformation B, −21 957.133 kJ/mol).

General Synthetic Procedures. Except where noted, solid-phase peptide synthesis was performed using NovaSyn TGR resin (Novabiochem, Inc., catalog number 01-64-0060) in 5 or 10 mL disposable polypropylene columns (Pierce) fitted with porous polyethylene filter disks. Resin mixing was achieved by means of 360° rotation on a Barnstead-Thermolyne Labquake shaker. Protected-peptide resins were manually constructed using Fmoc-based solid-phase protocols. Trityl protection was employed for Gln and Ser side chain protection. The pTyr residue was incorporated in its PO(NMe₂)₂ form. Fmoc amino acids were coupled in NMP by treatment with Fmoc amino acid (5 equiv) and coupling reagents [DIPCDI (5 equiv)/HOBt (5 equiv) for 2 h; PyBroP (5 equiv)/DIPEA (10 equiv) for 2 h]. Double couplings were used for secondary amines. After completion of the coupling reactions, resins were washed (DMF, 6 \times), and then Fmoc deprotection was achieved using 20% piperidine in DMF (20 min) followed by DMF wash (6 \times). Final N-terminal acetylation was performed in DMF by treating with 1-acetylimidazole (for primary amines) or acetic anhydride/DIPEA (for secondary amines). Introduction of N-terminal fluorescein was achieved by reaction of deprotected resins with fluorescein isothiocyanate (“isomer 1”, Sigma Aldrich Corp) in DMF containing 3 equiv of diisopropylamine (shaken overnight and shielded from light). Finished resins were washed with DMF, methanol, dichloromethane, and ether and then dried under vacuum. Peptides and peptoid–peptide hybrids were cleaved from the resin by treatment (3 h) with 5 mL of trifluoroacetic acid/EDT/H₂O (76:20:4). An additional 0.5 mL of H₂O was added, and mixing was continued (overnight). Resin was removed by filtration, and the filtrate was concentrated under vacuum. Crude peptide precipitated by the addition of diethyl ether, and the precipitate was washed with diethyl ether (2 \times). The resulting solids were dissolved in aqueous acetonitrile and purified by reverse phase preparative HPLC using a Phenomenex C₁₈ column (21.20 mm diameter \times 250 mm) with a linear gradient from 0% aqueous acetonitrile (0.1% trifluoroacetic acid) to 100% acetonitrile (0.1% trifluoroacetic acid) over 30 min at a flow rate of 10.0 mL/min. Lyophilization provided products as amorphous solids.

FITC-NH-(CH₂)₄-CO-pTyr-Gln-Gly-Leu-Ser-amide (8**).** ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.09 (s, 1H), 9.93 (s, 1H), 8.25 (br, 1H), 8.15 (d, 1H), 8.06 (t, 1H), 7.93 (d, 1H), 7.81 (d, $J = 4$ Hz, 1H), 7.73 (d, $J = 4$ Hz, 1H), 7.17 (d, 1H), 7.11 (d, 1H), 7.01 (d, $J = 4$ Hz, 2H), 6.73 (br, 1H), 6.62 (d, 2H), 6.54–6.50 (m, 3H), 4.50–4.45 (m, 1H), 4.33–4.27 (m, 1H), 4.18 (dd, $J_1 = 8$ Hz, $J_2 = 4$ Hz, 1H), 4.13–4.10 (m, 1H), 3.76–3.67 (m, 2H), 3.53–3.46 (m, 2H), 3.39 (s, 2H), 2.98 (d, $J = 8$ Hz, 1H), 2.68–2.62 (m, 1H), 2.06 (dd, $J_1 = 4$ Hz, $J_2 = 4$ Hz, 4H), 1.88–1.83 (m, 1H), 1.77–1.70 (m, 1H), 1.51–1.59 (m, 1H), 1.46–1.37 (m, 2H), 0.82 (d, $J = 4$ Hz, 3H), 0.79 (d, $J = 8$ Hz, 3H). MALDI-MS calcd for C₅₁H₆₀N₉O₁₇PS: 1134.11; found 1134.82.

FITC-[3-aminomethylbenzoyl]-pTyr-Gln-Gly-Leu-Ser-amide (23**).** ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.11 (s, 1H), 10.07 (s, 1H), 8.53 (d, $J = 8$ Hz, 1H), 8.29 (d, $J = 8$ Hz, 1H), 8.25 (s, 1H), 8.15 (t, $J = 8$ Hz, 1H), 8.00 (d, 1H), 7.86 (d, 1H), 7.76–7.79 (m, 3H), 7.41 (d, $J = 8$ Hz, 2H), 7.31 (d, $J = 8$ Hz, 2H), 7.25 (br, 1H), 7.19 (d, $J = 4$ Hz, 2H), 7.07 (s, 1H), 6.76 (br, 1H), 6.68 (d, $J = 4$ Hz, 2H), 6.58 (s, 1H), 6.55 (d, $J = 4$ Hz, 2H), 4.84 (d, $J = 4$ Hz, 2H), 4.72–4.66 (m, 1H), 4.38–4.25 (m, 2H), 4.17 (dd, $J_1 = 4$ Hz, $J_2 = 8$ Hz, 1H), 3.77 (d, 1H), 3.73 (d, 1H), 3.60 (t, $J = 4$ Hz, 2H), 3.51 (s, 2H), 3.13 (d, $J = 4$ Hz, 1H), 2.96 (t, $J = 16$ Hz, 1H), 2.66–2.67 (m, 1H), 2.33–2.32 (m, 1H), 2.14 (t, $J = 8$ Hz, 2H), 1.93–1.90 (m, 1H), 1.82–1.77 (m, 1H), 1.62–1.57 (m, 1H), 1.48 (br, 1H), 0.87 (d, $J = 4$ Hz, 3H), 0.84 (d, $J = 2$ Hz, 3H). MALDI-MS calcd for C₅₄H₅₈N₉O₁₇PS: 1168.13; found 1168.71.

Peptides Bearing C-Terminal N-Alkenylamides (31**–**65**).** Peptides bearing C-terminal N-alkenylamides were prepared by adding the appropriate N-alkenylamine (10 equiv) and NaBH(OAc)₃ (10 equiv) to a suspension of 4-(4-formyl)-3-methoxyphenoxy)bu-

tyrily-NovaGel HL resin (Novabiochem, Inc.) in dry 1,2-dichloroethane/trimethyl orthoformate (2:1) solution, followed by mixing at room temperature (12 h). The resin was washed successively with DMF, 10% *i*-Pr₂NEt/DMF, and DMF. Sequential coupling of amino acids was then achieved according to the general synthetic procedures outlined above. For peptides bearing N^α-alkenyl or N^α-alkyl residues, introduction of N^α-alkenyl and N^α-alkyl groups was conducted on solid-phase resin by reaction of the corresponding N-terminally deprotected peptide with the appropriate alkenyl or alkyl bromide (1.2 equiv) and powdered K₂CO₃ in anhydrous DMF at room temperature (overnight). Resins were washed with DMF (2×), 50% aqueous MeOH (5×), and CH₂Cl₂ (3×), followed by coupling of remaining amino acid residues. Cleavage of final peptides from the resin afforded side-chain-deprotected peptide products, which were purified by reverse phase preparative HPLC.

Ac-pTyr-Gln-Gly(N-allyl)-Leu-NH(homoallyl) (44). ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.24 (dd, *J* = 8, 12 Hz, 1H), 7.97–7.91 (m, 2H), 7.20 (d, 1H), 7.09–7.05 (m, 2H), 6.98 (d, *J* = 8 Hz, 2H), 5.75–5.56 (m, 2H), 5.16–5.08 (m, 1H), 5.04–5.01 (m, 1H), 4.96–4.91 (m, 1H), 4.65–4.61 (m, 1H), 4.47–4.44 (m, 2H), 4.18–4.16 (m, 2H), 4.06–4.02 (m, 2H), 3.63–3.55 (m, 2H), 3.12 (s, 1H), 3.10–2.99 (m, 1H), 2.65–2.62 (m, 1H), 2.04–1.98 (m, 1H), 1.73 (s, 3H), 1.53–1.46 (m, 1H), 1.40–1.38 (m, 2H), 0.82 (d, *J* = 4 Hz, 3H), 0.77 (d, *J* = 8 Hz, 3H). MALDI-MS calcd for C₃₁H₄₇N₆O₁₀P: 694.71; found 717.25.

Ac-pTyr-Gln-Gly(N-ethyl)-Leu-amide (46). ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.05 (d, *J* = 8 Hz, 2H), 6.98 (d, *J* = 8 Hz, 2H), 6.91 (br, 2H), 4.65–4.64 (m, 1H), 4.47–4.11 (q, *J* = 8 Hz, 2H), 4.09–4.07 (m, 1H), 3.68–3.62 (m, 1H), 3.55–3.52 (m, 1H), 3.26–3.19 (m, 2H), 2.85–2.83 (m, 1H), 2.65–2.62 (m, 2H), 2.33–2.32 (m, 1H), 2.08–2.06 (m, 2H), 1.93 (d, *J* = 8 Hz, 1H), 1.76 (d, 2H), 1.74 (s, 3H), 1.55–1.53 (m, 1H), 1.07 (t, *J* = 8 Hz, 3H), 0.88–0.87 (m, 1H), 0.84 (d, *J* = 4 Hz, 3H), 0.78 (d, *J* = 4 Hz, 3H). MALDI-MS calcd for C₂₆H₄₁N₆O₁₀P: 628.61; found 650.52.

Acknowledgment. Appreciation is expressed to Dr. Oleg Chertov for purification of Shc SH2 domain protein. The work was supported in part by the Intramural Research Program of the NIH, Center for Cancer Research, NCI–Frederick, and the National Cancer Institute, National Institutes of Health, under Contract N01-CO-12400.

Supporting Information Available: Mass spectral data for peptide products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Wells, J. A.; McClendon, C. L. Reaching for high-hanging fruit in drug discovery at protein–protein interfaces. *Nature* **2007**, *450*, 1001–1009.
- (2) Bradshaw, J. M.; Waksman, G. Molecular recognition by SH2 domains. *Adv. Protein Chem.* **2003**, *61*, 161–210.
- (3) Pawson, T. Specificity in signal transduction: from phosphotyrosine-SH2 domain interactions to complex cellular systems. *Cell* **2004**, *116*, 191–203.
- (4) Pawson, T.; Gish, G. D.; Nash, P. The SH2 domain: a prototype for protein interaction modules. *Modular Protein Domains* **2005**, 5–36.
- (5) Ladbury, J. E. Protein–protein recognition in phosphotyrosine-mediated intracellular signaling. *Protein Rev.* **2005**, *3*, 165–184.
- (6) Machida, K.; Mayer, B. J. The SH2 domain: versatile signaling module and pharmaceutical target. *Biochim. Biophys. Acta, Proteins Proteomics* **2005**, *1747*, 1–25.
- (7) Broadbridge, R. J.; Sharma, R. P. The Src homology-2 domains (SH2 domains) of the protein tyrosine kinase p56lck: structure, mechanism and drug design. *Curr. Drug Targets* **2000**, *1*, 365–386.
- (8) Shakespeare, W. C. SH2 domain inhibition: a problem solved. *Curr. Opin. Chem. Biol.* **2001**, *5*, 409–415.
- (9) Metcalf, C. A., III; Sawyer, T. Src homology-2 domains and structure-based, small-molecule library approaches to drug discovery. *Drug Discovery Strategies Methods* **2004**, 23–59.
- (10) Garcia-Echeverria, C. Inhibitors of signaling interfaces: targeting Src homology 2 domains in drug discovery. *Protein Tyrosine Kinases* **2006**, 31–52.
- (11) Burke, T. R., Jr. Development of Grb2 SH2 domain signaling antagonists: a potential new class of antiproliferative agents. *Int. J. Pept. Res. Ther.* **2006**, *12*, 33–48.
- (12) Batzer, A. G.; Rotin, D.; Urena, J. M.; Skolnik, E. Y.; Schlessinger, J. Hierarchy of binding sites for Grb2 and Shc on the epidermal growth factor receptor. *Mol. Cell. Biol.* **1994**, *14*, 5192–201.
- (13) Zhou, M. M.; Harlan, J. E.; Wade, W. S.; Crosby, S.; Ravichandran, K. S.; Burakoff, S. J.; Fesik, S. W. Binding affinities of tyrosine-phosphorylated peptides to the COOH-terminal SH2 and NH2-terminal phosphotyrosine binding domains of Shc. *J. Biol. Chem.* **1995**, *270*, 31119–31123.
- (14) Pelicci, G.; Lanfrancone, L.; Grignani, F.; McGlade, J.; Cavallo, F.; Forni, G.; Nicoletti, I.; Grignani, F.; Pawson, T.; Pelicci, P. G. A novel transforming protein (SHC) with an SH2 domain is implicated in mitogenic signal transduction. *Cell* **1992**, *70*, 93–104.
- (15) Ravichandran, K. S. Signaling via Shc family adapter proteins. *Oncogene* **2001**, *20*, 6322–6330.
- (16) Saucier, C.; Khoury, H.; Lai, K. M. V.; Peschard, P.; Dankort, D.; Naujokas, M. A.; Holash, J.; Yancopoulos, G. D.; Muller, W. J.; Pawson, T.; Park, M. The Shc adaptor protein is critical for VEGF induction by Met/HGF and ErbB2 receptors and for early onset of tumor angiogenesis. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 2345–2350.
- (17) Saucier, C.; Papavasiliou, V.; Palazzo, A.; Naujokas, M. A.; Kremer, R.; Park, M. Use of signal specific receptor tyrosine kinase oncoproteins reveals that pathways downstream from Grb2 or Shc are sufficient for cell transformation and metastasis. *Oncogene* **2002**, *21*, 1800–1811.
- (18) Ursini-Siegel, J.; Hardy, W. R.; Zuo, D.; Lam, S. H. L.; Sanguin-Gendreau, V.; Cardiff, R. D.; Pawson, T.; Muller, W. J. ShcA signalling is essential for tumour progression in mouse models of human breast cancer. *EMBO J.* **2008**, *27*, 910–920.
- (19) Zhou, M.-M.; Meadows, R. P.; Logan, T. M.; Yoon, H. S.; Wade, W. S.; Ravichandran, K. S.; Burakoff, S. J.; Fesik, S. W. Solution structure of the Shc SH2 domain complexed with a tyrosine-phosphorylated peptide from the T-cell receptor. *Proc. Natl. Acad. Sci. U.S.A.* **1995**, *92* (17), 7784–7788.
- (20) Protein Data Bank code 1TCE.
- (21) See *Novabiochem Letters*, January 2004 (catalog number 04-12-3911).
- (22) Kertesz, A.; Varadi, G.; Toth, G. K.; Fajka-Boja, R.; Monostori, E.; Sarmay, G. Optimization of the cellular import of functionally active SH2-domain-interacting phosphopeptides. *Cell. Mol. Life Sci.* **2006**, *63*, 2682–2693.
- (23) Garcia-Echeverria, C.; Jiang, L.; Ramsey, T. M.; Sharma, S. K.; Chen, Y. N. P. A new Antennapedia-derived vector for intracellular delivery of exogenous compounds. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 1363–1366.
- (24) Blackwell, H. E.; Sadowsky, J. D.; Howard, R. J.; Sampson, J. N.; Chao, J. A.; Steinmetz, W. E.; O'Leary, D. J.; Grubbs, R. H. Ring-closing metathesis of olefinic peptides: design, synthesis, and structural characterization of macrocyclic helical peptides. *J. Org. Chem.* **2001**, *66*, 5291–5302.
- (25) Oishi, S.; Shi, Z.-D.; Worthy, K. M.; Bindu, L. K.; Fisher, R. J.; Burke, T. R., Jr. Ring-closing metathesis of allylglycines with β-vinyl-substituted residues as an approach to novel macrocyclic β-bend mimetics. *ChemBioChem* **2005**, *6*, 668–674.
- (26) Scholl, M.; Ding, S.; Lee, C. W.; Grubbs, R. H. Synthesis and activity of a new generation of ruthenium-based olefin metathesis catalysts coordinated with 1,3-dimesityl-4,5-dihydroimidazol-2-ylidene ligands. *Org. Lett.* **1999**, *1*, 953–956.
- (27) Mikol, V.; Baumann, G.; Zurini, M. G. M.; Hommel, U. Crystal structure of the SH2 domain from the adaptor protein SHC: a model for peptide binding based on X-ray and NMR data. *J. Mol. Biol.* **1995**, *254*, 86–95.
- (28) Oishi, S.; Karki, R. G.; Shi, Z.-D.; Worthy, K. M.; Bindu, L.; Chertov, O.; Esposito, D.; Frank, P.; Gillette, W. K.; Maderia, M.; Hartley, J.; Nicklaus, M. C.; Barchi, J. J., Jr.; Fisher, R. J.; Burke, T. R., Jr. Evaluation of macrocyclic Grb2 SH2 domain-binding peptide mimetics prepared by ring-closing metathesis of C-terminal allylglycines with an N-terminal beta-vinyl-substituted phosphotyrosyl mimetic. *Bioorg. Med. Chem.* **2005**, *13*, 2431–2438.
- (29) Lakowicz, J. R. *Principles of Fluorescence Spectroscopy*, 2nd ed.; Kluwer Academic/Plenum Publishers: New York, 1999.
- (30) Boor, C. D. *Principles of Fluorescence Spectroscopy*, 2nd ed.; Kluwer Academic/Plenum Publishers: New York, 1999.

JM800789H