

Asymmetric Synthesis of Four Diastereomers of 3-Hydroxy-2,4,6-trimethylheptanoic Acid: Proof of Configurational Assignment

Jeffrey A. Turk, Gonzalo S. Visbal, and Mark A. Lipton*

Department of Chemistry, Purdue University, West Lafayette, Indiana 47907-2084

lipton@purdue.edu

Received May 30, 2003

Four unique diastereomers of 3-hydroxy-2,4,6-trimethylheptanoic acid—(2*R*,3*R*,4*R*), (2*S*,3*R*,4*S*), and (2*R*,3*R*,4*S*)—the fatty acid component of callipeltin A and D, have been synthesized from commercially available (+)- and (−)-pseudoephedrine propionamide in 6 steps and 59% average overall yield. Comparison of the ¹H and ¹³C NMR and optical rotation data of the resulting isomers with the natural fragment unambiguously verifies the configurational assignment of the natural isomer as (2*R*,3*R*,4*R*).

Introduction

The structure of the novel marine cyclic depsipeptide callipeltin A (**1**), isolated from the lithistid sponge *Callipelta* sp. collected off the East coast of New Caledonia, was originally published by Minale and co-workers, and recently revised by D'Auria.^{1,2} Callipeltin A displays potent cytotoxicity (10 µg/mL, ca. 80% inhibition) against a broad range of human carcinoma cell lines,³ and displays both antiviral activity against HIV-1 (Lai strain)-infected CEM4 lymphocytes and antifungal activity against *Fusarium oxysporum*, *Helminthosporium sativum*, *Phytophthora hevea*, and *Candida albicans*.¹ Additionally, callipeltin A was found to be a selective and powerful inhibitor of the Na⁺/Ca²⁺ exchanger and a positive inotropic agent in guinea pig atria.⁴ Callipeltin A has recently also been isolated from the marine sponge *Latrunculia* sp. in the Vanuatu Islands along with a new, truncated derivative named callipeltin D (**2**) (Figure 1).⁵

Two asymmetric syntheses of (2*R*,3*R*,4*S*)-3-hydroxy-2,4,6-trimethylheptanoic acid (TMHEA, (R,R,S)-**3**), the originally identified diastereomer of the fatty acid component of callipeltin A and D, have been published.⁶ Comparison of the ¹H NMR spectrum of (R,R,S)-**3** to that of the corresponding fragment obtained from the acid hydrosylate of callipeltin A, however, revealed that the original assignment of configuration was incorrect. D'Auria

et al. revised the configuration as (2*R*,3*R*,4*R*), and supported their revision by correlation of the natural fragment with synthetic (R,R,R)-**3**.² Comparison of the natural fragment to *all* diastereomeric possibilities, however, would provide a complete and unambiguous verification of the configurational assignment of TMHEA. The availability of multiple configurational isomers may also expedite future analogue studies. We herein report the efficient synthesis of four unique diastereoisomers of TMHEA, thereby confirming the configurational assignment of D'Auria.

Results and Discussion

Our approach to the synthesis of (R,R,R)-**3** and its diastereomers uses asymmetric crotylboration to stereoselectively form the key C-2/C-3 bond. The substrate aldehyde and its enantiomer could in turn be made in high enantiomeric purity using the diastereoselective alkylation chemistry of Myers.⁷ Accordingly, commercially available pseudoephedrine propionamide (S,S)-**4** was alkylated with 1-iodo-2-methylpropane, using Myers's conditions,⁷ to give (S,S,R)-**5** as a highly viscous oil in 98% yield and 99% de (Scheme 1). Reduction of the resulting tertiary amide to the aldehyde, (R)-**6**, was expected to proceed smoothly with lithium triethoxyaluminum hydride.⁸ The reaction initially gave inconsistent results, with yields ranging from 50% to 70%, but when all traces of residual ethyl acetate were removed from (S,S,R)-**5** by coevaporation with toluene, aldehyde (R)-**6** was obtained reproducibly in 86% yield. A convenient alternative, albeit one that affords lower yields, involves conversion of the tertiary amide to the primary alcohol with lithium amidotrihydroborate⁹ (LAB) and subsequent

(1) Zampella, A.; D'Auria, M. V.; Paloma, L. G.; Casapullo, A.; Minale, L.; Debitus, C.; Henin, Y. *J. Am. Chem. Soc.* **1996**, *118*, 6202–6209.

(2) D'Auria, M. V.; Zampella, A. *Tetrahedron: Asymmetry* **2002**, *13*, 1237–1239.

(3) D'Auria, M. V.; Zampella, A.; Paloma, L. G.; Minale, L.; Debitus, C.; Roussakis, C.; Le Bert, V. *Tetrahedron* **1996**, *52*, 9589–9596.

(4) Trevisi, L.; Bova, S.; Cargnelli, G.; Danielli-Betto, D.; Floreani, M.; Germinario, E.; D'Auria, M. V.; Lucianni, S. *Biochem. Biophys. Res. Commun.* **2000**, *279*, 219–222.

(5) Zampella, A.; Randazzo, A.; Borbone, N.; Lucianni, S.; Trevisi, L.; Debitus, C.; D'Auria, M. V. *Tetrahedron Lett.* **2002**, *43*, 6163–6166.

(6) Joullie, M. M.; Carroll, P. J.; Guerlavais, V. *Tetrahedron: Asymmetry* **2002**, *13*, 675–680. D'Auria, M. V.; Zampella, A.; Sorgente, M. *Tetrahedron: Asymmetry* **2002**, *13*, 681–685.

(7) Myers, A. G.; Yang, B. H.; Chen, H.; McKinstry, L.; Kopecky, D. J.; Gleason, J. L. *J. Am. Chem. Soc.* **1997**, *119*, 6496–6511.

(8) Brown, H. C.; Tsukamoto, A. *J. Am. Chem. Soc.* **1964**, *86*, 1089.

(9) Myers, A. G.; Yang, B. H.; Kopecky, D. J. *Tetrahedron Lett.* **1996**, *37*, 3623–3626.






FIGURE 1. Structures of callipeltin A (**1**) and D (**2**).

SCHEME 1^a



^a Reaction conditions: (a) LDA, LiCl, THF, -78 °C; 1-iodo-2-methylpropane, -78 °C, 98%; (b) LiAlH(OEt)₃, hexanes-THF, 0 °C; TFA, 1 M HCl, 86%; (c) LAB, THF; (d) TPAP, NMO, 4 Å MS, CH₂Cl₂, 77% (two steps).

SCHEME 2^a



^a Reaction conditions: (a) *t*-BuOK, *trans*-2-butene, *n*-BuLi, -78 to -57 °C; (-)-*B*-methoxydiisopinocampheylborane; BF₃·Et₂O, (*R*)-**6**, -78 °C, 75%; (b) benzyl 2,2,2-trichloroacetimidate, TfOH (cat.), cyclohexane-CH₂Cl₂, 96%; (c) TBSCl, imidazole, CH₂Cl₂, 98%; (d) OsO₄, NaIO₄, NMO, dioxane-H₂O; NaClO₂, H₂NSO₃H, 95%; (e) NaIO₄/RuCl₃·H₂O (cat.), CCl₄-CH₃CN-H₂O, 98%.

reoxidation with TPAP,¹⁰ giving *(R)*-**6** in 77% yield over two steps. In the latter case, the enantiomeric purity of the intermediate alcohol was determined to be 98% by employing Mosher's method.¹¹

Treatment of *(R)*-6 with the crotylborane derived from *(-)*-*B*-methoxydiisopinocampheylborane and *trans*-2-butene¹² provided alcohol *(S,R,R)*-7 in 72% yield and 96% de (Scheme 2). Following chromatography, the diastereomeric excess of *(S,R,R)*-7 was determined to be greater than 98%. Protection of *(S,R,R)*-7 as the benzyl ether, by reaction with benzyl-2,2,2-trichloroacetimidate and catalytic triflic acid,¹³ gave *(S,R,R)*-8a in 96% yield.

Alternatively, **(S,R,R)-7** could be protected as its silyl ether by reaction with *tert*-butyldimethylsilyl chloride and imidazole,¹⁴ giving **(S,R,R)-8b**.

Treatment of (S,R,R) -**8b** with ruthenium chloride and sodium periodate¹⁵ produced carboxylic acid (R,R,R) -**9b** in nearly quantitative yield, but partial deprotection of the benzyl protecting group of (S,R,R) -**8a** during this reaction necessitated the use of an alternative method. Instead, treatment of (S,R,R) -**8a** with catalytic osmium tetroxide and stoichiometric sodium periodate followed by in situ oxidation of the resultant aldehyde with sodium chlorite produced (R,R,R) -**9a** in excellent yield (95%). Finally, hydrogenolysis of (R,R,R) -**9a** proceeded in quantitative yield, giving (R,R,R) -**3** in 58% overall yield from (S,S) -**4** (Scheme 3).

The remaining diastereomers were synthesized with similar stereoselectivities and chemical yields. Accord-

(10) Ley, S. V.; Norman, J.; Griffith, W. D.; Marsden, S. D. *Synthesis* **1994**, 6, 639–666.


(11) Dale, J. A.; Mosher, H. S. *J. Am. Chem. Soc.* **1973**, *95*, 512–519. Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. *J. Am. Chem. Soc.* **1991**, *113*, 4092–4096. Kusumi, T.; Fukushima, T.; Ohtani, I.; Kakisawa, H. *Tetrahedron Lett.* **1991**, *32*, 2939–2942.

(12) Brown, H. C.; Bhat, K. S.; Randad, R. S. *J. Org. Chem.* **1989**, 54, 1570–1576.

(13) Iversen, T.; Bundle, D. R. *J. Chem. Soc., Chem. Commun.* **1981**, 23, 1240–1241.

(14) Corey, E. J.; Cho, H.; Rucker, C.; Hua, D. H. *Tetrahedron Lett.* **1981**, 22, 3455–3458.

(15) Carlsen, P. H. J.; Katsuki, T.; Martin, V. S.; Sharpless, B. *J. Org. Chem.* **1981**, *46*, 3936–3938.

SCHEME 3^a

^a Reaction conditions: (a) H₂/Pd(OH)₂-C, formic acid, MeOH.

TABLE 1. Comparison of the ¹H NMR Chemical Shifts^a (δ) of Synthetic 3-Hydroxy-(2,4,6)-trimethylheptanoic Acid Diastereomers with the Natural Fragment

natural fragment ^b δ , multiplicity (J), H	(R,R,R)-3 δ , multiplicity (J), H	(S,R,R)-3 δ , multiplicity (J), H	(S,R,S)-3 δ , multiplicity (J), H	(R,R,S)-3 δ , multiplicity (J), H
0.84, d (6.8), 3H	0.84, d (6.5), 3H	0.85, d (6.4), 3H	0.84, d (6.5), 3H	0.85, d (6.6), 3H
0.92, d (6.8), 3H	0.92, d (6.6), 3H	0.90, d (6.5), 6H	0.89, d (6.5), 3H	0.86, d (6.5), 3H
0.96, d (5.8), 3H	0.95, d (6.8), 3H		0.95, d (6.5), 3H	0.89, d (6.6), 3H
1.12–1.18, m, 2H	1.12–1.18, m, 2H	1.04, m, 1H	1.07, m, 2H	1.16, m, 1H
1.26, d (7.7), 3H	1.23, d (7.2), 3H	1.22, d (7.1), 3H	1.22, d (7.0), 3H	1.18, d (6.7), 3H
		1.49, m, 1H		1.24, m, 1H
1.64, m, 1H	1.64, m, 1H	1.62, m, 1H	1.66, m, 2H	1.66, m, 1H
1.70, m, 1H	1.69, m, 1H	1.66, m, 1H		1.70, m, 1H
2.73, quintet (7.7), 1H	2.74, dq (7.1, 7.1), 1H	2.74, dq (3.5, 3.5), 1H	2.72, dq (6.0, 6.0), 1H	2.66, dq (7.5, 7.5), 1H
3.48, dd (4.1, 7.7), 1H	3.48, dd (6.4, 6.4), 1H	3.67, dd (3.4, 7.7), 1H	3.68, dd (5.5, 5.5), 1H	3.61, dd (3.3, 8.3), 1H

^a All chemical shifts are reported in ppm and were measured in CDCl₃. Coupling constants (J) are expressed in Hz. ^b Spectral data for natural fragment are reproduced from ref 1.

ingly, (S,R,R)-3 was synthesized from (S,S)-4 and B-(Z)-2-butenyl-(-)-diisopinocampheylborane in 56% overall yield. (R,R,S)-3 (56%) and (S,R,S)-3 (62%) were obtained in a similar fashion from (R,R)-4 and the (E)- or (Z)-crotylboranes, respectively, derived from (-)-B-methoxyl-diisopinocampheylborane.

Proof of Configurational Assignment

The configuration of TMHEA at C-3 was determined by the modified Mosher method, and unambiguously assigned as *R*.^{1,11} Accordingly, (R,R,R)-3, (S,R,R)-3, (S,R,S)-3, and (R,R,S)-3 were synthesized for our study, as they represent the four possible diastereomers of 3-hydroxy-2,4,6-trimethylheptanoic acid (TMHEA) that have the *R* configuration at C-3.

Comparison of the ¹H NMR spectrum of the TMHEA obtained from the acid hydrolysis of callipeltin A with those of the four synthetic analogues unambiguously indicates the configurational assignment of the natural fragment as either *R,R,R* or *S,S,S* (Table 1). Aside from differences in the chemical shift values of the methyl signals of (R,R,S)-3 and the natural fragment, the ¹H NMR spectrum of (R,R,S)-3 displays two 1H signals (1.16 and 1.24 ppm) where the natural fragment displays a single broad 2H signal (1.12–1.18 ppm). The ¹H NMR spectrum of (S,R,R)-3 displays similar deviations in the methylene region, and the presence of two overlapping methyl signals at 0.90 ppm distinguishes this diastereomer as distinct from the natural isomer. (S,R,S)-3 differs

from that of the natural fragment by the presence of a 2H signal at 1.66 ppm instead of the two 1H signals (1.64 and 1.70 ppm) observed for the natural fragment. The most compelling evidence for the ¹H NMR assignment of the natural fragment is the chemical shift and coupling constant (J) differences observed for the C-2 methine proton. Despite minor differences in the vicinal coupling constants, the spectrum of (R,R,R)-3 matches that of the natural material far better than those of the other isomers.

Although ¹³C NMR spectroscopy should also confirm the configurational assignment, this comparison is not straightforward: the carbon chemical shift values for the natural TMHEA fragment were not available from the acid hydrolysis product, so instead were obtained from the ¹³C NMR spectrum of intact callipeltin A.¹ Because none of the spectra of the four isomers of 3 unambiguously match the chemical shifts of the natural fragment, it is impossible to make any conclusions regarding stereochemical assignment based on the ¹³C NMR data available.

Comparison of the specific rotation data of the four synthetic isomers of TMHEA (Scheme 3) with that of the natural fragments, $[\alpha]_D^{25} +14.9$ (c 0.4, CHCl₃), supports the configurational assignment of *R,R,R* or *S,R,R*.¹⁶

(16) The difference in the optical rotation of the natural fragment, $[\alpha]_D^{25} +14.9$ (c 0.4, CHCl₃), and the synthetic analogue, $[\alpha]_D^{25} +25.6$ (c 0.4, CHCl₃), indicated that (R,R,R)-3 was of a higher chemical purity than the crude acid hydrolysate, yet supporting ¹H NMR data verifies the assignment as (R,R,R).

Combining this fact with the conclusions drawn from the ¹H NMR spectra unambiguously defines the assignment of the 3-hydroxy-2,4,6-trimethylheptanoyl residue of calipeltin A as *R,R,R*.

Conclusion

In summary, we have synthesized four diastereomers of TMHEA (6 steps, 59% average yield), the fatty acid fragment of callipeltin A and D, using an efficient and flexible route. Comparison of the ¹H NMR, ¹³C NMR, and optical rotation data of the natural fragment with all its diastereomers unambiguously verifies the configurational assignment of the natural isomer as *R,R,R*.

Experimental Section

(R)-2,4-Dimethylpentanal (6). A 50-mL round-bottomed flask was charged with lithium aluminum hydride (95%, 262 mg, 6.55 mmol) and hexanes (6 mL), and the resulting suspension was cooled to 0 °C. Ethyl acetate (960 μ L, 9.83 mmol) was added dropwise over 1.5 h and the reaction mixture was cooled to -78 °C. A solution of (*S,S,R*)-5 (790 mg, 2.85 mmol) in THF (3 mL) was added over 5 min, and the reaction mixture was stirred at 0 °C for 1 h and transferred to a vigorously stirring solution of 1 M HCl (50 mL) and TFA (2.5 mL). After the mixture was stirred for 10 min at room temperature, 1 M HCl (65 mL) was added, and the solution was extracted with diethyl ether. The combined organic layers were neutralized by the slow addition of saturated aqueous NaHCO₃. The aqueous layer was separated and extracted with diethyl ether. The combined organic layers were dried (MgSO₄), passed through a short plug of silica, and concentrated to give (*R*)-6 (280 mg, 86%) as a volatile, colorless liquid that was used in the following step without further purification: ¹H NMR (500 MHz, CDCl₃) δ 9.60 (d, *J* = 1.9 Hz, 1H), 2.41 (ddq, *J* = 4.2, 4.2, 4.2 Hz), 1.63 (m, 1H), 1.59 (m, 1H), 1.20 (m, 1H), 1.08 (d, *J* = 6.9 Hz, 3H), 0.93 (d, *J* = 6.5 Hz, 3H), 0.90 (d, *J* = 6.5 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 205.4, 44.4, 39.7, 25.5, 22.9, 22.2, 20.6, 13.7.

(3*S,4R,5R*)-3,5,7-Trimethylocten-4-ol (7). A solution of potassium *tert*-butoxide in THF (1.0 M, 4.89 mL, 4.89 mmol) and THF (5 mL) was cooled to -78 °C, and *trans*-2-butene (0.60 mL, 6.3 mmol) was added, followed by *n*-butyllithium (2.50 M, 1.96 mL, 4.73 mmol). The resulting bright yellow solution was stirred at -78 °C for 2 min and -57 °C for 10 min. The reaction was re-cooled to -78 °C, and a solution of (-)-*B*-methoxydiisopinocampheylborane (1.55 g, 4.89 mmol) in THF (3 mL) was added. The reaction was stirred for 1 h, and BF₃·OEt₂ (0.614 mL, 4.89 mmol) was added followed by a pre-cooled (-78 °C) solution of (*R*)-6 (360 mg, 3.15 mmol) in THF (3 mL). The resulting solution was stirred at -78 °C for 4 h, and the reaction was quenched by the slow addition of aqueous

sodium hydroxide (3 M, 2 mL) and hydrogen peroxide (30%, 1 mL). The partially frozen mixture was removed from the cold bath and stirred for 18 h. The reaction was diluted with H₂O and extracted with diethyl ether. The combined organic extracts were dried and concentrated to give a colorless oil that was purified by flash chromatography to yield (*S,R,R*)-7 as a colorless oil (400 mg, 75%, 96% de): ¹H NMR (500 MHz, CDCl₃) δ 5.76 (m, 1H), 5.12 (m, 2H), 3.13 (t, *J* = 6.0 Hz, 1H), 2.37 (ddq, *J* = 7.0, 7.0, 7.0 Hz, 1H), 1.65 (m, 2H), 1.26 (m, 1H), 1.13 (m, 1H), 1.00 (d, *J* = 6.9 Hz, 3H), 0.92 (d, *J* = 6.7 Hz, 3H), 0.91 (d, *J* = 6.5 Hz, 3H), 0.86 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 140.7, 116.2, 79.5, 41.2, 39.5, 33.0, 25.3, 24.4, 21.3, 16.9, 16.8; IR (CHCl₃) 3048, 2959, 1705, 1462 cm⁻¹; $[\alpha]$ +21.8 (c 0.7, CHCl₃). Anal. Calcd for C₁₁H₂₂O·0.25H₂O: C, 75.58; H, 12.98. Found: C, 75.32; H, 13.35.

(2*R,3R,4R*)-2,4,6-Trimethyl-3-benzyloxyheptanoic Acid (9a). To a stirring solution of (*S,R,R*)-8a (617 mg, 2.37 mmol) in dioxane (3.5 mL) and H₂O (3.5 mL) was added 4-methylmorpholine N-oxide (416 mg, 3.55 mmol) and osmium tetroxide (4% in H₂O, 302 μ L, 95 μ mol). After 1 h, NaIO₄ (759 mg, 3.55 mmol) was added, and the suspension was stirred at room temperature for 1.5 h. The reaction mixture was cooled to 0 °C, and sodium chlorite (857 mg, 9.48 mmol) and sulfamic acid (920 mg, 9.48 mmol) were added, and the resulting bright yellow mixture was removed from the cold bath and stirred for 2 h. To the mixture was added 5% aqueous HCl, and the resulting solution was extracted with CH₂Cl₂. The combined organic extracts were washed with 5% aqueous HCl, dried over MgSO₄, concentrated, and purified by flash chromatography to give (*R,R,R*)-9a as a colorless oil (634 mg, 96%): ¹H NMR (500 MHz, CDCl₃) δ 7.31–7.26 (m, 5H), 4.58 (d, *J* = 11.2 Hz, 1H), 4.55 (d, *J* = 11.5 Hz, 1H), 3.48 (dd, *J* = 4.1, 6.8 Hz, 1H), 2.77 (dq, *J* = 7.1, 7.1 Hz, 1H), 1.81 (m, 1H), 1.59 (m, 1H), 1.17 (m, 2H), 1.17 (d, *J* = 7.1 Hz, 3H), 0.94 (d, *J* = 6.9 Hz, 3H), 0.87 (d, *J* = 6.5 Hz, 3H), 0.80 (d, *J* = 6.5 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 179.9, 138.3, 128.4, 128.3, 127.7, 127.7, 127.6, 86.1, 74.5, 42.3, 40.6, 32.9, 25.2, 24.0, 21.4, 16.3, 14.7; IR (CHCl₃) 3055, 2960, 1752, 1709, 1457 cm⁻¹; $[\alpha]$ ²⁵_D +14.9 (c 0.2, CHCl₃). Anal. Calcd for C₁₇H₂₆O₃·0.5H₂O: C, 71.49; H, 9.46. Found: C, 71.27, H, 9.71.

Acknowledgment. We thank Prof. Maria D'Auria for helpful discussions and spectra of the natural product. We also gratefully acknowledge the National Institute of Allergy and Infectious Diseases (NIH AI-50888) for financial support.

Supporting Information Available: Complete experimental descriptions of transformations not included in the Experimental Section and characterization for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO034738L