


Pyrrolidine and Piperidine Formation via Copper(II) Carboxylate-Promoted Intramolecular Carboamination of Unactivated Olefins: Diastereoselectivity and Mechanism

Eric S. Sherman, Peter H. Fuller, Dhanalakshmi Kasi, and Sherry R. Chemler*

Department of Chemistry, The University at Buffalo, The State University of New York,
Buffalo, New York 14260

schemler@buffalo.edu

Received February 15, 2007



An expanded substrate scope and in-depth analysis of the reaction mechanism of the copper(II) carboxylate-promoted intramolecular carboamination of unactivated alkenes is described. This method provides access to *N*-functionalized pyrrolidines and piperidines. Both aromatic and aliphatic γ - and δ -alkenyl *N*-arylsulfonamides undergo the oxidative cyclization reaction efficiently. *N*-Benzoyl-2-allylaniline also underwent the oxidative cyclization. The terminal olefin substrates examined were more reactive than those with internal olefins, and the latter terminated in elimination rather than carbon–carbon bond formation. The efficiency of the reaction was enhanced by the use of more organic soluble copper(II) carboxylate salts, copper(II) neodecanoate in particular. The reaction times were reduced by the use of microwave heating. High levels of diastereoselectivity were observed in the synthesis of 2,5-disubstituted pyrrolidines, wherein the *cis* substitution pattern predominates. The mechanism of the reaction is discussed in the context of the observed reactivity and in comparison to analogous reactions promoted by other reagents and conditions. Our evidence supports a mechanism wherein the N–C bond is formed via intramolecular syn aminocupration and the C–C bond is formed via intramolecular addition of a primary carbon radical to an aromatic ring.

Introduction

Nitrogen heterocycles make up a significant proportion of biologically active small organic molecules. Recently developed methods for the synthesis of nitrogen heterocycles by transition metal facilitated intramolecular amine additions onto unactivated alkenes have expanded the repertoire of tools available to the medicinal chemist.^{1–25} Methods that provide for concise build-up of functionality by installing two rings in a single operation


may prove especially useful in the concise synthesis of nitrogen heterocycles for natural product synthesis and drug discovery endeavors.^{11,14,15,26–32} Herein is reported an expansion of the substrate scope of the copper(II)-promoted intramolecular car-

- (1) Müller, T. E.; Beller, M. *Chem. Rev.* **1998**, 98, 675–703.
- (2) Zeni, G.; Larock, R. C. *Chem. Rev.* **2004**, 104, 2285–2309.
- (3) Nakamura, I.; Yamamoto, Y. *Chem. Rev.* **2004**, 104, 2127–2198.
- (4) Hong, S.; Marks, T. J. *Acc. Chem. Res.* **2004**, 37, 673–686.
- (5) Hegedus, L. S., In *Comprehensive Organic Synthesis*; Trost, B. M., Ed.; Pergamon: New York, 1991; Vol. 4, Chapter 3.1.


- (6) Larock, R. C.; Yang, H.; Weinreb, S. M.; Herr, R. J. *J. Org. Chem.* **1994**, 59, 4172–4178.
- (7) Overman, L. E.; Remarchuk, T. P. *J. Am. Chem. Soc.* **2002**, 124, 12–13.
- (8) Lutete, L. M.; Kadota, I.; Yamamoto, Y. *J. Am. Chem. Soc.* **2004**, 126, 1622–1623.
- (9) Fix, S. R.; Brice, J. L.; Stahl, S. S. *Angew. Chem., Int. Ed.* **2002**, 41, 164–166.
- (10) Trend, R. M.; Ramtohl, Y. K.; Ferreira, E. M.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2003**, 42, 2892–2895.
- (11) Sherman, E. S.; Chemler, S. R.; Tan, T. B.; Gerlits, O. *Org. Lett.* **2004**, 6 (10), 1573–1575.

boamination of unactivated alkenes,¹¹ a transformation that installs two new rings from acyclic *N*-substituted amines.

We recently reported that copper(II) acetate promotes the oxidative cyclization of *N*-arylsulfonyl-2-allylanilines **1** (eqs 1 and 2).¹¹ In these reactions, aryl sulfonamides with electron donating groups proved most reactive: 4-methyl, methoxy, chloro and bromoaryl sulfonamides reacted efficiently albeit the bromide was removed under the reaction conditions (eqs 1 and 2). 4-Nitro and 4-trifluoromethyl arylsulfonamides displayed




R	yield (%)
Me	73
OMe	63
Br	54 (R = H)
NO ₂	24
CF ₃	no rxn.



R	yield (%)	ortho : para
Me	67	2.3 : 1
OMe	62	2.7 : 1
Cl	60	1.8 : 1

significantly lower reactivity (eq 1). (Similar electronic effects have been observed in Brønsted acid catalyzed intramolecular hydroamination reactions.³³) Meta-substituted aryl sulfonamides demonstrated a preference (ca. 2:1) for the ortho addition

TABLE 1. Reaction Conditions: Carboxylate, Solvent, Temperature^a



entry	R	solvent	temp, time	yield (%) ^d
1	OAc	DMF	120 °C, ^b 24 h	69
2	OAc	CH ₃ CN	120 °C, ^b 24 h	73
3	OAc	<i>i</i> -PrOH	120 °C, ^b 24 h	16 ^e
4	OAc	<i>t</i> -amyl-OH	120 °C, ^b 24 h	33
5	OAc	EtOAc	120 °C, ^b 24 h	24 ^e
6	OAc	toluene	120 °C, ^b 24 h	51 ^e
7	OAc	DMA	120 °C, ^b 24 h	37
8	OPiv	DMF	120 °C, ^b 24 h	69
9	EH	DMF	120 °C, ^b 24 h	64
10	ND	DMF	120 °C, ^b 24 h	69
11	ND	toluene	120 °C, ^b 24 h	71
12	ND	DMF	120 °C, ^b 0.5 h	29 ^e
13	ND	DMF	120 °C, ^c 0.5 h	27 ^e
14	ND	DMF	160 °C, ^b 0.5 h	63
15	ND	DMF	160 °C, ^c 0.5 h	64

^a All reactions were run with 3 equiv of copper(II) carboxylate, 1 equiv of Cs₂CO₃ at 0.1 M **1a** concentration. ^b Oil bath heating, pressure tube. ^c Microwave heating. ^d Amount isolated after flash chromatography on SiO₂. ^e Remainder of the material is starting **1a**. In entry 3 another cyclization product (net aminoetherification, incorporation of *i*-PrOH in the product) was also formed in 21% yield (see the Supporting Information). Ac = COCH₃, Piv = COC(CH₃)₃, neodecanoate (ND) = OCO(CH₂)₅C(CH₃)₃, ethylhexanoate (EH) = OCOCH(C₂H₅)C₄H₉.

product over the para adduct (eq 2). This surprising ortho preference (the more sterically hindered site) indicated that C–C bond formation may occur via addition of a carbon radical to the aromatic ring [intermolecular additions of radicals to aromatics have shown a slight preference (ca. 2:1) for the ortho adduct].^{34,35}

Results and Discussion

To further expand the utility of this method, several copper(II) carboxylate salts, solvents, and reaction heating methods were examined. Our former optimized reaction conditions for the carboamination reaction of **1a** (R = Me) required DMF or CH₃CN solvent at 120 °C for 24 h (Table 1, entries 1 and 2). The need for polar solvents and additives (added DMSO increased the yield with some substrates) was attributed to the poor solubility of Cu(OAc)₂ in organic solvents. We therefore examined the reaction further by using more polar solvents with Cu(OAc)₂ and more organic soluble copper carboxylates with nonpolar solvents, respectively. The reaction did proceed in both *i*-PrOH and *tert*-amyl alcohol (Table 1, entries 4 and 5) although in the former case a substantial amount of another product was obtained (net aminoetherification, see the Supporting Information) while in the latter case a more efficient, albeit not optimal carboamination process occurred. Several copper(II) carboxylates were also compared. We found that while the use of Cu-

(31) Senboku, H.; Kajizuka, Y.; Hasegawa, H.; Fujita, H.; Sugimoto, H.; Orito, K.; Tokuda, M. *Tetrahedron* **1999**, *55*, 6465–6474.

(32) Hasegawa, H.; Senboku, H.; Kajizuka, Y.; Orito, K.; Tokuda, M. *Tetrahedron* **2003**, *59*, 827–832.

(33) Yin, Y.; Zhao, G. *Heterocycles* **2006**, *68* (1), 23–31.

(34) Ito, R.; Migita, T.; Morikawa, N.; Simamura, O. *Tetrahedron* **1965**, *21*, 955–961.

(35) Pryor, W. A.; Davis, W. H.; Gleaton, J. H. *J. Org. Chem.* **1975**, *40*, 2099–2102.

(12) Manzoni, M. R.; Zabawa, T. P.; Kasi, D.; Chemler, S. R. *Organometallics* **2004**, *23*, 5618–5621.

(13) Lei, A.; Lu, X.; Liu, G. *Tetrahedron Lett.* **2004**, *45*, 1785–1788.

(14) Zabawa, T. P.; Kasi, D.; Chemler, S. R. *J. Am. Chem. Soc.* **2005**, *127*, 11250–11251.

(15) Streuff, J.; Hovelmann, C. H.; Nieger, M.; Muniz, K. *J. Am. Chem. Soc.* **2005**, *127*, 14586–14587.

(16) Alexanian, E. J.; Lee, C.; Sorensen, E. J. *J. Am. Chem. Soc.* **2005**, *127*, 7690–7691.

(17) Donohoe, T. J.; Chughtai, M. J.; Klauber, D. J.; Griffin, D.; Campbell, A. D. *J. Am. Chem. Soc.* **2006**, *128*, 2514–2515.

(18) Donohoe, T. J.; Johnson, P. D.; Pye, R. J.; Keenan, M. *Org. Lett.* **2004**, *6*, 2583–2585.

(19) Lira, R.; Wolfe, J. P. *J. Am. Chem. Soc.* **2004**, *126*, 13906–13907.

(20) Ney, J. E.; Wolfe, J. P. *Angew. Chem., Int. Ed.* **2004**, *43*, 3605–3608.

(21) Bertrand, M. B.; Wolfe, J. P. *Tetrahedron* **2005**, *61* (26), 6447–6459.

(22) Hiroya, K.; Itoh, S.; Sakamoto, T. *Tetrahedron* **2005**, *61*, 10958–10964.

(23) Fallis, A. G.; Brinza, I. M. *Tetrahedron* **1997**, *53*, 17543–17594.

(24) Bender, C. F.; Widenhoefer, R. A. *Org. Lett.* **2006**, *8*, 5303–5305.

(25) Donohoe, T. J.; Churchill, G. H.; Wheelhouse, K. M. P.; Glossop, P. A. *Angew. Chem., Int. Ed.* **2006**, *45*, 8025–8028.

(26) Nakhla, J. S.; Kampf, J. W.; Wolfe, J. P. *J. Am. Chem. Soc.* **2006**, *128*, 2893–2901.

(27) Yip, K.-T.; Yang, M.; Law, K.-L.; Zhu, N.-Y.; Yang, D. *J. Am. Chem. Soc.* **2006**, *128*, 3130–3131.

(28) Molander, G. A.; Pack, S. K. *Tetrahedron* **2003**, *59*, 10581–10591.

(29) Sharp, L. A.; Zard, S. Z. *Org. Lett.* **2006**, *8*, 831–834.


(30) Banwell, M. G.; Lupton, D. W. *Heterocycles* **2006**, *68* (1), 71–92.

TABLE 2. Expanded Sulfonamide Substrate Scope^a

entry	substrate	product(s) ^b	temp; time	yield (%) ^c
1		 	120 °C; 24 h	49 exo (9) 46 endo (10)
2		 	160 °C; 72 h 160 °C; 72 h	23 CA (12) ^d 59 CA (12) + 21 HA (13)
4			160 °C; 72 h	87 ^d
5		 	210 °C, 72 h 210 °C; μ W, 3 h	56 CA (17 , <i>trans</i> : <i>cis</i> = 1 : 1), 16 HA (18 , <i>trans</i> : <i>cis</i> = 1 : 1) 52 CA (17 , <i>trans</i> : <i>cis</i> = 1 : 1), 20 HA (18 , <i>trans</i> : <i>cis</i> = 1 : 1)
7		 	180 °C, 72 h	65 CA (20 , <i>trans</i> : <i>cis</i> = 3 : 1) 33 HA (21 , <i>trans</i> : <i>cis</i> = 3 : 1)
8			200 °C; 72 h	48
9		 	200 °C; 72 h 210 °C, μ W, 3 h	53 CA (25 only) 55 CA (25) + 14 HA (26)
11			200 °C; 72 h	43

^a Reaction conditions: 3 equiv of Cu(ND)₂, 1 equiv of Cs₂CO₃, DMF (0.08–0.1 M), temp, time. ^b Selectivity determined by analysis of the crude ¹H NMR spectrum, and by the amount of isolated adducts. ^c Yield refers to the amount of product isolated by chromatography on silica gel. ^d Reaction run with Cu(OAc)₂ instead of Cu(ND)₂. CA = carboamination product, HA = hydroamination product, ND = neodecanoate.

SCHEME 1. Mechanism for Formation of the Carboamination and Hydroamination Adducts




(OAc)₂ in toluene did promote oxidative cyclization (51% yield + remaining starting material, entry 6, Table 1), the more organic soluble copper(II) neodecanoate [Cu(ND)₂] provided a more efficient reaction (71% yield, entry 11, Table 1). Copper(II) neodecanoate also provided a more efficient reaction with

more entropically challenging substrates (Table 2, *vide infra*). Copper(II) pivalate and copper(II) 2-ethylhexanoate also provided efficient reaction in DMF (entries 8 and 9) and under these conditions are comparable to Cu(ND)₂ and Cu(OAc)₂. On the basis of these examples, the relative steric demand of the

carboxylate alkyl chain does not affect the reactivity of the carboxylate salt in these reactions.

Increasing the reaction temperature decreased the time required for product formation. As judged by the crude ^1H NMR spectra, at 160 °C the reaction was complete after 0.5 h (63% isolated yield of **2**) while only a 29% yield of **2** was obtained at 120 °C for 0.5 h [using $\text{Cu}(\text{ND})_2$ in DMF], the remaining material being unreacted starting material (compare entries 12 and 14). At these reaction scales with substrate **1a** (ca. 30 mg of **1a**, 1 mL of solvent) we did not find a notable difference in reactivity between microwave and oil bath heating (compare entry 12 to 13 and entry 14 to 15). This may be due in part to the fact that both reactions are run in sealed tubes, thus, in both cases, the reaction solutions are subject to superheating and also the transfer of heat due to surface area heating is not a significant problem with small-scale reactions.³⁶

Effect of the Nitrogen Substituent. Although arylsulfonamides are superior substrates for this reaction (lower reaction temperatures required, higher yield), the *N*-benzoyl-2-allylaniline **6** also provides the oxidative cyclization product **7** (eq 3, ND = neodecanoate). Formation of fused carbon/nitrogen ring systems allows entry into a broader range of nitrogen heterocycles and further examination of the copper carboxylate-promoted reactions with such substrates is warranted.



Effect of Chain Length and Substitution. In an effort to expand the substrate scope of the copper(II) carboxylate-promoted carboamination reaction of sulfonamides as well as gain more insight into the reaction mechanism, the oxidative cyclizations of a number of substrates were explored, including arylsulfonamides derived from aliphatic amines, substrates with different olefin substitution patterns, and substrates containing chiral centers (Table 2). Both microwave and oil bath heating methods were used. Both five-membered-ring pyrrolidines and six-membered-ring piperidines can be formed through this oxidative cyclization protocol. All of the substrates examined in this reaction give exclusively the exo cyclization adduct except for the 1,1-disubstituted olefin **8**, which gave a ca. 1:1 mixture of the exo carboamination adduct **9** and the endo oxidative amination adduct **10** (Table 2, entry 1).


Sulfonamides **11**, **14**, **16**, and **19**, derived from primary aliphatic amines (entries 2–7), undergo cyclization efficiently although the *gem*-dimethyl-substituted substrate **14** demonstrated the highest reactivity.³⁷ The more organic soluble $\text{Cu}(\text{ND})_2$ had a significant effect on the reaction yields with less reactive substrates (compare entries 2 and 3, Table 2). The formation of hydroamination minor products was also observed with some substrates and the carboamination to hydroamination ratio is presumably a function of the relative rates of radical addition to the aromatic ring vs hydrogen atom abstraction from the reaction medium (vide infra). Reaction of the 2-phenyl-pent-

4-enyl sulfonamide **16** produced 1:1 diastereomeric mixtures of carboamination and hydroamination products, **17** and **18**, respectively (entries 5 and 6, ratio of **17**:**18** = 3.5:1 in an oil bath at 210 °C). In the case of substrate **19**, with a methylcyclohexyl group at the allylic position, a 3:1 mixture of 2,3-trans:2,3-cis diastereomers of carboamination and also 3:1 2,3-trans:2,3-cis hydroamination adducts, **20** and **21**, were formed (ratio of carboamination:hydroamination = 2:1). Cyclization reactions of substrates with allylic substitution under several other reaction conditions also favor the formation of the anti adduct, presumably due to reaction via a chairlike transition state that favors placement of the allylic alkyl substituent in a pseudoequatorial position.^{13,20} The need for high temperature in the copper(II) carboxylate-promoted cyclization may account for the modest level of asymmetric induction observed.

The carboamination adducts derived from styrenyl substrates **22** and **27** were obtained in modest yield upon heating at 200–210 °C (entries 8 and 11). These substrates are challenging due to the ground state resonance stabilization that is lost in the cyclization transition state. Few intramolecular carboamination protocols have been reported with styrenyl substrates.²⁸ Formation of six-membered rings is also possible via the copper(II)-promoted carboamination protocol (entries 9–11, Table 2).

The products formed in these oxidative cyclization reactions are consistent with a mechanism that involves formation of a radical intermediate (e.g., **29**) that may either perform intramolecular addition onto the neighboring aromatic ring or abstraction of a proton from the reaction medium (Scheme 1). In the case of endo cyclization adduct **10**, the secondary radical undergoes $\text{Cu}(\text{II})$ -facilitated elimination to form an alkene.^{38,39}

The mechanism was further probed by using the carboamination reaction of deuterioalkene **14-D** (eq 4). The partial conversion of **14-D** provides a 1:1 mixture of diastereomers **15-D**, indicating the presence of an sp^2 -hybridized carbon intermediate, likely a primary radical. The remaining deuterioalkene was recovered with complete retention of stereochemistry, indicating that the carbon radical intermediate does not revert back to the alkene.




When an unambiguous primary radical was generated from the primary bromide **31** (AIBN, HSnBu_3 , PhH, reflux), a mixture of carboamination adducts **4** and **5** (ortho:para = 2.7:1) and hydroamination adduct **32** was obtained (eq 5). The fact that the carboamination products were obtained in the same ortho:para ratio (2.7:1) as observed in the copper(II)-promoted reaction (see eq 2) is compelling evidence that a similar intermediate, a primary carbon radical, is involved in the C–C bond formation. No *ipso* (1,5-addition) substitution of the radical onto the aromatic ring was observed. Mixtures of *ipso* (1,5) addition products and direct (1,6) addition products along with the direct reduction products have previously been observed in the intramolecular radical reactions of *N*-arylsulfonyl-2-halometh-

(36) Larhed, M.; Moberg, C.; Hallberg, A. *Acc. Chem. Res.* **2002**, *35*, 717–727.

(37) Beesley, R. M.; Ingold, C. K.; Thorpe, J. F. *J. Chem. Soc.* **1915**, 107, 1080.


(38) Kochi, J. K. *Acc. Chem. Res.* **1974**, *7*, 351–360.

(39) Iqbal, J.; Bhatia, B.; Nayyar, N. K. *Chem. Rev.* **1994**, *94*, 519–564.



ylpiperidines (AIBN, HSnBu₃, PhH, reflux), and in many instances, the *ipso* product predominates.^{40–44} It has previously been observed, however, that the propensity for direct addition versus *ipso* substitution in the intramolecular addition reactions of carbon radicals to aromatic rings is highly dependent on the structure of the substrate.^{42,44,45}

In an additional experiment, we have trapped the copper(II) carboxylate-promoted carboamination reaction intermediate derived from sulfonamide **1** with TEMPO (2,2,6,6-tetramethylpiperidine-*N*-oxyl), a classical carbon radical trap (eq 6).⁴⁶



No reaction occurred under the same conditions but in the absence of Cu(OAc)₂. It should be noted that carbon radicals do react rapidly with copper(II) salts to form organocopper(III) intermediates.³⁸ This process might be reversible for primary carbon radicals under our reaction conditions. Although we cannot rule out the possibility that some of the reactions discussed above might take place via such an organocopper species, we believe the carbon radical mechanism is the simplest explanation and most consistent with the observed results (vide supra, eqs 5 and 6). Previously, organocopper(III) species have been shown to undergo oxidative elimination or reductive elimination reactions.³⁸

Substrates with internal disubstituted olefins require higher temperature conditions in the copper(II) carboxylate-promoted oxidative cyclization than their terminal olefin counterparts (Table 3). The reactions are also less efficient and the cyclization cascade concludes in elimination rather than radical addition to the aromatic ring of the sulfonamide (e.g., compare entry 4 in Table 2 to entries 2 and 3 in Table 3).


The reactivity pattern of the internal olefin substrates indicates that (1) the rate of the initial nitrogen addition to the alkene is retarded by substitution on the terminal carbon and (2) oxidation of the resulting carbon radical to the olefin is faster than addition

TABLE 3. 1,2-Disubstituted (Internal) Olefin Substrates

entry	substrate	product	temp, time, Cu(R) ₂	yield (%) ^c
1 ^a			200 °C, 78 h, Cu(ND) ₂	35
2 ^{a,b} 3 ^{a,b}			160 °C, 72 h, Cu(OAc) ₂ 160 °C, 72 h, Cu(ND) ₂	no rxn. 31

^a Remainder of the mass was recovered starting material. ^b A 5:1 *E:Z* mixture of olefin isomers of **36** was used. ^c Isolated yield. ND = neodecanoate.

SCHEME 2. Mechanism of Oxidative Cyclization of Substrates with Internal Olefins



of the radical to the aromatic ring in cases when the carbon radical is secondary rather than primary (Scheme 2). Copper(II)-promoted oxidative elimination reactions of carbon radicals have been previously invoked by Kochi; the elimination is not thought to occur via a carbocation intermediate unless a stable carbocation can be formed.⁴⁷

To aid in our mechanism investigation as well as further explore the scope of the carboamination reaction, we examined the oxidative cyclization of several α -substituted γ -alkenyl sulfonamides **39** for the synthesis of 2,5-disubstituted pyrrolidines (Table 4).⁴⁸ Sulfonamides **39** can be efficiently synthesized in enantiomerically enriched form via ring opening addition of substituted tosylaziridines with allylmagnesium bromide.⁴⁹ We found that *cis*-2,5-disubstituted pyrrolidines **40** can be formed with very high diastereoselectivity (>20:1) and in 31–51% yield in the copper(II)-promoted oxidative cyclization of substrates **39** using either oil bath heating (sealed tube, 170–200 °C, 72 h) or microwave irradiation (210 °C, 3 h). Heating in an oil bath at 200 °C for 3 h gave only slightly lower conversion (entry 3). The net hydroamination adducts **41** are formed in these reactions as well (15–28%) and are formed with high 2,5-*cis* selectivity (>20:1) in all cases examined. The products emerge as ca. 2:1 mixtures of carboamination and hydroamination adducts, **40** and **41**, respectively. A series of high boiling solvents [toluene, trifluorotoluene, *tert*-butyltoluene, *N,N*-dimethylacetamide (DMA), 1,3-dimethyl-2-imidazolidinone (DMI), and 1-methyl-2-pyrrolidinone (NMP)] and different reaction concentrations (0.01 and 0.5 M in DMF, reactions in Table 4 were run at 0.1 M substrate concentration) were tried with substrate **39b** in an attempt to improve the carboamination product yield; unfortunately no improved conditions have yet been identified.

When substrate **39b** was treated first with NaH (1.2 equiv) and then with Cu(ND)₂ (1.2 equiv) and subsequently heated, the reaction proceeded efficiently and gave similar results as when Cs₂CO₃ was used as base (compare entry 4 to 6). This

(40) Kohler, J. J.; Speckamp, W. N. *Tetrahedron Lett.* **1977**, 7, 631–634.

(41) Speckamp, W. N.; Kohler, J. J. *Chem. Commun.* **1978**, 166–167.

(42) Kohler, J. J.; Speckamp, W. N. *Tetrahedron Lett.* **1977**, 7, 635–638.

(43) Loven, R.; Speckamp, W. N. *Tetrahedron Lett.* **1972**, 1567–1570.

(44) da Mata, M. L. E. N.; Motherwell, W. B.; Ujjainwalla, F. *Tetrahedron Lett.* **1997**, 38, 137–140.

(45) Bonfand, E.; Motherwell, W. B.; Pennell, A. M. K.; Uddin, M. K.; Ujjainwalla, F. *Heterocycles* **1997**, 46, 523–534.

(46) Root, K. S.; Hill, C. L.; Lawrence, L. M.; Whitesides, G. M. *J. Am. Chem. Soc.* **1989**, 111, 5405–5412.

(47) Kochi, J. K.; Bacha, J. D. *J. Org. Chem.* **1968**, 33, 2746–2754.

(48) Pichon, M.; Figadere, B. *Tetrahedron: Asymmetry* **1996**, 7, 927–964.


(49) Akiyama, T.; Ishida, Y.; Kagoshima, H. *Tetrahedron Lett.* **1999**, 40, 4219–4222.

TABLE 4. Diastereoselective Formation of 2,5-*cis*-Pyrrolidines^a

Reaction scheme: Sulfonamide **39** (with R¹ and R² substituents) reacts with Cu(ND)₂ and Cs₂CO₃ in DMF to produce a mixture of *cis*-40 and *cis*-41.

entry	substrate	method, temp, time	yield (%) ^c 40	yield (%) ^c 41
1	39a , R ¹ = <i>i</i> -Pr, R ² = Me	oil bath, 190 °C, 72 h	49	25
2	39a	210 °C (μW), 3 h	51	28
3	39a	oil bath, 200 °C, 3 h	40	19
4	39b , R ¹ = <i>i</i> -Pr, R ² = OMe	oil bath, 190 °C, 72 h	49	23
5	39b	210 °C (μW), 3 h	51	23
6	39b	oil bath, 190 °C, 72 h ^b	33	24
7	39c , R ¹ = <i>t</i> -Bu, R ² = Me	oil bath, 170 °C, 72 h	34	15
8	39c	210 °C (μW), 3 h	31	17
9	39d , R ¹ = Me, R ² = Me	oil bath, 200 °C, 72 h	48	15
10	39d	210 °C (μW), 3 h	48	15
11	39e , R ¹ = <i>i</i> -PrCH ₂ , R ² = Me	oil bath, 200 °C, 72 h	49	20
12	39e	210 °C (μW), 3 h	49	19
13	39f , R ¹ = Bn, R ² = Me	oil bath, 200 °C, 72 h	50	22
14	39f	210 °C (μW), 3 h	47	21

^a Sulfonamides **39** were dissolved in DMF (0.1 M) and treated with Cs₂CO₃ (1 equiv) and Cu(ND)₂ (3 equiv) and were heated in the indicated manner at the indicated temperature and time. ^b Reaction run with NaH as base instead of Cs₂CO₃: sulfamide **39b** in DMF was treated with NaH (1.2 equiv) at 23 °C for 0.5 h, then Cu(ND)₂ (1.2 equiv) in DMF was added, the solution was stirred for 0.5 h, then the reaction was heated at 190 °C for 72 h. ^c Yields refer to isolated products; diastereomeric ratios (>20:1) were determined by analysis of the crude ¹H NMR and by isolated yields. ND = neodecanoate.

SCHEME 3. Possible Mechanisms for *cis*-Pyrrolidine Formation

indicates that a R₂N-CuL intermediate can productively convert to the observed products (vide infra).

Analysis of the C–N Bond Formation Mechanism. The level and direction of diastereoselectivity in the 2,5-disubstituted pyrrolidine formation (Table 4) provides insight into the initial steps of the reaction mechanism. Three mechanistic scenarios that lead to the carbon radical intermediate **43** were considered (Scheme 3): (1) trans aminocupration, generating an unstable organocopper(II) species^{50,51} that homolyzes to the carbon radical and Cu(I); (2) syn aminocupration, generating an unstable organocopper(II) species that homolyzes to the carbon radical and Cu(I); and (3) Cu(II) oxidation of the nitrogen to the nitrogen radical, followed by cyclization onto the olefin.

The predominance for the formation of the *cis* pyrrolidine diastereomer can provide insight into the reaction mechanism,

especially when compared to the stereochemical trends observed when other reagents are used to promote pyrrolidine formation (vide infra).

Consideration of a Nitrogen Radical Pathway. Pyrrolidines can be formed via cyclization of a nitrogen radical onto an olefin. Nitrogen radicals are usually formed via the homolytic cleavage of nitrogen–heteroatom bonds.^{23,52,53} For example, amine, amidyl, and sulfamidyl radicals have been generated by treatment of (1) xanthates (*N*-(*O*-ethyl thiocarbonylsulfanyl)-amides) with lauroyl peroxide and heat,⁵⁴ (2) *O*-acyl hydroxamic acids with Bu₃SnH and AIBN,^{55–57} or Cu(OTf)₂⁵⁵ (3) N–X

(52) Neale, R. S. *Synthesis* **1971**, 1, 1–15.

(53) Stella, L. *Angew. Chem., Int. Ed. Engl.* **1983**, 22, 337–422.

(54) Gagosz, F.; Moutrille, C.; Zard, S. Z. *Org. Lett.* **2002**, 4, 2707–2709.

(55) Clark, A. J.; Filik, R. P.; Peacock, J. L.; Thomas, G. H. *Synlett* **1999**, 441–443.


(56) Boivin, J.; Callier-Dublanche, A.-C.; Quiclet-Sire, B.; Schiana, A.-M.; Zard, S. Z. *Tetrahedron* **1995**, 51 (23), 6517–6528.

(57) Clark, A. J.; Deeth, R. J.; Samuel, C. J.; Wongtap, H. *Synlett* **1999**, 4, 444–446.


(50) Cotton, F. A.; Wilkinson, G.; Murillo, C. A.; Bochmann, M. *Advanced Inorganic Chemistry*, 6th ed.; John Wiley & Sons: New York, 1999.

(51) Chmielewski, P. J.; Latos-Grazynski, L.; Schmidt, I. *Inorg. Chem.* **2000**, 39, 5475–5482.

(X = Cl, Br, or I) with Et₃B, light, or a transition metal catalyst,^{52,58–65} and (4) *N*-acyltriazines with heat.⁶⁶ Typical products of nitrogen radical cyclizations onto olefins include net carboamination, hydroamination, and atom transfer (eqs 7 and 8).^{55,66} Typically, terminal, internal and trisubstituted olefins provide amine and amidyl radical cyclization adducts.^{54,63}




Clark has reported studies regarding the stereoselectivity of an amidyl radical cyclization where the substrate contained a stereocenter α to the amine.⁵⁷ He found that the trans pyrrolidine product is favored (dr = 3.3:1) (eq 9, remainder of isolated material is uncyclized amide resulting from reduction of the N–O bond).⁵⁷ Similar nitrogen radical reactions performed by Senboku and Tokuda involve homolysis of N–Cl substrates with AIBN and Bu₃SnH and provide the 2,5-trans pyrrolidines as the major diastereomeric products.^{31,32}



We further examined the stereoselectivity of a nitrogen radical reaction with a sulfonamide substrate. In our hands, of a number of methods attempted, the *N*-chlorosulfonamide **46** proved to be the most facile nitrogen radical precursor to synthesize. Hence, **46** was obtained uneventfully by treatment of the sulfonamide with NaH followed by *N*-chlorination with NCS.⁶⁴ The *N*-chlorosulfonamide was subjected to two conditions reported to induce nitrogen radical cyclization reactions.^{59,63,64} Treatment of **46** with Et₃B led to a 64:36 cis:trans mixture of chloromethylpyrrolidines **47** in 73% combined yield. When CuCl/CuCl₂ was used as the radical initiator,^{59,63,67} a 44:56 cis:trans mixture of **47** was obtained. Both reactions are relatively unselective (dr = <2:1). On the basis of these comparisons, it is unlikely that the highly diastereoselective copper(II) carboxylate-promoted oxidative cyclization proceeds via a nitrogen radical. A mechanism involving a copper(I)- or copper(II)-

coordinated nitrogen radical that promotes an organized addition to the alkene is also unlikely given the poor diastereoselectivity observed when CuCl/CuCl₂ was used to promote the radical reaction of chlorosulfonamide **46** (eq 10).




The oxidation of an amide or sulfonamide to the corresponding nitrogen radical by the agency of a copper salt is, in fact, unprecedented. (It should be noted, however, that the oxidation of *aliphatic* amines to nitriles, imines, and aldehydes by copper salts has been reported, and these reactions have been postulated to occur through both radical and two-electron processes.^{68–70}) The oxidation of *N*-aryl amides with *o*-iodoxybenzoic acid (IBX) has, however, been reported, and the derived radicals undergo intramolecular cyclization onto variously substituted olefins.⁷¹ Amides derived from aliphatic amines (non-aniline) are reported to be unreactive under these conditions.⁷¹ In our hands, subjection of *N*-arylsulfonyl-*o*-allylaniline **1a** to these reaction conditions resulted in no reaction by crude ¹H NMR analysis. In comparison to the IBX-promoted reaction, it is also noteworthy that the arylsulfonamides derived from aliphatic amines are not significantly less reactive than arylsulfonamides derived from anilines in the copper(II) carboxylate-promoted oxidative cyclization. This fact also argues against the formation of a nitrogen radical. In addition, the lower reactivity of the internal olefins [substrate **36** (Table 3) is less reactive than substrate **14** (Table 2)] indicates that steric hindrance at the terminal carbon impedes reactivity in the copper(II) carboxylate-promoted reactions; by comparison, amidyl radical reactions are not impeded by substitution on the terminal alkene carbon (e.g., eq 8).⁷² Transition metal-mediated additions of nitrogens to alkenes often suffer lower reactivity with internal alkenes.^{4,73} We therefore conclude that the reactivity pattern observed in the copper(II) carboxylate-promoted carboaminations is most consistent with a reaction mechanism that requires addition of [Cu] to the terminal alkene carbon.

Consideration of a Trans Aminocupration Mechanism. Intramolecular cyclization of amines onto olefins to form pyrrolidines can be promoted by metallic electrophiles such as mercury(II) and palladium as well as halogens and analogous electrophilic reagents.^{5,74,75} Elegant work by Harding and co-workers established that the trans-pyrrolidine is the kinetic


(58) Daoust, B.; Lessard, J. *Tetrahedron* **1999**, *55*, 3495–3514.
 (59) Heuger, G.; Kalsow, S.; Gottlich, R. *Eur. J. Org. Chem.* **2002**, 1848–1854.
 (60) Togo, H.; Hoshina, Y.; Muraki, T.; Nakayama, H.; Yokoyama, M. *J. Org. Chem.* **1998**, *63*, 5193–5200.
 (61) Martin, A.; Perez-Martin, I.; Suarez, E. *Org. Lett.* **2005**, *7*, 2027–2030.
 (62) Broka, C. A.; Gerlits, J. F. *J. Org. Chem.* **1988**, *53*, 2144–2150.
 (63) Bougeois, J.-L.; Stella, L.; Surzur, J.-M. *Tetrahedron Lett.* **1981**, *22*, 61–64.
 (64) Tsuritani, T.; Shinokubo, H.; Oshima, K. *Org. Lett.* **2001**, *3*, 2709–2711.
 (65) Tsuritani, T.; Shinokubo, H.; Oshima, K. *J. Org. Chem.* **2003**, *68*, 3246–3250.
 (66) Lu, H.; Li, C. *Tetrahedron Lett.* **2005**, *46*, 5983–5985.
 (67) Hemmerling, M.; Sjöholm, A.; Somfai, P. *Tetrahedron: Asymmetry* **1999**, *10*, 4091–4094.

(68) Yamaguchi, J.-i.; Takeda, T. *Chem. Lett.* **1992**, *10*, 1933–1936.
 (69) Minakata, S.; Ohshima, Y.; Takemiyu, A.; Ryu, I.; Komatsu, M.; Ohshiro, Y. *Chem. Lett.* **1997**, *4*, 311–312.
 (70) Maeda, Y.; Nishimura, T.; Uemura, S. *Bull. Chem. Soc. Jpn.* **2003**, *76*, 2399–2403.
 (71) Nicolaou, K. C.; Baran, P. S.; Zhong, Y.-L.; Barluega, S.; Hunt, K. W.; Kranich, R.; Vega, J. A. *J. Am. Chem. Soc.* **2002**, *124*, 2233–2244.
 (72) Horner, J. H.; Musa, O. M.; Bouvier, A.; Newcomb, M. *J. Am. Chem. Soc.* **1998**, *120*, 7738–7748.
 (73) Kim, J. Y.; Livinghouse, T. *Org. Lett.* **2005**, *7* (20), 4391–4393.
 (74) Williams, D. R.; Osterhout, M. H.; McGill, J. M. *Tetrahedron Lett.* **1989**, *30* (11), 1327–1330.
 (75) Takacs, J. M.; Helle, M. A.; Sanyal, B. J.; Eberspacher, T. A. *Tetrahedron Lett.* **1990**, *31* (47), 6765–6768.

product in the mercury(II)-promoted cyclization of *N*-Cbz γ -unsaturated amine (eq 11).⁷⁶ The proposed mechanism




involves a chairlike transition state and a *trans*-aminomercuriation of the olefin.⁷⁷ As proposed by Harding, the substituents adopt pseudoequatorial positions on the chairlike transition state. No evidence has been proposed to refute this mechanism. When sulfonamide **39a** was treated under modified⁷⁸ mercuriation–demercuration conditions, an 8:1 diastereomeric mixture favoring the *trans* hydroamination adduct *trans*-**41a** was obtained in 56% yield (eq 12). That the *trans* rather than *cis* adduct substantially predominates in this kinetically controlled reaction is a strong indication that the *cis*-selective copper(II) carboxylate oxidative cyclization reaction does not proceed by the same mechanism as the mercury(II) acetate reaction.





Consideration of a Syn Aminocupration Mechanism. A mechanism for the copper(II) carboxylate-promoted carboamination reaction that involves the intermediacy of a copper-coordinated nitrogen (cf. **44**, Scheme 3) is supported by the fact that when such an intermediate is generated at room temperature by deprotonation of the sulfonamide **39a** at 23 °C with NaH followed by treatment with 1 equiv of Cu(ND)₂ at 23 °C for 0.5 h, upon heating, it forms the oxidative cyclization products *cis*-**40b** and *cis*-**41b** (Table 4, entry 6). That internal olefins (Table 3) are less reactive than terminal olefins in this reaction also supports the hypothesis that a copper–carbon bond is formed in the rate-determining step.

In addition, palladium-catalyzed carboamination reactions that are thought to occur via syn aminopalladation favor formation of *cis*-2,5-disubstituted pyrrolidines.^{20,21,79} As reported by Wolfe and co-workers, *N*-Aryl, *N*-Boc, and *N*-acyl substrates all provide the *cis*-2,5-disubstituted pyrrolidine as the major adduct (eq 13). Examination of the reaction of cyclic disubstituted olefins led Wolfe to conclude that a syn insertion of the nitrogen and palladium into the olefin is operative in these reactions.²⁰ The nitrogen substitution pattern and *cis* pyrrolidine selectivity in this reaction is most analogous to the copper(II)-promoted




reactions described herein, suggestive of similar diastereocontrol elements in the two reactions.⁸⁰

It is also noteworthy that lanthanide-catalyzed hydroaminations of primary amines are routinely *trans* selective operations and are thought to occur via a syn addition mechanism (eq 14).^{4,73} Conversely, cationic titanocene and zirconocene catalyzed hydroaminations of *secondary* amines favor formation of the 2,5-*cis* pyrrolidines (eq 15).⁸¹ Hydroamination reactions of secondary amines catalyzed by *n*-BuLi also afford 2,5-*cis* pyrrolidines.⁸² Clearly, the nitrogen substituent (H vs Me, Ar, or Ts) significantly affects the diastereoselectivity of such reactions (vide infra).



Other Pyrrolidine Formation Methods. In addition to the methods described above, the Brønsted acid-catalyzed cyclization of γ -alkenyl sulfonamide **39e** provided a mixture of hydroamination products, pyrrolidines *trans*-**41e** and *cis*-**41e**, where the *trans* 2,5-pyrrolidine was slightly favored (eq 16).^{33,83} The direction of diastereoselectivity and level of stereocontrol in this reaction indicates that the mechanism of the copper(II) carboxylate reactions is not controlled by analogous factors.



Transition State Analysis for Copper(II) Carboxylate-Promoted Carboaminations. Syn addition transition state models that rationalize the observed diastereoselectivity of the copper(II) carboxylate-promoted carboamination reactions of substrates with stereocenters α to the amine are illustrated in Figure 1. The copper(II) is assumed to be tetracoordinate in these transition states. The top two transition states, **A** and **B**, are chairlike, where **A** leads to the 2,5-*cis*-disubstituted pyrrolidine and **B** leads to the 2,5-*trans*-disubstituted pyrrolidine. The boatlike transition state **C** also leads to the 2,5-*cis* pyrrolidine while the boatlike transition state **D** leads to the 2,5-*trans*

(76) Harding, K. E.; Marman, T. H. *J. Org. Chem.* **1984**, 49, 2838–2840.

(77) Harding, K. E.; Burks, S. R. *J. Org. Chem.* **1981**, 46, 3920–3922.

(78) Kang, S. H.; Lee, J. H.; Lee, S. B. *Tetrahedron Lett.* **1998**, 39, 59–62.

(79) Ney, J. E.; Hay, M. B.; Yang, Q. F.; Wolfe, J. P. *Adv. Synth. Catal.* **2005**, 347, 1614–1620.

(80) Bertrand, M. B.; Wolfe, J. P. *Org. Lett.* **2006**, 8 (11), 2353–2356.

(81) Gribkov, D. V.; Hultsch, K. C. *Angew. Chem., Int. Ed.* **2004**, 43, 5542–5546.

(82) Fujita, H.; Tokuda, M.; Nitta, M.; Sugimoto, H. *Tetrahedron Lett.* **1992**, 33, 6359–6362.

(83) Schlummer, B.; Hartwig, J. F. *Org. Lett.* **2002**, 4, 1471–1474.





FIGURE 1. Competing syn addition transition states that lead to the *cis*- and *trans*-pyrrolidines.

pyrrolidine. Both the chair and boat transition states **A** and **C** that lead to the *cis* pyrrolidine orient the SO₂Ar group *trans* to the α substituent, R. Both the chair and boat transition states **B** and **D** that lead to the *trans* pyrrolidine orient the SO₂Ar group *cis* with respect to substituent R. It appears that the steric interaction between these two groups is the dominant stereocontrol element. Chairlike transition state models of other intramolecular additions of amines to unactivated olefins are typically favored over the boatlike transition states.^{4,57,77,81}

Reductive Removal of Sulfur Dioxide. The use of sulfonamides and sultams as biologically active agents for medicinal chemistry purposes as well as herbicides is well established.⁸⁴ It is hoped that the methods described in this report for the direct synthesis of sultams from acyclic γ - and δ -alkenyl sulfonamides will prove useful for the synthesis of novel sultam small molecule scaffolds for medicinal chemistry endeavors. Should the free amine be desired, the SO₂ unit may be directly excised under dissolving metal conditions (Li, NH₃) (eq 17).⁸⁵



Additionally, methods for the use of sultams directly in Ni(acac)₂-catalyzed cross-coupling reactions with Grignard reagents have been reported and provide a method for subsequent C–C bond formation.⁸⁶

Conclusion

The low cost of copper and the wide range of valuable transformations enabled by this metal continue to stimulate significant research effort in the discovery, development, and refinement of copper-facilitated chemistry.^{22,38,87–104} The mech-

anism of these reactions is often complicated by the dual polar, organometallic, and redox properties of copper and it is often difficult to differentiate between radical and polar reaction mechanisms. We have reported the first copper-promoted addition of unfunctionalized nitrogens to unactivated alkenes forming a new sp³ stereocenter.^{11,14} As described herein, we have found these reactions to be highly diastereoselective in a number of cases and useful for the synthesis of a range of five- and six-membered nitrogen heterocycles. We have provided evidence for a syn aminocupration pathway for the formation of the N–C bond. A copper-facilitated N–C bond-forming process is required for asymmetric induction controlled by chiral ligands on copper(II) to be envisioned. We have also provided evidence for a carbon radical intermediate that can undergo either intramolecular addition to aryl rings or hydrogen abstrac-

- (88) Hassan, J.; Sevignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. *Chem. Rev.* **2002**, 102, 1359–1469.
- (89) Chen, X.; Hao, X.-S.; Goodhue, C. E.; Yu, J.-Q. *J. Am. Chem. Soc.* **2006**, 128, 6790–6791.
- (90) Baran, P. S.; Richter, J. M. *J. Am. Chem. Soc.* **2004**, 126, 7450–7451.
- (91) Li, X.; Hewgley, B.; Mulrooney, C. A.; Yang, J.; Kozlowski, M. C. *J. Org. Chem.* **2003**, 68, 5500–5511.
- (92) Taylor, J. G.; Whittall, N.; Hii, K. K. M. *Org. Lett.* **2006**, 8, 3561–3564.
- (93) Rovis, T.; Evans, D. A. *Prog. Inorg. Chem.* **2001**, 50, 1–150.
- (94) Wang, Q.; Chan, T. R.; Hilgraf, R.; Folkin, V. V.; Sharpless, K. B.; Finn, M. G. *J. Am. Chem. Soc.* **2003**, 125, 3192–3193.
- (95) Kanazawa, C.; Kamijo, S.; Yamamoto, Y. *J. Am. Chem. Soc.* **2006**, 128, 10662–10663.
- (96) Laiter, D. S.; Tsui, E. Y.; Sadighi, J. P. *J. Am. Chem. Soc.* **2006**, 128, 11036–11037.
- (97) Gooben, L. J.; Deng, G.; Levy, L. M. *Science* **2006**, 313, 662–664.
- (98) Barun, O.; Ila, H.; Junjappa, H.; Singh, O. M. *J. Org. Chem.* **2000**, 65, 1583–1587.
- (99) Zhu, J.; Grigoriadis, N. P.; Lee, J. P.; Porco, J. A. *J. Am. Chem. Soc.* **2005**, 127, 9342–9343.
- (100) Chemler, S. R.; Fuller, P. H. *Chem. Soc. Rev.* **2007**, DOI: 10.1039/B607819M.
- (101) Ezquerro, J.; Pedregal, C.; Lamas, C.; Barluenga, J.; Pérez, M.; García-Martin, M. A.; González, J. M. *J. Org. Chem.* **1996**, 61, 5804–5812.
- (102) Hiroya, K.; Itoh, S.; Sakamoto, T. *J. Org. Chem.* **2004**, 69, 1126–1136.
- (103) Xu, L.; Lewis, I. R.; Davidsen, S. K.; Summers, J. B. *Tetrahedron Lett.* **1998**, 39, 5159–5162.
- (104) Martin, R.; Rivero, M. R.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2006**, 45, 7079–7082.

(84) Sammes, P. G. In *Comprehensive Medicinal Chemistry*; Hansch, C., Sammes, P. G., Taylor, J. B., Eds.; Pergamon Press: Oxford, UK, 1990; Vol. 2, Chapter 7.1.

(85) Evans, P.; McCabe, T.; Morgan, B. S.; Reau, S. *Org. Lett.* **2005**, 7 (1), 43–46.

(86) Milburn, R. R.; Snieckus, V. *Angew. Chem., Int. Ed.* **2004**, 42, 888–891.

(87) Ley, S. V.; Thomas, A. W. *Angew. Chem., Int. Ed.* **2003**, 42, 5400–5449.

tion from the reaction medium. We have also observed oxidative amination with internal alkene substrates. Thus far we have demonstrated that both net intramolecular carboamination and diamination of unactivated alkenes can be efficiently facilitated by copper(II) carboxylates.^{11,14} This chemistry should also be amenable to the addition of other functional groups to alkenes via either radical or copper-facilitated mechanisms, and studies along those lines, as well as additional mechanistic studies and exploration of asymmetric catalysis methods, will be reported in due course.


Experimental Section

Representative Synthesis of α -Substituted *N*-Tosyl-4-pentenylamines (Substrates in Table 4): (*S*)-*N*-Tosyl-1-isopropyl-4-pentenylamine (**39a**). Magnesium metal (254 mg, 10.5 mmol, 5 equiv) was placed in a 25 mL 2-necked flask equipped with a reflux condenser and magnetic stirring bar, then suspended in 10 mL of Et₂O under Ar(g). Freshly distilled allyl bromide (1.24 g, 0.89 mL, 10.5 mmol, 5 equiv) was added dropwise at rt. The mixture was stirred for 2 h (until the magnesium was consumed). (*S*)-2-Isopropyl-*N*-tosylaziridine¹⁰⁵ (500 mg, 2.1 mmol, 1 equiv) was dissolved in 5 mL of Et₂O under Ar(g) and added dropwise. The mixture was stirred for an additional 16 h. The reaction was quenched with saturated NH₄Cl(aq) (15 mL), and extracted with ethyl acetate (2 \times 10 mL). The crude oil was purified by flash chromatography on SiO₂ (20% EtOAc in hexanes) to give 428 mg of (*S*)-*N*-tosyl-1-isopropyl-4-pentenylamine (**39a**) in a 73% yield as a white solid. Data for **39a**: mp 49–51 °C, [α]_D²⁰ –8.6 (c 1.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.74 (d, *J* = 8.0 Hz, 2 H), 7.28 (d, *J* = 8.4 Hz, 2 H), 5.64 (m, 1 H), 4.85–4.95 (m, 2 H), 4.26 (d, *J* = 9.2 Hz, 1 H), 3.12 (m, 1 H), 2.41 (s, 3 H), 1.85–1.95 (m, 2 H), 1.73 (m, 1 H), 1.48 (m, 1 H), 1.30 (m, 1 H), 0.78 (d, *J* = 3.6 Hz, 3 H), 0.76 (d, *J* = 2.8 Hz, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 143.2, 138.9, 138.3, 130.0, 127.6, 115.5, 59.3, 31.6, 31.5, 30.3, 22.0, 18.7, 18.1; IR (neat, thin film) ν 2961, 1641, 1599, 1429, 1323, 1156, 1094 cm^{–1}; HRMS (EI) calcd for C₁₅H₂₃O₂NS [M]⁺ 281.1444, found 281.1447.

Representative Procedure (Table 4) for Carboamination Reactions in a Microwave: (3*S*,10*aS*)-3-isopropyl-8-methyl-2,3,10,10*a*-tetrahydro-1*H*-pyrrolo[1,2-*b*]^{1,2}benzothiazine 5,5-Dioxide (**40a**) and (2*S*,5*S*)-*N*-Tosyl-2-isopropyl-5-methylpyrrolidine (**41a**). (*S*)-*N*-Tosyl-1-isopropyl-4-pentenylamine **39a** (35.1 mg, 0.125 mmol) in a microwave vial equipped with a magnetic stir bar was treated with copper(II) neodecanoate (60% by wt in toluene, 152 mg, 0.275 mmol, 3 equiv) and Cs₂CO₃ (40.7 mg, 0.125 mmol, 1 equiv) and dissolved in DMF (1.3 mL) under Ar(g). The vial was sealed and the reaction mixture was heated at 210 °C for 1.5 h in a microwave. After cooling to rt, it was heated again at 210 °C for 1.5 h in a microwave. The mixture was cooled to rt, diluted with Et₂O, and washed with saturated EDTANa₂(aq). The organic layer was dried over Na₂SO₄ and filtered, and the solvents were removed in vacuo. The mixture was purified by flash chromatography on SiO₂ (10–30% EtOAc in hexanes gradient) providing (*S*,*S*)-*N*-tosyl-2-isopropyl-5-methylpyrrolidine **41a** (9.8 mg, 0.0351 mmol) in 28% yield (*R*_f 0.5 in 10% EtOAc in hexanes) and (*S*,*S*)-3-isopropyl-8-methyl-2,3,10,10*a*-tetrahydro-1*H*-pyrrolo[1,2-*b*]^{1,2}benzothiazine 5,5-dioxide **40a** (17.8 mg, 0.0639 mmol) in 51% yield (*R*_f 0.3 in 10% EtOAc in hexanes).

The relative stereochemistry of the carboamination product **40a** was determined by a 1D NOE experiment that showed a signal between H_a and H_b.


Data for **40a**: mp 85–88 °C; [α]_D²⁰ +198.9 (c 1.2, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.77 (d, *J* = 8.5 Hz, 1 H), 7.19 (d, *J* = 8.0 Hz, 1 H), 7.02 (s, 1 H), 4.11 (m, 1 H), 3.94 (m, 1 H), 2.88–3.09 (m, 2 H), 2.36 (s, 3 H), 2.24 (m, 1 H), 2.19 (m, 1 H), 1.86–



1.90 (m, 2 H), 1.76 (m, 1 H), 0.93 (d, *J* = 7.0 Hz, 3 H), 0.88 (d, *J* = 6.5 Hz, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 142.8, 136.3, 135.8, 130.0, 128.7, 124.3, 62.5, 60.8, 36.7, 32.9, 32.6, 29.6, 21.9, 20.4, 16.5; IR (neat, thin film) ν 2962, 1698, 1465, 1323, 11578, 1094, 909 cm^{–1}; HRMS (EI) calcd for C₁₅H₂₁O₂NSNa [M + Na]⁺ 302.1185, found 302.1192.

The relative stereochemistry of the hydroamination product **41a** was determined by X-ray crystallography.

X-ray Structure of (2*S*,5*S*)-*N*-Tosyl-2-isopropyl-5-methylpyrrolidine (41a**).** Data for **41a**: mp 83–86 °C, [α]_D²⁰ +130.0 (c 0.28,



CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ 7.72 (d, *J* = 8.1 Hz, 2 H), 7.30 (d, *J* = 7.8 Hz, 2 H), 3.65 (m, 1 H), 3.40 (m, 1 H), 2.42 (s, 3 H), 2.10 (m, 1 H), 1.30–1.65 (m, 4 H), 1.29 (d, *J* = 6.6 Hz, 3 H), 0.98 (d, *J* = 7.2 Hz, 3 H), 0.92 (d, *J* = 6.9 Hz, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 143.0, 135.2, 129.5, 127.6, 67.4, 57.3, 31.9, 31.5, 25.3, 23.1, 21.5, 20.0, 17.3; IR (neat, thin film) ν 2962, 1734, 1592, 1459, 1341, 1153, 1096, 1014 cm^{–1}; HRMS (ESI) calcd for C₁₅H₂₄O₂NS [M + H]⁺ 282.1522, found 282.1524.

Representative Procedure for Carboamination in an Oil Bath (Table 4). (*S*)-*N*-Tosyl-1-isopropyl-4-pentenylamine **39a** (92.6 mg, 0.330 mmol) was treated with copper(II) neodecanoate (60% by wt in toluene, 402 mg, 0.990 mmol, 3 equiv), and cesium carbonate (108 mg, 0.330 mmol, 1 equiv) in DMF (3.3 mL) under Ar(g) in a pressure tube equipped with a magnetic stir bar. The tube was capped and the mixture was heated at 190 °C for 72 h. Workup and chromatography as described above provided the carboamination adduct **40a** (45.2 mg, 0.162 mmol) in 49% yield, and the hydroamination adduct **41a** (23.2 mg, 0.083 mmol) in 25% yield.

Acknowledgment. The authors thank Dr. Cara L. Nygren [CCDC 614781 (*trans*-**17**) and CCDC 296132 (**41a**)] and Dr. Marc Messerschmidt [CCDC 628996 (*trans*-**20**)] for X-ray crystallographic analysis. The authors thank Prof. Dennis Curran and Dr. Maria Manzoni for helpful discussions. The authors thank Mr. Thomas Zabawa and Mr. Thomas Hayes for their efforts in the synthesis of substrates **14-D** and **22**, respectively. The authors gratefully acknowledge the financial support of the NIH (NIGMS 1R01-GM07838301) and the donors of the Petroleum Research Fund (PRF 40968-G1).

Supporting Information Available: Procedures and characterization data and NMR spectra for all new products and X-ray data in the form of CIF files. This material is available free of charge via the Internet at <http://pubs.acs.org>.