


Pd–C-Induced Catalytic Transfer Hydrogenation with Triethylsilane

Pijus K. Mandal and John S. McMurray*

Department of Experimental Therapeutics, The University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd., Houston, Texas 77030

jsmcmur@mdanderson.org

Received March 29, 2007



In situ generation of molecular hydrogen by addition of triethylsilane to palladium–charcoal catalyst results in rapid and efficient reduction of multiple bonds, azides, imines, and nitro groups, as well as benzyl group and allyl group deprotection under mild, neutral conditions.

The reduction of alkenes, alkynes, nitrates, etc. is often carried out by hydrogenation reactions using metal catalysts. Catalytic transfer of hydrogen (CTH) is a widely accepted alternative method that does not require the use of potentially dangerous hydrogen gas.¹ For this simple operation H_2 is replaced by a hydrogen donor such as 1,4-cyclohexadiene,² hydrazine,³ formic acid,⁴ ammonium formate,⁵ phosphinic acid,⁶ or sodium hypophosphite⁷ with Pd–C as the catalyst. Though these methods are elegant, some transfer agents require high temperature, and some are not applicable to acid-sensitive or base-sensitive substrates. Following the lead of Fukuyama et al.,⁸ we found that TES/Pd–C is a widely applicable, very convenient CTH reagent. The reactions are carried out at room temperature and are rapid, often complete in 10 min or less using excess TES and 10–20% Pd–C (by weight) in MeOH. The conditions are neutral, and thus acid- or base-sensitive substrates can be

(1) Johnstone, R. A. W.; Wilby, A. H.; Entwistle, I. D. *Chem. Rev.* **1985**, 85, 129–170.

(2) Felix, A. M.; Heimer, E. P.; Lambros, T. J.; Tzougraki, C.; Meienhofer, J. *J. Org. Chem.* **1978**, 43, 4194–4196.

(3) (a) Anwer, M. K.; Khan, S. A.; Sivanandaiah, K. M. *Synthesis* **1978**, 751. (b) Furst, A.; Berlo, R. C.; Hooton, S. *Chem. Rev.* **1965**, 65, 51–68.

(4) (a) Gray, B. D.; Jeffs, P. W. *J. Chem. Soc., Chem. Commun.* **1987**, 1329. (b) Elamin, B.; Anantharamaiah, G. M.; Royer, G. P.; Means, G. E. *J. Org. Chem.* **1979**, 44, 3442–3444.

(5) (a) Anwer M. K.; Spatola, A. F. *Synthesis* **1980**, 929–931 (b) Adger, B. M.; O'Farrell, C.; Lewis, N. J.; Mitchell, M. B. *Synthesis* **1987**, 53–54. (c) Carpino, L.; Tunga, A. *J. Org. Chem.* **1986**, 51, 1930–1932. (d) Ram, S.; Ehrenkaufer, R. E. *Synthesis* **1986**, 133–135. (e) For a recent review, see: Ranu, B. C.; Sarkar, A.; Guchhait, S. K.; Ghosh, K. *J. Ind. Chem. Soc.* **1998**, 75, 690–694.

(6) Entwistle, I. D.; Gilkerson, T.; Johnstone, R. A. W.; Telford, R. P. *Tetrahedron* **1978**, 34, 213–215.

(7) Sala, R.; Doria, G.; Passarotti, C. *Tetrahedron Lett.* **1984**, 25, 4565–4568.

(8) Fukuyama, T.; Lin S. C.; Li, L. *J. Am. Chem. Soc.* **1990**, 112, 7050–7051.

reduced without harm. In this communication, we present several examples of TES/Pd–C-mediated CTH of a wide variety of substrates.

Table 1 illustrates the utility of the reaction. Compounds **1–6** show that reduction of simple alkene, alkyne, and conjugate double and triple bonds occurs rapidly and in high yield. Mirza-Aghayan et al. demonstrated that treatment of 1-alkenes with 1 equiv of TES with Pd–C catalysis resulted in partial reduction and double bond migration.⁹ However, with 2 equiv of the silane, only the alkane was observed. In keeping with this work, our results show that excess TES leads to complete reduction of unsaturated bonds.

The Fmoc group is stable to these conditions (compounds **5**, **9**, **15**, and **21**). This protecting group, highly used in peptide synthesis, is subject to hydrogenolysis using H_2 and Pd–C.¹⁰

Interestingly, substrates **7a** and **8a**, containing both benzylic alcohol functionality and multiple carbon–carbon bonds, were reduced completely to alkyl arenes. Benzyl esters and *N*-benzyloxycarbonyl (Z) groups are removed in 5–10 min (entries 9 and 10). Previous studies using TES and PdCl_2 or $\text{Pd}(\text{OAc})_2$ required extended reaction times at elevated temperatures to cleave benzyl-based protecting groups in peptide applications.¹¹ Transprotection to a Boc-amino acid¹² was achieved by the TES/Pd–C-mediated hydrogenolysis of the Z group in **11a** in the presence of di-*tert*-butyl dicarbonate (**11b**). Cleavage of benzyl ethers requires longer reaction times than benzyl esters or the Z group (**12** and **13**). Aromatic benzyl ethers are removed more rapidly than aliphatic ones (compare **14** versus **12** and **13**). Azetidine-2-ones bearing 4-aryl substituents are susceptible to 1,4-ring cleavage reactions by hydrogenolysis.¹³ As shown in **14**, the β -lactam ring is stable to TES/Pd–C, whereas the benzyl ether was cleaved.

Allylic protecting groups such as the allyl ester (All) for protection of carboxylic acids and the allyloxycarbonyl (Alloc) group for amine protection are widely used in synthetic organic chemistry and in peptide synthesis. These groups can be removed with catalytic palladium ($\text{Pd}(\text{PPh}_3)_4$ or $\text{Pd}(\text{OAc})_2$) π -allyl methodology using hydride donors such as HCO_2H , Bu_3SnH , NaBH_4 , PhSiH_3 , and TES.¹⁴ We were able to deprotect both All and Alloc groups using simple TES/Pd–C (**15** and **17**). For substrate **16a** simple reduction of the triple bond to the alkane was observed. We did not detect formation of allyl esters or cleavage of the presumed allyl ester intermediate. Alloc-to-Boc transprotection was also achieved by treating allyloxycarbamates with TES/Pd–C in the presence of di-*tert*-butyl dicarbonate (**18**).

(9) Mirza-Aghayan, M.; Boukherroub, R.; Bolourchian, M. *Appl. Organomet. Chem.* **2006**, 20, 214–219.

(10) Martinez, J.; Tolle, J. C.; Bodanszky, M. *J. Org. Chem.* **1979**, 44, 3596–3598.

(11) (a) Watanabe, Y.; Maki, Y.; Kikuchi, K.; Sugiyama, H. *Chem. Ind.-London* **1984**, 7, 272. (b) Birkofner, L.; Bierwirth, E.; Ritter, A. *Chem. Ber.* **1961**, 94, 821–824.

(12) Bajwa, J. S. *Tetrahedron Lett.* **1992**, 3, 2955–2956.

(13) (a) Ojima, I. *Acc. Chem. Res.* **1995**, 28, 383–389 and references therein. (b) Banik, B. K.; Barakat, K. J.; Wagle, D. R.; Manhas, M. S.; Bose, A. K. *J. Org. Chem.* **1999**, 64, 5746–5753. (c) Bertha, F.; Fetter, J.; Kajtá-Pérez, M.; Lempert, K. *Tetrahedron* **1999**, 55, 5567–5580. (d) Srirajan, V.; Desmukh, A. R. A. S.; Puranik, V. G.; Bhawal, B. M. *Tetrahedron: Asymmetry* **1996**, 7, 2733–2738.

(14) Guibe, F. *Tetrahedron* **1998**, 54, 2967–3042.

TABLE 1. Substrates and Products of TES/Pd-C-Mediated Catalytic Transfer Hydrogenation Reactions

Substrate	Product	%wt Pd-C	Time (min)	Yield ^a (%)
1a	1b	10	10	90
2a	2b	10	10	91
3a	3b	20	10	88
4a	4b	15	15	91
5a	5b	15	5	94
6a	6b	15	10	93
7a	7b	10	5	90
8a	8b	10	5	91
9a	9b	20	5	90
10a	10b	20	10	quantitative ^b
11a	11b	20	50	88
12a	12b	20	60	91
13a	13b	20	72	90
14a	14b	20	20	92
15a	15b	20	5	92
16a	16b	10	10	86
17a	17b	20	5	quantitative ^b
18a	18b	20	30	88
19a	19b	10	10	quantitative ^b
20a	20b	10	30	quantitative ^b
21a	21b	20	5	quantitative ^b
22a	22b	10	5	92

^a Isolated yields after silica gel column chromatography. ^b Reactions were performed in 10:1 MeOH–CHCl₃. Products were isolated as HCl salts and titrated with hexane–ether.

Reduction of both aromatic and aliphatic nitro groups, imines, and azide moieties is also readily achieved using TES/Pd-C (**19–22**). Note that the benzylamine in **22a** was stable to the short reaction conditions.

Chemoslectivity can be achieved by altering the solvent. 4-Benzylbenzaldehyde, on treatment with TES/Pd-C in acetone, resulted in reduction of the carbonyl to a methyl group. In chloroform the reduction halted at the alcohol stage. In both cases the benzyl group remained untouched. However, using methanol as the solvent, all functional groups were reduced and *p*-hydroxy toluene was obtained.

The deprotection of benzyl carboxylates and carbamates, benzyl ethers, and reduction of multiple bond, nitro, azide, and imine groups suggests that Et₃SiH/Pd-C behaves as a hydrogenation reagent through in situ generation of H₂.⁹ Indeed, effervescence was observed on adding TES to the Pd-C catalyst. On the other hand, it is likely that removal of allyl carboxylates and carbamates proceeds via palladium activation of the π -allyl system with triethylsilane as a hydride donor.

It should be noted that catalytic transfer hydrogenation using TES as a hydrogen donor has been used with other metal catalysts. For example, a combined metathesis and olefin hydrogenation using ruthenium Grubbs-type catalyst^{15a} and dehalogenation of halides using PdCl₂ were recently reported.^{15b,c} However, owing to the rapid reaction times and neutral conditions and simplicity of method, the combination of TES and Pd-C has great potential for use in all types of organic synthesis.

Experimental Section

General Procedure. To a stirred solution of substrate (1 mmol) and 10%Pd-C (10–20% by weight) in MeOH (2–3 mL) was added neat TES (10 mmol) dropwise from a pressure-equalizing dropping funnel under an argon-filled balloon. When the reaction was complete (TLC), the mixture was filtered through celite and the solvent was removed in vacuo. The product was chromatographically purified on a short silica gel column. Note: Although

(15) (a) Menozzi, C.; Dalko, P. I.; Cossy, J. *Synlett* **2005**, 2449–2452. (b) Boukherroub, R.; Chatgilialoglu, C.; Manuel, G. *Organometallics* **1996**, 15, 1508 – 1510. (c) Beshore, D. C.; Smith, A. B., III. *J. Am. Chem. Soc.* **2007**, 129, 4148–4149.

we ran all of our reactions in MeOH, this methodology is likely to be successful using ethanol, which has a better safety profile with Pd-C reactions.

Preparation of *N*-(9-Fluorenylmethoxycarbonyl)-4-methyl-4-aminobutyric Acid (3b). To a stirred suspension of 4(S)-[9-fluorenylmethoxycarbonyl]amino]-2-pentenoic acid benzyl ester (**3a**) (0.5 g, 1.2 mmol) and 10% Pd-C (100 mg) in 5 mL of methanol was added triethylsilane (1.8 mL, 11.7 mmol) dropwise under argon. The reaction started with evolution of gas. After completion of the reaction (TLC), the mixture was filtered through celite, and the solvent was removed in vacuo. The crude product was purified by short silica gel column chromatography, eluting with 2% hexane–ethylacetate, followed by hexane–ethylacetate–methanol (3:6:1) to obtain 0.348 g of **3b**, a white solid. ¹H NMR (DMSO-*d*₆) δ 1.24 (d, *J* = 6.6 Hz, 2H), 1.62 (m, 2H), 2.2 (m, 2H), 3.51 (m, 1H), 4.18–4.35 (m, 3H), 7.15 (d, *J* = 6.6 Hz, 1H), 7.3–7.43 (m, 4H), 7.69 (d, *J* = 7.2 Hz, 2H), 7.88 (d, *J* = 7.2 Hz, 2H). HRMS (M + H) calcd for C₂₀H₂₂NO₄ 340.1559, found 340.1556.

Transprotection of Alloc-Leu-OMe (17a) to Boc-Leu-OMe (18b). To a stirred suspension of Alloc-Leu-OMe (**17a**) (0.3 g, 1.3 mmol), 10% Pd-C (60 mg), and Boc-anhydride (0.340 g, 1.56 mmol) in 5 mL of methanol was added triethylsilane (2.1 mL, 13 mmol) dropwise under argon. After completion of reaction (TLC), the mixture was filtered through celite, and the filtrate was concentrated under vaccum. The residue was diluted with 30 mL of ether, and the organic layer was then washed with 10% NaHCO₃ solution followed by brine and dried over MgSO₄. The crude product was purified by short silica gel column chromatography, eluting with 10% hexane–ethylacetate, to get 0.280 g of **18b** as an oil. ¹H NMR (CDCl₃) δ 0.93 (m, 6H), 1.38 (s, 9H), 1.44–1.66 (m, 3H), 3.67 (s, 3H), 4.25 (m, 1H), 4.9 (brs, 1H).

Acknowledgment. We are grateful to the National Cancer Institute (CA096652) and the M. D. Anderson Cancer University Cancer Fund for support of this work. We also acknowledge the NCI Cancer Center Support Grant CA016672 for the support of our NMR facility.

Supporting Information Available: Procedures for the synthesis of substrates **1a–22a**, characterizorion of products **1b–22b**, and ¹H and ¹³C NMR spectra for compounds **3a**, **3b**, **5a**, **8a**, **14a**, **14b**, **15a**, **16a**, and **16b**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO0706123