


Nucleophilic Addition/Double Cyclization Cascade Processes between Enynyl Fischer Carbene Complexes and Alkynyl Malonates

Ana Álvarez-Fernández, Tatiana Suárez-Rodríguez, and Ángel L. Suárez-Sobrino*

Departamento de Química Orgánica e Inorgánica, Universidad de Oviedo, Julián Clavería 8, 33071 Oviedo, Spain

Supporting Information

ABSTRACT: Two new selective cascade processes for enynyl Fischer carbene complexes **1** are described in their reaction with alkynyl malonates. When carbene complexes **1** react with the sodium enolate of homopropargyl malonates **3** a consecutive Michael-type addition/cyclopentannulation/6-*exo* cyclization takes place leading, in a regio- and stereoselective way, to *n*/5/6 angular tricyclic compounds **5**. Furthermore, when propargylic malonates are used, a delayed protonation of the reaction mixture allows intermediate 1,4-addition adduct **Ia** to evolve through a 5-*exo* cyclization, consisting of an intramolecular nucleophilic attack from the central carbon of the allenylmetallate over the triple C–C bond. Further spontaneous cyclopentannulation of the resulting metallatriene gives rise to bicyclic and linear polycyclic compounds **6** and **7**, some of them bearing a polyquinane framework.



INTRODUCTION

Cascade reactions are among the most advantageous tools in organic synthesis.¹ They include multiple benefits, such as atom economy, reduction of workup and purification steps, as well as decreased production of waste. It has been demonstrated that Fischer carbene complexes (FCCs) can participate in many cascade and multicomponent reactions² due to the ability of the metal pentacarbonyl moiety to activate consecutive reaction steps. Thus, enynyl Fischer metal carbenes **1** are the popular substrates for these kinds of reactions, which may generate at least two rings in a single operation. In this context, a well-developed strategy comprises first a cycloaddition involving the activated C–C triple bond followed by ring closure through the C–metal and C–C double bonds. Figure 1 illustrates three different processes, (i) [6 + 2] cycloaddition/cyclopentannu-

lation (compound A),³ (ii) [4 + 2] cycloaddition/benzannulation (compound B),⁴ and (iii) triple [4 + 2] cycloaddition/cyclopentannulation (compound C).⁵

In contrast, cascade reactions initiated by 1,4-conjugate addition of nucleophiles to enynyl FCCs have been less studied.⁶ Recently, we described a new entry to cyclopentenones consisting of a 1,4-addition of enolates to enynylcarbenes followed by cyclopentannulation.⁷ We considered that the use of appropriately functionalized nucleophiles, e.g., readily accessible alkyne-containing malonates **2** and **3**, might provide an alternative pathway. Thus, we observed that cyclopentene derivatives **4** were isolated by mixing enynyl carbenes **1** and dimethyl propargylmalonates **2** at low temperature followed by rapid quenching (30 min) and further stirring at room temperature (Scheme 1).⁸ The complete process involves (i) 1,4-addition of enolate, (ii) protonation and regeneration of the metal carbene function, (iii) cyclopentannulation (cyclization a), (iv) chromium-promoted intramolecular nucleophilic addition (cyclization b), and (v) retrocyclization.

Considering the participation of the key allenyl chromium metallate intermediate **Ia**, other reaction pathways might be realized by appropriate variation on its structure, such as using a longer enolate–alkyne tether like an homopropargylmalonate derivative. Moreover, delaying the protonation would allow intermediate **I** to survive and likely attack an electrophilic center. Herein are reported new cascade polycyclizations from enynyl chromium carbenes **1** and dimethyl propargyl and homopropargylmalonates **2** and **3** leading to bicyclic and tricyclic systems of potential interest (Figure 2). Specifically, compounds **5** were formed from homopropargyl malonates **3**




Figure 1. Cascade reactions of enynyl FCCs **1**.

Received: February 17, 2014

Published: June 17, 2014

Scheme 1. Reaction between Enynyl FCCs 1 and Propargylmalonates 2 (Ref 8)





Figure 2. Cascade cyclizations reported in this work.

(cyclizations a and c), while compounds **6** resulted from the reaction of **1** and propargyl malonates **2** followed by late quenching (cyclizations d and a).


RESULTS AND DISCUSSION

Reaction of Enynyl FCCs 1a–c and Homopropargylmalonates 3: 1,4-Nucleophilic Addition/Double Cyclization Process. We started our study by reacting the corresponding enolate of homopropargylmalonates **3** with enynyl carbene complexes **1** (Scheme 2). Thus, carbene complexes **1a–c** dissolved in THF were added to a THF solution of sodium enolates **3**. After the mixture was stirred for 30 min, the reaction was quenched with an aqueous saturated solution of NH₄Cl. Purification gave tricycles **5** in 89%–76% yield with complete diastereoselectivity.⁹


The structural arrangement and stereochemistry of compounds **5** are in accordance with the spectroscopic data (¹H and ¹³C NMR, HRMS), and the full structure was unambiguously determined for **5c** by X-ray analysis of crystals grown from hexanes/chloroform.¹⁰ The process tolerates well the bromo-C(sp) function (**5**; R³ = Br)¹¹ affording stereoselectively the *Z*-bromoalkenyl derivative **5d** whose structure was also confirmed by X-ray analysis.¹⁰

The consecutive processes leading to tricycles **5** can be explained as depicted in Scheme 3. Thus, Michael-type addition of the homopropargylmalonate enolate **3** to carbenes **1a–c**

Scheme 2. Synthesis of Tricycles 5 from Enynyl FCCs 1a–c and Homopropargylmalonate Enolates 3



Scheme 3. Mechanism Proposed for the Formation of Tricycles 5




would produce allenyl intermediate **Ib** (*n* = 2), which then would undergo protonation on addition of NH₄Cl to regenerate the metal carbene functionality. The resulting metallatriene **IIb** would provide the intermediate **III** upon cyclopentannulation (cyclization a), metal fragment liberation and alkyne–Cr(CO)₅ coordination. Finally, the formation of compounds **5** is thought to occur via 6-*exo* cyclization (cyclization c) by attack from the C4 of the methoxydiene function over the C–C triple bond (intermediate **IV**) followed by protodemetalation.¹²

It is worth noting that this 1,4-addition/double cyclization cascade sequence represents a stereoselective access to functionalized 5/5/6, 6/5/6, and 7/5/6 angular tricyclic frameworks¹³ in a single operation from simple starting materials. Some of these structures are present in the skeleton of natural and bioactive products.¹⁴

Reaction of Propargylmalonates 2 and Enynyl FCCs 1. 1,4-Nucleophilic Addition/Double Cyclization Process. Keeping in mind the experimental conditions and the associated mechanisms shown in the cycloaddition reactions of propargylmalonates (see Scheme 1, ref 8) and homopropargylmalonates (see Schemes 2,3), it seems that the rapid


quenching of intermediates **Ia**⁸ (Scheme 1) and **Ib** (Scheme 3) resulted in the generation of reactive dienylcarbene species **II**. If the quenching of these intermediates **I** were delayed, it could allow new modes of cyclization to develop. Thus, a THF solution of carbenes **1** was added at 0 °C to a THF solution of sodium propargylic enolates **2**. After being stirred for 12 h, the reaction was quenched with an aqueous saturated solution of NH_4Cl . Purification gave enol ethers **6** and ketones **7** in good yields (62–92%) (Scheme 4). The structures of compounds **6** and **7** were confirmed by mono- and bidimensional NMR experiments and by X-ray analysis of compound **6e**.¹⁰

Scheme 4. Reaction between Enynyl FCCs **1** and Propargylic Enolates **2** To Give Polycycles **6** and **7**



As shown in Scheme 4, a variety of polycyclic compounds, depending on the structure of the enynyl FCCs, were readily accessible by taking advantage of this strategy. Thus, tricyclic (**6a–c**, Scheme 4, eq 1) and bicyclic (**6d,e**, Scheme 4, eq 2) structures, some of them with an interesting functionalized polyquinane framework,¹⁵ were formed in very satisfactory yields. The reaction also worked well for indole-substituted alkynylcarbene, where the C2–C3 of 1-methylindole is able to participate as the alkenyl moiety ($\text{R}^1\text{C}=\text{CR}^2$ in **1**) providing a complex tetracyclic indol framework (compound **7a**, Scheme 4, eq 3).¹⁶ Finally, TMS-substituted propargylmalonate **2b** ($\text{R}^3 = \text{TMS}$) has been successfully used to generate the linear triquinane **7b** in high yield (Scheme 4, eq 4).¹⁶ A plausible mechanism for this new process involving a Michael-type addition and a double cyclization is depicted in Scheme 5. In this case, the allenylmetallate **Ia**, resulting from the addition of the propargylmalonate enolate to the Fischer carbene, would undergo an intramolecular 5-*exo* cyclization by nucleophilic attack over the C–C triple bond to give cyclic metallatriene **V**

Scheme 5. Mechanism Proposed for the Formation of **6**



(cyclization d).¹⁷ The latter readily would undergo cyclopentannulation and proton rearrangement to cyclopentadienyl anion intermediate **VI** (cyclization a), and then protonation would give rise to compounds **6** and **7**.¹⁸

CONCLUSIONS

We have described two new 1,4-nucleophilic addition/double-cyclization cascade processes involving enynyl FCCs. Functionalized tricyclic angular compounds **5** can be obtained selectively from cyclic enynyl FCCs **1a–c** and homopropargylmalonates **3** by a 1,4-nucleophilic addition/cyclopentannulation/6-*exo* cyclization process. This reaction represents a simple approach to a variety of 5/5/5, 5/5/6, and 5/5/7 angular tricyclic compounds, the basic structure of some of them being present in bioactive molecules.¹⁴ Moreover, new transformations can be achieved when the protonation of the allenylmetallate addition adducts **Ia** is delayed. Thus, the intramolecular 5-*exo* nucleophilic addition to the triple bond from the central carbon of the allenyl moiety can be favored to produce metallatriene intermediates that spontaneously cyclize to a variety of polycyclic compounds (**6** and **7**), several of them bearing an interesting polyquinane structure.¹⁵

EXPERIMENTAL SECTION

General Methods. All reactions involving air-sensitive compounds were carried out under a N_2 atmosphere (99.99%). All glassware was oven-dried (120 °C), evacuated, and purged with nitrogen. Enynyl Fischer carbene complexes **1** were prepared following described procedures.⁵ Enynes were purchased (2-methyl-1-buten-3-yne, 1-ethynylcyclohexene) or synthesized following the described procedures¹⁹ (1-ethynylcyclopentene and 1-ethynylcycloheptene). Dimethyl 2-(2-propynyl)malonate was purchased; dimethyl 2-(3-trimethylsilyl-2-propynyl)malonate,²⁰ dimethyl 2-(3-butynyl)malonate,²¹ and dimethyl 2-(4-bromo-3-butynyl)malonate²¹ were synthesized. Solvents were dried by standard methods and distilled prior to use. Flash column chromatography was carried out on silica gel 60, 230–240 mesh. Deactivated silica gel was prepared by treatment with a 4% potassium hydrogenophosphate aqueous solution. NMR experiments were recorded on 300, 400, and 600 MHz spectrometers with tetramethylsilane ($\delta = 0.0$) as the internal standard. NMR chemical shifts are reported in ppm. ¹H NMR splitting pattern abbreviations are as follows: s, singlet; d, doublet; m, multiplet; brs, broad singlet. ¹³C NMR spectra were recorded with complete proton decoupling, and the multiplicities were determined by DEPT. High-resolution mass spectra (HRMS) were obtained with electron ionization techniques (EI) (70 eV) using a magnetic sector analyzer.

General Procedure for the 1,4-Nucleophilic Addition/Cyclopentannulation/6-Exocyclization Process. Preparation of Compounds **5**. The corresponding homopropargylmalonate (1.3 mmol) was added to a THF (3 mL) suspension of NaH (1.6 mmol) at 0 °C and stirred for 10 min. Then, a THF solution (2 mL) of chromium enynyl carbene **1** (1 mmol) was added dropwise at the same temperature. The mixture was warmed to rt, stirred for 30 min, and quenched with a saturated aq solution of NH₄Cl (10 mL). The mixture was stirred for 4 h and extracted with diethyl ether (2 × 10 mL). After solvent removal, the resulting crude material was dissolved in CH₂Cl₂ (5 mL) and treated with concd HCl (1 mL); after 12 h at rt, the mixture was extracted with CH₂Cl₂ and the solvent removed. Purification of the crude by column chromatography (silica gel, hexanes/ethyl acetate, 5:1) gave tricycles **5** as solids (76–89%).

cis-Dimethyl 9-methylene-4-oxo-2,3,3a,4,8,9-hexahydro-1H-cyclopenta[c]indene-6,6(7H)-dicarboxylate (**5a**): white solid; 271 mg, 89% yield; mp 119–121 °C dec; ¹H NMR (400 MHz, CDCl₃) δ 1.49–1.52 (m, 2H), 1.67–1.84 (m, 3H), 1.95–2.0 (m, 1H), 2.11–2.15 (m, 1H), 2.35–2.59 (m, 2H), 2.62–2.66 (m, 1H), 2.95–2.98 (m, 1H), 3.79 (s, 6H), 4.75 (brs, 1H), 4.80 (brs, 1H), 5.97 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 25.9 (CH₂), 27.9 (CH₂), 29.7 (CH₂), 33.2 (CH₂), 39.7 (CH₂), 53.1 (CH₃), 53.3 (CH₃), 55.8 (CH), 59.3 (C), 61.8 (C), 107.0 (CH₂), 130.9 (CH), 150.2 (C), 168.7 (C), 169.8 (C), 174.4 (C), 209.9 (C); HRMS calcd for C₁₇H₂₀O₅ (M⁺) 304.1311, found 304.1313.

cis-Dimethyl 1-methylene-6-oxo-2,3,6,6a,7,8,9,10-octahydrobenzo[c]indene-4,4(1H)-dicarboxylate (**5b**): white solid; 248 mg, 78% yield; mp 121–124 °C dec; ¹H NMR (400 MHz, CDCl₃) δ 1.11–1.24 (m, 3H), 1.30–1.41 (m, 1H), 1.58–1.71 (m, 2H), 1.87–1.92 (m, 1H), 2.01–2.13 (m, 1H), 2.28–2.43 (m, 2H), 2.43–2.75 (m, 3H), 3.77 (s, 6H), 4.85 (brs, 1H), 5.00 (brs, 1H), 6.05 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 20.3 (CH₂), 22.0 (CH₂), 23.0 (CH₂), 28.9 (CH₂), 33.8 (CH₂), 37.7 (CH₂), 52.5 (2 × CH₃), 53.0 (CH), 53.2 (C), 58.7 (C), 111.0 (CH₂), 129.7 (CH), 148.9 (C), 168.5 (C), 170.1 (C), 176.0 (C), 206.8 (C); HRMS calcd for C₁₈H₂₂O₅ (M⁺) 318.1467, found 318.1461.

cis-Dimethyl 1-methylene-6-oxo-2,3,6a,7,8,9,10,11-octahydro-1H-benzo[c]azulene-4,4(6H)-dicarboxylate (**5c**): white solid; 252 mg, 76% yield; mp 129–130 °C dec; ¹H NMR (400 MHz, CDCl₃) δ 0.93–1.05 (m, 1H), 1.17–1.25 (m, 1H), 1.35–1.46 (m, 1H), 1.50–1.65 (m, 4H), 1.75–1.86 (m, 1H), 1.87–1.95 (m, 1H), 2.08–2.23 (m, 2H), 2.26–2.42 (m, 1H), 2.51–2.63 (m, 1H), 2.70–2.73 (m, 1H), 2.82–2.92 (m, 1H), 3.76 (s, 3H), 3.78 (s, 3H), 4.80 (brs, 1H), 4.90 (brs, 1H), 6.12 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 22.7 (CH₂), 25.0 (CH₂), 25.5 (CH₂), 29.2 (CH₂), 30.6 (CH₂), 33.7 (CH₂), 37.7 (CH₂), 53.1 (CH₃), 53.3 (CH₃), 55.6 (CH), 58.5 (C), 58.8 (C), 109.2 (CH₂), 132.7 (CH), 152.6 (C), 168.8 (C), 170.2 (C), 173.3 (C), 207.9 (C); HRMS calcd for C₁₉H₂₄O₅ (M⁺) 332.1622, found 332.1627. Anal. Calcd for C₁₉H₂₄O₅: C, 68.66; H, 7.28. Found: C, 68.39; H, 7.41.

(3a*R*/5*aR*/S*Z*)-Dimethyl-9-(bromomethylene)-4-oxo-2,3,3a,4,8,9-hexahydro-1H-cyclopenta[c]indene-6,6(7H)-dicarboxylate (**5d**): white solid; 310 mg, 81% yield; mp 143–145 °C dec; ¹H NMR (400 MHz, CDCl₃) δ 1.30–1.45 (m, 1H), 1.64–1.79 (m, 2H), 1.89–2.09 (m, 2H), 2.20–2.35 (m, 1H), 2.40–2.52 (m, 4H), 3.39 (d, *J* = 10 Hz, 1H), 3.77 (s, 3H), 3.81 (s, 3H), 6.03 (s, 1H), 6.28 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 24.5 (CH₂), 29.1 (CH₂), 31.1 (CH₂), 31.8 (CH₂), 36.8 (CH₂), 53.3 (CH₃), 53.4 (CH₃), 55.7 (CH), 58.1 (C), 59.7 (C), 99.1 (CH), 135.7 (CH), 145.7 (C), 169.2 (C), 169.8 (C), 171.9 (C), 210.5 (C); HRMS calcd for C₁₇H₁₉BrO₅ (M⁺) 382.0421, found 382.0416.

General Procedure for the 1,4-Nucleophilic Addition/5-Exocyclization/Cyclopentannulation Process. Preparation of Compounds **6** and **7**. A THF solution (2 mL) of chromium enynyl carbene **1** (1 mmol) was added dropwise at 0 °C to a THF (3 mL) solution of sodium enolate **2**, prepared in situ by the addition of the corresponding propargylmalonate (1.3 mmol) to a NaH (1.6 mmol) THF suspension. The mixture was stirred at rt for 12 h and then quenched with a saturated aqueous solution of NH₄Cl (10 mL). The resulting crude was isolated by extraction with diethyl ether (2 × 10 mL), followed by solvent removal, and then purified by column

chromatography (deactivated SiO₂, hexanes/ethyl acetate, 5:1) to get compounds **6** and **7** (72–92%).

Dimethyl 7-methoxy-6-methyl-2,3-dihydro-1H-cyclopenta[a]pentalene-4,4(5H)-dicarboxylate (**6a**): colorless oil; 280 mg, 92% yield; ¹H NMR (300 MHz, CDCl₃) δ 2.12 (s, 3H), 2.15–2.25 (m, 2H), 2.36–2.47 (m, 2H), 2.67–2.76 (m, 2H), 3.52 (s, 2H), 3.76 (s, 6H), 3.84 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 15.1 (CH₃), 24.4 (CH₂), 26.4 (CH₂), 30.5 (CH₂), 52.4 (CH₂), 52.7 (2 × CH₃), 57.2 (C), 57.8 (CH₃), 120.1 (C), 130.6 (C), 140.4 (C), 141.1 (C), 145.0 (C), 145.7 (C), 170.6 (2 × C); HRMS calcd for C₁₇H₂₀O₅ (M⁺) 304.1311, found 304.1325.

Dimethyl 8-methoxy-1-methyl-4,5,6,7-tetrahydrocyclopenta[a]indene-3,3(2H)-dicarboxylate (**6b**): colorless oil; 264 mg, 83% yield; ¹H NMR (300 MHz, CDCl₃) δ 1.58–1.64 (m, 4H), 2.15 (s, 3H), 2.49–2.55 (m, 2H), 2.58–2.65 (m, 2H), 3.50–3.54 (s, 2H), 3.75 (s, 6H), 3.76 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 15.6 (CH₃), 23.0 (CH₂), 23.1 (CH₂), 23.8 (CH₂), 24.4 (CH₂), 52.7 (2 × CH₃), 53.0 (CH₂), 57.8 (C), 60.5 (CH₃), 126.2 (C), 130.1 (C), 131.4 (C), 140.1 (C), 142.7 (C), 146.2 (C), 170.4 (2 × C); HRMS calcd for C₁₈H₂₂O₅ (M⁺) 318.1467, found 318.1471.

Dimethyl 9-methoxy-1-methyl-5,6,7,8-tetrahydro-2H-cyclopenta[a]azulene-3,3(4H)-dicarboxylate (**6c**): colorless oil; 239 mg, 72% yield; ¹H NMR (300 MHz, CDCl₃) δ 1.54–1.57 (m, 6H), 2.17 (s, 3H), 2.44–2.57 (m, 4H), 3.55 (s, 2H), 3.70 (s, 3H), 3.76 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 15.7 (CH₃), 26.1 (CH₂), 28.5 (CH₂), 29.4 (CH₂), 29.8 (CH₂), 32.8 (CH₂), 52.8 (2 × CH₃), 53.2 (CH₂), 57.8 (C), 62.4 (CH₃), 127.6 (C), 135.7 (C), 140.0 (C), 140.7 (C), 142.7 (C), 145.3 (C), 170.7 (2 × C); HRMS calcd for C₁₉H₂₄O₅ (M⁺) 332.1624, found 332.1627.

Dimethyl 4-methoxy-3,6-dimethylpentalene-1,1(2H)-dicarboxylate (**6d**): colorless oil; 178 mg, 64% yield; ¹H NMR (400 MHz, CDCl₃) δ 1.98 (s, 3H), 2.15 (s, 3H), 3.55 (s, 2H), 3.69 (s, 3H), 3.76 (s, 6H), 5.39 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 13.7 (CH₃), 15.4 (CH₃), 52.7 (2 × CH₃), 53.2 (CH₂), 56.8 (CH₃), 57.7 (C), 110.8 (CH), 125.9 (C), 129.3 (C), 140.2 (C), 147.3 (C), 151.3 (C), 170.7 (2 × C); HRMS calcd for C₁₅H₁₈O₅ (M⁺) 278.1154, found 278.1160.

Dimethyl 4-methoxy-3-methyl-5-phenylpentalene-1,1(2H)-dicarboxylate (**6e**): white solid; 272 mg, 80% yield; mp 103–105 °C; ¹H NMR (400 MHz, CDCl₃) δ 2.31 (s, 3H), 3.66 (s, 2H), 3.73 (s, 3H), 3.79 (s, 6H), 6.46 (s, 1H), 7.21–7.29 (m, 1H), 7.32–7.41 (m, 2H), 7.71–7.78 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 16.7 (CH₃), 53.5 (CH₂), 53.6 (2 × CH₃), 58.5 (C), 61.6 (CH₃), 118.2 (CH), 127.2 (CH), 127.5 (2 × CH), 128.7 (2 × CH), 134.1 (C), 135.3 (C), 136.7 (C), 141.6 (C), 152.5 (C), 170.5 (2 × C); HRMS calcd for C₂₀H₂₀O₅ (M⁺) 340.1311, found 340.1315. Anal. Calcd for C₂₀H₂₀O₅: C, 70.57; H, 5.92. Found: C, 70.35; H, 6.15.

cis-Dimethyl 5-methyl-3-methylene-4-oxo-2,3,3a,4-tetrahydro-1H-pentaleno[2,1-*b*]indole-1,1(5H,9*CH*)-dicarboxylate (**7a**): colorless oil; 219 mg, 62% yield; ¹H NMR (300 MHz, CDCl₃) δ 2.74 (d, *J* = 15.3 Hz, 1H), 3.06 (d, *J* = 15.3 Hz, 1H), 3.60 (s, 3H), 3.80 (s, 3H), 3.90 (m, 1H), 3.93 (s, 3H), 4.85 (m, 1H), 5.11 (brs, 1H), 5.33 (brs, 1H), 7.10–7.21 (m, 1H), 7.26–7.30 (m, 2H), 7.36–7.42 (m, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 30.5 (CH₃), 39.9 (CH₂), 45.1 (CH), 53.3 (CH₃), 53.6 (CH₃), 61.7 (C), 63.6 (CH₃), 111.50 (CH₂), 111.6 (CH), 121.0 (CH), 122.9 (CH), 123.1 (C), 127.5 (CH), 139.6 (C), 140.8 (C), 142.9 (C), 145.6 (C), 170.5 (C), 171.9 (C), 191.9 (C); HRMS calcd for C₂₀H₁₉NO₅ (M⁺) 353.1263, found 353.1271.

cis-Dimethyl 7-oxo-6-((trimethylsilyl)methyl)-3,3a,7,7a-tetrahydro-1H-cyclopenta[a]pentalene-4,4(2H)-dicarboxylate (**7b**): colorless oil; 297 mg, 82% yield; ¹H NMR (300 MHz, CDCl₃) δ 0.12 (s, 9H), 1.33–1.43 (m, 1H), 1.58–1.78 (m, 4H), 1.86–1.93 (m, 1H), 2.81–2.89 (m, 1H), 3.10 (s, 2H), 3.54–3.58 (m, 1H), 3.79 (s, 3H), 3.80 (s, 3H), 6.36 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 0.2 (3 × CH₃), 25.5 (CH₂), 27.4 (CH₂), 29.4 (CH₂), 30.3 (CH₂), 48.2 (CH), 52.2 (CH), 53.6 (CH₃), 53.7 (CH₃), 61.5 (C), 99.7 (C), 101.1 (C), 135.1 (CH), 169.5 (C), 173.8 (2 × C), 212.0 (C); HRMS calcd for C₁₉H₂₆O₅Si (M⁺) 362.1550, found 362.1556.

■ ASSOCIATED CONTENT

■ Supporting Information

¹H and ¹³C NMR spectra for all compounds; X-ray data and CIF for compounds **5c**, **5d**, and **6e**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Phone: +34-98-510-4952. Fax +34-98-510-3446. E-mail: alss@uniovi.es.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was partially supported by the Governments of Spain (CTQ2010-20517-C02-01; undergraduate fellowship to T.S.-R.) and Principado de Asturias (IB08-088; Severo Ochoa-PCTI predoctoral fellowship to A.A.-F).

■ REFERENCES

- (1) (a) Tietze, L. F.; Brasche, G.; Gericke, K. *Domino Reactions in Organic Synthesis*; Wiley-VCH: Weinheim, 2006; p 672. (b) Nicolau, K. C.; Edmonds, D. J.; Bulger, P. G. *Angew. Chem., Int. Ed.* **2006**, *45*, 7134–7186. (c) Poulin, J.; Grisé-Bard, C. M.; Barriault, L. *Chem. Soc. Rev.* **2009**, *38*, 3092–3101. (d) Nicolau, K. C.; Chen, J. S. *Chem. Soc. Rev.* **2009**, *38*, 2993–3009. (e) Ruiz, M.; López-Alvarado, P.; Giorgi, G.; Menéndez, J. C. *Chem. Soc. Rev.* **2011**, *40*, 3445–3454.
- (2) (a) Barluenga, J. *Pure Appl. Chem.* **2002**, *74*, 1317–1325. (b) Barluenga, J.; Rodríguez, F.; Fañanás, F.; Flórez, J. *Top. Organomet. Chem.* **2004**, *13*, 59–121. (c) Barluenga, J.; Fernández-Rodríguez, M. A.; Aguilar, E. J. *Organomet. Chem.* **2005**, *690*, 539–587. (d) Dötz, K. H.; Stendel, J. *Chem. Rev.* **2009**, *109*, 3227–3274.
- (3) Barluenga, J.; García-Rodríguez, J.; Martínez, S.; Suárez-Sobrinó, A. L.; Tomás, M. *Chem.—Eur. J.* **2006**, *12*, 3201–3210.
- (4) (a) Barluenga, J.; Aznar, F.; Palomero, M. A. *Angew. Chem., Int. Ed.* **2000**, *39*, 4346–4348. (b) Barluenga, J.; Aznar, F.; Palomero, M. A. *Chem.—Eur. J.* **2002**, *8*, 4149–4163. (c) Barluenga, J.; Martínez, S.; Suárez-Sobrinó, A. L.; Tomás, M. *Org. Lett.* **2008**, *10*, 677–679.
- (5) Barluenga, J.; Aznar, F.; Barluenga, S.; Fernández, M.; Martín, A.; García-Granda, S.; Piñera-Nicolás, A. *Chem.—Eur. J.* **1998**, *4*, 2280–2298.
- (6) For cascade processes involving 1,4-addition of heteroatom nucleophiles to enynyl Fischer carbenes, see: (a) Wu, H.-P.; Auman, R.; Frölich, R.; Wegelius, E.; Saavenketo, P. *Organometallics* **2000**, *18*, 2373–2381. (b) Wu, H.-P.; Aumann, R.; Venne-Dunker, S.; Saarenketo, P. *Eur. J. Org. Chem.* **2000**, 463–3473. (c) Wu, H.-P.; Auman, R.; Frölich, R.; Saarenketo, P. *Chem.—Eur. J.* **2001**, *7*, 700–710.
- (7) Barluenga, J.; Álvarez-Fernández, A.; Martínez, S.; Suárez-Sobrinó, A. L.; Tomás, M. *Tetrahedron Lett.* **2009**, *50*, 3606–3608.
- (8) Barluenga, J.; Álvarez-Fernández, A.; Suárez-Rodríguez, T.; Suárez-Sobrinó, A. L.; Tomás, M. *Org. Lett.* **2013**, *15*, 488–491.
- (9) All attempts using acyclic enynyl FFCs gave rise to a complex mixture of products.
- (10) See the Supporting Information
- (11) Unfortunately, in the case of homopropargyl derivatives with R³ ≠ H, Br the reaction did not work, but a mixture of decomposition products was always formed. Probably, the severe steric hindrance imposed by the congested framework along with the resulting *cis* stereochemistry makes the final cyclization unlikely to occur.
- (12) 5-*exo*-W(CO)₅-amine catalyzed cyclizations of ω -alkynyl silyl enol ethers were reported; see: Kusama, H.; Yamabe, H.; Iwasawa, N. *Org. Lett.* **2002**, *4*, 2569–2571.
- (13) For a recent one-step construction of angular tricyclic 5/5/6 compounds, see: Lin, M.; Li, F.; Jiao, L.; Yu, Z.-X. *J. Am. Chem. Soc.* **2011**, *133*, 1690–1693.
- (14) For instance, the 5/5/6 angular system is present in the lycopodium alkaloid family; for recent total syntheses, see: (a) Sha, C.-K.; Lee, F.-K.; Chang, C.-J. *J. Am. Chem. Soc.* **1999**, *121*, 9875–9876. (b) Yen, C.-F.; Liao, C.-C. *Angew. Chem., Int. Ed.* **2002**, *41*, 4090–4093. (c) Kozaka, T.; Miyakoshi, N.; Mukai, C. *J. Org. Chem.* **2007**, *72*, 10147–10154.
- (15) For a review on polyquinanes, see: Mehta, G.; Srikrishna, A. *Chem. Rev.* **1997**, *97*, 651–719.
- (16) The corresponding methoxycyclopentadiene derivatives **6** cannot be isolated, but they suffer hydrolysis to the ketones **7a,b** under the standard quenching and purification steps.
- (17) A 5-*exo* cyclization of an allenolate intermediate was proposed in a recent example of an intramolecular allenolate Rauhut–Currier reaction; see: MacKay, A. J. A.; Landis, Z. C.; Motika, S. E.; Kench, M. H. *J. Org. Chem.* **2012**, *77*, 7768–7774.
- (18) It should be mentioned that all attempts to reproduce this delayed-protonation strategy to the case of enynyl carbenes **1** and homopropargylicmalonates **3** (see Scheme 2) were unsuccessful. Most probably, the resulting allenyl metalate intermediate **1b** does not overwhelm the reaction conditions and evolves to decomposition products.
- (19) Arai, S.; Koike, Y.; Hade, H.; Nishida, A. *J. Am. Chem. Soc.* **2010**, *132*, 4522–4523.
- (20) Buisine, O.; Aubert, C.; Malacria, M. *Chem.—Eur. J.* **2001**, *7*, 3517–3525.
- (21) (a) Finaru, A.; Berthault, A.; Besson, T.; Guillaumet, G.; Bertaina-Raboin, S. *Tetrahedron Lett.* **2002**, *43*, 787–790. (b) Pardo-Rodríguez, V.; Marco-Martínez, J.; Buñuel, E.; Cárdenas, D. J. *Org. Lett.* **2009**, *11*, 4548–4551.