

Hard-soft interactions in early transition-metal thioether macrocyclic chemistry: spectroscopic and extended X-ray absorption fine structure studies on chromium(III) thioether complexes


Simon J. A. Pope, Neil R. Champness and Gillian Reid *

Department of Chemistry, University of Southampton, Highfield, Southampton SO17 1BJ, UK

A series of chromium(III) thioether macrocyclic complexes of the form $[\text{CrX}_3(\text{[9]aneS}_3)]$, $[\text{CrX}_3(\text{[10]aneS}_3)]$, $[(\text{CrX}_3)_2(\mu\text{-[18]aneS}_6)]$ ($\text{X} = \text{Cl}$ or Br), $[\text{CrX}_2(\text{[14]aneS}_4)]\text{PF}_6$ and $[\text{CrX}_2(\text{[16]aneS}_4)]\text{PF}_6$ ($\text{X} = \text{Cl}$, Br or I) ($\text{[9]aneS}_3 = 1,4,7\text{-trithiacyclononane}$, $\text{[10]aneS}_3 = 1,4,7\text{-trithiacyclodecane}$, $\text{[18]aneS}_6 = 1,4,7,10,13,16\text{-hexathiacyclooctadecane}$, $\text{[14]aneS}_4 = 1,4,8,11\text{-tetrathiacyclotetradecane}$, $\text{[16]aneS}_4 = 1,5,9,13\text{-tetrathiacyclohexadecane}$) has been prepared and characterised by IR and UV/VIS spectroscopy, microanalyses, and in some cases electrospray mass spectrometry. Chromium K-edge extended X-ray absorption fine structure studies have provided information on the Cr–X and Cr–S bond-length distributions in a selection of these compounds, and these, together with the relatively low Dq values derived from UV/VIS spectroscopy, confirm rather weak chromium(III)–thioether interactions in these unusual species involving hard metal and soft ligand combinations.

In contrast to the vast number of chromium(III) complexes with hard nitrogen- and oxygen-donor ligands,¹ there are very few examples of complexes with soft donor ligands such as phosphines^{2–5} or thioethers.^{6–9} This is principally due to the severe incompatibility of the hard chromium(III) centre with the soft phosphine or thioether functions.

Thioether macrocycles have been intensively studied in recent years since the ability of these ligands to stabilise unusual oxidation states and co-ordination geometries was noted. Notably, however, to date thioether macrocyclic chemistry has been dominated by middle and late d- and p-block ions.¹⁰ Most significantly, it has been noted that while acyclic thioethers act as relatively poor ligands to metal ions, when the thioether functions are included within a macrocyclic framework they tend to co-ordinate much more readily often to yield quite stable complexes. We have become interested in the possibility of utilising the additional stability characteristic of macrocyclic complexes to stabilise metal/ligand combinations which are usually considered incompatible. In this context we have been investigating the reactions of Cr^{III} with various thioether macrocycles to determine whether the incorporation of macrocyclic environments would render such hard-soft interactions more accessible and to establish the nature of the S (thioether) to Cr^{III} interaction. A few examples of macrocyclic thioether complexes of early transition-metal ions have been reported. These include carbonyl derivatives of Mo^0 and W^0 ,^{11,12} $[\text{MoCl}_3(\text{[9]aneS}_3)]$ ($\text{[9]aneS}_3 = 1,4,7\text{-trithiacyclononane}$),¹² the metal(II) species $[\text{M}(\text{CO})_3(\text{[9]aneS}_3)]^+$ ($\text{M} = \text{Mo}$ or W),¹³ $[\text{CrCl}_3(\text{[9]aneS}_3)]$, $[\text{Cr}(\text{SO}_3\text{CF}_3)_3(\text{[9]aneS}_3)]^7$ and some examples with the formula $[\text{CrCl}_3\text{L}]$ ($\text{L} = \text{S}_4$, S_5 or S_6 donor macrocycle) which are thought to involve tridentate co-ordination.⁸ The crystal structure of the vanadium(IV) species $[\text{VOCl}_2(\text{[9]aneS}_3)]$ has also been reported, and shows *fac* tridentate [9]aneS_3 ligation, giving V–S (*trans* to O) 2.634(5), (*trans* to Cl) 2.470(5) (mean), V–Cl 2.295(5) (mean) and V–O 1.579(4) Å.¹⁴ Very recently we have also reported the synthesis and crystal structure of the corresponding vanadium(III) species $[\text{VCl}_3(\text{[9]aneS}_3)]$ which also involves *fac* tridentate [9]aneS_3 co-ordination, giving V–S 2.486(3), 2.515(2), 2.500(2), V–Cl 2.279(3), 2.310(3), 2.281(3) Å.¹⁵ The only structurally characterised examples of chromium(III) thioether macrocyclic species are $[\text{CrCl}_3(\text{[18]aneS}_6)]$ ($\text{[18]aneS}_6 = 1,4,7,10,13,16\text{-hexathiacyclooctadecane}$) which involves *fac* tridentate co-ordination of the hexathia crown to Cr^{III} , leaving three free thioether donors (see text below)⁸ and *cis*-



$[\text{CrCl}_2(\text{[14]aneS}_4)]\text{PF}_6$ ($\text{[14]aneS}_4 = 1,4,8,11\text{-tetrathiacyclotetradecane}$), the structure of which we have reported, involving S_4Cl_2 co-ordination at Cr^{III} with Cr–S 2.393(1), 2.407(2), Cr–Cl 2.295(1) Å.⁹ We now report the preparation and spectroscopic characterisation of a series of chromium(III) thioether macrocyclic complexes incorporating [9]aneS_3 , [10]aneS_3 (1,4,7-trithiacyclodecane), [14]aneS_4 , [16]aneS_4 (1,5,9,13-tetrathiacyclohexadecane) and [18]aneS_6 , together with chromium K-edge extended X-ray absorption fine structure (EXAFS) studies on selected examples.

Results and Discussion

Reaction of $[\text{CrX}_3(\text{thf})_3]$ ($\text{X} = \text{Cl}$ or Br , $\text{thf} = \text{tetrahydrofuran}$) with [9]aneS_3 or [10]aneS_3 in a 1:1 molar ratio in CH_2Cl_2 solution gives the products $[\text{CrX}_3(\text{[9]aneS}_3)]$ or $[\text{CrX}_3(\text{[10]aneS}_3)]$ as powdered solids in high yield. Similarly, using $[\text{CrX}_3(\text{thf})_3]$ and [18]aneS_6 in a 2:1 molar ratio yields the binuclear species $[(\text{CrX}_3)_2(\mu\text{-[18]aneS}_6)]$. We found that in order to promote co-ordination through the thioether donors it was necessary to remove the thf from the reaction mixture, otherwise the harder O-donor thf ligand would bind preferentially. These neutral, highly coloured products proved to be very poorly soluble in both chlorinated and organic solvents, and attempts to dissolve them in dimethyl sulfoxide (dmsO) resulted in decomposition.

They are relatively stable in the solid state when stored in a dry atmosphere, hence all were stored in an N₂-purged glove-box. The IR spectra of these species provided evidence for the presence of co-ordinated thioether, and in the case of the chloro and bromo derivatives peaks were observed in most cases in the region 200–400 cm⁻¹, which might be assigned tentatively to $\nu(\text{Cr-X})$. However, given that in some cases these peaks were rather weak, and since $\nu(\text{Cr-S})$ would also be expected to appear in this region, definitive assignments were not possible. Owing to their poor solubilities and sensitivity to moisture, FAB and electrospray mass spectra proved unhelpful. However, microanalytical data and UV/VIS spectroscopic data consistent with the proposed formulations were obtained in all cases.

Reaction of $[\text{CrX}_3(\text{thf})_3]$ (X = Cl or Br) with $[\text{14}] \text{aneS}_4$ or $[\text{16}] \text{aneS}_4$ in dry MeNO₂ solution in the presence of 1 molar equivalent of TlPF₆ resulted in the formation of an intense blue-green or green solution for $[\text{14}] \text{aneS}_4$ and $[\text{16}] \text{aneS}_4$ respectively, and a white precipitate of TlCl. Following removal of the TlCl by filtration, the complexes $[\text{CrX}_2([\text{14}] \text{aneS}_4)]\text{PF}_6$ and $[\text{CrX}_2([\text{16}] \text{aneS}_4)]\text{PF}_6$ were isolated by reducing the volume of solvent to ca. 3 cm³ *in vacuo* and addition of dry diethyl ether. The complexes $[\text{CrI}_2([\text{14}] \text{aneS}_4)]\text{BF}_4$ and $[\text{CrI}_2([\text{16}] \text{aneS}_4)]\text{BF}_4$ were obtained by addition of the tetrathioether macrocycle to a solution of $[\text{Cr}(\text{thf})_6][\text{BF}_4]_3$ in dry MeNO₂. This precursor was generated *in situ* by treatment of $[\text{CrCl}_3(\text{thf})_3]$ with 3 molar equivalents of AgBF₄ in thf solution, followed by removal of the AgCl by filtration, addition of MeNO₂ and removal of the thf *in vacuo*.¹⁶ Treatment of this MeNO₂ solution with 2 molar equivalents of NEt₄I gave a green-brown solution from which the products were obtained by addition of diethyl ether. These cationic tetrathioether compounds are considerably more soluble than the neutral species described above, dissolving readily in dry MeCN and MeNO₂ and partially in dry acetone. Also, the ionic products are noticeably more sensitive to hydrolysis, especially in solution. Attempts to prepare similar cationic species using the smaller tetrathioether macrocycle $[\text{12}] \text{aneS}_4$ (1,4,7,10-tetrathiacyclododecane) under similar conditions were not successful, yielding only a purple insoluble product thought to be $[\text{CrX}_3([\text{12}] \text{aneS}_4)]$ involving tridentate co-ordination of the macrocycle.

The IR spectra of the cationic tetrathioether complexes confirmed the presence of PF₆⁻ anion [$\nu(\text{PF}_6^-)$ 840, $\delta(\text{PF}_6^-)$ 558 cm⁻¹] and co-ordinated macrocycle in each case. For X = Cl and Br the spectra also revealed weak features in the range 200–400 cm⁻¹ which can tentatively be assigned to $\nu(\text{Cr-X})$ [$\nu(\text{Cr-I})$ is expected to occur below 200 cm⁻¹]. The fast atom bombardment (FAB) or electrospray mass spectra showed that the most intense peaks had the correct isotopic distributions at m/z = 480 corresponding to $[\text{Cr}^{79}\text{Br}^{81}\text{Br}([\text{14}] \text{aneS}_4)]^+$, 418 to $[\text{Cr}^{35}\text{Cl}_2([\text{16}] \text{aneS}_4)]^+$ and 508 to $[\text{Cr}^{79}\text{Br}^{81}\text{Br}([\text{16}] \text{aneS}_4)]^+$, as well as peaks at lower m/z due to loss of Cl or Br. No useful information was obtained from the mass spectra of the iodo derivatives, possible due to the increased sensitivity of these particular species to hydrolysis. Satisfactory microanalytical data were obtained for each of these compounds, confirming the proposed formulations.

There are very few structurally characterised examples of chromium(III) complexes involving thioether co-ordination. Gahan and co-workers¹⁷ reported the structure of a Cr^{III}-S (thioether) complex. In their system the metal ion is encapsulated by an N₅S donor caging ligand which enforces the thioether co-ordination [Cr-S 2.399(2) Å]. Chakravorty and co-workers¹⁸ have reported the preparation of a series of octahedral chromium(III) compounds involving S₂N₂O₂ chelating ligands, including a crystal structure of one example which shows $d(\text{Cr-S}) = 2.417(3)$ and $2.445(3)$ Å. Both of these systems involve much harder donor sets than those of the present macrocyclic thioether complexes. Of more direct relevance to our work, the structure of the mononuclear complex $[\text{CrCl}_3-$




Fig. 1 View of the structure of $\text{cis-}[\text{CrBr}_2([\text{14}] \text{aneS}_4)]^+$ with the numbering scheme adopted (atoms marked with * are related by a crystallographic two-fold axis) [Cr-Br(1) 2.441(8), Cr-S(1) 2.41(1), Cr-S(2) 2.39(1) Å; Br(1)-Cr-Br(1*) 95.9(3), S(1)-Cr-S(1*) 179.7(8), S(1)-Cr-S(2) 84.5(4), S(1)-Cr-S(2*) 95.2(5), S(2)-Cr-S(2*) 80.5(6)°]

$[\text{18}] \text{aneS}_6]$ which involves an S₃Cl₃ donor set has been reported, Cr-S 2.459(3), 2.442(5), 2.440(5), Cr-Cl 2.305(5), 2.279(5), 2.291(5) Å.⁸ The Cr-S and Cr-Cl bond lengths compare well with those in $\text{cis-}[\text{CrCl}_2([\text{14}] \text{aneS}_4)]\text{PF}_6$, the structure of which we have reported, in which the geometry at Cr^{III} is a distorted octahedron, with *cis*-chloride ligands and the tetrathia macrocycle adopting a folded conformation, Cr-Cl(1) 2.295(1), Cr-S(1) 2.393(1), Cr-S(2) 2.407(2) Å.⁹ We noted previously that the trend in $d(\text{M-Cl})$ vs. $d(\text{M-S})$ in this compound is opposite to that observed in complexes of $[\text{14}] \text{aneS}_4$ with late transition-metal ions. Thus, in $\text{cis-}[\text{CrCl}_2([\text{14}] \text{aneS}_4)]^+$ $d(\text{Cr-S}) > d(\text{Cr-Cl})$ (by ca. 0.1 Å), whereas in $\text{cis-}[\text{RhCl}_2([\text{14}] \text{aneS}_4)]^+$,¹⁹ $\text{cis-}[\text{IrCl}_2([\text{14}] \text{aneS}_4)]^+$ ²⁰ and $\text{cis-}[\text{RuCl}_2([\text{14}] \text{aneS}_4)]^+$ ²¹ $d(\text{M-S}) < d(\text{M-Cl})$ (by ca. 0.1 Å), suggesting rather more favourable M-S interactions in the later transition-metal complexes compared to the chromium species.

In an attempt to confirm the stereochemistry at the Cr^{III} single-crystal X-ray diffraction data were collected† on a small, very weakly diffracting crystal of $[\text{CrBr}_2([\text{14}] \text{aneS}_4)]\text{PF}_6$. The data were very weak and hence the final residuals rather high and it was not possible to refine the F and C atoms anisotropically due to the low data: parameters ratio. However, the analysis was sufficient to establish the stereochemistry of the $[\text{CrBr}_2([\text{14}] \text{aneS}_4)]^+$ cation. The structure of this species shows (Fig. 1) a very similar arrangement to that seen previously for $[\text{CrCl}_2([\text{14}] \text{aneS}_4)]^+$, with distorted-octahedral co-ordination at Cr^{III} via two mutually *cis*-bromide ligands and the four

† C₁₀H₂₀Br₂CrF₆PS₄, $M = 625.3$, monoclinic $C2/c$, $a = 10.808(3)$, $b = 11.514(2)$, $c = 15.879(3)$ Å, $\beta = 92.71(2)^\circ$, $U = 1974$ Å³, $Z = 4$, $D_c = 2.104$ g cm⁻³, $T = 150$ K, small dark blue block, $0.30 \times 0.20 \times 0.10$ mm, $\mu = 51.91$ cm⁻¹, $F(000) = 1228$. Data collection used a Rigaku AFC7S four-circle diffractometer equipped with an Oxford Systems low-temperature attachment. ω -2 θ Scans; 4571 unique reflections measured ($2\theta_{\text{max}} = 50^\circ$), 515 with $I > 2\sigma(I)$ used in all calculations. The structure was solved by direct methods²² and refined (based on F) using iterative cycles of full-matrix least squares²³ which revealed one half cation (with the Cr atom lying on a two-fold axis) and one half PF₆⁻ anion (with the P atom occupying a crystallographic inversion centre) in the asymmetric unit. At isotropic convergence the data were corrected for absorption using DIFABS²⁴ (maximum, minimum transmission factors = 1.000, 0.582), and the Cr, Br, P and S atoms were then refined anisotropically and H atoms included in fixed, calculated positions. The weighting scheme $w^{-1} = \sigma^2(F)$ gave satisfactory agreement analyses. This model refined to R , $R' = 0.089$, 0.090 respectively and $S = 2.23$ for 71 parameters and the final Fourier-difference map showed residual electron-density peaks of 1.19 and -1.18 e Å⁻³.

Table 1 Chromium K-edge EXAFS structural data^a for chromium(III) thioether macrocyclic compounds

Complex	$d(\text{Cr-S})$ ^b /Å	$2\sigma^2/\text{\AA}^2$	$d(\text{Cr-X})$ ^b /Å	$2\sigma^2/\text{\AA}^2$	R^d	Fit index ^e
[CrCl ₃ ([9]aneS ₃)]	2.396(11)	0.0121(20)	2.306(9)	0.0119(16)	22.5	2.9
[CrBr ₃ ([9]aneS ₃)]	2.401(4)	0.0064(14)	2.450(3)	0.0095(6)	21.2	3.5
[CrCl ₃ ([10]aneS ₃)]	2.361(3)	0.0076(6)	2.239(8)	0.0257(23)	20.1	3.8
<i>cis</i> -[CrCl ₂ ([14]aneS ₄)]PF ₆	2.394(5)	0.0199(11)	2.291(4)	0.0086(6)	17.8	1.9
	2.40 (average) ^f		2.295(1) ^f			
<i>cis</i> -[CrBr ₂ ([14]aneS ₄)]PF ₆	2.409(3)	0.0133(7)	2.446(1)	0.0072(2)	16.3	1.4
[Cr ₂ Cl ₆ ([18]aneS ₆)]	2.424(4)	0.0162(10)	2.291(2)	0.0094(4)	19.8	1.9
[CrCl ₃ (thf) ₃]	2.009(13) ^g	0.0245(20)	2.295(4)	0.0092(41)	23.3	4.1
[CrBr ₃ (thf) ₃]	1.998(4) ^g	0.0105(8)	2.458(2)	0.0110(2)	18.6	3.3

^a Recorded in transmission mode on station 7.1 or 8.1, using powdered samples diluted with BN; AFAC (a factor compensating for the reduction in amplitude due to multi-electron processes) = 0.80 for all refinements. ^b Standard deviations in parentheses. Note that the systematic errors in bond distances arising from data collection and analysis procedures are ± 0.02 – 0.03 Å for well defined co-ordination shells. ^c Debye–Waller factor. ^d Defined as $[\int(\chi^T - \chi^E)^2 dk / \int \chi^E k^3 dk]$ (where T is theoretical and E is experimental) 100%. ^e Defined as $\Sigma[(\chi^T - \chi^E)/k_i]^2$. ^f X-Ray crystallographic data from ref. 9. ^g $d(\text{Cr-O})$.




Fig. 2 Background-subtracted chromium K-edge EXAFS data (a) and the corresponding Fourier transform (b) for *cis*-[CrBr₂([14]aneS₄)]PF₆ (solid line, experimental; dashed line, calculated data)

thioether donors of the macrocycle, which adopts a folded arrangement, Cr–Br(1) 2.441(8), Cr–S(1) 2.41(1), Cr–S(2) 2.39(1) Å. As a result of the poor crystal quality the geometric parameters have high associated estimated standard deviations, and hence comparisons with related species require caution. However, the Cr–S distances are similar to those in both *cis*-[CrCl₂([14]aneS₄)]⁹ and [CrCl₃([18]aneS₆)],⁸ while $d(\text{Cr-Br})$ is *ca.* 0.15 Å longer than $d(\text{Cr-Cl})$ in these compounds. The angles subtended at Cr^{III} show a marked deviation from the 90 and 180° expected for a regular octahedron. Again this is similar to the case in *cis*-[CrCl₂([14]aneS₄)]⁹.

Chromium K-edge EXAFS studies

Owing to the limited solubilities displayed by many of the compounds isolated, and their sensitivity to moisture, in most cases crystals suitable for an X-ray study could not be obtained. However, given the lack of structural data on compounds of

this type, we proposed that chromium K-edge EXAFS data would provide useful information regarding the metal–ligand bond lengths for the first co-ordination sphere, *i.e.* $d(\text{Cr-S})$ and $d(\text{Cr-X})$. Importantly, the spectroscopic studies carried out in parallel provided key information concerning the donor sets involved in these products. Details of the refined EXAFS data for the complexes are given in Table 1 and Fig. 2 shows a typical example. Chromium K-edge EXAFS spectra were also recorded for the [CrX₃(thf)₃] model compounds and for [CrCl₂([14]aneS₄)]PF₆ in order to compare the Cr–S and Cr–X distances derived from this method with the *average* values obtained from X-ray crystallographic studies. The EXAFS data for the thf adducts, [CrX₃(thf)₃], were satisfactorily modelled to a first shell of three oxygens giving Cr–O distances of 2.01 and 2.00 Å for X = Cl and Br respectively, with a second shell of three halides giving $d(\text{Cr-Cl}) = 2.29$ and $d(\text{Cr-Br}) = 2.46$ Å. In the case of *cis*-[CrCl₂([14]aneS₄)]PF₆ the data were satisfactorily modelled to an S₄Cl₂ donor set with $d(\text{Cr-Cl}) = 2.29$, $d(\text{Cr-S}) = 2.39$ Å. These results correlate very well with the *average* Cr–S, Cr–O or Cr–X distances derived from the X-ray crystallographic studies.^{9,25}

For [CrX₃([9]aneS₃)], [CrCl₃([10]aneS₃)] and [(CrCl₃)₂(μ-[18]aneS₆)] the EXAFS data were modelled for three sulfurs and three halogen atoms (Cl or Br as appropriate), while for *cis*-[CrBr₂([14]aneS₄)]PF₆ two shells were modelled, one with four sulfurs and one with two bromides. For the complexes *cis*-[CrX₂([14]aneS₄)]PF₆ no attempt was made to split the sulfur shell to take account of the slightly different distances expected for the Cr–S (*trans* to X) and Cr–S (*trans* to S) since X-ray structural studies have shown that the difference is very small (in this case only *ca.* 0.01 Å). In all cases the EXAFS data gave $d(\text{Cr-S})$ values very similar to those obtained crystallographically for chromium(III)–thioether interactions (*ca.* 2.4 Å) and significantly longer than $d(\text{Cr-Cl})$ (*ca.* 2.3 Å). This is consistent with the thioether interaction being rather weak, as might be anticipated for such a hard metal–soft ligand combination. The Cr–Br bond lengths of *ca.* 2.45 Å are in accord with the trend expected on the basis of the increased size of bromine over chlorine and also agree very well with $d(\text{Cr-Br})$ derived from the X-ray crystallographic study on *cis*-[CrBr₂([14]aneS₄)]PF₆ described above.

Electronic spectra

The UV/VIS spectra were recorded by diffuse reflectance (and also in solution for the soluble species), and the data for all of the new compounds are presented in Table 2. In an *O_h* environment, three spin-allowed d–d bands are predicted for chromium(III) species, $^4A_{2g} \longrightarrow ^4T_{2g}(v_1)$, $^4A_{2g} \longrightarrow ^4T_{1g}(F)(v_2)$ and $^4A_{2g} \longrightarrow ^4T_{1g}(P)(v_3)$, of which the latter is rarely observed as it is often obscured by intense charge-transfer bands.²⁶

Table 2 Electronic spectroscopic data (cm⁻¹)^a for chromium(III) thioether macrocyclic compounds

Compound	Colour	ν_1	ν_2	Dq	B
[CrCl ₃ ([9]aneS ₃)]	Purple	14 450	20 080	1 445	556
[CrBr ₃ ([9]aneS ₃)]	Blue-purple	14 350	19 230	1 435	463
[CrCl ₃ ([10]aneS ₃)]	Purple	14 750	20 240	1 473	590
[CrBr ₃ ([10]aneS ₃)]	Dark purple	14 140	19 120	1 414	505
[(CrCl ₃) ₂ ([18]aneS ₆)]	Pink-purple	14 450	19 530	1 445	516
[(CrBr ₃) ₂ ([18]aneS ₆)]	Blue-purple	14 250	18 620	1 425	491
[CrCl ₂ ([14]aneS ₄)]PF ₆ ^b	Blue-green	16 500	22 400	1 650	536
[CrBr ₂ ([14]aneS ₄)]PF ₆ ^b	Dark blue	16 500	21 550	1 650	485
[CrI ₂ ([14]aneS ₄)]PF ₆ ^b	Olive green	16 700 (sh)	—	—	—
[CrCl ₂ ([16]aneS ₄)]PF ₆ ^b	Green	15 820	21 790	1 582	546
[CrBr ₂ ([16]aneS ₄)]PF ₆ ^b	Dark green	15 510	20 940	1 551	554
[CrI ₂ ([16]aneS ₄)]PF ₆ ^b	Light green	16 100	—	—	—

^a Recorded for solid diluted with BaSO₄ by diffuse reflectance, except where indicated; sh = shoulder; $C = 4.5B$. ^b Recorded in MeNO₂ solution.

Noticeable splitting of these bands usually occurs upon reducing the symmetry from O_h to D_{4h} (i.e. a *trans*-L₄X₂ donor set), whereas reducing the symmetry to C_{2v} (i.e. a *cis*-L₄X₂ donor set) usually just results in some broadening of the transitions.¹⁸ Also, reducing the symmetry from O_h to C_{3v} (i.e. a *fac*-L₃X₃ donor set as in the trithia and binuclear hexathia species) should theoretically result in splitting of the bands, although in practice only broadening of the bands is observed usually. Owing to the geometric constraints of the ligands used only the *fac* isomer is expected in each of the trithia species.

The UV/VIS spectra of the chromium(III) thioether species in this work are typical of chromium(III) complexes with an L₄Cl₂ donor set, showing two d-d transitions. The absence of any noticeable splitting in the UV/VIS spectra of [CrX₂([14]aneS₄)]PF₆ or [CrX₂([16]aneS₄)]PF₆ either in solution or the solid state suggests that these exist as the *cis* isomers. This has been confirmed for [CrX₂([14]aneS₄)]⁺ through single-crystal structure analyses on the chloride and bromide derivatives, however in the absence of such data there is some uncertainty over the geometric arrangement (*cis* vs. *trans*) in the complexes of [16]aneS₄.

For each of the chromium(III) compounds Dq is obtained directly from ν_1 , while the Racah parameter (B) can be extracted from the appropriate Tanabe–Sugano diagram.²⁶ It should be noted, however, that there is a considerable error associated with B of ca. ± 50 cm⁻¹. From the values of Dq obtained it can be seen that for the neutral chromium(III) species with S₃X₃ donor sets Dq is smaller than for the species with S₄X₂ donor sets, consistent with the replacement of π -donor halogen ligands with thioether donors (probably weak σ donors only in these compounds). Consistent with this, Dq is even smaller for the hexachloro species [CrCl₆]³⁻ (1318).²⁷ The values of Dq for the thioether compounds are considerably smaller than for other chromium(III) species with harder N- and O-donor ligands, e.g. [Cr(H₂NCH₂CH₂NH₂)₃]³⁺ ($Dq = 2180$), [Cr(bipy)₃]³⁺ (bipy = 2,2'-bipyridine) ($Dq = 2340$), and [Cr(H₂O)₆]³⁺ ($Dq = 1740$).^{27,28} For chromium(III) compounds with other soft donor sets such as phosphines the Dq values generally compare well with those for the thioether species in this work, e.g. for [CrCl₃{MeC(CH₂PPh₂)₃}] (P₃Cl₃ donor set) $Dq = 1600$.³ The values are also very similar to those for the other chromium(III)–thioether species known, e.g. [CrCl₃{MeC(CH₂SMe)₃}] (S₃Cl₃ donor set) $Dq = 1470$ cm⁻¹.³ These trends are consistent with only weak interactions between the Cr^{III} and the soft thioether functions.

For all of the compounds studied B lies in the range 463–590 cm⁻¹. These values are rather smaller than those observed for [Cr(H₂O)₆]³⁺ ($B = 710$) and [Cr(dmsO)₆]³⁺ ($B = 685$ cm⁻¹).²⁷

Conclusion

These results confirm that, despite the anticipated incompatibility between hard chromium(III) centres and soft thioether func-

tions, a series of relatively stable mono- and di-nuclear complexes involving S₃X₃ and S₄X₂ donor sets has been successfully synthesized, adding considerably to the previously very limited range of such compounds known. Chromium K-edge EXAFS studies have proved to be extremely useful, providing structural data on the Cr–S and Cr–X bond distances in this unusual series of compounds. The values derived from the EXAFS measurements compare very well with those from X-ray crystallographic studies where such information is available. In particular, the Cr–S bond lengths are apparently largely independent of the macrocycle ring size or number of donor atoms available, and the Cr–S bond lengths, together with the Dq values, indicate relatively weak Cr^{III}–S (thioether) interactions in these compounds. Current work investigating the possibility of substituting the halide ligand to generate organometallic chromium–thioether macrocyclic derivatives is underway.

Experimental

Infrared spectra were measured as KBr or CsI discs or as Nujol mulls between CsI plates using a Perkin-Elmer 983 spectrometer over the range 200–4000 cm⁻¹, mass spectra by FAB using 3-nitrobenzyl alcohol as matrix on a VG Analytical 70-250-SE normal-geometry double-focusing spectrometer or by electrospray using a Hewlett-Packard Series 1050 spectrometer and UV/VIS spectra in solution using 1 cm path length quartz cells or by diffuse reflectance using samples diluted with BaSO₄ on a Perkin-Elmer Lambda19 spectrophotometer. Microanalytical data were obtained using a F & M model 185 C, H and N analyser. The EXAFS measurements were made at the Daresbury Laboratory, operating at 2.0 GeV (ca. 3.2×10^{-7} J) with typical currents of 200 mA. Chromium K-edge data were collected on stations 7.1 and 8.1 using a silicon(111) order-sorting monochromator, with harmonic rejection achieved by stepping off the peak of the rocking curve by 50% of the full height level. Data were collected in transmission mode from samples diluted with boron nitride and mounted between Sellotape in 1 mm aluminium holders.

Diethyl ether was dried over sodium wire. The compounds [CrCl₃(thf)₃],²⁹ [CrBr₃(thf)₃], [Cr(thf)₆][BF₄]₃¹⁶ and *cis*-[CrCl₂([14]aneS₄)]PF₆⁹ were prepared by the literature methods.

Preparations

(a) [CrCl₃([9]aneS₃)]. To a solution of [CrCl₃(thf)₃] (0.104 g, 0.28 mmol) in dry, degassed CH₂Cl₂ (15 cm³) was added [9]aneS₃ (0.05 g, 0.28 mmol). The resulting purple solution was stirred at room temperature under N₂ for 2 h and Et₂O was added to afford a purple solid which was filtered off and dried *in vacuo*. Yield = 0.066 g, 70% (Found: C, 21.5; H, 3.7. C₆H₁₂Cl₃CrS₃ requires C, 21.3; H, 3.5%). IR (CsI disc): 2970w, 1445s, 1405s, 826m, 349 (br) m and 293w cm⁻¹.

(b) [CrBr₃([9]aneS₃)]. Method as for (a), using [CrBr₃(thf)₃] (0.101 g, 0.20 mmol) and [9]aneS₃ (0.036 g, 0.20 mmol), giving a blue-purple solid. Yield = 0.059 g, 62% (Found: C, 14.9; H, 2.7. C₆H₁₂Br₃CrS₃ requires C, 15.3; H, 2.6%). IR (Nujol mull): 346m and 324m cm⁻¹.

(c) [CrCl₃([10]aneS₃)]. Method as for (a), using [CrCl₃(thf)₃] (0.108 g, 0.29 mmol) and [10]aneS₃ (0.056 g, 0.29 mmol), giving a purple solid. Yield = 0.077 g, 75% (Found: C, 24.0; H, 4.1. C₇H₁₄Cl₃CrS₃ requires C, 23.9; H, 4.0%). IR (Nujol mull): 359m and 346m cm⁻¹.

(d) [CrBr₃([10]aneS₃)]. Method as for (a), using [CrBr₃(thf)₃] (0.091 g, 0.18 mmol) and [10]aneS₃ (0.035 g, 0.18 mmol), giving a dark purple solid. Yield = 0.037 g, 68% (Found: C, 17.1; H, 3.0. C₇H₁₄Br₃CrS₃ requires C, 17.3; H, 2.9%). IR (Nujol mull): 342w, 325m and 300m cm⁻¹.

(e) [(CrCl₃)₂(μ-[18]aneS₆)]. Method as for (a), using [CrCl₃(thf)₃] (0.066 g, 0.18 mmol) and [18]aneS₆ (0.03 g, 0.09 mmol), giving a pink-purple solid. Yield = 0.080 g, 66% (Found: C, 21.6; H, 3.5. C₁₂H₂₄Cl₆Cr₂S₆ requires C, 21.3; H, 3.5%). IR (Nujol mull): 340 (br) m and 298w cm⁻¹.

(f) [(CrBr₃)₂(μ-[18]aneS₆)]. Method as for (a), using [CrBr₃(thf)₃] (0.07 g, 0.14 mmol) and [18]aneS₆ (0.025 g, 0.07 mmol), giving a blue-purple solid. Yield = 0.07 g, 52% (Found: C, 15.1; H, 2.6. C₁₂H₂₄Br₆Cr₂S₆ requires C, 15.3; H, 2.5%). IR (Nujol mull): 310m and 301w cm⁻¹.

(g) [CrBr₂([14]aneS₄)]PF₆. To a solution of [CrBr₃(thf)₃] (0.12 g, 0.24 mmol) in dry, degassed MeNO₂ (15 cm³) was added TIPF₆ (0.085 g, 0.24 mmol) and [14]aneS₄ (0.065 g, 0.24 mmol). After stirring for ca. 12 h the white TiCl₄ precipitate was filtered off, leaving a blue-green solution. Addition of Et₂O gave a deep blue solid which was filtered off, washed with Et₂O and dried *in vacuo*. Yield = 0.06 g, 41% (Found: C, 18.8; H, 3.4. C₁₀H₂₀Br₂CrF₆PS₄ requires C, 19.2; H, 3.2%). Electrospray mass spectrum (MeCN solution): *m/z* = 480 {calc. for [Cr⁷⁹Br⁸¹Br([14]aneS₄)]⁺ *m/z* = 480}. IR (Nujol mull): 839vs, 557s, 346w and 323w cm⁻¹.

(h) [CrI₂([14]aneS₄)]BF₄. The salt AgBF₄ (0.156 g, 0.80 mmol) and [CrCl₃(thf)₃] (0.10 g, 0.27 mmol) were stirred in dry thf (25 cm³) for ca. 2 h to give a white AgCl precipitate which was filtered off to leave a pale green solution of [Cr(thf)₆][BF₄]₃. Dry MeNO₂ (25 cm³) was then added and the thf removed *in vacuo*. The compounds [14]aneS₄ (0.072 g, 0.27 mmol) and NEt₄I (0.139 g, 0.54 mmol) were added and the deep green reaction mixture stirred for 1 h. The volume was then reduced *in vacuo* to <3 cm³. Addition of Et₂O yielded an olive-green solid which was filtered off and dried. Yield = 0.04 g, 22% (Found: C, 21.1; H, 3.9. C₁₀H₂₀BCrF₁₂I₂S₄ requires C, 21.8; H, 3.6%). IR (Nujol mull): 1077s, 839m, 564m and 351w cm⁻¹.

(i) [CrCl₂([16]aneS₄)]PF₆. As for (g), using [CrCl₃(thf)₃] (0.06 g, 0.16 mmol), TIPF₆ (0.056 g, 0.16 mmol) and [16]aneS₄ (0.048 g, 0.016 mmol), giving a light green solid. Yield = 0.03 g, 30% (Found: C, 25.0; H, 4.6. C₁₂H₂₄Cl₂CrF₆PS₄ requires C, 25.5; H, 4.3%). Electrospray mass spectrum (MeCN solution): *m/z* = 418 {calc. for [Cr³⁵Cl₂([16]aneS₄)]⁺ *m/z* = 418}. IR (Nujol mull): 840vs, 558s and 366 (br) w cm⁻¹.

(j) [CrBr₂([16]aneS₄)]PF₆. As for (g), using [CrBr₃(thf)₃] (0.08 g, 0.16 mmol), TIPF₆ (0.056 g, 0.16 mmol) and [16]aneS₄ (0.048 g, 0.016 mmol), giving a dark green solid. Yield = 0.03 g, 29% (Found: C, 22.4; H, 3.9. C₁₂H₂₄Br₂CrF₆PS₄ requires C, 22.0; H, 3.7%). Electrospray mass spectrum (MeCN solution): *m/z* = 508 {calc. for [Cr⁷⁹Br⁸¹Br([16]aneS₄)]⁺ *m/z* = 508}. IR (Nujol mull): 841vs, 558s and 302w cm⁻¹.

(k) [CrI₂([16]aneS₄)]BF₄. Method as for (h), using [Cr(thf)₆][BF₄]₃ (0.25 mmol), [16]aneS₄ (0.08 g, 0.27 mmol) and NEt₄I (0.139 g, 0.54 mmol), giving a green solid. Yield = 0.03 g, 16% (Found: C, 21.5; H, 3.8. C₁₂H₂₄BCrF₁₂I₂S₄ requires C, 20.9; H, 3.5%). IR (Nujol mull): 1056s, 846m, 562m and 359w cm⁻¹.

EXAFS Refinements

Typically two or three data sets were collected for each complex and the analyses were carried out on the averaged spectra. The raw data were background-subtracted using the program PAXAS³⁰ by fitting a sixth- or eighth-order split polynomial to the pre-edge subtracted spectrum between *k* = 2 and to 13–15 Å⁻¹. Curve fitting was carried out using the program EXCURV 92.³¹ Ground-state potentials of the atoms were calculated using Von Barth theory and phase shifts using Hedin–Lundqvist potentials. Two shells (either 3 S and 3 X or 4 S and 2 X) were fitted in each case. Refinements were also carried out using 6 S or 6 X, as well as other combinations, and in the case of X = Br the results clearly supported the S₃Br₃ or S₄Br₂ donor sets expected. In the case of the chloro derivatives the very similar backscattering from S and Cl made the assignment of the donor set difficult on the basis of the EXAFS data alone, although a better fit was obtained using two shells (either 3 S and 3 Cl or 4 S and 2 Cl as appropriate) and, in addition, the UV/VIS, IR and mass spectrometric data provide very strong evidence for the donor sets chosen. The distances and Debye–Waller factors were refined for all the shells, as well as the Fermi energy difference. No attempt was made to refine the carbons of the ligand backbones since these occur over a range of distances and are not expected to be well defined.

Acknowledgements

We thank the EPSRC for an Earmarked Studentship (S. J. A. P.) and for funding the X-ray diffractometer, and the University of Southampton for support. We also thank the Director of the Synchrotron Radiation Source at Daresbury for the use of the facilities and we are indebted to Dr. W. Levason and Professor J. Evans (University of Southampton) for help in collecting the EXAFS data and the former for microanalytical measurements.

References

- 1 D. A. House and C. S. Garner, *Transition Met. Chem.*, 1970, **6**, 59; *Comprehensive Coordination Chemistry*, eds. G. Wilkinson, R. D. Gillard and J. A. McCleverty, Pergamon, Oxford 1987, vol. 3.
- 2 See, for example, A. L. Hale, W. Levason and F. P. McCullough, *Inorg. Chem.*, 1982, **21**, 3570; L. R. Gray, A. L. Hale, W. Levason, F. P. McCullough and M. Webster, *J. Chem. Soc., Dalton Trans.*, 1983, 2573.
- 3 L. R. Gray, A. L. Hale, W. Levason, F. P. McCullough and M. Webster, *J. Chem. Soc., Dalton Trans.*, 1984, 47.
- 4 A. L. Hale and W. Levason, *J. Chem. Soc., Dalton Trans.*, 1983, 2569.
- 5 M. A. Bennett, R. J. H. Clark and A. D. J. Goodwin, *J. Chem. Soc. A*, 1970, 541.
- 6 R. J. H. Clark and G. Natile, *Inorg. Chim. Acta*, 1970, **4**, 533.
- 7 H.-J. Kuppers and K. Wiegardt, *Polyhedron*, 1989, **8**, 1770.
- 8 G. J. Grant, K. E. Grant, W. N. Setzer and D. G. VanDerveer, *Inorg. Chim. Acta*, 1995, **234**, 35.
- 9 N. R. Champness, S. R. Jacob, G. Reid and C. S. Frampton, *Inorg. Chem.*, 1995, **34**, 396.
- 10 A. J. Blake and M. Schröder, *Adv. Inorg. Chem.*, 1990, **35**, 1; S. R. Cooper and S. C. Rawle, *Struct. Bonding (Berlin)*, 1990, **72**, 1.
- 11 See, for example, M. T. Ashby and D. L. Lichtenberger, *Inorg. Chem.*, 1985, **24**, 636; D. Sellman and L. Zapf, *Angew. Chem., Int. Ed. Engl.*, 1984, **23**, 807; D. Sevdic, M. Curic and Lj. Tusak-Bozic, *Polyhedron*, 1989, **8**, 505; T. Adachi, N. Sasaki, T. Ueda, M. Kaminaka and T. Yoshida, *J. Chem. Soc., Chem. Commun.*, 1989, 1320; T. Yoshida, T. Adachi, T. Ueda, M. Kaminaka, N. Sasaki, T. Higuchi, T. Aoshima, I. Mega, Y. Mizobe and M. Hidai, *Angew. Chem., Int. Ed. Engl.*, 1989, **28**, 1040.
- 12 D. Sellmann and L. Zapf, *J. Organomet. Chem.*, 1985, **289**, 57.

- 13 P. K. Baker, S. J. Coles, M. C. Durrant, S. D. Harris, D. L. Hughes, M. B. Hursthouse and R. L. Richards, *J. Chem. Soc., Dalton Trans.*, 1996, 4003.
- 14 G. R. Willey, M. T. Lakin and N. W. Alcock, *J. Chem. Soc., Chem. Commun.*, 1991, 1414.
- 15 M. C. Durrant, S. Davies, D. L. Hughes, C. Le Floch, R. L. Richards, J. R. Sanders, N. R. Champness, S. J. Pope and G. Reid, *Inorg. Chim. Acta*, 1996, **251**, 13.
- 16 P. J. Jones, A. L. Hale, W. Levason and F. P. McCullough, *Inorg. Chem.*, 1983, **22**, 2642.
- 17 J. I. Bruce, L. R. Gahan, T. W. Hambley and R. Stranger, *J. Chem. Soc., Chem. Commun.*, 1993, 702.
- 18 S. Pattanayak, D. Kumar Das, P. Chakravorty and A. Chakravorty, *Inorg. Chem.*, 1995, **34**, 6556.
- 19 A. J. Blake, G. Reid and M. Schröder, *J. Chem. Soc., Dalton Trans.*, 1989, 1675.
- 20 A. J. Blake, R. O. Gould, G. Reid and M. Schröder, *J. Organomet. Chem.*, 1988, **356**, 389.
- 21 T.-F. Lai and C.-K. Poon, *J. Chem. Soc., Dalton Trans.*, 1982, 1465.
- 22 SHELXS 86, program for crystal structure solution, G. M. Sheldrick, *Acta Crystallogr., Sect. A*, 1990, **46**, 467.
- 23 TEXSAN, Crystal Structure Analysis Package, Molecular Structure Corporation, Houston, TX, 1992.
- 24 N. Walker and D. Stuart, *Acta Crystallogr., Sect. A*, 1983, **39**, 158.
- 25 F. A. Cotton, S. A. Duraj, G. L. Powell and W. J. Roth, *Inorg. Chim. Acta*, 1986, **113**, 81.
- 26 A. P. B. Lever, *Inorganic Electronic Spectroscopy*, Elsevier, Amsterdam, 2nd edn., 1984.
- 27 L. S. Forster, *Transition Metal Chemistry*, Marcel Dekker, New York, 1969, vol. 5, p. 1.
- 28 N. Serpone, M. A. Jamieson, M. S. Henry, M. Z. Hoffman, F. Bolletta and M. Maestri, *J. Am. Chem. Soc.*, 1979, **101**, 2907.
- 29 W. Herzig and H. H. Zeiss, *J. Org. Chem.*, 1958, **23**, 1404.
- 30 N. Binsted, PAXAS, Program for the analysis of X-ray absorption spectra, University of Southampton, 1988.
- 31 N. Binsted, J. W. Campbell, S. J. Gurman and P. C. Stephenson, EXCURV 92, SERC Daresbury Laboratory, 1992.

Received 11th December 1996; Paper 6/08332C