

# Ultrafast reductive elimination of hydrogen from a metal carbonyl dihydride complex; a study by time-resolved IR and visible spectroscopy †


Mirco Colombo,<sup>a</sup> Michael W. George,<sup>b</sup> John N. Moore,<sup>a</sup> David I. Pattison,<sup>a</sup> Robin N. Perutz,<sup>\*,a</sup> Ian G. Virrels<sup>b</sup> and Tian-Qing Ye<sup>a</sup>

<sup>a</sup> Department of Chemistry, University of York, York, UK YO1 5DD

<sup>b</sup> Department of Chemistry, University of Nottingham, Nottingham, UK NG7 2RD

Laser flash photolysis of  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})(\text{H})_2]$  **1** in benzene solution yielded transient  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$  which was observed by both microsecond UV/VIS and IR spectroscopy [ $\lambda_{\text{max}} = 380$  nm,  $\nu(\text{CO}) = 1845 \text{ cm}^{-1}$ ] and reacted with  $\text{H}_2$  to reform **1** [ $k_2 = (8.4 \pm 0.4) \times 10^7 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ ]; photolysis of **1** with an ultrafast laser equipped with IR detection demonstrates that reductive elimination of  $\text{H}_2$  and formation of  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$  is complete within 6 ps.

Reductive elimination of hydrogen and the reverse reaction, oxidative addition, occupy fundamental positions in transition-metal chemistry. When initiated thermally, such reactions play a crucial role in catalysis. However, it is often the photo-induced reactions which offer a gateway to mechanistic studies, since numerous metal dihydride complexes undergo reductive elimination of hydrogen on ultraviolet irradiation.<sup>1–5</sup> When placed under a hydrogen atmosphere, the photoproduct may react back with hydrogen to regenerate the precursor [equation (1)].<sup>3–5</sup> We have used flash photolysis with UV/VIS detection to



observe the primary photoproducts,  $[\text{M}]$ , and to determine the kinetics of the back reaction with hydrogen, notably with  $[\text{Ru}(\text{drpe})_2]$  (drpe =  $\text{R}_2\text{PCH}_2\text{CH}_2\text{PR}_2$ , dmpe when  $\text{R} = \text{Me}$ , depe when  $\text{R} = \text{Et}$ , etc.).<sup>3–5</sup> Such measurements are made with typical 'nanosecond apparatus' with an instrumental risetime of ca. 50 ns. It is invariably found that formation of the 16-electron transient is complete within the risetime. The early steps of the photochemical reaction, including formation of excited state(s), M–H bond breaking, H–H bond making and rearrangement of the  $\text{MP}_4$  skeleton to its new equilibrium geometry must occur within this time.

Ultrafast spectroscopy (time-scale  $10^{-13}$ – $10^{-9}$  s) offers the opportunity of studying the early stages of reaction but is less straightforward and requires very different equipment. In our first investigation of reductive elimination on the picosecond time-scale, we irradiated  $[\text{Ru}(\text{dmpe})_2\text{H}_2]$  with an ultrafast laser and followed the transient UV/VIS absorption over time.<sup>6</sup> We obtained evidence that reductive elimination was already complete ca. 16 ps after the laser pulse, but these experiments were confined to a few probe wavelengths and were limited by the lack of conclusive evidence of oxidation states from UV/VIS spectroscopy.

In this communication, we report ultrafast experiments on a

related metal carbonyl dihydride complex in which we have followed the reaction from excitation to 2000 ps after the laser pulse by time-resolved infrared (TRIR) spectroscopy using the CO stretching vibration as a reporter.<sup>7</sup> These experiments form the first application of ultrafast IR spectroscopy to study reductive elimination, and provide conclusive evidence that reductive elimination is complete within 6 ps.

Our precursor,  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})(\text{H})_2]$  **1**, has been shown to lose  $\text{H}_2$  upon photolysis in steady-state experiments.<sup>8</sup> It is of particular interest because it catalyses insertion of alkenes into C–H bonds at unsaturated carbon (alkene or aromatic) in a  $\beta$ -position relative to a carbonyl group.<sup>9–11</sup> In one of the putative reaction mechanisms, the thermal reaction is initiated by hydrogenation of the alkene to form  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$ , the same intermediate as would be expected to form upon photolysis.<sup>9</sup>


The white precursor **1** is soluble in benzene and exhibits a shoulder at 325 nm ( $\epsilon \approx 9000 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$ ) in its UV/VIS absorption spectrum and a  $\nu(\text{CO})$  band at  $1939 \text{ cm}^{-1}$ . On laser flash photolysis† (instrumental risetime ca. 50 ns), a conspicuous transient is observed with  $\lambda_{\text{max}} \approx 380$  nm [Fig. 1(a)]. The transient decays with pseudo-first-order kinetics ( $k_{\text{obs}} = 3 \times 10^3 \text{ s}^{-1}$ ) under an argon atmosphere. The rate constant increases linearly with concentration of dissolved hydrogen [Fig. 1(b)]. The slope of this plot yields the second-order rate constant for reaction of the transient with  $\text{H}_2$  in benzene of  $(8.4 \pm 0.4) \times 10^7 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ . Transient absorption experiments with **1** dissolved under 1 atm  $\text{H}_2$  in tetrahydrofuran (thf) yield a comparable value of  $k_{\text{obs}}$  ( $2.1 \times 10^5 \text{ s}^{-1}$ ) to the experiments with benzene as solvent ( $2.5 \times 10^5 \text{ s}^{-1}$ ).‡

Similar experiments were performed with **1** dissolved in  $\text{C}_6\text{D}_6$  under  $\text{H}_2$ , but with IR detection.§ A spectrum measured 1  $\mu\text{s}$  after the laser flash revealed loss of precursor at  $1940 \text{ cm}^{-1}$  and

† [1] typically ca.  $10^{-4} \text{ mol dm}^{-3}$  in  $\text{C}_6\text{H}_6$ , 10 mm pathlength, laser at 308 nm, pulse energy at sample ca. 5 mJ. Samples were degassed by freeze–pump–thaw methods and  $\text{H}_2$ –Ar gas mixtures admitted with a total pressure of 1 atm. One measurement was made at higher pressure of  $\text{H}_2$ , viz. 3.2 atm. No significant change in the rate constant under 1 atm  $\text{H}_2$  was found when  $\text{C}_6\text{H}_6$  solvent was replaced by  $\text{C}_6\text{D}_6$ , nor when the concentration of **1** was increased to  $5 \times 10^{-4}$  and  $1 \times 10^{-3} \text{ mol dm}^{-3}$  (2 and 1 mm pathlengths, respectively). The concentration of dissolved hydrogen was taken as  $2.85 \times 10^{-3} \text{ mol dm}^{-3} \text{ atm}^{-1}$ .<sup>12a</sup> The solubility of  $\text{H}_2$  in thf has not been documented, but is ca.  $6 \times 10^{-3} \text{ mol dm}^{-3} \text{ atm}^{-1}$  in  $\text{Et}_2\text{O}$ .<sup>12b</sup> The value derived for the rate constant for reaction of the transient with  $\text{H}_2$  in thf is ca.  $3.6 \times 10^7 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ .

§ Pump laser at 355 or 308 nm, detection with diode IR laser, sample  $2 \times 10^{-3}$  or  $3 \times 10^{-3} \text{ mol dm}^{-3}$  dissolved in  $\text{C}_6\text{D}_6$  (for IR transparency) under 1 atm hydrogen and flowed through an IR cell ( $\text{CaF}_2$  windows, 1 mm pathlength) within a closed circulating system (glass with PTFE connections and stainless steel micropump). Notice that the kinetics were sometimes monitored away from the band maxima, in order to use the optimum laser output. The instrument was set up to maximise the signal-to-noise ratio, such that the risetime was ca. 500 ns.

† Non-SI unit employed: atm = 101 325 Pa.




**Fig. 1** (a) Transient absorption, monitored at 410 nm, following laser flash photolysis ( $\lambda_{\text{ex}} = 308$  nm) of **1** dissolved in benzene under 1 atm  $\text{H}_2$ ; (●) experimental points, (—) exponential fit. (b) Plot of the variation of  $k_{\text{obs}}$  vs. concentration of dissolved hydrogen. The slope yields the second-order rate constant for reaction of  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$  with hydrogen to reform **1**

formation of product bands with maxima at 1845 and 1974  $\text{cm}^{-1}$  [Fig. 2(a)]. The kinetics of the first product band were monitored at 1840  $\text{cm}^{-1}$ ; the band rose within the risetime of the instrument and decayed to about 30% of its initial absorbance with first-order kinetics with a rate constant,  $k_{\text{obs}}$ , of  $2.0 \times 10^5 \text{ s}^{-1}$  [Fig. 2(b)]. The same value of  $k_{\text{obs}}$  was obtained at two concentrations of **1** and is consistent with the kinetics from the UV experiments. The second transient band at 1973  $\text{cm}^{-1}$  showed a significant risetime of ca. 5  $\mu\text{s}$  with rate constants for growth dependent on **1** and did not show any subsequent decay over 50  $\mu\text{s}$ . The bleach signal at 1940  $\text{cm}^{-1}$  recovered partially (to ca. 80% of initial change in absorbance).


The transient species with  $\lambda_{\text{max}}$  at 380 nm and  $\nu(\text{CO})$  at 1845  $\text{cm}^{-1}$  is assigned unequivocally as  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$  on the basis of its rapid rise, the kinetics of its reaction with  $\text{H}_2$ , the partial recovery of starting material and the position of the IR band (Scheme 1). The very large shift of ca. 100  $\text{cm}^{-1}$  in  $\nu(\text{CO})$  from  $\text{Ru}^{\text{II}}$  in **1** to  $\text{Ru}^0$  in  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$  is consistent with shifts recently determined for related ruthenium complexes in matrices.<sup>13,14</sup> The lack of kinetic stabilisation by thf provides no support for solvent co-ordination to  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$ . The co-ordination geometry at ruthenium in  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$  cannot be determined from these experiments. In contrast, the relative intensity of the  $\nu(\text{CO})$  bands provides bond angle information for  $[\text{Ru}(\text{CO})_2(\text{dmpe})]$ .<sup>14</sup> The long-lived species with  $\nu(\text{CO})$  at 1973  $\text{cm}^{-1}$  must be a secondary dinuclear product since it rose relatively slowly with kinetics dependent on **1**.

The photolysis of **1** has also been monitored by TRIR spec-

† The observed rate constants for the growth of the 1973  $\text{cm}^{-1}$  band were  $6.8 \times 10^5 \text{ s}^{-1}$  and  $9.3 \times 10^5 \text{ s}^{-1}$  at  $[\text{1}] = 2 \times 10^{-3}$  and  $3 \times 10^{-3} \text{ mol dm}^{-3}$ , respectively. The kinetics of the bleach at 1940  $\text{cm}^{-1}$  could be fitted satisfactorily to a model comprising a finite rise and a competing decay component corresponding to the rate constants of the two product bands.




**Fig. 2** Microsecond TRIR spectrum recorded 1  $\mu\text{s}$  after laser flash photolysis ( $\lambda_{\text{ex}} = 355$  nm) of **1** dissolved in  $\text{C}_6\text{D}_6$  under 1 atm  $\text{H}_2$ . The negative feature is due to bleach of **1**, the positive features are due to photoproducts. (b) Microsecond kinetics of the lower wavenumber product band monitored at 1840  $\text{cm}^{-1}$  ( $\lambda_{\text{ex}} = 308$  nm)



**Scheme 1** Photochemical reactions of complex **1**

troscopy on the picosecond time-scale. An IR spectrum measured 25 ps after the laser pulse revealed two features: a negative peak at  $1940 \pm 3 \text{ cm}^{-1}$  corresponding to the loss of the precursor and a positive peak at  $1843 \pm 3 \text{ cm}^{-1}$  corresponding to formation of  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$ . No other bands were observed between 1800 and 2050  $\text{cm}^{-1}$ . A kinetic trace measured at 1843  $\text{cm}^{-1}$  showed a very fast rise and decay within the instrumental response function of ca. 6 ps. After this initial period, the signal maintained its amplitude over 2 ns [Fig. 3(a)]. The very fast decay corresponds to the solvent response, since a positive signal with these kinetics was observed at all wavenumbers for the solvent alone. The long-lived signal arises from  $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$ ; curve fitting shows that this signal appeared within ca. 6 ps and remained constant [Fig. 3(a)]. A kinetic trace measured at 1941  $\text{cm}^{-1}$  showed that the bleaching signal of **1** also rose within 6 ps and remained steady for the succeeding 2 ns [Fig. 3(b)]; again, a signal was observed which decayed with the instrument response function.

†† Ultrafast setup;<sup>7</sup> pump pulses at 304 nm of 4 ps full width at half maximum (6 ps autocorrelation time) duration and 1.05 kHz repetition rate, pulse energy ca. 1–2  $\mu\text{J}$ . IR absorption was measured with a CO laser (1800–1960  $\text{cm}^{-1}$  at 4  $\text{cm}^{-1}$  intervals) or with a diode laser (1965–2050  $\text{cm}^{-1}$ ) via a detection system employing upconversion. The solutions were handled in the same flow system as for the microsecond experiments, but with a faster flow rate. Complex **1** (ca.  $10^{-3} \text{ mol dm}^{-3}$ ) dissolved in  $\text{C}_6\text{D}_6$  under 1 atm  $\text{H}_2$ . IR cell of pathlength 1 mm giving an absorbance ca. 1 at the pump wavelength and ca. 0.3 at 1940  $\text{cm}^{-1}$ .



**Fig. 3** Ultrafast TRIR kinetics following laser photolysis of **1** dissolved in  $C_6D_6$  under 1 atm  $H_2$  ( $\lambda_{ex} = 304$  nm, laser pulse duration = 4 ps) (a) at  $1843\text{ cm}^{-1}$  showing formation of  $[Ru(PPh_3)_3(CO)]$  (●), (b) at  $1941\text{ cm}^{-1}$  showing bleaching of **1** (■). The very fast component in both traces lasting *ca.* 6 ps lies within the instrumental response function

Ultrafast experiments on binary metal carbonyls usually show the absorptions of metal carbonyls in *vibrational* excited states of the electronic ground state which decay over tens or hundreds of picoseconds.<sup>15</sup> The vibrational excitation is detected through broadening and shifts in the  $\nu(CO)$  bands. However,  $[Ru(PPh_3)_3(CO)]$  shows no signs of vibrational relaxation although the photon energy absorbed by **1** greatly exceeds the energy needed to break the bonds, just as when photodissociating CO from metal carbonyls. When expelling CO, the CO bond length is altered little, so the CO carries away relatively little vibrational energy leaving much of the excess of energy in the photoproduct. When  $H_2$  is dissociated from a metal dihydride, the  $H\cdots H$  distance is necessarily compressed with the result that much of the excess of energy may be removed by vibrational excitation of expelled  $H_2$  (*cf.*  $CH_2O$ ).<sup>16</sup> Almost all of the excess of energy would be removed if  $H_2$  leaves in its  $v = 7$  state.<sup>\*\*</sup> In reality, some of the excess of energy will go into rotation of  $H_2$  and translation of the photofragments, as well as vibrations of  $[Ru(PPh_3)_3(CO)]$ . However, the lack of observable vibrational relaxation of  $[Ru(PPh_3)_3(CO)]$  is understandable.

The ultrafast experiments provide unambiguous evidence for removal of **1** and formation of  $[Ru(PPh_3)_3(CO)]$  within the experimental time-scale. The use of IR detection has removed any possibility that the signals are due to excited states of precursor or photoproducts. We conclude that photoelimination of  $H_2$  from **1** is a process in which H–H bond formation, Ru–H

bond cleavage and any reorganisation of the co-ordination geometry at ruthenium occur within 6 ps.

## Acknowledgements

We are grateful to EPSRC for support for equipment, studentships and an Advanced Fellowship (to J. N. M.). We also acknowledge the support of the European Commission and British Gas.

## References

- 1 M. Berry, K. Elmitt, M. L. H. Green and S. J. Simpson, *J. Chem. Soc., Dalton Trans.*, 1979, 1950; P. Grebenik, R. Grinter and R. N. Perutz, *Chem. Soc. Rev.*, 1988, **17**, 453.
- 2 W. D. Jones and F. J. Feher, *Acc. Chem. Res.*, 1989, **22**, 91; R. M. Chin, L. Dong, S. B. Duckett, M. G. Partridge, W. D. Jones and R. N. Perutz, *ibid.*, 1993, **115**, 7685; A. H. Janowicz and R. G. Bergman, *J. Am. Chem. Soc.*, 1983, **105**, 3929; J. M. Buchanan, J. M. Stryker and R. G. Bergman, *ibid.*, 1986, **108**, 1537; B. M. Sponsler, B. H. Weiller, P. O. Stoutland and R. G. Bergman, *ibid.*, 1989, **111**, 6841; P. E. Bloyce, A. J. Rest and I. Whitwell, *J. Chem. Soc., Dalton Trans.*, 1990, 813; M. G. Partridge, A. McCamley and R. N. Perutz, *ibid.*, 1994, 3519; W. A. Kiel, R. G. Ball and W. A. G. Graham, *J. Organomet. Chem.*, 1990, **383**, 481; S. A. Brough, C. Hall, A. McCamley, R. N. Perutz, S. Stahl, U. Wecker and H. Werner, *ibid.*, 1995, **504**, 33; H. Azizian and R. H. Morris, *Inorg. Chem.*, 1983, **22**, 6.
- 3 R. N. Perutz, *Chem. Soc. Rev.*, 1993, 361; R. Osman, D. I. Pattison, R. N. Perutz, C. Bianchini and M. Peruzzini, *J. Chem. Soc., Chem. Commun.*, 1994, 513.
- 4 L. Cronin, M. C. Nicasio, R. N. Perutz, R. G. Peters, D. M. Roddick and M. K. Whittlesey, *J. Am. Chem. Soc.*, 1995, **117**, 10 047.
- 5 C. Hall, W. D. Jones, R. J. Mawby, R. Osman, R. N. Perutz and M. K. Whittlesey, *J. Am. Chem. Soc.*, 1992, **114**, 7425.
- 6 R. Osman, R. N. Perutz, A. D. Rooney and A. J. Langley, *J. Phys. Chem.*, 1994, **98**, 3562.
- 7 T. Q. Ye, C. J. Arnold, D. I. Pattison, C. L. Anderton, D. Dukic, R. N. Perutz, R. E. Hester and J. N. Moore, *Appl. Spectrosc.*, 1996, **50**, 597.
- 8 G. L. Geoffroy and M. G. Bradley, *Inorg. Chem.*, 1977, **16**, 744.
- 9 F. Kakiuchi, S. Sekine, Y. Tanaka, A. Kamatani, M. Sonoda, N. Chatani and S. Murai, *Bull. Chem. Soc. Jpn.*, 1995, **68**, 62 and refs. therein.
- 10 M. Sonoda, F. Kakiuchi, A. Kamatani and S. Murai, *Chem. Lett.*, 1996, 109.
- 11 B. M. Trost, K. Imi and I. W. Davies, *J. Am. Chem. Soc.*, 1995, **117**, 5371; F. Kakiuchi, Y. Tanaka, T. Sato, N. Chatani and S. Murai, *Chem. Lett.*, 1995, 679.
- 12 (a) E. Wilhelm and R. Battino, *Chem. Rev.*, 1973, **73**, 1; (b) P. G. T. Fogg and W. Gerrard, *Solubility of Gases in Liquids*, Wiley, Chichester, 1991.
- 13 R. J. Mawby, R. N. Perutz and M. K. Whittlesey, *Organometallics*, 1995, **14**, 3268.
- 14 M. K. Whittlesey, R. N. Perutz, I. G. Virrels and M. W. George, *Organometallics*, 1997, **16**, 268.
- 15 M. Lee and C. B. Harris, *J. Am. Chem. Soc.*, 1989, **111**, 8963; J. C. King, J. Z. Zhang, B. J. Schwartz and C. B. Harris, *J. Chem. Phys.*, 1993, **99**, 7595; T. P. Dougherty and E. J. Heilweil, *Chem. Phys. Lett.*, 1994, **227**, 19; S. M. Arrivo, T. P. Dougherty, W. T. Grubbs and E. J. Heilweil, *Chem. Phys. Lett.*, 1995, **235**, 247.
- 16 C. B. Moore, *Ann. Rev. Phys. Chem.*, 1983, **34**, 525.
- 17 S. T. Belt, J. C. Scaiano and M. K. Whittlesey, *J. Am. Chem. Soc.*, 1993, **115**, 1921.
- 18 G. Herzberg and L. L. Howe, *Can. J. Phys.*, 1959, **37**, 636.

Received 25th June 1997; Communication 7/04484D

<sup>\*\*</sup> At 304 nm,  $h\nu = 393\text{ kJ mol}^{-1}$ . Enthalpy<sup>17</sup> of  $[Ru(dmpe)_2H_2] \longrightarrow [Ru(dmpe)_2] + H_2 = 95\text{ kJ mol}^{-1}$ . If enthalpy change for **1**  $\longrightarrow [Ru(PPh_3)_3(CO)] + H_2$  is the same, excess of energy =  $298\text{ kJ mol}^{-1}$ . Energy of  $H_2$  ( $v = 7$ )  $\approx 290\text{ kJ mol}^{-1}$ .<sup>18</sup>