

Bis(diphosphino)methanide and hydride derivatives of Ir₄(CO)₁₂

Serena Detti,^a Tito Lumini,^a Raymond Roulet,^{*a} Kurt Schenk,^b Renzo Ros^{*c} and Augusto Tassan^c

^a Institut de Chimie Minérale et Analytique de l'Université, BCH, CH-1015 Lausanne, Switzerland

^b Institut de Crystallographie de l'Université, BSP, CH-1015 Lausanne, Switzerland

^c Dipartimento di Processi Chimici dell'Ingegneria e Centro di Chimica Metallorganica del CNR, Via Marzolo 9, I-35131 Padova, Italy

Received 16th December 1999, Accepted 31st March 2000

Published on the Web 27th April 2000

The reaction of [Ir₄(CO)₁₀(μ-(Ph₂P)₂CH₂)] with dried KOH forms [Ir₄(CO)₁₀(μ-(Ph₂P)₂CH)][−] which subsequently converts into [Ir₄(CO)₉(μ₃-(Ph₂P)₂CH)][−], a rare example of an Ir cluster with an Ir–C bond (average 2.274(8) Å). The same reaction with the weaker base 1,8-diazobicyclo[5.4.0]undec-7-ene or with K₂CO₃ in wet CH₂Cl₂ under CO affords the anion [HIr₄(CO)₉(μ-(Ph₂P)₂CH₂)][−], a hydride-cluster with an unusually long Ir–H distance (2.08(6) Å at 90 K). Intramolecular CO scrambling is observed in CD₂Cl₂ solution above 200 K, preserving the terminal–axial position of the hydride ligand.


Introduction

The hydrido-derivatives of Ir₄(CO)₁₂ reported to date are the anions [HIr₄(CO)₁₁][−],¹ and [H₂Ir₄(CO)₁₀]^{2−},² as well as the neutral orthometalated species [HIr₄(CO)₇(Ph₂PCH=CHPPh₂)-(PhC₆H₄PCH=CHPPh₂)].³ The examples of clusters containing Ir–C(alkyl) bonds are [Ir₄(CO)₁₁(CH₂CO₂CH₃)][−] and [Ir₄(CO)₁₀(CH₂CO₂CH₃)₂]^{2−}.⁴ Clusters containing Ir=C bonds are the carbene complexes [Ir₄(CO)_{12−n}(C(=O)CH₂CH₂O)_n] (n = 1–3) derived from the nucleophilic attack of the coordinated CO ligands of Ir₄(CO)₁₂ by oxirane.⁵ The classical organometallic reactions used for the preparation of metal alkyls failed to give any well defined derivatives from Ir₄(CO)₁₂. An idea for obtaining new clusters with Ir–C bonds is therefore to deprotonate a methylene group which is part of a bidentate ligand already bonded to Ir and form an Ir–C bond by intramolecular displacement of carbon monoxide.


The bidentate ligand bis(diphenylphosphino)methane (dppm) has widely been used in coordination and organometallic chemistry.⁶ Its deprotonated form, bis(diphenylphosphino)methanide, [(Ph₂P)₂CH][−], has also received some attention because of its ability to act as a two-, four- or six-electron donor ligand, being a versatile building block for the construction of heterometallic species.⁷ We report here a simple way to obtain a new class of methanide complexes derived from deprotonation of the dppm ligand of [Ir₄(CO)₁₀(μ-dppm)]⁸ or alternatively, depending on the choice of the base, a way to obtain new hydrido derivatives of Ir₄ clusters.

Results and discussion

The reaction of [Ir₄(CO)₁₀(μ-dppm)] (1) with two equivalents of bis(triphenylphosphoranylidene)ammonium chloride (PPNCl) and an excess of KOH (dry powder) in CH₂Cl₂ at −20 °C gives rapidly a first product which could only be observed in solution, and was identified as [Ir₄(CO)₁₀(μ-(Ph₂P)₂CH)][−] on the basis of its NMR characteristics (see Experimental section). After ca. 30 minutes, intramolecular coordination of the methanide moiety is completed, affording the yellow product [PPN][Ir₄(CO)₉(μ₃-(Ph₂P)₂CH)] (2) (yield 76%) by the overall equation:



The same reaction, but with a large excess of the weaker base 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (or more slowly with K₂CO₃) in wet CH₂Cl₂ under CO gave orange crystals of the hydrido-cluster [PPN][HIr₄(CO)₉(μ-dppm)] (3) (yield 86%) by the overall reaction:



The crystal structures of 2 and 3 were determined by X-ray analysis (see below).

Crystal structures of [PPN][Ir₄(CO)₉(μ₃-(Ph₂P)₂CH)]·CH₂Cl₂ and [PPN][HIr₄(CO)₉(μ-dppm)]

The structure of the anionic part of 2 and the labelling scheme are shown in Fig. 1. Some relevant interatomic distances are reported in Table 1. Crystallographic data for 2·CH₂Cl₂ and 3 are reported in Table 2.

The anion of 2 has approximate C_s symmetry with all terminal CO's. The [(Ph₂P)₂CH][−] ligand is face bridging with the plane P(3)–C(32)–P(2) eclipsed relative to the plane Ir(4)–Ir(2)–Ir(3). The mean value of the Ir–Ir distances (2.700(1) Å) is signif-





Fig. 1 ORTEP¹⁵ representation at the 30% probability level of [Ir₄(CO)₉(μ₃-(Ph₂P)₂CH)][−].

Table 1 Bond lengths [Å] for **2** and **3**

	2	3
Ir(1)–C(1)	1.864(12)	1.889(6)
Ir(1)–C(1A)	1.879(14)	1.888(7)
Ir(1)–C(1B)	1.917(13)	1.889(7)
Ir(1)–Ir(2)	2.7066(5)	2.7448(3)
Ir(1)–Ir(3)	2.7080(5)	2.7598(4)
Ir(1)–Ir(4)	2.7179(6)	2.7860(4)
Ir(2)–C(2C)	1.892(13)	
Ir(2)–C(2)	1.899(12)	1.856(7)
Ir(2)–C(23)		2.053(6)
Ir(2)–C(24)		2.176(7)
Ir(2)–P(2)	2.287(2)	2.275(2)
Ir(2)–Ir(4)	2.7203(6)	2.7426(4)
Ir(2)–Ir(3)	2.7359(5)	2.6847(3)
Ir(3)–C(3)	1.872(12)	1.859(7)
Ir(3)–C(3C)	1.873(12)	
Ir(3)–C(23)		2.103(6)
Ir(3)–C(34)		2.151(7)
Ir(3)–P(3)	2.299(2)	2.291(2)
Ir(3)–Ir(4)	2.6620(6)	2.7276(4)
Ir(4)–H(4)		2.08(6)
Ir(4)–C(4)		1.855(7)
Ir(4)–C(4B)	1.869(12)	
Ir(4)–C(4A)	1.871(12)	
Ir(4)–C(24)		2.046(7)
Ir(4)–C(34)		2.064(6)
Ir(4)–C(32)	2.274(8)	
Ir(4)–P(3)	2.931(2)	
Ir(4)–P(2)	2.956(2)	
O(1)–C(1)	1.164(12)	1.168(8)
O(1A)–C(1A)	1.178(13)	1.152(8)
O(1B)–C(1B)	1.141(12)	1.131(8)
O(2)–C(2)	1.137(12)	1.170(8)
O(2C)–C(2C)	1.147(12)	
O(23)–C(23)		1.173(7)
O(24)–C(24)		1.167(8)
O(3)–C(3)	1.157(12)	1.160(8)
O(3C)–C(3C)	1.168(12)	
O(34)–C(34)		1.172(8)
O(4)–C(4)		1.153(9)
O(4A)–C(4A)	1.150(12)	
O(4B)–C(4B)	1.142(11)	
P(2)–C(32)	1.786(9)	1.843(6)
C(32)–H(32A)		0.97
C(32)–H(32B)		0.97
P(2)–C(200)	1.818(10)	1.829(6)
P(2)–C(210)	1.840(10)	1.817(6)
P(3)–C(32)	1.792(9)	1.829(6)
P(3)–C(310)	1.822(9)	1.836(6)
P(3)–C(300)	1.841(9)	1.841(6)
C(32)–H(32)	1.01	

Table 2 Crystallographic data for compounds **2**·CH₂Cl₂ and **3**

	2·CH ₂ Cl ₂	3
Formula	C ₇₁ H ₅₃ Cl ₂ Ir ₄ NO ₉ P ₄	C ₇₀ H ₅₃ Ir ₄ NO ₉ P ₄
<i>M</i>	2027.72	1944.81
Crystal size/mm		0.5 × 0.4 × 0.3
<i>T</i> /K	293(2)	90(3)
Crystal system	Monoclinic	Monoclinic
Space group	<i>P</i> 2 ₁ / <i>n</i>	<i>P</i> 2 ₁ / <i>c</i>
<i>a</i> /Å	12.6694(8)	10.2149(8)
<i>b</i> /Å	26.479(2)	18.4724(13)
<i>c</i> /Å	20.7990(13)	33.772(3)
β /°	94.0210(10)	94.578(9)
<i>V</i> /Å ³	6960.4(8)	6352.3(8)
<i>Z</i>	4	4
<i>D</i> _c /g cm ^{−3}	1.935	2.034
<i>F</i> (000)	3840	3680
μ /mm ^{−1}	7.848	8.259
2 θ _{max} /°	51.0	51.88
Reflections collected/unique	28587/11402	49085/12165
Variables	821	797
<i>R</i> ₁	0.0554	0.0412
<i>wR</i> ₂	0.823	0.838

**Fig. 2** ORTEP representation at the 50% probability level of [Ir₄(CO)₉(μ-dppm)][−].

icantly shorter than that found for [Ir₄(CO)₇(μ-CO)₃-(μ-(Ph₂P)₂CHMe)] (**1'**) (2.729(1) Å).⁹ A DFT calculation on Rh₄(CO)₁₂ and IrRh₃(CO)₁₂ has shown that unbridged metal-metal bonds are shorter than CO-bridged ones.¹⁰ This conclusion seems to be also valid for the present, substituted clusters. The metal tetrahedron is not regular, as the three Ir–Ir distances of the Ir(2)–Ir(3)–Ir(4) face bearing the [(Ph₂P)₂CH][−] ligand differ by more than 0.06 Å, the longest being that between the Ir-atoms bearing the P-atoms and the shortest being the other two bonds of that face. All the bond distances (and bond angles) of the Ir₂P₂C ring are smaller than the corresponding ones in **1'**, probably denoting the better donor properties of the deprotonated ligand relative to the neutral dppm ligand. As expected, the average Ir–C distance (2.274(8) Å) is longer than the Ir=C(carbene) distances of 1.94–2.06 Å in [Ir₄(CO)₁₂−*n*](CO-CH₂CH₂O)_{*n*}] (*n* = 1–3),⁵ but also longer than the Ir–C distances found in [Ir₄(CO)₁₁(CH₂CO₂Me)][−] and [Ir₄(CO)₁₀(CH₂CO₂Me)₂]^{2−} (2.15–2.19 Å).⁴

The anionic structure of **3** and the labelling scheme are shown in Fig. 2. Some relevant interatomic distances are reported in Table 1. The anion has approximate C_s symmetry with the mirror plane defined by atoms Ir(1) and Ir(4), and the methylene group. Three CO's are edge-bridging, defining the basal face Ir(2)–Ir(3)–Ir(4). The P-atoms of the dppm ligand are in axial positions relative to the basal face. The comparison between corresponding bond distances of **3** and **2** gives the same trends as that described above between **1'** and **2**. Due to the presence of 3 edge-bridging CO's, the mean value of the Ir–Ir distance (2.769(3) Å) is longer than that of **2**. Likewise, the interatomic distances in the Ir₂P₂C ring (mean Ir–P 2.283, mean P–C 1.836 Å) in **3** are longer than those in **2** (2.293 and 1.789 Å, respectively). The value for the Ir–H distance (2.08(6) Å at 90 K) is unusually large, both relative to the range found in monometallic complexes (1.5–1.9 Å)¹¹ and to the calculated range (1.54–1.60 Å).¹² The temperature was low enough to locate this H-atom during the refinement down to 4.16%. However, a neutron diffraction experiment will be undertaken to verify this value.

Behaviour of clusters **2** and **3** in solution

Complex **2** is thermally stable in CH₂Cl₂ solution up to room temperature. It reverts back to **1** upon adding 1 equivalent of HCl under CO. It converts to **3** upon adding excess DBU and 1 equivalent of H₂O under CO, or to the deuterido-analog of **3** when replacing H₂O by D₂O.

Its geometry in solution is similar to that found in the solid, as deduced from IR, ¹H-, ³¹P- and ¹³C-NMR spectroscopy (see Experimental section). A 2D-COSY spectrum of an enriched




Fig. 3 2D-COSY spectrum of **2** in CD_2Cl_2 at 193 K (see Fig. 4 for the carbonyl labelling scheme).

sample of **2** in ^{13}CO (ca. 30%) at 193 K (Fig. 3) shows the pseudo-*trans* coupling between carbonyls d and f, the signal at 177.2 ppm being due to carbonyls d closest to the CH^- group, and shows also that the apical CO's e and g (respectively at 163.4 and 160.4 ppm) are already in the slow exchange domain.

The only exchange mechanism in CD_2Cl_2 for compound **2**, noticeable above 203 K, is the rotation of the apical carbonyls, as shown by the variable temperature ^{13}C -NMR spectra (Fig. 4, left) which were simulated using a two-site exchange matrix (Fig. 4, right). Regression of the Eyring plot $\ln(k/T)$ vs. $1/T$ gave a value of $43.1 \pm 0.6 \text{ kJ mol}^{-1}$ for the free enthalpy of activation at 298 K for the rotation of the apical carbonyls about the local C_3 axis.

The geometry of **3** in solution is similar to that found in the solid, as deduced from IR, ^1H -, ^{31}P - and ^{13}C -NMR spectroscopy (see Experimental section). The ^1H -NMR spectrum at 210 K shows the CH_2 group of the dppm ligand as an ABX_2 spin system due to the inequivalence of the two methylene protons: whereas the H_B proton shows the expected chemical shift at 2.69 ppm, the H_A proton is abnormally deshielded at 6.03 ppm. This pattern is quite similar to that found for **1**;⁸ therefore, the conformation of the dimetallacyclopentane ring must be that found in solid **1**⁹ (where a methyl group replaces the H_B proton of dppm). The hydride ligand is coordinated in an axial position, replacing the unique axial CO in **1** (−15.7 ppm). This is deduced from the presence of three resonances in the ^{13}C -NMR limiting spectrum of a ^{13}CO enriched sample of **3** (ca. 30% enrichment) in the region of radial CO's (the two P-atoms and the H-atom must therefore occupy axial positions). Cluster **3** is fluxional in solution, as shown by the variable temperature ^{13}C -NMR spectra (Fig. 5, left). The first




Fig. 4 Experimental (left) and calculated (right) VT ^{13}C -NMR spectra of **2** in CD_2Cl_2 .

process of CO scrambling, starting at ca. 200 K, is the merry-go-round of the basal carbonyls a, b, d and f.

The second process is the rotation of apical carbonyls, which could not be simulated due to the similar δ values of carbonyls g and e. Simulation of the merry-go-round was effected with a four sites exchange matrix (Fig. 5, right). Regression of the function $\ln(k/T)$ vs. $1/T$ gave a value of $39.4 \pm 1.4 \text{ kJ mol}^{-1}$ for the free enthalpy of activation at 298 K, which is quite similar to that found for anionic Ir_4 clusters such as $[\text{Ir}_4(\text{CO})_{11}\text{X}]^-$ ($\text{X} = \text{Br}, \text{I}, \text{NCS}, \text{NO}_2$).¹³

Experimental

Unless otherwise stated, all the operations were carried out under nitrogen, all solvents were dried by conventional methods prior to use. PPNCl (Fluka) was used as received. KOH (Fluka) was dried at 150°C under reduced pressure (10^{-4} mm Hg). DBU (Aldrich) was dried with KOH and distilled. $[\text{Ir}_4(\text{CO})_{10}(\mu\text{-dppm})]$ was prepared as described in the literature,⁸ as well as ^{13}CO enriched derivatives of **2** and **3** (ca. 30% enrichment).¹⁴ Infrared spectra were recorded from 0.1 mm CaF_2 cells previously purged with N_2 on a Perkin-Elmer FT-IR 2000 spectrometer. Elemental analyses were performed at the Analytische Laboratorien, Lindlar, Germany. ^1H -, ^{31}P - and ^{13}C -NMR spectra were recorded on a Bruker AC200 spectrometer, using the residual proton peaks and carbon resonances of deuterated solvents as chemical shift references for ^1H - and ^{13}C -NMR spectra. ^{31}P -NMR spectra were referenced to external 85% H_3PO_4 . ^{13}C -NMR and 2D- ^{13}C -NMR spectra were recorded in 10 mm o.d. tubes on a Bruker WH360 spectrometer. COSY experiment: 256 t_1 increments with 2K transients, spectral




Fig. 5 Experimental (left) and calculated (right) VT ^{13}C -NMR spectra of **3** in CD_2Cl_2 .

width 2840.9 Hz in the F_2 domain and 1420.5 Hz in the F_1 domain.

Bis(triphenylphosphoranylidene)ammonium tetrahedrononacarbonyl $\{\mu_3$ -bis(diphenylphosphino)methanido}tetrairidate (2**) ([PPN][$\text{Ir}_4(\text{CO})_9(\mu_3\text{-(Ph}_2\text{P)}_2\text{CH})$])**

Dry KOH powder (0.20 g) and PPNCl (0.37 g, 0.64 mmol) were added to a solution of **1** (0.40 g, 0.28 mmol) in CH_2Cl_2 (15 cm^3) under N_2 at -20°C . After a few minutes the deprotonated intermediate $[\text{Ir}_4(\text{CO})_{10}(\mu\text{-(Ph}_2\text{P)}_2\text{CH})]^-$ was formed but could not be isolated. δ_{C} (CD_2Cl_2 , 210 K): 231.2 (1 CO, s), 214.7 (2 CO, m), 185.2 (2 CO, radial, s), 174.3 (1 CO, radial, s), 166.4 (1 CO, s), 159.8 (1 CO, s), 159.0 (2 CO, m). δ_{P} (CD_2Cl_2 , 210 K): -59.2 (PPh_2). The suspension was stirred for 35 minutes, filtered, and cold isopropanol (80 cm^3) added. The yellow precipitate was recrystallised at -30°C from CH_2Cl_2 /isopropanol, and the crystals of **2** washed with cold isopropanol and pentane (0.41 g, 76%) (Found: C, 43.14; H, 2.69; N, 0.71; P, 6.32; Ir, 39.75%. $\text{C}_{70}\text{H}_{51}\text{Ir}_4\text{NO}_9\text{P}_4$ requires C, 43.27; H, 2.65; N, 0.72; P, 6.38; Ir, 39.57%). $\nu_{\text{max}}/\text{cm}^{-1}$ (CH_2Cl_2): 2034s, 1983vs, 1924w $\nu(\text{CO})$. δ_{H} (CD_2Cl_2 , 298 K): 7.7–6.6 (50 H, m, Ph); 3.95 [1 H, t, $^2J(\text{P,H})$ 5.5 Hz, CH]. δ_{P} (CD_2Cl_2 , 298 K): 21.0 (PPN^+), -44.4 (PPh_2). δ_{C} (CD_2Cl_2 , 173 K): 177.1 (2 CO, s, d), 172.8 (2 CO, s, f), 167.0 (2 CO, s, c), 163.3 (1 CO, s, e), 160.2 (2 CO, m, g). Suitable crystals of **2** for an X-ray diffraction study were obtained by slow diffusion of hexane in a CH_2Cl_2 solution at 0°C .

Bis(triphenylphosphoranylidene)ammonium tetrahedrononacarbonyl(hydrido) $\{\mu$ -bis(diphenylphosphino)methane}tetrairidate (3**) ([PPN][$\text{Ir}_4(\text{CO})_9(\mu\text{-dppm})$])**

A solution of **1** (0.27 g, 0.19 mmol) and PPNCl (0.21 g, 0.37

mmol) in CH_2Cl_2 (20 cm^3) under CO was cooled to -10°C and DBU (600 μl , 4 mmol) was added. The solution was stirred for 2 h, then cold isopropanol (20 cm^3) was added and the volume of the mixture reduced by half. The orange precipitate was recrystallised at -30°C from CH_2Cl_2 (+PPNCl)/isopropanol and the crystals of **3** were washed with cold isopropanol and pentane (0.32 g, 87%) (Found: C, 43.08; H, 2.71; N, 0.69; P, 6.28; Ir, 39.70%. $\text{C}_{70}\text{H}_{53}\text{Ir}_4\text{NO}_9\text{P}_4$ requires C, 43.23; H, 2.75; N, 0.72; P, 6.37; Ir, 39.53%). $\nu_{\text{max}}/\text{cm}^{-1}$ (CH_2Cl_2): 2010m, 1960vs, 1827vs, 1766 m $\nu(\text{CO})$. δ_{H} (CD_2Cl_2 , 210 K): 7.6–7.0 (50 H; m, Ph); 6.03 [1 H, dt, $^2J(\text{H}_\text{A},\text{H}_\text{B})$ 12.8 Hz, $^2J(\text{P,H})$ 10.3 Hz, H_A], 2.69 [1 H, dt, $^2J(\text{H}_\text{A},\text{H}_\text{B})$ 12.8 Hz, $^2J(\text{P,H})$ 12.8 Hz, H_B], -15.7 (Ir-H). δ_{P} (CD_2Cl_2 , 210 K): 20.1 (PPN^+), -52.9 (PPh_2). δ_{C} (CD_2Cl_2 , 200 K): 225.0 (2 CO; s, a), 224.5 (1 CO, s, b), 186.7 (1 CO, s, d), 185.1 (2 CO, s, f), 165.0 (1 CO, s, e), 164.9 (2 CO, m, g). Suitable crystals of **3** for an X-ray diffraction study were obtained by slow diffusion of hexane in a CH_2Cl_2 solution at 0°C .

X-Ray crystallography

Compound 2. A yellow crystal was brought onto a Bruker SMART CCD system equipped with Mo radiation. All non-hydrogen atoms were refined anisotropically, and all hydrogens were made to ride on their associated carbons.

Compound 3. An orange crystal of which the habitus consisted of $\{100\}$, $\{001\}$ pinacoids, $(10, \bar{2})$, $(0, \bar{1} 0)$ pedions and a (161) fracture plane was cooled, using an Oxford Cryostream, to 90 K, the temperature of the data collection which took place on a Stoe IPDS system equipped with Mo radiation. All non-hydrogen atoms were refined anisotropically, and all hydrogens but H_4 were made to ride on their associated carbons. The hydrogen H_4 was difficult to find in difference maps and isotropically refined to a position surprisingly far away from Ir₄ [2.08(6) Å].

CCDC reference number 186/1923.

See <http://www.rsc.org/suppdata/dt/a9/a909881j/> for crystallographic files in .cif format.

Acknowledgements

We thank the Swiss National Science Foundation for financial support.

References

- 1 L. Malatesta and C. Caglio, *Chem. Commun.*, 1967, 420; R. Bau, M. Y. Chiang, C.-Y. Wei, L. Garlaschelli, S. Martinengo and T. F. Koetzle, *Inorg. Chim. Acta*, 1984, **23**, 4758; M. J. Davis and R. Roulet, *Inorg. Chim. Acta*, 1992, **197**, 15.
- 2 G. Ciani, M. Manassero, V. G. Albano, F. Canziani, G. Giordano, S. Martinengo and P. Chini, *J. Organomet. Chem.*, 1978, **150**, C17.
- 3 V. G. Albano, D. Braga, R. Ros and A. Scrivanti, *J. Chem. Soc., Chem. Commun.*, 1985, 866.
- 4 F. Ragaini, F. Porta and F. Demartin, *Organometallics*, 1991, **10**, 185.
- 5 G. Bondietti, R. Ros, R. Roulet, F. Musso and G. Gervasio, *Inorg. Chim. Acta*, 1993, **213**, 301.
- 6 R. J. Puddephatt, *Chem. Soc. Rev.*, 1983, **12**, 99.
- 7 A. Laguna and M. Laguna, *J. Organomet. Chem.*, 1990, **394**, 743.
- 8 R. Ros, A. Scrivanti, V. G. Albano, D. Braga and L. Garlaschelli, *J. Chem. Soc., Dalton Trans.*, 1986, 2411; A. Strawczynski, G. Suardi, R. Ros and R. Roulet, *Helv. Chim. Acta*, 1993, **76**, 2210.
- 9 T. Lumini, G. Laurenczy, R. Roulet, A. Tassan, R. Ros, K. Schenk and G. Gervasio, *Helv. Chim. Acta*, 1998, **81**, 781.
- 10 K. Besançon, G. Laurenczy, T. Lumini, R. Roulet, R. Bruyndonckx and C. Daul, *Inorg. Chem.*, 1998, **37**, 5634.
- 11 R. G. Teller and R. Bau (Editors), in *Structure and Bonding*, Springer-Verlag, Berlin, 1981, vol. 44, pp. 1–82.
- 12 M. Casarrubios and L. Seijo, *J. Chem. Phys.*, 1999, **110**, 784.
- 13 R. Roulet, in *The Synergy Between Dynamics and Reactivity at Clusters and Surfaces*, ed. L. J. Farrugia, Kluwer Academic Publishers, Dordrecht, 1995, pp. 159–173.
- 14 R. Ros and A. Tassan, *Inorg. Chim. Acta*, 1997, **260**, 89.
- 15 C. K. Johnson, ORTEP, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, TN, 1976.