

Ether functionalised aminophosphines: synthesis and co-ordination chemistry of palladium(II) and platinum(II) complexes†

Andrew D. Burrows,* Mary F. Mahon and Mark T. Palmer

Department of Chemistry, University of Bath, Claverton Down, Bath, UK BA2 7AY.
E-mail: a.d.burrows@bath.ac.uk

Received 22nd February 2000, Accepted 5th April 2000

Ether-functionalised aminophosphines Ph_2PNHR [$\text{R} = \text{CH}_2\text{CH}_2\text{OCH}_3$ L^1 , $\text{CH}_2\text{CH}(\text{OCH}_3)_2$ L^2 , $\text{CH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$ L^3 or $\text{C}_6\text{H}_4\text{OCH}_3$ -2 L^4] were prepared from the reaction of Ph_2PCl and the amine RNH_2 in the presence of triethylamine. Reaction of L^{1-4} with $[\text{MCl}_2(\text{cod})]$ ($\text{M} = \text{Pd}$ or Pt) or $[\text{PtBr}_2(\text{cod})]$ gave the complexes $[\text{MCl}_2\text{L}_2]$ (**1**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^1$; **2**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^2$; **3**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^3$; **4**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^4$; **5**, $\text{M} = \text{Pt}$, $\text{L} = \text{L}^1$; **6**, $\text{M} = \text{Pt}$, $\text{L} = \text{L}^2$; **7**, $\text{M} = \text{Pt}$, $\text{L} = \text{L}^3$) and $[\text{PtBr}_2(\text{L}^1)_2]$ **8**. X-Ray crystallographic analyses of complexes **1** and **2** indicated the presence of bifurcated hydrogen bonds between the NH protons and both the chloride ligands and the ether oxygen atoms. Reaction of **1**, **5** and **7** with two equivalents of AgBF_4 led to formation of *trans*- $[\text{Pd}(\text{L}^1)_2][\text{BF}_4]_2$ **9** and *cis*- $[\text{PtL}_2][\text{BF}_4]_2$ ($\text{L} = \text{L}^1$; **10** or L^4 **11**) containing bidentate P,O-co-ordinated ligands whereas reaction of **1** with one equivalent of AgBF_4 gave exclusively $[\{\text{Pd}(\mu\text{-Cl})(\text{L}^1)_2\}_2][\text{BF}_4]_2$ **12**, containing only unidentate aminophosphine ligands. Reaction of **5** and **8** with NaNO_2 led to formation of *cis*- $[\text{PtX}(\text{NO}_2)(\text{L}^1)_2]$ ($\text{X} = \text{Cl}$ **13** or Br **14**). Shielding of one of the NH protons with respect to **5** and **8** suggested loss of $\text{N-H} \cdots \text{X}$ hydrogen bonding and this was confirmed by an X-ray crystallographic analysis of **14**. Reaction of L^{1-4} with $[\{\text{M}(\text{dmba})(\mu\text{-Cl})\}_2]$ ($\text{M} = \text{Pd}$ or Pt ; $\text{Hdmba} = N,N$ -dimethylbenzylamine) gave $[\text{M}(\text{dmba})\text{ClL}]$ (**16**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^1$; **17**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^2$; **18**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^3$; **19**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^4$; **20**, $\text{M} = \text{Pt}$, $\text{L} = \text{L}^1$; **21**, $\text{M} = \text{Pt}$, $\text{L} = \text{L}^2$; **22**, $\text{M} = \text{Pt}$, $\text{L} = \text{L}^4$). The crystal structure of **16** reveals a similar bifurcated hydrogen bond to those observed in **1** and **2**, whereas the structure of **18** shows the $\text{N-H} \cdots \text{Cl}$ interaction to be present but the $\text{N-H} \cdots \text{O}$ interaction absent. Reaction of **16**, **20** and **22** with AgBF_4 gave the bidentate P,O-co-ordinated complexes $[\text{M}(\text{dmba})\text{L}]\text{BF}_4$ (**23**, $\text{M} = \text{Pd}$, $\text{L} = \text{L}^1$; **24**, $\text{M} = \text{Pt}$, $\text{L} = \text{L}^1$; **25**, $\text{M} = \text{Pt}$, $\text{L} = \text{L}^4$). Displacement of the co-ordinated ether oxygen was achieved by reaction of **24** and **25** with CO and of **25** with xylol isocyanide and acetonitrile. Attempts to recrystallise **24** led instead to isolation of the dinuclear species $[\{\text{Pt}(\text{dmba})(\mu\text{-PPh}_2\text{O})\}_2]$ **30** which was characterised by X-ray crystallography.

Introduction

Phosphines lie at the heart of modern co-ordination and organometallic chemistry, with their transition metal complexes of primary importance as homogeneous catalysts for many processes. Although there now exists a huge body of work on the chemistry of both phosphines and phosphites, aminophosphines (or phosphinamines) containing P–N bonds instead of P–C or P–O bonds have been relatively neglected as ligands. One potential reason for the underdevelopment of this chemistry is that the P–N bond is envisaged as being susceptible to relatively easy cleavage,¹ although a similar lack of stability to hydrolysis has not prevented the widespread use of phosphites.² Despite the relative lack of development, there are indications of the potential utility of transition metal complexes of aminophosphines. For example, rhodium(i)³ and platinum(ii)⁴ complexes of chiral aminophosphines have proved to be efficient catalysts for asymmetric hydrogenation and hydroformylation reactions respectively, and nickel complexes have been employed⁵ in the cyclodimerisation of buta-1,3-diene. In addition, Woollins and co-workers have reported a number of examples of aminophosphines including those derived from 1,2-diaminobenzene⁶ and $\text{Ph}_2\text{PNHCOPh}$,⁷ and aminophosphines of the type $\text{Ph}_2\text{PNHC}_6\text{H}_4\text{COR}$ have also recently been reported.⁸

The chemistry of ether phosphines has received considerable

attention, most notably from Lindner and co-workers,⁹ and the hemilabile co-ordination by the ether oxygen in ligands such as $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{OCH}_3$ to late transition metals has important catalytic implications.¹⁰ In this paper we explore the synthesis and reactivity of aminophosphines that contain ether functionalities.

Results and discussion

(i) Ligand synthesis

Since NH bonds are generally more acidic than CH bonds, the synthesis of aminophosphines from Ph_2PCl requires milder conditions than the analogous synthesis of phosphines. Reaction of Ph_2PCl with an equimolar amount of the appropriate primary amine in the presence of triethylamine leads to good yields of the aminophosphines $\text{Ph}_2\text{PNHCH}_2\text{CH}_2\text{OCH}_3$ (L^1), $\text{Ph}_2\text{PNHCH}_2\text{CH}(\text{OCH}_3)_2$ (L^2), $\text{Ph}_2\text{PNHCH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$ (L^3) and $\text{Ph}_2\text{PNHC}_6\text{H}_4\text{OCH}_3$ -2 (L^4) with precipitation of NEt_3HCl the main driving force. Care needs to be taken during the syntheses to exclude water, in order to prevent the formation of $\text{Ph}_2\text{PP}(\text{O})\text{Ph}_2$, which is catalysed by NEt_3 .¹¹ The aminophosphines L^{1-4} were isolated as colourless moisture-sensitive oils, and characterised on the basis of multinuclear NMR and IR spectroscopy (see Table 1) and microanalysis.

Compounds L^{1-4} are susceptible to alcoholysis, the reaction with methanol leading to cleavage of the P–N bond and formation of Ph_2POMe . The influence of the alkyl and aryl substituents on this reaction was studied by comparing the relative rates of methanolysis of L^1 and L^4 . 20 equivalents of methanol

† Electronic supplementary information (ESI) available: rotatable 3-D crystal structure diagram in CHIME format. See <http://www.rsc.org/suppdata/dt/b0/b001436m/>

Table 1 ^{31}P - $\{^1\text{H}\}$ NMR data, IR data and $\delta(\text{NH})$ from the ^1H NMR spectrum for ligands **L**^{1–4} and complexes **1–29**

Compound		$\delta(\text{P})$	$^1J(\text{Pt,P})/\text{Hz}$	$\delta(\text{NH})$	$\nu(\text{NH})/\text{cm}^{-1}$	Other data
L ¹	$\text{Ph}_2\text{PNHCH}_2\text{CH}_2\text{OCH}_3$	42.0	—	2.46	3370	
L ²	$\text{Ph}_2\text{PNHCH}_2\text{CH}(\text{OCH}_3)_2$	42.9	—	2.23	3368	
L ³	$\text{Ph}_2\text{PNHCH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$	41.9	—	2.20	3370	
L ⁴	$\text{Ph}_2\text{PNHC}_6\text{H}_4\text{OCH}_3$ -2	27.2	—	5.32	3381	
<i>cis</i> - 1	<i>cis</i> - $[\text{PdCl}_2(\text{L}^1)_2]$	59.0	—	4.40		
<i>trans</i> - 1	<i>trans</i> - $[\text{PdCl}_2(\text{L}^1)_2]$	46.4	—	4.28	3351	
<i>cis</i> - 2	<i>cis</i> - $[\text{PdCl}_2(\text{L}^2)_2]$	59.2	—	4.42		
<i>trans</i> - 2	<i>trans</i> - $[\text{PdCl}_2(\text{L}^2)_2]$	46.9	—	4.29	3331	
<i>cis</i> - 3	<i>cis</i> - $[\text{PdCl}_2(\text{L}^3)_2]$	58.8	—	4.47		
<i>trans</i> - 3	<i>trans</i> - $[\text{PdCl}_2(\text{L}^3)_2]$	46.2	—	4.11	3332	
4	<i>trans</i> - $[\text{PdCl}_2(\text{L}^4)_2]$	—	—	—	3300	
5	<i>cis</i> - $[\text{PtCl}_2(\text{L}^1)_2]$	35.5	3940	4.08	3380, 3293	
6	<i>cis</i> - $[\text{PtCl}_2(\text{L}^2)_2]$	35.8	3940	4.09	3350, 3247	
7	<i>cis</i> - $[\text{PtCl}_2(\text{L}^4)_2]$	30.1	3934	6.82	3370, 3199	
8	<i>cis</i> - $[\text{PtBr}_2(\text{L}^1)_2]$	36.7	3904	3.97	3399, 3293	
9	<i>trans</i> - $[\text{Pd}(\text{L}^1)_2][\text{BF}_4]_2$	80.1	—	4.07	3316	
10	<i>cis</i> - $[\text{Pt}(\text{L}^1)_2][\text{BF}_4]_2$	39.6	4346	3.87	3326	
11	<i>cis</i> - $[\text{Pt}(\text{L}^4)_2][\text{BF}_4]_2$	27.7	4162	6.48	3296	
12	$[\{\text{Pd}(\mu\text{-Cl})_2(\text{L}^1)_2\}_3][\text{BF}_4]_2$	61.4	—	4.39	3274	
13	<i>cis</i> - $[\text{PtCl}(\text{NO}_2)(\text{L}^1)_2]$	33.8	4115	4.40	3396, 3264	$^2J(\text{P,P})$ 24
		25.5	3186	3.50		
14	<i>cis</i> - $[\text{PtBr}(\text{NO}_2)(\text{L}^1)_2]$	33.8	4101	4.20	3397, 3311	$^2J(\text{P,P})$ 23
		25.5	3166	3.30		
15	<i>cis</i> - $[\text{Pt}(\text{NO}_2)_2(\text{L}^1)_2]$	26.8	3370	3.54		
16	$[\text{Pd}(\text{dmba})\text{Cl}(\text{L}^1)]$	68.4	—	4.40	3285	
17	$[\text{Pd}(\text{dmba})\text{Cl}(\text{L}^2)]$	68.4	—	4.48	3284	
18	$[\text{Pd}(\text{dmba})\text{Cl}(\text{L}^3)]$	67.6	—	4.40	3303	
19	$[\text{Pd}(\text{dmba})\text{Cl}(\text{L}^4)]$	59.7	—	6.60	3197	
20	$[\text{Pt}(\text{dmba})\text{Cl}(\text{L}^1)]$	42.1	4460	4.23	3308	
21	$[\text{Pt}(\text{dmba})\text{Cl}(\text{L}^2)]$	42.3	4470	4.24	3294	
22	$[\text{Pt}(\text{dmba})\text{Cl}(\text{L}^4)]$	34.8	4464	6.37	3214	
23	$[\text{Pd}(\text{dmba})(\text{L}^1)]\text{BF}_4$	75.2	—	3.40		
24	$[\text{Pt}(\text{dmba})(\text{L}^1)]\text{BF}_4$	48.8	4396	3.15	3399	
25	$[\text{Pt}(\text{dmba})(\text{L}^4)]\text{BF}_4$	60.9	4393	6.00	3390	
26	$[\text{Pt}(\text{dmba})(\text{L}^1)(\text{CO})]\text{BF}_4$	38.8	3666	3.57	3334	$\nu(\text{CO})$ 2114
27	$[\text{Pt}(\text{dmba})(\text{L}^4)(\text{CO})]\text{BF}_4$	35.4	3696	5.90		$\nu(\text{CO})$ 2105
28	$[\text{Pt}(\text{dmba})(\text{L}^4)(\text{CNXyl})]\text{BF}_4$	36.7	3897	5.85	3376	$\nu(\text{NC})$ 2179
29	$[\text{Pt}(\text{dmba})(\text{L}^4)(\text{NCCH}_3)]\text{BF}_4$	38.5	4256	5.74	3370	

were added to **L**¹ and **L**⁴ in CDCl_3 and the solutions monitored by ^{31}P - $\{^1\text{H}\}$ NMR spectroscopy. **L**¹ reacted slowly, with the reaction taking 24 h to reach completion, whereas in contrast **L**⁴ was fully converted into Ph_2POMe after 30 min. These results were somewhat surprising since it has previously been reported that aminophosphines with aryl substituents such as $(\text{Ph}_2\text{PNH})_2\text{C}_6\text{H}_4$ -1,2 are air-stable indefinitely.⁶

(ii) Palladium(II) and platinum(II) complexes $[\text{MCl}_2\text{L}_2]$

The reaction of two equivalents of **L**^{1–4} with $[\text{PdCl}_2(\text{cod})]$ in dichloromethane gave virtually quantitative yields of $[\text{PdCl}_2\text{L}_2]$ (**L** = **L**¹ **1**, **L**² **2**, **L**³ **3** or **L**⁴ **4**). The ^{31}P - $\{^1\text{H}\}$ NMR spectra of the palladium complexes **1–3** showed two phosphorus resonances which were assigned to the *cis* and *trans* isomers (Table 1). In accordance with this, two sets of resonances were also observed in the ^1H NMR spectra. Crystallisation of the mixture of geometric isomers gave exclusively *trans*- $[\text{PdCl}_2\text{L}_2]$ in the solid state (see below). The formation of an equilibrium mixture of the *cis* and *trans* isomers was followed by observing the slow isomerisation of the crystallographically characterised isomers *trans*- $[\text{PdCl}_2(\text{L}^1)_2]$ and *trans*- $[\text{PdCl}_2(\text{L}^2)_2]$ over a period of 24 hours. In both cases there was a 1 : 1 mixture of the *cis* and *trans* isomers at equilibrium in CDCl_3 . These studies enabled the ^{31}P - $\{^1\text{H}\}$ and ^1H resonances for the two isomers to be assigned. The complex $[\text{PdCl}_2\{\text{PPh}_2\text{NHP}(\text{O})\text{Ph}_2\}_2]$ also exists in the *trans* form in the solid state and as a *cis*–*trans* mixture in solution, though in this case there is a 1 : 3.7 ratio of the *cis* and *trans* isomers at equilibrium in CDCl_3 .¹² The complex $[\text{PdCl}_2(\text{L}^4)_2]$ **4** proved to be insoluble in common solvents so could not be characterised using NMR spectroscopy. However, the IR spectrum showed one $\nu(\text{NH})$ band as observed for

1–3 suggesting that **4** also exists in the solid state as the *trans* isomer.

The reaction of two equivalents of **L**¹, **L**² or **L**⁴ with $[\text{PtCl}_2(\text{cod})]$ in dichloromethane gave virtually quantitative yields of *cis*- $[\text{PtCl}_2\text{L}_2]$ (**L** = **L**¹ **5**, **L**² **6** or **L**⁴ **7**). In the ^{31}P - $\{^1\text{H}\}$ NMR spectra of complexes **5–7** $^1J(\text{Pt,P})$ is ca. 3940 Hz (Table 1), typical for phosphorus *trans* to chloride. The IR spectra for the *cis* complexes **5–7** all show two bands for $\nu(\text{NH})$, in contrast to the *trans* complexes **1–4** which show only one band. Reaction of **L**¹ with $[\text{PtBr}_2(\text{cod})]$ in an analogous manner gave *cis*- $[\text{PtBr}_2(\text{L}^1)_2]$ **8**.

Complexes **1** and **2** were recrystallised from dichloromethane–diethyl ether as yellow needle-shaped crystals, suitable for single crystal X-ray analyses. The crystal structures confirmed the *trans* orientation of the ligands; selected bond lengths and angles are given in Table 2 and the molecular structures are shown in Figs. 1 and 2. In both cases the asymmetric unit contains only half the molecule, with the remaining half being generated by inversion through a centre of symmetry on which the palladium atom resides with half-site occupancy. The palladium(II) centres are distorted square-planar, with *cis* angles of 88.49(4) and 91.51(4)° in **1** and 88.23(4) and 91.77(4)° in **2**. The Pd–P and Pd–Cl bond distances in **1** and **2** are unremarkable. The P–N distances [1.643(3) Å for **1**, 1.644(4) Å for **2**] are shorter than the generally accepted range for P–N single bonds (e.g. 1.689–1.727 Å in *N*-piperidinophosphines¹³), suggesting a degree of double bond character. This is generally observed in complexes of aminophosphines, and a search of the Cambridge Structural Database¹⁴ revealed the average P–N distance for aminophosphines of the type Ph_2PNHR (R = alkyl or aryl), unconstrained by chelation, to be 1.67 Å (range 1.63–1.73 Å). The sums of angles around the nitrogen atoms in **1** and **2**

Table 2 Selected bond lengths (Å) and angles (°) for complexes *trans*-1 and *trans*-2

	<i>trans</i> -1	<i>trans</i> -2
Pd(1)–Cl(1)	2.2963(10)	2.2957(12)
Pd(1)–P(1)	2.3158(10)	2.3304(12)
P(1)–N(1)	1.643(3)	1.644(4)
N(1)–C(13)	1.464(5)	1.456(5)
Cl(1)–Pd(1)–P(1)	91.51(4)	88.23(4)
Cl(1)–Pd(1)–P(1)′	88.49(4)	91.77(4)
N(1)–P(1)–Pd(1)	111.59(13)	111.31(14)
C(13)–N(1)–Pd(1)	127.6(3)	127.9(3)

Primed atoms generated by symmetry operations $-x, -y, -z + 1$ (*trans*-1), $-x, -y, -z$ (*trans*-2).




Fig. 1 Molecular structure of *trans*-[PdCl₂(L¹)₂] **1** with thermal ellipsoids at the 30% probability level. Unprimed atoms represent those in the asymmetric unit.




Fig. 2 Molecular structure of *trans*-[PdCl₂(L²)₂] **2**. Details as in Fig. 1.

are 354 and 360° respectively, consistent with significant sp² character.

The structure of complex **1** shows the presence of an intramolecular hydrogen bond between the N(1)–H(1) proton and the chloride ligand Cl(1) [N⋯Cl 3.170(3), H⋯Cl 2.54 Å, N–H⋯Cl 126°]. N–H⋯Cl hydrogen bonds have been

observed in a number of other complexes containing mutually *cis* aminophosphine and chloride ligands,^{12,15–20} and in these previous examples N⋯Cl distances range from 3.00 to 3.12 Å and H⋯Cl distances from 2.1 to 2.5 Å. The orientation of the ether chain suggests that the hydrogen bond is more accurately described as bifurcated, with the N–H⋯O interaction being the minor component of the 3-centre hydrogen bond. Vicinal intramolecular hydrogen bonding such as this is generally only observed in this manner due to the unfavourable angular geometry of the 5-membered hydrogen-bond ring, and the parameters [N⋯O 2.777(4), H⋯O 2.42 Å, N–H⋯O 104°] are similar to those observed for minor components of bifurcated hydrogen bonds in the crystal structures of nucleosides and nucleotides.^{21,22} A similar intramolecular bifurcated hydrogen bond is observed in the structure of **2** [N⋯Cl 3.139(3), H⋯Cl 2.49 Å, N–H⋯Cl 131°; N⋯O 2.861(4), H⋯O 2.56 Å, N–H⋯O 100°].

Unlike the free ligands L^{1–4}, complexes **1–7** do not readily react with methanol. The addition of an excess of wet methanol to a dichloromethane solution of **1** led to no change in the ³¹P-{¹H} NMR spectrum after 24 hours. The stabilisation of the aminophosphine ligands to methanolysis and hydrolysis on co-ordination is in direct contrast to diphosphazanes (Ph₂PNRPPH₂) for which the ligand is stable in methanol for several days at room temperature, but on co-ordination the P–N bond is readily cleaved.¹⁸ Similarly, Ph₂PNHP(O)Ph₂ is activated to methanolysis on co-ordination to platinum.¹⁶

In order to examine the potential bidentate co-ordination of the ether functionalised aminophosphines, the complexes *trans*-[PdCl₂(L¹)₂] **1** and *cis*-[PtCl₂L₂] (L = L¹ **5** or L⁴ **7**) were treated with two equivalents of AgBF₄. Reaction in dichloromethane led to the rapid precipitation of silver chloride, and the products, *trans*-[Pd(L¹)₂][BF₄]₂ **9** and *cis*-[Pt(L¹)₂][BF₄]₂ (L = L¹ **10** or L⁴ **11**), were characterised on the basis of multinuclear NMR spectroscopy and microanalysis. The platinum complexes **10** and **11** show a significant increase in ¹J(Pt,P), as expected on replacing chloride with a weaker *trans* influence ligand such as an ether oxygen. The ¹H NMR spectra too show evidence of co-ordination of the ether oxygen atoms, with a deshielding of the methoxy protons (Δδ = 0.43 ppm for **9** with respect to **1**, 0.63 ppm for **10** with respect to **5**, 0.14 ppm for **11** with respect to **7**) and, for **9** and **10**, deshielding of the methylene protons, now attached to a 6-membered chelate ring. It is notable also that the NH protons become shielded on formation of the chelate rings in **9**, **10** and **11**. This is a likely consequence of the loss of hydrogen bonding to the NH proton.


Reaction of **1** with one equivalent of AgBF₄ led exclusively to the chloro-bridged dimer [{Pd(μ-Cl)(L¹)₂]₂}(BF₄)₂ **12** which was characterised on the basis of multinuclear NMR spectroscopy, microanalysis and FAB-MS. Lindner and co-workers²³ have previously shown that a range of ether phosphines lead, under similar conditions, to complexes of the type [PdClL₂]⁺ in which one phosphine is bidentate and P,O-co-ordinated, whereas the other is unidentate and P-co-ordinated. Only for ether phosphines in which the oxygen atom is less basic were dimers observed. In our case, it is likely to be the unfavorable formation of a 6-membered ring that favours the formation of **12** over [PdCl(L¹)₂]⁺.

(iii) Nitrito complexes

In order to probe further the N–H⋯X hydrogen bonding in this system, halide substitution reactions were carried out. The reaction of *cis*-[PtX₂(L¹)₂] (X = Cl **5** or Br **8**) with sodium nitrite in aqueous acetone was followed by ³¹P-{¹H} NMR spectroscopy, and showed that in both cases the formation of *cis*-[PtX(NO₂)(L¹)₂] (X = Cl **13** or Br **14**) reached a maximum yield after ca. 4 hours. Continuation of the reaction for 24 hours using an excess of sodium nitrite led to *cis*-[Pt(NO₂)₂(L¹)₂] **15** in both cases, though this complex was not isolated from the

Table 3 Selected bond lengths (Å) and angles (°) for complex **14**

Pt(1)–P(1)	2.2564(15)	P(2)–N(2)	1.661(5)
Pt(1)–P(2)	2.2684(14)	N(1)–C(13)	1.471(8)
Pt(1)–Br(1)	2.4889(8)	N(2)–C(28)	1.475(8)
Pt(1)–N(3)	2.150(6)	N(3)–O(1)	1.2235(14)
P(1)–N(1)	1.653(5)	N(3)–O(2)	1.2246(10)
N(3)–Pt(1)–P(1)	91.91(12)	N(1)–P(1)–Pt(1)	110.9(2)
N(3)–Pt(1)–P(2)	173.86(13)	C(13)–N(1)–P(1)	127.8(4)
N(3)–Pt(1)–Br(1)	83.19(12)	N(2)–P(2)–Pt(1)	110.1(2)
P(1)–Pt(1)–P(2)	93.78(5)	C(28)–N(2)–P(2)	126.6(4)
P(1)–Pt(1)–Br(1)	174.96(4)	O(1)–N(3)–O(2)	130.4(7)
P(2)–Pt(1)–Br(1)	91.07(4)		

**Fig. 3** Molecular structure of *cis*-[PtBr(NO₂)(L)₂] **14** with thermal ellipsoids at the 30% probability level.

reaction mixture. The ³¹P-¹H} NMR spectra of **13** and **14** show two sets of doublets, each with ¹⁹⁵Pt satellites, as expected for complexes containing two inequivalent phosphorus atoms. The doublets can be assigned on the basis of ¹J(Pt,P), the value of which is greater for the phosphorus atom *trans* to the halide. The effect of the different electronic effects of halide and nitrite on ¹J(Pt,P) is also reflected in the coupling constant for **15** which at 3370 Hz is 570 Hz less than that for **5**.

The ¹H NMR spectra of complexes **13** and **14** show two resonances for δ(NH), separated by approximately 0.9 ppm. Comparison of these spectra with those for **5** and **8** show that one of the amine hydrogen atoms has become significantly more shielded. This shielding is probably a consequence of the loss of the intramolecular N–H···X hydrogen bonding for this proton. The ¹H NMR spectrum of **15** shows that both amine hydrogen atoms are shielded with respect to the amine protons in **5** and **8**, thus δ(NH) provides a useful probe of intramolecular hydrogen bonding in these compounds.

In order to assess whether the same pattern was observed in the solid state, a single crystal X-ray analysis was carried out. Suitable crystals of complex **14** were grown from dichloromethane–hexane as colourless blocks. Selected bond lengths and angles are given in Table 3 and the molecular structure is shown in Fig. 3. The platinum(II) centre in **14** is distorted square planar with *cis* angles between 83.19(12) and 93.78(5)°. The Pt–P and Pt–Br distances are unremarkable, and as with the structures of **1** and **2** the P–N distances [1.653(5) and 1.661(5) Å] are shorter than would be expected for single P–N bonds. The sums of angles round the nitrogen atoms are 352 and 355° respectively, showing significant sp² character. An intramolecular hydrogen bond is observed between the N(2)–H(2A) proton and the bromine atom Br(1) [N(2)···Br(1) 3.175(4), H(2A)···Br(1) 2.48 Å, N(2)–H(2A)···Br(1) 128°], and as with **1** and **2** the orientation of the ether chain suggests that there is a significant N–H···O interaction as the minor com-

Table 4 Selected bond lengths (Å) and angles (°) for complexes **16** and **18**

16		18	
Pd(1)–C(13)	1.955(6)	Pd(1)–C(17)	2.034(4)
Pd(1)–N(2)	2.141(5)	Pd(1)–N(2)	2.156(3)
Pd(1)–P(1)	2.247(2)	Pd(1)–P(1)	2.2472(8)
Pd(1)–Cl(1)	2.365(2)	Pd(1)–Cl(1)	2.3882(11)
P(1)–N(1)	1.616(6)	P(1)–N(1)	1.648(3)
N(1)–C(19)	1.470(8)	N(1)–C(13)	1.455(4)
C(13)–Pd(1)–N(2)	80.1(2)	C(17)–Pd(1)–N(2)	82.72(13)
C(13)–Pd(1)–P(1)	96.7(2)	C(17)–Pd(1)–P(1)	96.80(10)
N(2)–Pd(1)–P(1)	169.53(14)	N(2)–Pd(1)–P(1)	177.49(8)
C(13)–Pd(1)–Cl(1)	166.0(2)	C(17)–Pd(1)–Cl(1)	170.91(10)
N(2)–Pd(1)–Cl(1)	92.5(2)	N(2)–Pd(1)–Cl(1)	90.16(9)
P(1)–Pd(1)–Cl(1)	92.51(6)	P(1)–Pd(1)–Cl(1)	90.07(3)
N(1)–P(1)–Pd(1)	111.2(2)	N(1)–P(1)–Pd(1)	109.37(10)
C(19)–N(1)–P(1)	124.9(4)	C(13)–N(1)–P(1)	128.1(2)

ponent of a bifurcated hydrogen bond [N···O 2.791(7), H···O 2.67 Å, N–H···O 87°]. In contrast, no close contact is made by the amine hydrogen on the second aminophosphine ligand [N(1)–H(1A)] to either the nitrite or the ether oxygen atom, suggesting the solid state structure is the same as that observed in solution.

(iv) Palladium(II) and platinum(II) complexes [M(dmmba)ClL]

Complexes **9–11** were found readily to decompose in solution over several hours to give complex mixtures, so in an attempt to increase the stability of compounds containing L^{1–4} in a bidentate co-ordination mode complexes containing only one phosphine ligand were prepared. *N,N*-Dimethylbenzylamine (Hdmmba) was used as a bidentate monoanionic C,N-donor ligand to block two co-ordination sites of the metal complex. The reaction of two equivalents of the appropriate phosphine with [{M(dmmba)(μ-Cl)}₂] (M = Pd or Pt) gave the complexes [Pd(dmmba)ClL] (L = L¹ **16**, L² **17**, L³ **18** or L⁴ **19**) or [Pt(dmmba)ClL] (L = L¹ **20**, L² **21** or L⁴ **22**) in virtually quantitative yield.


The ³¹P-¹H} NMR spectra of complexes **16–19** showed single phosphorus resonances, suggesting that, in contrast to **1–3**, no isomerisation occurs in solution. The orientation of L^{1–4} *trans* to nitrogen can be inferred from the observation of ⁴J(P,H) coupling constants in the dimethylbenzylamine methyl and methylene resonances. This arrangement of phosphine *trans* to nitrogen was also observed for the complex [Pd(dmmba)ClL] (L = PPh₂CH₂CO₂Et)²⁴ and the non-observation of the other potential isomer was, in this case, ascribed to the lability of a phosphine *trans* to a σ-bonded carbon.

The ³¹P-¹H} NMR spectra of complexes **20–22** show single phosphorus resonances with ¹J(Pt,P) in the range 4460–4470 Hz. The high value is consistent with the phosphine *trans* to the nitrogen atom of the cyclometallated *N,N*-dimethylbenzylamine, and as for complexes **16–19**, the observation of ⁴J(P,H) couplings also supports this. ³¹P-¹H} NMR spectra of crude samples of **20** showed the presence of an additional compound [δ 53.9, ¹J(Pt,P) 2170]. The lower magnitude of ¹J(Pt,P) is consistent with this compound being the isomer with the phosphorus atom *trans* to carbon. If the reaction mixture is left overnight this compound is converted into **20**. It has previously been demonstrated that the reaction of cyclometallated platinum chloro-bridged dimers with pyridines can give either *N,N*-*cis* or *N,N*-*trans* products, depending on substituents.²⁵

Recrystallisation of complexes **16** and **18** from dichloromethane–pentane and dichloromethane–diethyl ether respectively produced single crystals suitable for X-ray analyses. Selected bond lengths and angles are given in Table 4 and the molecular structures are shown in Figs. 4 and 5. The palladium(II) centres in **16** and **18** are distorted square planar, with *cis* angles between 80.1(2) and 96.7(2)° and 82.73(13) and

Table 5 Selected bond lengths (Å) and angles (°) for complex **30**

Pt(1)–C(13)	2.00(2)	Pt(2)–O(2)	2.098(11)
Pt(1)–O(1)	2.116(11)	Pt(2)–N(2)	2.15(2)
Pt(1)–N(1)	2.160(15)	Pt(2)–P(2)	2.217(5)
Pt(1)–P(1)	2.205(5)	P(1)–O(2)	1.531(13)
Pt(2)–C(34)	2.01(2)	P(2)–O(1)	1.556(12)
C(13)–Pt(1)–O(1)	170.4(7)	O(2)–Pt(2)–N(2)	88.8(8)
C(13)–Pt(1)–N(1)	82.1(7)	C(34)–Pt(2)–P(2)	98.7(7)
O(1)–Pt(1)–N(1)	88.9(5)	O(2)–Pt(2)–P(2)	89.9(4)
C(13)–Pt(1)–P(1)	98.9(6)	N(2)–Pt(2)–P(2)	176.2(4)
O(1)–Pt(1)–P(1)	90.1(3)	O(2)–P(1)–Pt(1)	115.4(5)
N(1)–Pt(1)–P(1)	179.0(4)	O(1)–P(2)–Pt(1)	115.5(5)
C(34)–Pt(2)–O(2)	171.3(8)	P(2)–O(1)–Pt(1)	126.1(7)
C(34)–Pt(2)–N(2)	82.6(8)	P(1)–O(2)–Pt(2)	128.6(7)

**Fig. 6** Molecular structure of $[\{\text{Pt}(\text{dmmba})(\mu\text{-PPh}_2\text{O})\}_2]$ **30** with thermal ellipsoids at the 30% probability level. Hydrogen atoms and the CH_2Cl_2 of solvation have been omitted for clarity.

ation of the 6-membered ring in **30** can best be described as two distorted square planar units linked at an $\text{O} \cdots \text{O}$ hinge, with the angle between the $\text{Pt}(1)\text{--P}(1)\text{--O}(1)\text{--O}(2)$ and $\text{Pt}(2)\text{--P}(2)\text{--O}(1)\text{--O}(2)$ least squares planes being 48° . This conformation is in marked contrast to that observed in $[\{\text{Pd}[\text{PPh}_2\text{CH}=\text{C}(\text{O})\text{Ph}](\mu\text{-PPh}_2\text{O})\}_2]$ which is best described as a boat with the palladium atoms in the prows.²⁹

Experimental

Reactions were routinely carried out using Schlenk-line techniques under pure dry dinitrogen using dry dioxygen-free solvents unless noted otherwise. Microanalyses (C, H and N) were by Mr Alan Carver (University of Bath Microanalytical Service). Infrared spectra were recorded on a Nicolet 510P spectrometer as KBr pellets, Nujol mulls or dichloromethane solutions in KBr cells, ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR spectra on a JEOL JNM-EX270 spectrometer operating at 270 MHz referenced to TMS and 109.4 MHz referenced to H_3PO_4 , respectively, coupling constants are given in Hz and FAB mass spectra on a VG AutoSpec-Q spectrometer using 3-nitrobenzyl alcohol as the matrix. The complexes $[\text{PtX}_2(\text{cod})]$ ($\text{X} = \text{Cl}$ or Br),³¹ $[\text{PdCl}_2(\text{cod})]$,³² and $[\{\text{M}(\text{dmmba})(\mu\text{-Cl})\}_2]$ ($\text{M} = \text{Pd}$ or Pt)³³ were prepared by standard literature methods. $\text{L}^1\text{--L}^4$ were isolated in 80–90% yield, the palladium and platinum complexes in 70–90% yield unless noted otherwise below.

Syntheses

$\text{Ph}_2\text{PNHCH}_2\text{CH}_2\text{OCH}_3$ (L^1). Triethylamine (0.566 g, 5.5 mmol) and PPh_2Cl (1.23 g, 5.5 mmol) were added sequentially with stirring to a solution of 2-methoxyethylamine (0.41 g, 5.5 mmol) in THF (20 cm^3). The reaction mixture was stirred for 30 minutes and the solution filtered to remove NEt_3HCl , then evaporated under reduced pressure to give a pale yellow, viscous

oil (Found: C, 69.3; H, 6.96; N, 5.15. $\text{C}_{15}\text{H}_{18}\text{NOP}$ requires C, 69.5; H, 7.00; N, 5.40%). ^1H NMR (CDCl_3): δ 7.47–7.40 (m, 4 H, Ar), 7.33–7.24 (m, 6 H, Ar), 3.32 (t, 2 H, CH_2O), 3.23 (s, 3 H, CH_3), 3.07 (dt, 2 H, CH_2N) and 2.46 (m, 1 H, NH). ^{13}C NMR (CDCl_3): δ 141.2 (d, $^1J(\text{CP})$ 12, C_{ipso}), 130.8 (d, $^2J(\text{CP})$ 20, C_{ortho}), 128.0 (s, C_{para}), 127.7 (d, $^3J(\text{CP})$ 6, C_{meta}), 73.5 (d, $^3J(\text{CP})$ 6, CH_2O), 58.1 (s, CH_3) and 45.3 (d, $^2J(\text{CP})$ 14, CH_2N).

L^2 . As for L^1 using NEt_3 (1.13 g, 11 mmol), PPh_2Cl (2.46 g, 11 mmol) and 2,2-dimethoxyethylamine (1.17 g, 11 mmol). Product extracted with diethyl ether at -78°C (Found: C, 66.4; H, 6.86; N, 4.70. $\text{C}_{16}\text{H}_{20}\text{NO}_2\text{P}$ requires C, 66.4; H, 6.97; N, 4.84%). ^1H NMR (CDCl_3): δ 7.39–7.23 (m, Ar), 4.13 (t, 1 H, CH), 3.22 (s, 6 H, CH_3), 3.00 (m, 2 H, CH_2) and 2.23 (m, 1 H, NH). ^{13}C NMR (CD_2Cl_2): δ 141.4 (d, $^1J(\text{CP})$ 12, C_{ipso}), 130.7 (d, $^2J(\text{CP})$ 20, C_{ortho}), 127.8 (s, C_{para}), 127.6 (d, $^3J(\text{CP})$ 6, C_{meta}), 104.3 (d, $^3J(\text{CP})$ 6, CH), 53.0 (s, CH_3) and 47.2 (d, $^2J(\text{CP})$ 14, CH_2).

L^3 . As for L^1 using NEt_3 (1.11 g, 11 mmol), PPh_2Cl (2.41 g, 11 mmol) and 3-methoxypropylamine (0.98 g, 11 mmol). Product extracted with diethyl ether at -78°C followed by hexane (Found: C, 70.8; H, 7.20; N, 4.67. $\text{C}_{16}\text{H}_{20}\text{NOP}$ requires C, 70.3; H, 7.38; N, 5.12%). ^1H NMR (CDCl_3): δ 7.52–7.46 (m, 4 H, Ar), 7.40–7.33 (m, 6 H, Ar), 3.44 (t, 2 H, CH_2O), 3.32 (s, 3 H, CH_3), 3.07 (m, 2 H, CH_2N), 2.20 (m, 1 H, NH) and 1.76 (q, 2 H, CH_2C_2). ^{13}C NMR (CDCl_3): δ 141.3 (d, C_{ipso}), 130.9 (d, $^2J(\text{CP})$ 18, C_{ortho}), 127.8 (s, C_{para}), 127.7 (s, C_{meta}), 70.2 (s, CH_2O), 58.0 (s, CH_3), 43.1 (d, $^2J(\text{CP})$ 11, CH_2N) and 32.1 (s, CCH_2C).

L^4 . As for L^1 using NEt_3 (1.13 g, 11 mmol), PPh_2Cl (2.46 g, 11 mmol) and *o*-anisidine (1.38 g, 11 mmol). Product extracted with diethyl ether at -78°C (Found: C, 73.3; H, 6.11; N, 4.95. $\text{C}_{19}\text{H}_{18}\text{NOP}$ requires C, 74.2; H, 5.90; N, 4.56%). ^1H NMR (CDCl_3): δ 7.70–7.64 (m, 4 H, Ar), 7.55–7.44 (m, 6 H, Ar), 7.02 (m, 2 H), 6.93 (m, 2 H), 5.32 (d, 1 H, $^2J(\text{PH})$ 7, NH) and 3.93 (s, 3 H, CH_3). ^{13}C NMR (CDCl_3): δ 147.3 (d, $^3J(\text{CP})$ 4, C2), 139.8 (d, $^2J(\text{CP})$ 12, C_{ipso}), 135.7 (d, $^2J(\text{CP})$ 18, C1), 130.6 (d, $^2J(\text{CP})$ 21, C_{ortho}), 128.5 (s, C_{para}), 128.0 (d, $^3J(\text{CP})$ 7, C_{meta}), 120.7 (s, C5), 118.3 (s, C4), 113.6 (d, $^3J(\text{CP})$ 21, C6), 109.6 (s, C3) and 54.9 (s, CH_3).

$[\text{PdCl}_2(\text{L}^1)_2]$ **1.** The complex $[\text{PdCl}_2(\text{cod})]$ (0.100 g, 0.35 mmol) was added with stirring to a solution of L^1 (0.182 g, 0.70 mmol) in dichloromethane (40 cm^3). After 30 min the solution was concentrated under reduced pressure, and diethyl ether added to give yellow crystals of complex **1** (Found: C, 50.9; H, 5.20; N, 3.87. $\text{C}_{30}\text{H}_{36}\text{Cl}_2\text{N}_2\text{O}_2\text{P}_2\text{Pd}$ requires C, 50.7; H, 5.13; N, 3.91%). ^1H NMR (CDCl_3): *trans* isomer, δ 7.85–7.75 (m, Ar), 7.47–7.36 (m, Ar), 4.26 (m (br), 1 H, NH), 3.22 (s, 3 H, CH_3), 3.22 (t, 2 H, CH_2O) and 2.76 (m, 2 H, CH_2N); *cis* isomer, δ 7.8–7.3 (m, Ar), 4.45 (m (br), 1 H, NH), 3.28 (t, 2 H, CH_2O), 3.16 (s, 3 H, CH_3) and 2.55 (m, 2 H, CH_2N). FAB-MS: m/z 697, $[\text{M} + \text{H}]^+$; and 661, $[\text{M} - \text{Cl}]^+$.

Complex 2. As for complex **1** using $[\text{PdCl}_2(\text{cod})]$ (0.100 g, 0.35 mmol) and L^2 (0.203 g, 0.70 mmol) (Found: C, 50.6; H, 5.31; N, 3.68. $\text{C}_{32}\text{H}_{40}\text{Cl}_2\text{N}_2\text{O}_4\text{P}_2\text{Pd}$ requires C, 50.8; H, 5.33; N, 3.71%). ^1H NMR (CDCl_3): *trans* isomer, δ 7.90–7.82 (m, 4 H, Ar), 7.53–7.44 (m, 6 H, Ar), 4.29 (m (br), 1 H, NH), 4.05 (t, 1 H, CH), 3.25 (s, 6 H, CH_3) and 2.75 (m, 2 H, CH_2N); *cis* isomer, δ 7.94–7.86 (m, Ar), 7.58–7.35 (m, Ar), 4.42 (m, 1 H, NH), 4.02 (m, 1 H, CH), 3.23 (s, 6 H, CH_3) and 2.52 (m, 2 H, CH_2N). FAB-MS: m/z 756, $[\text{M} + \text{H}]^+$; and 721, $[\text{M} - \text{Cl}]^+$. Yield 0.177 g (67%).

Complex 3. As for complex **1** using $[\text{PdCl}_2(\text{cod})]$ (0.103 g, 0.36 mmol) and L^3 (0.198 g, 0.73 mmol) (Found: C, 53.0; H, 5.58; N, 3.82. $\text{C}_{32}\text{H}_{40}\text{Cl}_2\text{N}_2\text{O}_2\text{P}_2\text{Pd}$ requires C, 53.1; H, 5.57; N, 3.87%). ^1H NMR (CDCl_3): *cis-trans* mixture, δ 7.86–7.80 (m, Ar), 7.60–7.36 (m, Ar), 7.31–7.25 (m, Ar), 4.49 (br, NH), 4.11 (br, NH),

3.32 (t, CH₂O), 3.25 (m, CH₂O), 3.23 (s, CH₃), 3.19 (s, CH₃), 2.73 (m, CH₂N), 2.48 (m, CH₂N), 1.58 (q, CH₂C₂) and 1.50 (q, CH₂C₂).

Complex 4. As for complex **1** using [PdCl₂(cod)] (0.100 g, 0.35 mmol) and **L**⁴ (0.216 g, 0.70 mmol) (Found: C, 57.3; H, 4.66; N, 3.46. C₃₈H₃₆Cl₂N₂O₂P₂Pd requires C, 57.6; H, 4.58; N, 3.54%).

Complex 5. As for complex **1** using [PtCl₂(cod)] (0.100 g, 0.27 mmol) and **L**¹ (0.138 g, 0.53 mmol). Colourless crystals (Found: C, 44.2; H, 4.57; N, 3.37. C₃₀H₃₆Cl₂N₂O₂P₂Pt·½CH₂Cl₂ requires C, 44.3; H, 4.51; N, 3.39%). ¹H NMR (CDCl₃): δ 7.57 (m, 4 H, Ar), 7.43 (m, 2 H, Ar), 7.30 (m, 4 H, Ar), 4.08 (m (br), 1 H, NH), 3.14 (s, 3 H, CH₃), 3.09 (t, 2 H, CH₂O) and 2.52 (m, 2 H, CH₂N). FAB-MS: *m/z* 785, [M + H]⁺; and 749, [M – Cl]⁺.

Complex 6. As for complex **1** using [PtCl₂(cod)] (0.136 g, 0.36 mmol) and **L**² (0.210 g, 0.73 mmol). Colourless crystals (Found: C, 44.1; H, 4.72; N, 3.07. C₃₂H₄₀Cl₂N₂O₂P₂Pt·½CH₂Cl₂ requires C, 44.0; H, 4.66; N, 3.16%). ¹H NMR (CDCl₃): δ 7.56 (m, 4 H, Ar), 7.45 (m, 2 H, Ar), 7.32 (m, 4 H, Ar), 4.09 (m (br), 1 H, NH), 3.93 (t, 1 H, CH), 3.14 (s, 6 H, CH₃) and 2.45 (m, 2 H, CH₂).

Complex 7. As for complex **1** using [PtCl₂(cod)] (0.100 g, 0.27 mmol) and **L**⁴ (0.164 g, 0.53 mmol). Colourless crystals (Found: C, 50.1; H, 4.09; N, 2.72. C₃₈H₃₆Cl₂N₂O₂P₂Pt·½CH₂Cl₂ requires C, 50.1; H, 4.04; N, 3.03%). ¹H NMR (CDCl₃): δ 7.62 (m, 4 H, Ar), 7.46 (m, 2 H, Ar), 7.30 (m, 4 H, Ar), 6.82 (m, 1 H, NH), 6.74 (m, Ar), 6.72 (m, Ar), 6.33 (m, Ar), 6.07 (d, 1 H, Ar) and 3.65 (s, 3 H, CH₃). FAB-MS: *m/z* 880, [M]⁺; 844, [M – Cl]⁺; and 808 [M – 2Cl]⁺.

Complex 8. As for complex **1** using [PtBr₂(cod)] (0.100 g, 0.21 mmol) and **L**¹ (0.112 g, 0.43 mmol). Colourless crystals (Found: C, 40.3; H, 4.01; N, 2.75. C₃₀H₃₆Br₂N₂O₂P₂Pt·½CH₂Cl₂ requires C, 40.0; H, 4.07; N, 3.06%). ¹H NMR (CDCl₃): δ 7.62 (m, 4 H, Ar), 7.45 (m, 2 H, Ar), 7.35 (m, 4 H, Ar), 3.97 (m (br), 1 H, NH), 3.16 (s, 3 H, CH₃), 3.12 (t, 2 H, CH₂O) and 2.55 (m, 2 H, CH₂N).

Reaction of complex 1 with AgBF₄ giving [Pd(L¹)]₂[BF₄]₂ **9.** The salt AgBF₄ (0.093 g, 0.48 mmol) was added to a solution of complex **1** (0.150 g, 0.22 mmol) in dichloromethane (30 cm³) and the mixture stirred in darkness for 1 hour. The resulting solution was filtered, the residue washed with dichloromethane, and the filtrate and washings were combined. The solvent was then evaporated under reduced pressure to give **9** as an orange solid. ¹H NMR (CDCl₃): δ 7.59 (m, Ar), 7.39 (m, Ar), 4.07 (m (br), 1 H, NH), 3.93 (m, 2 H, CH₂O), 3.65 (s, 3 H, CH₃) and 3.20 (m, 2 H, CH₂N).

Complex 10. As for complex **9** using AgBF₄ (0.060 g, 0.31 mmol) and **5** (0.100 g, 0.13 mmol). Recrystallisation from dichloromethane–hexane gave colourless microcrystals (Found: C, 38.1; H, 4.15; N, 2.56. C₃₀H₃₆B₂F₈N₂O₂P₂Pt·CH₂Cl₂ requires C, 38.3; H, 3.94; N, 2.88%). ¹H NMR (CDCl₃): δ 7.61 (m, Ar), 7.52 (m, Ar), 7.35 (m, Ar), 4.20 (m, 2 H, CH₂O), 3.87 (m (br), 1 H, NH), 3.77 (s, 3 H, CH₃) and 3.22 (m, 2 H, CH₂N).

Complex 11. As for complex **9** using AgBF₄ (0.117 g, 0.60 mmol) and **7** (0.213 g, 0.30 mmol). Recrystallisation from dichloromethane–hexane gave yellow microcrystals (Found: C, 46.4; H, 3.79; N, 2.48. C₃₈H₃₆B₂F₈N₂O₂P₂Pt requires C, 46.4; H, 3.69; N, 2.85%). ¹H NMR (CD₂Cl₂): δ 7.44 (m, Ar), 7.22 (m, Ar), 6.82 (m, 2 H, Ar), 6.48 (d, ²J(PH) 8, NH), 6.40 (m, 2 H, Ar) and 3.79 (s, 3 H, CH₃). Yield 0.076 g (32%).

Complex 12. As for complex **9** using AgBF₄ (0.030 g, 0.15 mmol) and **1** (0.107 g, 0.15 mmol) (Found: C, 47.1; H, 4.83; N,

3.66. C₆₀H₇₂B₂Cl₂F₈N₄O₄P₄Pd₂·½CH₂Cl₂ requires C, 47.3; H, 4.79; N, 3.65%). ¹H NMR (CDCl₃): δ 7.35 (m, Ar), 4.39 (br, NH), 3.32 (t, CH₂O), 3.26 (s, CH₃) and 2.72 (t, CH₂N). FAB-MS: *m/z* 1407, [M – BF₄]⁺; 1321, [M – 2BF₄]⁺; and 660, [PdCl(L¹)₂]⁺.

[PtCl(NO₂)(L¹)₂] **13.** A solution of NaNO₂ (0.070 g, 1.0 mmol) in water (4 cm³) was added with stirring to a solution of complex **5** (0.150 g, 0.19 mmol) in acetone (20 cm³). The mixture was stirred for 4 hours, after which the solvent was removed under reduced pressure, the product extracted with dichloromethane (3 × 20 cm³) and recrystallised from dichloromethane–diethyl ether, followed by dichloromethane–hexane to give colourless crystals (Found: C, 42.2; H, 4.24; N, 4.82. C₃₀H₃₆ClN₃O₄P₂Pt·CH₂Cl₂ requires C, 42.3; H, 4.35; N, 4.77%). ¹H NMR (CDCl₃): δ 7.68–7.28 (m, br, Ar), 4.40 (m, br, NH), 3.50 (m, br, NH), 3.20–3.17 (m, CH₂O/CH₃O), 2.72 (m, CH₂N) and 2.58 (m, CH₂N).

Complex 14. As for complex **13** using NaNO₂ (0.300 g, 4.35 mmol) and **8** (0.377 g, 0.43 mmol). Recrystallisation from dichloromethane–hexane gave colourless crystals (Found: C, 42.8; H, 4.34; N, 4.92. C₃₀H₃₆BrN₃O₄P₂Pt requires C, 42.9; H, 4.32; N, 5.01%). ¹H NMR (CDCl₃): δ 7.60 (m, Ar), 7.43 (m, Ar), 7.32 (m, Ar), 4.20 (m, NH), 3.30 (m, NH), 3.21 (s, CH₃), 3.19 (s, CH₃), 3.16 (m, CH₂O), 2.73 (m, CH₂N) and 2.56 (m, CH₂N). FAB-MS: *m/z* 793, [M – NO₂]⁺; and 712, [M – NO₂ – Br]⁺. Yield 0.162 g (45%).

[Pd(dmba)Cl(L¹)] **16.** The complex [{Pd(dmba)(μ-Cl)}₂] (0.400 g, 0.75 mmol) was added to a solution of **L**¹ (0.375 g, 1.45 mmol) in dichloromethane (30 cm³). The solution was stirred for 1 hour, after which it was concentrated under reduced pressure, filtered and pentane added, to give yellow crystals of **16** (Found: C, 53.5; H, 5.66; N, 5.19. C₂₄H₃₀ClN₂OPPd requires C, 53.8; H, 5.65; N, 5.23%). ¹H NMR (CDCl₃): δ 7.85 (m, Ar), 7.35 (m, Ar), 6.93 (d, Ar), 6.82 (t, Ar), 6.53 (m, Ar), 4.41 (m, br, NH), 3.91 (m, br, CH₂N (dmba)), 3.23 (m, br, CH₂O/CH₃O) and 2.76 (m, NCH₂, CH₃N). FAB-MS: *m/z* 535, [M + H]⁺; and 499, [M – Cl]⁺.

Complex 17. As for complex **16** using [{Pd(dmba)(μ-Cl)}₂] (0.200 g, 0.36 mmol) and **L**² (0.210 g, 0.73 mmol). Recrystallisation from dichloromethane–pentane gave yellow crystals (Found: C, 52.4; H, 5.61; N, 4.80. C₂₅H₃₁ClN₂O₂PPd requires C, 53.2; H, 5.54; N, 4.96%). ¹H NMR (CDCl₃): δ 7.85 (m, 4 H, Ar), 7.42 (m, 6 H, Ar), 6.95 (d, 1 H, Ar), 6.82 (m, 1 H, Ar), 6.55 (m, 2 H, Ar), 4.48 (m, 1 H, NH), 4.14 (t, 1 H, CH), 3.93 (s, 2 H, CH₂N (dmba)), 3.25 (s, 6 H, CH₃O), 2.78 (d, J(CP) 3, 6 H, CH₃N) and 2.74 (m, 2 H, CH₂).

Complex 18. As for complex **16** using [{Pd(dmba)(μ-Cl)}₂] (0.100 g, 0.18 mmol) and **L**³ (0.099 g, 0.36 mmol). Recrystallisation from dichloromethane–diethyl ether gave yellow crystals (Found: C, 54.2; H, 5.85; N, 5.01. C₂₅H₃₂ClN₂OPPd requires C, 54.7; H, 5.87; N, 5.10%). ¹H NMR (CDCl₃): δ 7.80 (m, Ar), 7.40 (m, Ar), 6.89 (d, 1 H, Ar), 6.78 (m, 1 H, Ar), 6.50 (m, 2 H, Ar), 4.40 (m, 1 H, NH), 3.89 (s, 2 H, CH₂N (dmba)), 3.30 (t, 2 H, CH₂O), 3.19 (s, 3 H, CH₃O), 2.74 (m, CH₂N/CH₃N) and 1.57 (m, 2 H, CH₂C₂).

Complex 19. As for complex **16** using [{Pd(dmba)(μ-Cl)}₂] (0.275 g, 0.90 mmol) and **L**⁴ (0.247 g, 0.45 mmol). Recrystallisation from dichloromethane–pentane gave yellow microcrystals (Found: C, 49.9; H, 4.53; N, 3.91. C₂₈H₃₀ClN₂OPPd·½CH₂Cl₂ requires C, 49.9; H, 4.68; N, 3.94%). ¹H NMR (CDCl₃): δ 7.85 (m, 4 H, Ar), 7.39 (m, 7 H, Ar), 6.98 (m, 1 H, Ar), 6.88 (m, 1 H, Ar), 6.71 (m, 2 H, Ar), 6.59 (m, 2 H, Ar), 6.42 (m, 1 H, Ar), 6.30 (d, ²J(PH) 8, 1 H, NH), 3.94 (d, ⁴J(PH) 2, 2 H, CH₂N (dmba)), 3.81 (s, 3 H, CH₃O) and 2.77 (d, 6 H ⁴J(PH) 3, CH₃N).

Table 6 Crystallographic data for complexes **1**, **2**, **14**, **16**, **18** and **30**

	1	2	14	16	18	30
Empirical formula	C ₃₀ H ₃₆ Cl ₂ N ₂ O ₂ ·P ₂ Pd	C ₃₂ H ₄₀ Cl ₂ N ₂ O ₄ ·P ₂ Pd	C ₃₀ H ₃₆ BrN ₃ O ₄ ·P ₂ Pt	C ₂₄ H ₃₀ ClN ₂ ·OPPd	C ₂₅ H ₃₂ ClN ₂ ·OPPd	C ₄₃ H ₄₆ Cl ₂ N ₃ O ₂ ·P ₂ Pt ₂
<i>M</i>	695.84	755.90	839.56	535.32	549.35	1145.84
<i>T</i> /K	150(2)	293(2)	293(2)	293(2)	293(2)	293(2)
Crystal system	Monoclinic	Triclinic	Monoclinic	Orthorhombic	Monoclinic	Monoclinic
Space group	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> $\bar{1}$	<i>P</i> 2 ₁ / <i>n</i>	<i>Pbca</i>	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> 2 ₁ / <i>c</i>
<i>a</i> /Å	10.456(2)	8.396(2)	9.576(2)	14.070(2)	16.835(2)	13.877(5)
<i>b</i> /Å	7.553(1)	10.146(2)	24.119(5)	17.080(3)	8.3350(9)	20.984(7)
<i>c</i> /Å	19.935(2)	10.849(3)	14.306(2)	18.899(4)	19.590(2)	15.345(5)
<i>a</i> /°		94.97(2)				
<i>β</i> /°	100.641(13)	111.01(2)	105.09(2)		111.922(2)	107.09(3)
<i>γ</i> /°		94.71(2)				
<i>U</i> /Å ³	1547.3(4)	853.1(4)	3190.2(10)	4541.7(14)	2550.1(5)	4271(3)
<i>Z</i>	2	1	4	8	4	4
<i>μ</i> /mm ^{−1}	0.906	0.832	5.790	1.024	0.914	6.781
Reflections collected	6352	3413	5596	3785	14866	6620
Independent reflections	2395	2973	5587	3715	5207	6620
	[<i>R</i> (int) = 0.0904]	[<i>R</i> (int) = 0.0537]	[<i>R</i> (int) = 0.0329 ^a]	[<i>R</i> (int) = 0.002]	[<i>R</i> (int) = 0.0385]	[<i>R</i> (int) = 0.0267 ^a]
<i>R</i> 1, <i>wR</i> 2 [<i>I</i> > 2σ(<i>I</i>)]	0.0383, 0.1084	0.0518, 0.1298	0.0325, 0.0887	0.0534, 0.1366	0.0392, 0.0942	0.0715, 0.1449
(all data)	0.0451, 0.1098	0.0620, 0.1458	0.0454, 0.0949	0.0874, 0.1446	0.058, 0.1028	0.1541, 0.1776

^a Pre DIFABS.

Complex 20. As for complex **16** using [{Pt(dmba)(μ-Cl)]₂ (0.141 g, 0.19 mmol) and **L**¹ (0.100 g, 0.39 mmol). Recrystallisation from dichloromethane–diethyl ether gave colourless crystals (Found: C, 45.9; H, 4.85; N, 4.34. C₂₄H₃₀ClN₂OPPt requires C, 46.2; H, 4.85; N, 4.49%). ¹H NMR (CDCl₃): δ 7.86 (m, Ar), 7.40 (m, Ar), 6.98 (m, Ar), 6.86 (m, Ar), 6.75 (m, Ar), 6.52 (m, Ar), 4.23 (m, br, NH), 3.95 (m, CH₂N (dmba)), 3.28 (t, CH₂O), 3.25 (s, CH₃O), 2.89 (m, CH₃N) and 2.86 (m, CH₂N).

Complex 21. As for complex **16** using [{Pt(dmba)(μ-Cl)]₂ (0.189 g, 0.26 mmol) and **L**² (0.150 g, 0.52 mmol). Recrystallisation from dichloromethane–hexane gave a pale yellow powder (Found: C, 46.0; H, 5.06; N, 4.21. C₂₅H₃₁ClN₂O₂Pt requires C, 45.9; H, 4.93; N, 4.28%). ¹H NMR (CDCl₃): δ 7.88 (m, 4 H, Ar), 7.42 (m, 6 H, Ar), 7.02 (d, 1 H, Ar), 6.89 (t, 1 H, Ar), 6.78 (m, 1 H, Ar), 6.55 (m, 1 H, Ar), 4.24 (m, 1 H, NH), 4.18 (t, 1 H, CH), 3.99 (m, 2 H, CH₂N (dmba)), 3.28 (s, 6 H, CH₃O), 2.94 (m, 6 H, CH₃N) and 2.82 (m, 2 H, CH₂).

Complex 22. As for complex **16** using [{Pt(dmba)(μ-Cl)]₂ (0.257 g, 0.35 mmol) and **L**⁴ (0.217 g, 0.71 mmol). Recrystallisation from THF–hexane, followed by dichloromethane–hexane gave yellow microcrystals (Found: C, 46.6; H, 4.40; N, 3.64. C₂₈H₃₀ClN₂OPPt· $\frac{3}{4}$ CH₂Cl₂ requires C, 46.9; H, 4.32; N, 3.81%). ¹H NMR (CDCl₃): δ 7.88 (m, 4 H, Ar), 7.39 (m, 6 H, Ar), 7.27 (m, 1 H, Ar), 7.02 (m, 1 H, Ar), 6.88 (m, 2 H, Ar), 6.71 (m, 2 H, Ar), 6.58 (m, 1 H, Ar), 6.46 (m, 1 H, Ar), 6.37 (d, ²*J*(PH) 8, 1 H, NH), 3.96 (m, ³*J*(PtH) 28, ⁴*J*(PH) 3, 2 H, CH₂N (dmba)), 3.80 (s, 3 H, CH₃O) and 2.90 (m, 6 H, ³*J*(PtH) 23, ⁴*J*(PH) 3, CH₃N). Yield 0.218 g (46%).

Reaction of complex 16 with AgBF₄ giving [Pd(dmba)(L¹)]BF₄ **23.** The salt AgBF₄ (0.122 g, 0.63 mmol) was added to a solution of complex **16** (0.200 g, 0.37 mmol) in dichloromethane (20 cm³) and the mixture stirred in darkness for 30 min. The resulting solution was filtered, the residue washed with dichloromethane, and the filtrate and washings were combined. This solvent was then evaporated under reduced pressure to give **23** as a brown solid. ¹H NMR (CDCl₃): δ 7.79 (m, 4 H, Ar), 7.47 (m, 6 H, Ar), 6.94 (d, 1 H, Ar), 6.84 (t, 1 H, Ar), 6.43 (t, 1 H, Ar), 6.19 (t, 1 H, Ar), 4.01 (m, 2 H, CH₂N (dmba)), 3.85 (m, 2 H, CH₂O), 3.40 (br, 1 H, NH), 3.33 (m, 5 H, CH₃O/CH₂N) and 2.73 (d, 6 H, CH₃N).

Complex 24. As for complex **23** using AgBF₄ (0.052 g, 0.27 mmol) and **20** (0.100 g, 0.16 mmol). Recrystallisation from

dichloromethane–pentane gave colourless crystals (Found: C, 42.9; H, 4.65; N, 3.96. C₂₄H₃₀BF₄N₂OPPt requires C, 42.7; H, 4.48; N, 4.15%). ¹H NMR (CDCl₃): δ 7.83 (m, Ar), 7.48 (m, Ar), 6.92 (d, Ar), 6.77 (t, Ar), 6.32 (t, Ar), 6.15 (m, Ar), 4.11 (m, CH₂O), 4.02 (m, CH₂N (dmba)), 3.52 (s, CH₃O), 3.38 (s, CH₂N), 3.15 (m, br, NH) and 2.83 (m, CH₃N).

Complex 25. As for complex **23** using AgBF₄ (0.030 g, 0.16 mmol) and **22** (0.104 g, 0.16 mmol). ¹H NMR (CDCl₃): δ 7.70 (m, Ar), 7.10 (m, Ar), 6.50 (m, Ar), 6.40 (m, Ar), 6.00 (NH), 4.17 (s, CH₃O), 4.09 (CH₂) and 3.01 (CH₃N).

Reaction of complex 24 with CO giving [Pt(dmba)(L¹)(CO)]·BF₄ **26.** Carbon monoxide was bubbled through a solution of complex **24** formed *in situ* from AgBF₄ (0.052 g, 0.27 mmol) and **20** (0.100 g, 0.16 mmol). Recrystallisation from dichloromethane–diethyl ether gave colourless crystals of **26** (Found: C, 42.6; H, 4.44; N, 3.95. C₂₅H₃₀BF₄N₂O₂Pt· $\frac{3}{4}$ CH₂Cl₂ requires C, 42.3; H, 4.51; N, 3.80%). ¹H NMR (CD₂Cl₂): δ 7.74 (m, Ar), 7.54 (m, Ar), 7.25 (m, Ar), 7.17 (m, Ar), 7.10 (m, Ar), 6.85 (m, Ar), 4.35 (m, CH₂N (dmba)), 3.57 (m, br, NH), 3.48 (t, CH₂O), 3.26 (m, CH₃O/CH₃N) and 3.20 (m, CH₂N).

Complex 27. As for complex **26** using AgBF₄ (0.052 g, 0.27 mmol) and **22** (0.100 g, 0.16 mmol). ¹H NMR (CD₂Cl₂): δ 7.86 (m, Ar), 7.57 (m, Ar), 7.31 (m, Ar), 7.18 (m, Ar), 6.88 (m, Ar), 6.73 (m, Ar), 5.90 (m, NH), 4.33 (m, CH₂), 3.76 (s, CH₃O) and 3.25 (m CH₃N).

Reaction of complex 25 with XylNC giving [Pt(dmba)(L⁴)(CNXyl)]BF₄ **28.** The compound CNXyl (0.050 g, 0.38 mmol) was added to a solution of complex **25**, formed *in situ* from AgBF₄ (0.015 g, 0.08 mmol) and **22** (0.050 g, 0.08 mmol). ¹H NMR (CDCl₃): δ 7.86 (m, Ar), 7.45 (m, Ar), 7.17 (m, Ar), 6.82 (m, Ar), 5.85 (m, NH), 4.37 (d, CH₂, ⁴*J*(PH) 3), 3.66 (s, CH₃O), 3.24 (d, CH₃N, ⁴*J*(PH) 3) and 2.19 (s, CH₃C).

Reaction of complex 25 with acetonitrile giving [Pt(dmba)(L⁴)(NCCH₃)]BF₄ **29.** An excess of acetonitrile (1 cm³) was added to a solution of complex **25**, formed *in situ* from AgBF₄ (0.015 g, 0.08 mmol) and **22** (0.050 g, 0.08 mmol). ¹H NMR (CDCl₃): δ 7.76 (m, Ar), 7.49 (m, Ar), 7.10 (d, Ar), 7.01 (m, Ar), 6.89 (m, Ar), 6.73 (m, Ar), 5.74 (m, NH), 4.07 (m, CH₂), 3.77 (s, CH₃O), 2.98 (m, CH₃N) and 1.83 (s, CH₃CN).

Crystallography

Data for complexes **2**, **14**, **16** and **30** were collected on a CAD4 automatic 4-circle diffractometer, and for **1** and **18** on the EPSRC FAST system and a Bruker SMART 1000 CCD diffractometer, respectively. Hydrogen atoms were included in calculated positions on carbon centres for all structures. Hydrogens attached to nitrogen atoms were readily located throughout in the penultimate Fourier difference electron density maps, and included in the final stage of refinement with various restraints. In **1** the hydrogen attached to N(1) was refined at a fixed distance of 0.90 Å from the parent atom. The corresponding hydrogens in compounds **2**, **14** and **18** were treated similarly. In **16**, H(1A) and H(2A) were refined at a fixed distance of 0.98 Å from N(1) and N(2) respectively. In the structural refinement of complex **14** the maximum residual electron density peak was noted to be very close to the nitrite moiety, which points to possible disorder and librational effects in this region of the electron density map, given the large thermal vibration of the atoms therein. This disorder could not be successfully modelled and in the latter stages of refinement the N–O distances in the nitrite group were constrained to be similar to each other.

The crystal used for the structural determination of complex **30** was not a strong diffractor. This is exemplified by the statistics on the data and the larger than desirable ESDs for the geometric parameters associated with this crystal structure. In the final stages of the refinement the components of the anisotropic displacement parameters for atoms in the phenyl ring containing C(7) were refined subject to 'rigid bond' restraints. In addition the thermal parameters for C(5) and C(6) were treated similarly using SHELXL.³⁴

An empirical absorption correction (DIFABS³⁵) was applied to the data for compounds **14** and **30**. The associated maximum and minimum transmission factors for **14** were 1.000 and 0.860 respectively. The corresponding values for **30** were 1.000 and 0.219.

CCDC reference number 186/1924.

See <http://www.rsc.org/suppdata/dt/b0/b001436m/> for crystallographic files in .cif format.

Acknowledgements

We would like to thank the University of Bath for a studentship (to M. T. P.), Professor M. B. Hursthouse and the EPSRC National Crystallography Service for providing the data set for compound **1**, and Dr Michael Ruf of Bruker AXS GmbH Analytical X-ray Systems for collecting data on compound **18**.

References

- 1 L. A. Hamilton and P. S. Landis, in *Organic Phosphorus Compounds*, eds. G. M. Kosolapoff and L. Maier, Wiley-Interscience, New York, 1972, vol. 4, p. 504.
- 2 D. Thompson (Editor), *Insights into Speciality Inorganic Chemicals*, RSC, Cambridge, 1995.
- 3 K. Osakada, T. Ikariya, M. Saburi and S. Yoshikawa, *Chem. Lett.*, 1981, 1691.
- 4 S. Naili, J.-F. Carpentier, F. Agbossou and A. Mortreux, *Organometallics*, 1995, **14**, 401.
- 5 I. Suisse, H. Bricout and A. Mortreux, *Tetrahedron Lett.*, 1994, **35**, 413.
- 6 T. Q. Ly, A. M. Z. Slawin and J. D. Woollins, *J. Chem. Soc., Dalton Trans.*, 1997, 1611.
- 7 D. J. Birdsall, J. Green, T. Q. Ly, J. Novosaf, M. Necas, A. M. Z. Slawin and J. D. Woollins, *Eur. J. Inorg. Chem.*, 1999, 1445.
- 8 K. G. Gaw, A. M. Z. Slawin and M. B. Smith, *Organometallics*, 1999, **18**, 3255.
- 9 A. Bader and E. Lindner, *Coord. Chem. Rev.*, 1991, **108**, 27.
- 10 E. Lindner and E. Glaser, *J. Organomet. Chem.*, 1990, **391**, C37.
- 11 J. McKechnie, D. S. Payne and W. Sim, *J. Chem. Soc.*, 1965, 3500.
- 12 P. Bhattacharyya, A. M. Z. Slawin, M. B. Smith and J. D. Woollins, *Inorg. Chem.*, 1996, **35**, 3675.
- 13 R. A. Burrow, D. H. Farrar and C. H. Honeyman, *Acta Crystallogr., Sect. C*, 1994, **50**, 681.
- 14 D. A. Fletcher, R. F. McMeeking and D. Parkin, *J. Chem. Inf. Comput. Sci.*, 1996, **36**, 746; F. H. Allen and O. Kennard, *Chem. Des. Autom. News*, 1993, **8**, 1; 31.
- 15 A. M. Z. Slawin, M. B. Smith and D. J. Woollins, *J. Chem. Soc., Dalton Trans.*, 1996, 1283.
- 16 A. M. Z. Slawin, M. B. Smith and D. J. Woollins, *J. Chem. Soc., Dalton Trans.*, 1996, 4567.
- 17 A. Badia, L. R. Falvello, R. Navarro and E. P. Urriolabeitia, *J. Organomet. Chem.*, 1997, **547**, 121.
- 18 R. P. Kamalesh Babu, S. S. Krishnamurthy and M. Nethaji, *Polyhedron*, 1996, **15**, 2689.
- 19 A. Fischer, I. Neda, P. G. Jones and R. Schmutzler, *Z. Naturforsch., Teil B*, 1994, **49**, 1481.
- 20 M. Paul and H. Schmidbaur, *Chem. Ber.*, 1996, **129**, 77.
- 21 G. A. Jeffrey, *An Introduction to Hydrogen Bonding*, O.U.P., New York, 1997 and references therein.
- 22 G. A. Jeffrey, H. Maluszynska and J. Mitra, *Int. J. Biol. Macromol.*, 1985, **7**, 336.
- 23 E. Lindner, J. Dettinger, H. A. Mayer, H. Kuhbauch, R. Fawzi and M. Steimann, *Chem. Ber.*, 1993, **126**, 1317.
- 24 P. Braunstein, D. Matt, Y. Dusaosoy, J. Fischer, A. Mitschler and L. Ricard, *J. Am. Chem. Soc.*, 1981, **103**, 5115.
- 25 A. D. Ryabov, L. G. Kuz'mina, V. A. Polyakov, G. M. Kazankov, E. S. Ryabova, M. Pfeffer and R. van Eldik, *J. Chem. Soc., Dalton Trans.*, 1995, 999.
- 26 G. K. Anderson and R. Kumar, *Inorg. Chem.*, 1984, **23**, 4064.
- 27 A. D. Burrows and M. T. Palmer, unpublished results.
- 28 D. E. Berry, K. A. Beveridge, J. Browning, G. W. Bushnell and K. R. Dixon, *Can. J. Chem.*, 1986, **64**, 1903.
- 29 D. Matt, F. Ingold, F. Balegroune and D. Grandjean, *J. Organomet. Chem.*, 1990, **399**, 349.
- 30 N. W. Alcock, P. Bergamini, T. M. Gomes-Carniero, R. D. Jackson, J. Nicholls, A. G. Orpen, P. G. Pringle, S. Sostero and O. Traverso, *J. Chem. Soc., Chem. Commun.*, 1990, 980.
- 31 D. Drew and J. R. Doyle, *Inorg. Synth.*, 1972, **13**, 48.
- 32 J. Chatt, L. M. Vallarino and L. M. Venanzi, *J. Chem. Soc.*, 1957, 3413.
- 33 A. C. Cope and E. C. Friedrich, *J. Am. Chem. Soc.*, 1968, **90**, 909.
- 34 G. M. Sheldrick, SHELXL 93, a computer program for crystal structure refinement, University of Göttingen, 1993.
- 35 N. Walker and D. Stuart, *Acta Crystallogr., Sect. A*, 1983, **39**, 158.