

Ferrocenium salt induced ring expansion of a 2*H*-azaphosphirene complex: Synthesis of the first Δ^3 -1,4,2-oxazaphospholene complexes†

Rainer Streubel,* Christoph Neumann and Peter G. Jones


Institut für Anorganische und Analytische Chemie der Technischen Universität Braunschweig, Postfach 3329, 38023-Braunschweig, Germany. E-mail: r.streubel@tu-bs.de

Received 15th June 2000, Accepted 19th June 2000

Published on the Web 4th July 2000

Regioselective ring expansion of the 2*H*-azaphosphirene tungsten complex **1** in CH_2Cl_2 is performed using substoichiometric amounts of ferrocenium hexafluorophosphate and benzaldehyde or cyclohexanone, thus yielding Δ^3 -1,4,2-oxazaphospholene complexes **2a,c**; the molecular structure of complex **2c** was established by a single crystal X-ray diffraction study.

There is considerable current interest in the development of sophisticated ligands including either sp^2 - and/or sp^3 -hybridised phosphorus atoms in heterocyclic rings.¹ In the last few years, we have focussed our research on the synthesis of so-called biomimetic P-heterocycles of the diazole- and triazole-type, such as 2*H*-1,2-azaphospholes,² 2*H*-1,3,2-diazaphospholes³ and 2*H*-1,4,2-diazaphospholes.⁴ These were obtained as transition metal complexes using (i) selective P–C cleaving ring expansions of the 2*H*-azaphosphirene ring system as a new synthetic methodology (Scheme 1). Five-membered heterocycles of




Scheme 1 Ring expansion reactions of 2*H*-azaphosphirene complexes ($a=b$ denotes a π -system).

these or related types are especially interesting with regard to the intrinsic electronic ambiguity of a tri-coordinate phosphorus centre, which depends significantly on the degree of pyramidalisation.⁵ Recently, the first striking example of a phosphole-type ring system was published, 1-[bis(trimethylsilyl)methyl]-3,5-bis(trimethylsilyl)-1,2,4-triphosphole,⁶ bearing a phosphorus centre that is tri-coordinated and planar.

During our current study of oxidation and reduction reactions of 2*H*-azaphosphirene metal complexes, we have observed that such complexes undergo selective P–N cleaving ring expansion reactions (ii) at ambient temperature with various π -systems, when ferrocenium hexafluorophosphate (Fc^+PF_6^-) is present (Scheme 1). We now report the synthesis of the first Δ^3 -1,4,2-oxazaphospholene complexes under very mild conditions using Fc^+PF_6^- , benzaldehyde and cyclohexanone and a 2*H*-azaphosphirene tungsten complex.

The reaction of 2*H*-azaphosphirene tungsten complex **1** with 2 equivalents of either benzaldehyde or cyclohexanone and 0.2 equivalents of Fc^+PF_6^- in dichloromethane at ambient temperature afforded regioselectively the Δ^3 -1,4,2-oxazaphospholene complexes **2a** (together with **2b**, most probably a diastereoisomer) and **2c** (Scheme 2) in reasonable yields (45 and 44%, respectively); ferrocene was isolated in both reactions in



Scheme 2 Reaction of complex **1** with Fc^+PF_6^- , benzaldehyde and cyclohexanone. *Reagents and conditions:* to a solution of 1 mmol of complex **1**, 2 mmol of benzaldehyde or cyclohexanone in 3 ml of dichloromethane was added 0.2 mmol Fc^+PF_6^- and stirred at ambient temperature until **1** was consumed (^{31}P NMR control); work-up by column chromatography (SiO_2) at low temperature and crystallization from *n*-pentane yielded complexes **2a,c** as pale yellow solids (**2a**: 45%, mp 92 °C, decomp.; **2c**: 44%, mp 140 °C, decomp.).

ca. 10% yield. If one equivalent of the ferrocenium salt was added the reaction time decreased without affecting the selectivity of the reaction. In the case of benzaldehyde, ^{31}P NMR spectroscopic monitoring showed the formation of two reactive intermediates (<5–10% of all phosphorus-containing products) having resonances at δ 108 and 119. Because of the need for Fc^+PF_6^- in these reactions or other oxidizing agents such as Cu(II) salts or elemental sulfur, in the absence of oxidizing agents no reactions occurred at ambient temperature, we propose that the ring expansion proceeds *via* electron transfer catalysis,⁹ which, to the best of our knowledge, would be unprecedented in the chemistry of phosphorus-containing heterocycles.¹⁰ At higher temperature the reactions yielded selectively 2*H*-1,3,2-oxazaphospholene complexes *via* insertion of the C–O π -system into the P–C bond.^{3b}

The complexes **2a,c** were isolated by low-temperature chromatography and crystallisation; the constitutions of the complexes **2a,c** are unambiguously established by their NMR spectroscopic data‡ (^{13}C , ^{31}P) in solution and their MS data.§ Furthermore, the ring constitution of complex **2c** was confirmed by X-ray structure analysis (Fig. 1).§

Complexes **2a,c** show resonances for the imino carbon atoms at δ 173.6 and 169.7 with coupling constant magnitudes $|J(^{31}\text{P},^{13}\text{C})|$ of ca. 13–16 Hz, and for the sp^3 -hybridized carbon atoms at δ 108.1 and 113.1 with coupling constant magnitudes $|J(^{31}\text{P},^{13}\text{C})|$ of ca. 4–7 Hz. ^{31}P resonances are observed in a small range [δ 135.3 (**2a**), 141.5 (**2b**) and 136.4 (**2c**)] with characteristic coupling constants $|J(^{183}\text{W},^{31}\text{P})|$ of ca. 270–280 Hz. The molecular structure of complex **2c** (Fig. 1) confirms the constitution of the heterocyclic ring system, which is almost planar (mean

† Dedicated to Professor Herbert W. Roesky on the occasion of his 65th birthday.




Fig. 1 Molecular structure of complex **2c** (ellipsoids represent 30% probability levels; solvent and hydrogen atoms are omitted for clarity). Selected bond lengths (Å) and angles (°): W–C1 2.011(3), W–P 2.5209(6), P–C8 1.821(2), P–O6 1.6309(17), P–C6 1.877(2), N–C6 1.273(3), N–C7 1.454(3), C7–O6 1.465(3); C8–P–W 117.45(8), O6–P–C6 88.92(9), P–C6–N 111.51(17), C6–N–C7 115.21(9), N–C7–O6 108.02(18).

deviation 0.042 Å) and shows, in comparison to {[2-bis(trimethylsilyl)methyl-5-phenylbenz[*c*]-1,2-oxaphospholane]penta-carbonyltungsten(0)}¹¹ **3** and 1,1-diphenyl-3,3,5,5,8,8-hexakis-(trifluoromethyl)-2,4,9-trioxa-7-aza-1λ⁵-phosphabicyclo[4.3.0]-non-6-ene¹² **4**, C–O and P–O bond distances [C7–O6 1.465(3) and P–O6 1.6309(17) Å], which are very similar to those in **3** [C–O 1.463(4) and P–O 1.641(2)¹¹ Å], but differ significantly from those in **4** [C–O 1.374(3) and P–O 1.732(2)¹² Å]; the C–N double bond distances of **2c** and **4** are also slightly different [**2c**: N–C6 1.273(3) and **4**: N–C 1.291(4)¹² Å].

Currently, electrochemical and ESR studies to elucidate the reaction mechanism are in progress, furthermore, we are exploiting this new synthetic methodology by employing nitriles instead of carbonyl derivatives.¹³

Acknowledgements

We are grateful to the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie for financial support and Mr Andreas Weinkauff for the crystallographic measurement.

Notes and references

‡ Satisfactory elemental analyses were obtained for complexes **2a,c**. NMR data were recorded in CDCl₃ solutions at 50.3 MHz (¹³C) and 81.0 MHz (³¹P), using SiMe₄ and 85% H₃PO₄ as standard references; *J*/Hz. Selected spectroscopic data for **2a** (≅8:1 mixture of **2a,b**): ¹³C NMR: δ 2.9 [d, ³*J*_{PC} 1.9, Si(CH₃)₃], 3.3 [d, ³*J*_{PC} 2.8, Si(CH₃)₃], 33.3 (d, ¹*J*_{PC} 16.7, CH), 108.1 (d, ²*J*_{PC} 6.3 POC), 126.5 (s, Ph), 128.6 (s, Ph),

130.6 (s, Ph), 130.7 (s, Ph), 132.7 (d, *J*_{PC} 25.2, Ph), 137.8 (d, *J*_{PC} 3.5, Ph) 173.6 (d, *J*_{PC} 13.3, PCN), 197.1 (d, ³*J*_{PC} 7.4, *cis*-CO), 198.9 (d, ³*J*_{PC} 28.2, *trans*-CO); ³¹P NMR: δ 135.3 (s, ¹*J*_{PW} 280.2); *m/z* (EI) 723 (M⁺, 28). **2c**: ¹³C NMR: δ 1.9 [d, ³*J*_{PC} 1.6, Si(CH₃)₃], 2.2 [d, ³*J*_{PC} 2.6, Si(CH₃)₃], 21.8 (s, CH₂), 22.0 (s, CH₂), 23.9 (s, CH₂), 35.4 (d, ¹*J*_{PC} 13.3, CH), 36.1 (s, CH₂), 36.3 (s, CH₂), 113.1 (d, ²*J*_{PC} 4.7 POC), 127.5 (s, Ph), 129.3 (s, Ph), 130.2 (s, Ph), 132.7 (d, *J*_{PC} 25.2, Ph), 169.7 (d, *J*_{PC} 15.4, PCN), 198.2 (d, ³*J*_{PC} 7.2, *cis*-CO), 199.4 (d, ³*J*_{PC} 26.3, *trans*-CO); ³¹P NMR: δ 136.4 (s, ¹*J*_{PW} 273.4); *m/z* (EI) 715 (M⁺, 8).

§ Crystal structure analysis of **2c**·0.5C₃H₁₂: empirical formula C_{27.5}H₄₀NO₆PSi₂W, *M* = 751.6; monoclinic, space group C2/c; *a* = 36.058(4), *b* = 10.4819(12), *c* = 19.602(2) Å, β = 119.330(6)°, *V* = 6458.9(13) Å³, *Z* = 8, *D*_c = 1.546 Mg m^{−3}; λ = 0.71073 Å, *T* = 143(2) K. The crystal (0.27 × 0.24 × 0.17 mm) was mounted in inert oil. 38692 intensities were measured (ω and θ scans, 2θ 3–60°) using using Mo-Kα radiation on a Bruker SMART 1000 CCD diffractometer. After absorption correction (multiple scans) 9462 were unique (*R*_{int} = 0.0454) and used for all calculations (SHELXL-97¹⁴). All hydrogen atoms were refined with a riding model or as rigid methyl groups. Final *wR*(*F*²) was 0.0589, with conventional *R*(*F*) 0.0247, for 186 parameters and 351 restraints; highest peak 1.479, hole −0.861 e Å^{−3}.

CCDC reference number 186/2042.

See <http://www.rsc.org/suppdata/dt/b0/b004795n/> for crystallographic files in .cif format.

- 1 K. B. Dillon, F. Mathey and J. F. Nixon, *Phosphorus: The Carbon Copy*, Wiley, Chichester, 1998, p. 203.
- 2 R. Streubel, H. Wilkens, A. Ostrowski, C. Neumann, F. Ruthe and P. G. Jones, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1492; U. Rohde, F. Ruthe, P. G. Jones and R. Streubel, *Angew. Chem., Int. Ed.*, 1999, **38**, 215; H. Wilkens, A. Ostrowski, J. Jeske, F. Ruthe, P. G. Jones and R. Streubel, *Organometallics*, 1999, **18**, 5627.
- 3 (a) H. Wilkens, J. Jeske, P. G. Jones and R. Streubel, *Chem. Commun.*, 1997, 2317; (b) H. Wilkens, F. Ruthe, P. G. Jones and R. Streubel, *Chem. Eur. J.*, 1998, **4**, 1542.
- 4 R. Streubel, H. Wilkens, F. Ruthe and P. G. Jones, *Chem. Commun.*, 1999, 2127.
- 5 For a more recent work on aromaticity in phospholes and polyphosphaphospholes, see: A. Dransfeld, L. Nyulászi and P. v. R. Schleyer, *Inorg. Chem.*, 1998, **37**, 4413.
- 6 F. G. N. Cloke, P. B. Hitchcock, P. Hunnabell, J. F. Nixon, L. Nyulászi, E. Niecke and V. Thelen, *Angew. Chem., Int. Ed.*, 1998, **37**, 1083.
- 7 R. Streubel, U. Rohde, J. Jeske, F. Ruthe and P. G. Jones, *Eur. J. Inorg. Chem.*, 1998, 2005.
- 8 N. G. Connelly and W. E. Geiger, *Chem. Rev.*, 1996, **96**, 877.
- 9 Reviews: H. Taube, *J. Chem. Ed.*, 1968, **45**, 452; M. Chanon, *Acc. Chem. Res.*, 1987, **20**, 214; R. A. Marcus, *Angew. Chem., Int. Ed. Engl.*, 1993, **32**, 1111; D. Astruc, *Electron-Transfer Processes in Transition Metal Chemistry*, VCH Publishers, New York, 1995.
- 10 For ET-catalyzed ring expansions of oxiranes and aziridines in organic synthesis, see: L. Ebersson, *Electron Transfer Reactions in Organic Chemistry*, Springer Verlag, Berlin, 1987.
- 11 R. Streubel, A. Ostrowski, H. Wilkens, F. Ruthe, J. Jeske and P. G. Jones, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 378.
- 12 H. W. Roesky, V. W. Pogatzki, K. S. Dhathathreyan, A. Thiel, H. G. Schmidt, M. Dyrbusch, M. Noltemeyer and G. M. Sheldrick, *Chem. Ber.*, 1986, **119**, 2687.
- 13 R. Streubel, H. Wilkens, C. Neumann, F. Ruthe and P. G. Jones, unpublished work.
- 14 G. M. Sheldrick, SHELXL-97, program for crystal structure refinement, Universität Göttingen, 1997.