


Synthesis of Daunorubicin Analogues with Novel 9-Acyl Substituents¹

Thomas H. Smith,* Allan N. Fujiwara, and David W. Henry

Bio-Organic Chemistry Department, SRI International, Menlo Park, California 94025. Received July 20, 1978

Synthetic approaches to anthracyclines bearing novel 9-acyl substituents were investigated. Reaction of the lithium enolate of *N*-(trifluoroacetyl)daunorubicin (9) with methyl iodide in tetrahydrofuran afforded only the 9-propionyl derivative 10 in high yield. Reaction of 10 under identical conditions cleanly afforded the 9-isobutyryl derivative 11. Extension of this procedure to other alkylating agents (ethyl iodide, benzyl bromide, and heptyl iodide) required hexamethylphosphoramide as cosolvent and afforded mixtures of mono- and dialkylated products as well as recovered 9. The amino group was deblocked with NaOH in aqueous tetrahydrofuran, except in the case of the dibenzyl derivative 13 which was inert under these conditions. The 9-formyl analogue 23 was prepared via NaIO₄ cleavage of 13-dihydroadriamycin (21). Antitumor evaluation against P388 leukemia in mice showed 23 to have activity comparable to the parent compounds, while the C-alkylated analogues were less active.

The anthracycline antibiotics daunorubicin² (1) and




adriamycin³ (2) are clinically useful antineoplastic agents, with adriamycin having an especially broad spectrum of activity that includes various solid tumors that are normally resistant to most modes of chemotherapy. However, clinical use of these drugs is hampered by a number of undesirable side effects, the most serious being a dose-related and irreversible cardiotoxicity.⁴ As part of our efforts to prepare analogues of 1 and 2 having improved therapeutic properties, we now report the synthesis and biological evaluation of a number of anthracyclines bearing novel acyl groups at C-9.

The apparent clinical superiority of adriamycin over daunorubicin, from which it differs only by the presence of a hydroxyl group at C-14, led us to investigate the side chain as a site for structural modification. Recent reports indicate that 1 and 2 can tolerate substantial modification and even removal of the 9-acyl group while retaining antitumor activity. These prior modifications involved either degradation of the side chain,⁵ modifying the oxidation level or derivatization of the 13-carbonyl,⁶ or the introduction of a novel heterosubstituent at C-14.⁷ Our present approach to novel 9-acylanthracyclines involved the generation of new carbon-carbon bonds at C-14, except for the 9-formyl derivative 23 obtained by a degradative sequence.

Chemistry. The methodology developed in our laboratory for the synthesis of [14-¹³C]daunorubicin and adriamycin⁸ seemed appropriate as a general synthesis of 9-acylanthracyclines. In this approach, adriamycinone (3) is reacted with the appropriate Grignard reagent, and the resulting glycol 4 is cleaved with NaIO₄ to afford the 13-ketone. While this sequence afforded daunomycinone (5) in a 56% chemical yield from 3, it failed in several cases when R ≠ Me. A 10% yield of 6 from 3 was the next best result obtained by this method, with Grignard addition to the quinone carbonyls appearing to be a serious side reaction.


The ready availability of 14-bromodaunomycinone (7)¹²



led us to evaluate it as a 9-acylanthracycline synthon. Reaction of 7 with lithiodimethyl cuprate afforded only a low yield of the desired 9-propionyl aglycon 8, with daunomycinone being the major product. The formation of 5 can be attributed to halogen-metal exchange forming the enolate, which decomposes to 5 on aqueous workup. Treatment of the reaction mixture with excess methyl iodide prior to workup⁹ increased the yield of 8 to 49%. Moreover, the absence of any O-methylated products, despite the presence of four hydroxyl groups, indicated that direct alkylation of the 14-enolate may be a feasible route to C-alkylated aglycons such as 8.

Indeed, reaction of the enolate of 5 (generated with 12 equiv of lithium diisopropylamide (LDA) in THF at -78 °C) with excess methyl iodide afforded 8 in 76% yield. In light of the demonstrated susceptibility of the phenolic hydroxyls to O-methylation (dimethyl sulfate, K₂CO₃),¹⁰ the chemospecificity observed in the C-alkylation procedure is impressive. The absence of O-methylation is probably due to the greater association of the lithium cation to the oxygen atoms, as compared with the larger potassium ion, thus enhancing the extent of C-alkylation.¹¹

Pilot experiments showed the glycoside bond to be stable to the C-alkylation conditions, and extension of this procedure to the intact glycosides went smoothly, thus obviating the need for the relatively laborious aglycon-sugar condensation. The amino group of 1 was protected via conversion to the trifluoroacetamide 9¹² with *S*-ethyl



- 9, R = CH₃; R' = COCF₃
 10, R = CH₂CH₃; R' = COCF₃
 11, R = CH(CH₃)₂; R' = COCF₃
 12, R = CH₂CH₂Ph; R' = COCF₃
 13, R = CH(CH₂Ph)₂; R' = COCF₃
 14, R = (CH₂)₂CH₃; R' = COCF₃
 15, R = CH(CH₂CH₃)₂; R' = COCF₃
 16, R = (CH₂)₇CH₃; R' = COCF₃
 17, R = CH(*n*-C₈H₁₇)₂; R' = COCF₃
 18, R = C₂H₅; R' = H
 19, R = CH(CH₃)₂; R' = H
 20, R = CH₂CH₂Ph; R' = H

trifluoroacetate and sodium methoxide. Generation of the enolate of **9** with LDA in THF followed by excess methyl iodide afforded the 9-propionyl glycoside **10** in 77% yield. Although **9** could not be dimethylated directly, **10** was smoothly alkylated under the above conditions to afford **11** in high yield.


Table I summarizes the results of our C-alkylation studies. Extension of the scope of the glycoside C-alkylation to permit the use of reagents other than methyl iodide required hexamethylphosphoramide (HMPA) as cosolvent. In the absence of HMPA, ethyl iodide was inert, and only low yields of alkylated products could be obtained with benzyl bromide and heptyl iodide. Although monoalkylation predominated, significant amounts of dialkylated products and unreacted **9** were obtained.


The amino group was deprotected with NaOH in aqueous THF at 0 °C to afford the amino glycosides **18–20**, which were isolated as the HCl salts. However, this procedure was unsuccessful with the dibenzyl derivative **13**, which was recovered unchanged after prolonged reaction under the above and more rigorous conditions. An examination of a Drieding model of **13** offers a possible explanation of this result. The second phenyl group can afford appreciable steric hindrance to the approach of base to the trifluoroacetamide moiety. The side chain of **12** has no effect on the reactivity of the trifluoroacetamido moiety, since it can readily assume a conformation minimizing such interaction. However, in **13** one of the phenyl groups is forced into close proximity to the ring A and sugar portions of the molecule.


The 9-formyl analogue **23** was obtained in a manner conceptually similar to the Grignard approach (vide supra). Adriamycin (**2**) was reduced with NaCNBH₃, and the resulting alcohol **21**¹³ was treated with sodium metaperiodate to afford **23**, which was isolated as the hydrate. Since **23** showed both a weak aldehyde proton signal in its ¹H NMR and weak aldehyde carbonyl absorbance in its IR spectra, additional confirmation of its structure was sought. Acid hydrolysis of **23** afforded an aglycon (**24**) identical (TLC, IR) with that obtained from periodate cleavage of 13-dihydroadriamycinone (**22**). Further characterization of **24** was obtained by its conversion to the 2,4-dinitrophenylhydrazone **25**.


Results and Discussion

Table II presents a comparison of test data for the parent antibiotics and the 9-acylanthracyclines prepared



- 21, R = 
 22, R = H



- 23, R = ; R' = O
 24, R = H; R' = O
 25, R = H; R' = NNHC₆H₃-*o*, *p*-(NO₂)₂

in this study. All of the compounds bind to DNA to a significant degree, as indicated by their ability to stabilize helical DNA to thermal denaturation (ΔT_m). The ΔT_m value is the difference in degree Celsius between the melting temperature of calf thymus DNA at pH 7 and the same DNA in the presence of drug at a drug-DNA phosphate ratio of 1:10. Each new analogue also displays potent inhibition of DNA and RNA syntheses in cultured L1210 cells. The most complete set of in vivo antitumor data is in the advanced P388 test (QD5, **9**, **13**). In this test, daunorubicin (**1**) effects only a very modest increase in survival time. Along with daunorubicin, all C-14 alkylated analogues (**18–20**) show weak but significant antitumor activity in this test, branched analogue **19** being the most borderline case. Compound **19** was inactive in the P388 QD1-9 test according to NCI criteria. Neither **18** nor **20** was tested in this latter system. Formyl derivative **23** showed efficacy intermediate between those of daunorubicin and adriamycin in both in vivo tests.

The available data indicate that acyl groups on the 9 position larger than acetyl do not interfere substantially with DNA binding or with nucleic acid synthesis inhibition in cultured cells. Similarly, antitumor efficacy in vivo is not grossly affected by the larger and increasingly lipophilic acyl moieties of compounds **18–20** in the advanced P388 test when compared with daunorubicin. Potency in this three-compound group is reduced substantially, however, and roughly according to the lipophilicity of the side chain. These data taken together suggest that distribution of the drugs is being adversely affected by these structural modifications. 9-Formyl analogue **23** is the only member of the group with a small acyl group and should be less lipophilic than daunorubicin. It provides equal or somewhat enhanced efficacy relative to daunorubicin but at much larger doses, thus not falling into the lipophilicity-potency correlation seen with the other analogues. The existence of **23** as the *gem*-diol hydrate, with tetrahedral stereochemistry at C-13, might be expected to give it unique status in the group. This is also reflected by its relatively poor ranking in ΔT_m and nucleic acid synthesis inhibition.

Although none of the new anthracycline compounds prepared in this study appear especially interesting as clinical antitumor drug candidates, the new chemistry involved in their synthesis provides a potentially useful

Table I. Summary of Enolate C-Alkylation

substrate	alkylating agent	solvent	products ^a (% yield)		
			monoalkyl	dialkyl	recov substr
5	CH ₃ I	THF	8 (76)		
9	CH ₃ I	THF	10 (77)		
10	CH ₃ I	THF	11 (82)		
9	C ₆ H ₅ CH ₂ Br	2:1 THF-HMPA	12 (36)	13 (27)	9 (16)
9	CH ₃ CH ₂ I	2:1 THF-HMPA	14 (31)	15 (11)*	9 (5)
9	CH ₃ (CH ₂) ₆ I	2:1 THF-HMPA	16 (43)	17 (12)*	9 (25)

^a Yields refer to isolated products, homogenous by TLC and having expected IR and NMR spectra. Acceptable elemental analyses were obtained for all new compounds except where, as noted by an asterisk, sufficient material was not available.

Table II. Comparison of Biological Data for Daunorubicin, Adriamycin, and 9-Acyl Analogues Prepared in This Study

compd	ΔT_m , ^a °C	nucleic acid synth inhibn, ^a		antitumor activity in mice ^b			
		ED ₅₀ , μ M		P388, qd 5, 9, 13		P388, qd 1-9	
		DNA	RNA	opt dose, mg/kg	T/C ^c (<i>n</i>) ^d	opt dose, mg/kg	T/C (<i>n</i>)
1	11.2	0.66	0.33	8	132 · 4 (29)	0.78	160 · 27 (8)
2	13.4	1.5	0.58	8	159 · 20 (29)	0.78	197 · 26 (8)
18	11.0	0.67	0.29	12.5	128 · 8 (2)		
19	10.3	1.0	0.40	37.5	122 · 6 (2)	3.13	117
20	8.6	1.5	1.6	50	129 · 6 (2)		
23	6.5	4.1	2.0	50	154	4	169

^a ΔT_m and ED₅₀ values were determined by the method of G. Tong, W. W. Lee, D. R. Black, and D. W. Henry, *J. Med. Chem.*, **19**, 395 (1976), except that the drugs were initially dissolved in a volume of Me₂SO that resulted in a final Me₂SO concentration of 5 and 1%, respectively, in the assay medium. This modification greatly aided solubilization and did not affect assay results according to extensive control experiments. ^b Assays were arranged through the Drug Research and Development Program, Division of Cancer Treatment, National Cancer Institute. RDF or CDF mice are injected ip with 10⁶ P388 lymphocytic leukemia cells on day 0 and are treated ip on days 1-9 or 5, 9, and 13 with the specified drug dose. For detailed protocols, see R. I. Geran, N. H. Greenberg, M. M. MacDonald, A. M. Schumacher, and B. J. Abbott, *Cancer Chemother. Rep., Part 3*, **3**(2), 9 (1972). ^c Ratio of average survival times of treated mice to untreated controls in percent. The average survival time of untreated controls is approximately 11 days. Activity is defined as values of T/C > 120 in the qd 5, 9, 13 protocol and T/C > 125 in the qd 1-9 protocol. ^d *n* = number of tests.

tool for more complex structural manipulations of the 9-acyl group of daunorubicin.

Experimental Section

Solvent extracts of aqueous solutions were dried over anhydrous Na₂SO₄. Solutions were concentrated under reduced pressure using a rotary evaporator. All melting points are uncorrected. HMPA was stored over 13X molecular sieves, and THF was distilled from LiAlH₄ immediately prior to use. Measurements of 100-MHz NMR were performed by Mr. L. Cary using a Varian XL-100 spectrometer in CDCl₃, unless specified otherwise, with Me₄Si as an internal standard. Elemental microanalyses were provided by the microanalytical laboratory of Stanford University.

Thin-layer chromatograms (TLC) were obtained on silica gel GF 250- μ m plates (Analtech). Preparative layer chromatograms (PLC) were obtained on 20 × 20 × 0.2 cm silica gel 60 F-254 plates (E. Merck). Preparative liquid chromatography was performed on Bio-Sila 200 325 mesh or E. Merck prepacked silica gel 60 columns.

N-(Trifluoroacetyl)daunorubicin (9). Daunorubicin hydrochloride (1.10 g, 1.95 mmol) was dissolved in MeOH (10 mL) and cooled to 5 °C. Sodium methoxide (108.0 mg, 1.95 mmol) was added and the solution was stirred at 5 °C for 5 min. *N*-Ethyl trifluoroacetate (10.80 mL) was added, and the mixture was stirred at 23 °C for 16 h. Additional sodium methoxide (~20 mg) was added and stirring was continued for 24 h (total reaction time 36 h). The solvent was removed and the residue partitioned between CHCl₃ (200 mL) and 0.5 M citric acid (20 mL). The organic phase was separated, washed with saturated NaCl (20 mL), dried, and evaporated. The residue was precipitated from CHCl₃/petroleum ether to afford 1.15 g (94%) of 9.

(7S,9S)-7,8,9,10-Tetrahydro-6,9,11-trihydroxy-4-methoxy-9-propionyl-7[[2,3,6-trideoxy-3-(trifluoroacetamido)- α -L-lyxo-hexopyranosyl]oxy]-5,12-naphthacenedione (10). Methylolithium (24.5 mL of 0.90 M solution in ether, 22.0 mmol) was added to diisopropylamine (6.5 mL) in THF (6.50 mL) at 0 °C under N₂. The solution was cooled to -78 °C. *N*-(trifluoroacetyl)daunorubicin (9, 500 mg, 0.80 mmol) in THF (8.0 mL) was

added dropwise with stirring, and the mixture was maintained at -78 °C for 15 min. Methyl iodide (6.5 mL) was added, and the mixture was allowed to warm to 0 °C over 30 min and kept at 0 °C for 2 h. The reaction mixture was poured into saturated NH₄Cl ice (150 mL) and extracted with CHCl₃ (4 × 40 mL). The extracts were combined, dried, and evaporated. The residue was chromatographed (E. Merck prepacked silica gel 60 column, size B, 98:2 CHCl₃/MeOH) to afford 394.5 mg (77%) of 10: IR 2.85 (OH, NH), 5.75 (sh), 5.80 (C=O), 6.15, 6.30 (H-bonded quinone); NMR δ 1.16 (t, 3, 15-H₂), 1.32 (d, 3, 6'-H₂), 1.85-2.35 (m, 4, 8- and 2'-H₂'s), 2.88 (q, 3, 10 β -H and 14-H₂), 3.28 (d, 1, *J* = 17 Hz, 10 α -H), 3.68 (br s, 1, 4'-H), 4.11 (s, 3, OMe), 4.57 (m, 2, 3'- and 5'-H's), 5.27 (br s, 1, 7-H), 5.54 (br s, 1, 1'-H), 6.65 (d, 1, NH), 7.41 (dd, 1, *J* = 8 and 1 Hz, 2-H), 7.79 (t, 1, *J* = 8 Hz, 2-H), 8.06 (dd, 1, *J* = 8 and 1 Hz, 1-H), 13.30 (s, 1, 11-OH), 14.02 (s, 1, 6-OH); MS *m/e* 637 M⁺ (45), 412 (61), 376 (38), 337 (100), 57 (48); UV visible λ_{max} (MeOH) 233 nm (ϵ 37,665), 252 (26,976), 289 (8901), 480 (11961), 496 (12,216), 531 (6617); [α]_D²⁵ (MeOH) c 0.05; TLC (95:5 CHCl₃/MeOH) *R*_f 0.35. Anal. (C₂₉H₃₀F₃NO₁₁·H₂O) C, H, N.

(7S,9S)-9-Isobutyryl-7,8,9,10-tetrahydro-6,9,11-trihydroxy-4-methoxy-7-[[2,3,6-trideoxy-3-(trifluoroacetamido)- α -L-lyxo-hexopyranosyl]oxy]-5,12-naphthacenedione (11). By the procedure described in the above experiment, 10 (634.0 mg, 0.994 mmol) was converted to 531.3 mg (82%) of 11: IR 2.85 (OH, NH), 5.81 (sh), 5.85 (C=O), 6.15, 6.30 (H-bonded quinone), 9.55, 10.10 μ m; NMR δ 1.22 [d, 6, CH(Me)₂], 1.32 (d, 3, 6'-H₂), 1.95 (m, 2, 2'-H₂), 2.27 (m, 2, 8-H₂), 2.96 (d, 1, *J* = 19 Hz, 10 β -H), 3.29 (d, 1, *J* = 19 Hz, 10 α -H), 3.59 (m, 2, 4'- and 14-H's), 4.12 (s, 3, OMe), 4.29 (m, 2, 3'- and 5'-H's), 5.35 (br s, 1, 7-H), 5.51 (br s, 1, 1'-H), 6.66 (d, 1, NH), 7.40 (dd, 1, *J* = 8 and 1 Hz, 2-H), 7.79 (t, 1, *J* = 8 Hz, 2-H), 8.06 (dd, 1, *J* = 8 and 1 Hz, 1-H), 13.30 (s, 1, 11-OH), 14.03 (s, 1, 6-OH); MS *m/e* 651 M⁺ (16), 426 (27), 390 (13), 337 (100); TLC (95:5 CHCl₃/MeOH) *R*_f 0.41. Anal. (C₃₃H₃₂F₃NO₁₁·0.5H₂O) C, H, N.

Alkylation of the Lithium Enolate of 9 with Benzyl Bromide. Methylolithium (2.0 mL of 0.9 M solution in ether, 2.56 mmol) was added to a stirred solution of diisopropylamine (1.0

mL) in 2:1 THF-HMPA (15 mL) under N_2 . The solution was cooled to -78°C and **9** (50 mg, 0.080 mmol) in 2:1 THF-HMPA (2.0 mL) was added. After 10 min at -78°C , benzyl bromide (0.5 mL) was added, and the mixture was allowed to warm to 0°C over 30 min and kept at 0°C for 1.75 h. The reaction was quenched with saturated NH_4Cl -ice (20 mL) and extracted with ethyl acetate (20 mL). The extract was washed with saturated NaHCO_3 (10 mL), water (4×15 mL), and saturated NaCl (10 mL), dried, and evaporated. The residue was precipitated from CHCl_3 -petroleum ether, collected, and chromatographed (E. Merck prepacked silica gel 60, column-size A-99:1, CHCl_3 -MeOH) to afford 17.5 mg (27%) of **(7S,9S)-9-(2-benzyl-3-phenylpropionyl)-7,8,9,10-tetrahydro-6,9,11-trihydroxy-4-methoxy-7-[[2,3,6-trideoxy-3-(trifluoroacetamido)- α -L-lyxo-hexopyranosyl]oxy]-5,12-naphthacenedione (13)**: IR 2.85 (OH, NH), 5.80, 5.82 (C=O), 6.15, 6.30 (H-bonded quinone), 9.85, 10.10 μm ; NMR δ 1.17 (d, 3, 6'- H_3), 1.84 (m, 4, 8- and 2'- H_2 's), 2.72 (m, 4, 15- and 16- H_2 's), 3.08 (m, 2, 10- H_2), 3.62 (d, 1, 4'-H), 4.09 (s and m, 6, OMe, 14-, 3'-, and 5'-H's), 5.13 (br s, 1, 7-H), 5.43 (br s, 1, 1'-H), 6.66 (d, 1, NH), 7.30 (m, 11, 3-H and Ar-H's), 7.76 (t, 1, $J = 8$ Hz, 2-H), 8.04 (dd, 1, $J = 8$ and 1 Hz, 1-H), 13.18 (s, 1, 11-OH), 14.03 (s, 1, 6-OH); UV-vis λ_{max} (MeOH) 235 nm (ϵ 38778), 253 (29006), 289 (9807), 480 (13381), 498 (13467), 531 (7666); $[\alpha]_D^{245}$ (MeOH) (c 0.05); TLC (95:5 CHCl_3 -MeOH) R_f 0.42. Anal. ($\text{C}_{43}\text{H}_{40}\text{F}_3\text{NO}_{11} \cdot 0.5\text{H}_2\text{O}$) C, H, N.

Continued elution afforded 20.5 mg (36%) of **(7S,9S)-7,8,9,10-tetrahydro-6,9,11-trihydroxy-4-methoxy-9-(3-phenylpropionyl)-7-[[2,3,6-trideoxy-3-(trifluoroacetamido)- α -L-lyxo-hexopyranosyl]oxy]-5,12-naphthacenedione (12)**: IR 2.85 (OH, NH), 5.80, 5.80 (C=O), 6.15, 6.30 (H-bonded quinone), 9.85, 10.10 μm ; NMR δ 1.27 (d, 3, 6'- H_3), 1.87 (m, 2, 2'- H_2), 2.18 (m, 2, 8- H_2), 2.8-3.4 (m, 6, 10-, 14- and 15- H_2 's), 3.66 [d (br s on D_2O exchange), 1, 4'-H], 4.12 (s, 3, OMe), 4.03 (m, 2, 3'-, and 5'-H's), 5.25 (br s, 1, 7-H), 5.51 (br s, 1, 1'-H), 6.63 (d, 1, NH), 7.30 (s, 5, Ar-H's), 7.38 (dd, 1, $J = 8$ and 1 Hz, 3-H), 7.78 (t, 1, $J = 8$ Hz, 2-H), 8.04 (dd, 1, $J = 8$ and 1 Hz, 1-H), 13.29 (s, 1, 11-OH), 14.02 (s, 1, 6-OH); UV-vis λ_{max} (MeOH) 234 nm (ϵ 37035), 252 (27082), 289 (9067), 479 (12478), 495 (12552), 530 (6911); $[\alpha]_D^{216}$ (MeOH) (c 0.05); TLC (95:5 CHCl_3 -MeOH) R_f 0.38. Anal. ($\text{C}_{36}\text{H}_{34}\text{F}_3\text{NO}_{11}$) C, H, N. Further elution afforded 7.5 mg (16%) of **9**.

(7S,9S)-7-[(3-Amino-2,3,6-trideoxy- α -L-lyxo-hexopyranosyl)oxy]-7,8,9,10-tetrahydro-6,9,11-trihydroxy-4-methoxy-9-propionyl-5,12-naphthacenedione Hydrochloride (18). To a solution of **10** (325 mg, 0.510 mmol) in THF (36 mL) at 0°C was added 0.1 N NaOH (36 mL). The solution was kept at 5°C for 16 h, quenched with 0.5 M citric acid (~ 10 mL), neutralized with NaHCO_3 , and extracted with 9:1 CHCl_3 -MeOH (4×40 mL). The extracts were combined, washed with saturated NaCl, dried, and evaporated. The residue was dissolved in MeOH (2 mL) and cooled to 0°C . Cold methanolic HCl (2 mL of 0.276 M solution) was added followed by ether (100 mL). The precipitate was collected to afford 247.2 mg (84%) of **18**: IR 2.90 (OH, NH), 5.81 (C=O), 6.15, 6.30 μm (H-bonded quinone); NMR (free base) δ 1.17 (t, 3, 15- H_3), 1.36 (d, 3, 6'- H_3), 1.73 (m, 2, 8- H_2), 2.25 (m, 2, 8- H_2), 2.75-3.40 (m, 5, 3'-H, 10- and 14- H_2 's), 3.50 (br s, 1, 4'-H), 4.12 (s, 4, OMe and 5'-H), 5.30 (br s, 1, 7-H), 5.52 (br s, 1, 1'-H), 7.39 (dd, 1, $J = 8$ and 1 Hz, 3-H), 7.78 (t, 1, $J = 8$ Hz, 2-H), 8.03 (dd, 1, $J = 8$ and 1 Hz, 1-H); UV-vis λ_{max} (MeOH), 234 nm (ϵ 33864), 252 (23333), 290 (7949), 478 (10513), 495 (10513), 530 (5641); $[\alpha]_D^{234}$ (MeOH) (c 0.06); TLC (30:10:1, CHCl_3 -MeOH- H_2O) R_f 0.36. Anal. ($\text{C}_{28}\text{H}_{33}\text{NO}_{10} \cdot \text{HCl} \cdot 1.7\text{H}_2\text{O}$) C, H, N.

13-Dihydroadriamycin Hydrochloride (21). To a solution of **2** (1.97 g, 3.39 mmol) in water (90 mL) was added NaCNBH_3 (0.64 g, 10.1 mmol) in water (45 mL). The pH was adjusted and maintained at 3.52 with 1 N HOAc while the solution was stirred at 23°C for 6 h. The reaction mixture was basified to pH 9.52 with 1 M NaOH and lyophilized. The residue was dissolved in water (200 mL) and extracted with 6:4 CHCl_3 -MeOH (10×100 mL). The extracts were combined and evaporated. The residue was dissolved in CHCl_3 (200 mL) and washed with H_2O (50 mL). The aqueous phase was reextracted with CHCl_3 (5×50 mL), and the organic solutions were combined, dried, and evaporated. The residue was dissolved in 4:1 CHCl_3 -MeOH (75 mL), filtered, and cooled to $\sim 5^\circ\text{C}$. HCl (8.25 mL of 0.40 M solution in ether, 3.30

mmol) was added and the HCl salt precipitated with ether (175 mL) to afford 1.79 g (91%) of **21**.

(7S,9S)-7-[(3-Amino-2,3,6-trideoxy- α -L-lyxo-hexopyranosyl)oxy]-9-formyl-7,8,9,10-tetrahydro-6,9,11-trihydroxy-4-methoxy-5,12-naphthacenedione Hydrochloride (23). To a solution of **21** (58 mg, 0.095 mmol) in water (3 mL) was added NaIO_4 (26 mg, 0.120 mmol) in water (2 mL). The solution was stirred at 0°C for 5.5 h, basified to pH 9.4 with 1 M NaOH, and extracted with 6:4 CHCl_3 -MeOH (4×25 mL). The extracts were combined and washed with water (35 mL), and the aqueous phase was extracted with CHCl_3 (3×10 mL). The organic solutions were combined, dried, and evaporated. The residue was dissolved in 6:1 CHCl_3 -MeOH (5 mL), filtered, and cooled to $\sim 5^\circ\text{C}$. HCl (4.57 mL of 0.02 M solution in ether, 0.092 mmol) was added followed by ether (40 mL). The precipitate was collected to afford 36 mg (64%) of **23** essentially homogeneously by TLC. An analytical sample was prepared via chromatography (E. Merck silica gel 60 prepacked column, size A, 30:10:1 CHCl_3 -MeOH- H_2O), which afforded 17 mg of analytically pure **23**: IR 3.0 (OH, NH), 5.80 (wk, C=O), 6.20, 6.35 (H-bonded quinone), 7.80, 8.25, 8.95, 10.10 μm ; NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.20 (d, 3, 6'- H_3), 1.5-2.3 (m, 4, 8-, and 2'- H_2 's), 2.7-3.1 (m, 2, 10- H_2), 4.00 (s, 3, OMe), 5.00 (br s, 1, 7-H), 5.30 (br s, 1, 1'-H), 7.5-8.0 (m, 3, 1-, 2-, and 3-H's), 9.6 (s, CHO); TLC (30:10:1 CHCl_3 -MeOH- H_2O) R_f 0.35. Anal. ($\text{C}_{26}\text{H}_{27}\text{NO}_{10} \cdot \text{HCl} \cdot 1.6\text{H}_2\text{O}$) C, H, N.

Acknowledgment. This work was supported by Contract No. NO1-CM-33742 from the Division of Cancer Treatment, National Cancer Institute, Department of Health, Education, and Welfare. We are grateful to Dr. Harry B. Wood, Jr., of the National Cancer Institute for arranging the acquisition of in vivo antitumor test data and to Dorris Taylor, Nancy Charbeneau, and Keith Hohlfeldt of SRI for providing the in vitro test data.

References and Notes

- (1) A preliminary account of portions of this work was presented by T. H. Smith, A. N. Fujiwara, and D. W. Henry at the 2nd Joint Conference CIC/ACS, Montreal, Quebec, May, 1977, abstract MEDI 20.
- (2) (a) F. Arcamone, G. Franceschi, P. Orezzi, G. Cassinelli, W. Barbieri, and R. Mondelli, *J. Am. Chem. Soc.*, **86**, 5334 (1964); (b) J. Bernard, R. Paul, M. Boiron, C. Jacquillat, and R. Maral, Ed., "Rubidomycin", Springer-Verlag, New York, N.Y., 1969.
- (3) (a) F. Arcamone, G. Franceschi, S. Penco, and A. Selva, *Tetrahedron Lett.*, 1007 (1969); (b) R. H. Blum and S. K. Carter, *Ann. Intern. Med.*, **80**, 249 (1974); (c) S. K. Carter, *J. Natl. Cancer Inst.*, **55**, 1265 (1975); (d) D. W. Henry, *ACS Symp. Ser.*, **No. 30**, 15-57 (1976).
- (4) (a) E. A. Lefrak, J. Pitha, S. Rosenheim, and J. A. Gottlieb, *Cancer*, **32**, 302 (1973); (b) L. Lenaz and J. A. Page, *Cancer Treat. Rev.*, **3**, 111 (1976).
- (5) (a) G. Tong, W. W. Lee, D. R. Black, and D. W. Henry, *J. Med. Chem.*, **19**, 395 (1976); (b) S. Penco, F. Angelucci, A. Vigevani, E. Arlandini, and F. Arcamone, *J. Antibiot.*, **30**, 764 (1977).
- (6) (a) K. Yamamoto, E. M. Acton, and D. W. Henry, *J. Med. Chem.*, **15**, 872 (1972); (b) G. Jolles, R. Maral, M. Messer, and G. Ponsinet, *Chemotherapy*, **8**, 237; (c) T. H. Smith, A. N. Fujiwara, and D. W. Henry, *J. Med. Chem.*, **21**, 280 (1978); (d) G. Tong, M. Cory, W. W. Lee, D. W. Henry, and G. Zbinden, *J. Med. Chem.*, **21**, 732 (1978).
- (7) (a) F. Arcamone, G. Franceschi, A. Minghetti, S. Penco, S. Redaelli, A. DiMarco, A. M. Casazza, T. Dasdia, G. DiFronzo, F. Guilianni, L. Lenaz, A. Necco, and C. Soranzo, *J. Med. Chem.*, **17**, 335 (1974); (b) M. Israel, E. J. Modest, and E. Frei III, *Cancer Res.*, **35**, 1365 (1975); (c) F. Arcamone, *Lloydia*, **40**, 45 (1977); (d) R. Seshadri, W. J. Pegg, and M. Israel, 175th National Meeting of the American Chemical Society, Anaheim, Calif., March 1978, abstract no. MEDI 47.
- (8) C. R. Chen, M. T. Fong, A. N. Fujiwara, D. W. Henry, M. A. Leaffer, W. W. Lee, and T. H. Smith, *J. Labelled Compd. Radiopharm.*, **14**, 111 (1978).
- (9) J.-E. DuBois and C. Lion, *Tetrahedron*, **31**, 1227 (1975).

(10) (a) F. Arcamone, G. Franceschi, P. Orezzi, G. Cassinelli, W. Barbieri, and R. Mondelli, *J. Am. Chem. Soc.*, 86, 5334 (1964); (b) D. W. Henry, G. Tong, A. N. Fujiwara, and W. W. Lee, 172nd National Meeting of the American Chemical Society, San Francisco, Calif., 1976, abstract no. MF1D190.

(11) L. M. Jackinan and B. C. Lange, *Tetrahedron*, 33, 2737 (1977).

(12) F. Arcamone, G. Franceschi, and S. Penco, U.S. Patent 3 803 124, 1974.

(13) S. A. Rhône-Poulenc, U.K. Patent 1 322 872, 1973.


Development of Ethyl 3,4-Dihydro-4-oxopyrimido[4,5-*b*]quinoline-2-carboxylate, a New Prototype with Oral Antiallergy Activity¹

T. H. Althuis,* P. F. Moore, and H.-J. Hess

Medicinal Research Laboratories, Central Research, Pfizer Inc., Graton, Connecticut 06340. Received July 17, 1978

Structural modification of 3,4-dihydro-4-oxoquinazoline-2-carboxylic acid leading to ethyl 3,4-dihydro-4-oxopyrimido[4,5-*b*]quinoline-2-carboxylate, a new prototype with oral antiallergy activity of the disodium cromoglycate type, is described. This prototype is 10 times more potent than disodium cromoglycate in the rat passive cutaneous anaphylaxis test. Structure activity studies indicate that a carboxylic acid moiety directly attached to the 2 position of the pyrimidine ring is most favorable for intravenous activity while esters of this acid are preferred for oral activity. The oral activity of ethyl 3,4-dihydro-4-oxopyrimido[4,5-*b*]quinoline-2-carboxylate (ED₅₀ = 3 mg/kg) places this ester among the more potent orally active antiallergy agents reported to date.


The discovery about a decade ago that inhalation of disodium cromoglycate (1, DSCG) provides protection



against antigen-induced seasonal allergic asthma² has stimulated considerable research seeking agents acting by a similar pharmacological mechanism.


Since DSCG is not absorbed orally this effort has focused on finding orally active agents. In the past few years a number of different chemical series have been reported to possess oral activity in the rat passive cutaneous anaphylaxis (PCA) procedure, a test capable of identifying antiallergy agents pharmacologically related to DSCG. Examples of these orally active series include xanthone-2-carboxylic acids,³ 2-nitroindan-1,3-diones,⁴ 3-(5-tetra-*o*lyl)thioxanthone 10,10-dioxides,⁵ 8-azapurinones,⁶ 1,4,6,9-tetrahydro-4,6-dioxypyrido[3,2-*g*]quinoline-2,8-dicarboxylic acids,⁷ cinnoline-3-propionic acids,⁸ 2- and 3-substituted chromones,⁹⁻¹² and aryl oxamates.¹³

We wish to report the development of a novel prototype, ethyl 3,4-dihydro-4-oxopyrimido[4,5-*b*]quinoline-2-carboxylate, which exhibits oral DSCG-type antiallergy activity. This compound was developed by successive molecular modification of the carbocyclic ring of 3,4-dihydro-4-oxoquinazoline-2-carboxylic acid (3), leading to



acids 5, 11, and then 16. Initial attempts to increase potency by substitution of the carbocyclic ring of 3 were unsuccessful. However, either incorporation of a nitrogen atom into this ring or fusion of another ring onto 3, af-

Scheme 1. Synthesis of 3,4-Dihydro-4-oxobenzo[*g*]quinazoline-2-carboxylic Acids (X = CH) and 3,4-Dihydro-4-oxopyrimido[4,5-*b*]quinoline-2-carboxylic Acids (X = N)



fording 5 and 11, respectively, did increase potency. A further, marked potency enhancement was achieved with compound 16 which incorporates both of these changes. More significantly, the ethyl ester of 16 displayed potent oral activity.

Chemistry. Although a few 3,4-dihydro-4-oxoquinazoline-2-carboxylic acids (3) have previously been reported in the literature,¹⁴ this communication reports the first synthesis of 3,4-dihydro-4-oxopyrimido[2,3-*d*]pyrimidine-2-carboxylic acids (5), 3,4-dihydro-4-oxobenzo[*g*]quinazoline-2-carboxylic acids (11), and 3,4-dihydro-4-oxopyrimido[4,5-*b*]quinoline-2-carboxylic acids (16). Esters of these four series of 3,4-dihydro-4-oxopyrimi-