

The fermentation was carried out in a 400-liter resin-lined tank using the synthetic medium described by Stone and Farrell¹ and culture Q-176 of *Penicillium notatum*. The precursor, γ -chlorocrotylmercaptoacetic acid, was employed at a concentration of 250 mg. per liter. The assay of the beer at harvest (sixty-four hours) was 250 Oxford units per ml. When processed by the carbon-acetone method,² 44 g. of crude sodium salt was obtained; bio-assay, 625 Oxford units per mg.; purity indicated by hydroxylamine assay,³ 24%. The crude penicillin was purified by partition chromatography^{4,5,6} of the free acid on an ether-silica column using pH 6.2 potassium phosphate buffer. A peak fraction in the eluates⁷ which included 70% of the activity was titrated to pH 7.0 with 1% potassium hydroxide solution and dried from the frozen state. The resulting amorphous potassium salt crystallized when treated with dry acetone. It was recrystallized by dissolving it in 90% acetone and adding 3 volumes of dry acetone. The yield of recrystallized potassium salt was 4.2 g.; bio-assay, 1900 Oxford units per mg.

Anal. Calcd. for $C_{14}H_{18}N_2O_4ClS_2K$: C, 40.32; H, 4.35; S, 15.38; Cl, 8.50. Found: C, 40.32; H, 4.35; S, 15.22; Cl, 8.37.

(1) Stone and Farrell, *Science*, **104**, 445 (1946).

(2) Whitmore, Wagner, Noll, Bassler, Fleming, Carnahan, Weisgerber, Oakwood, Herr, Patterson, Haggard, Mraz, Hoover, Di-Giorgio, Weisel, Lovell, Walter and Ropp, *Ind. Eng. Chem.*, **38**, 942 (1946).

(3) Ford, *Anal. Chem.*, **19**, 1004 (1947).

(4) Gordon, Martin and Syngé, *Biochem. J.*, **37**, 79 (1943).

(5) Harris and Wick, *Ind. Eng. Chem., Anal. Ed.*, **18**, 276 (1946).

(6) Behrens, Corse, Edwards, Garrison, Jones, Soper, Van Abeele and Whitehead, *J. Biol. Chem.*, **175**, 793 (1948).

(7) The optical rotation was found to be a convenient measurement for use in grouping the eluate fractions.

JARED H. FORD

HARRISON A. NELSON

UPJOHN RESEARCH LABORATORIES WILLIAM H. DeVRIES
KALAMAZOO 99, MICHIGAN ROBERTA HANSON

RECEIVED JUNE 14, 1948

Some New Derivatives of 4-Methylimidazole

During a search for compounds which possess antihistaminic activity, a brief study was made of substances which contained the imidazole nucleus. A few compounds were synthesized by treatment of 4-(chloromethyl)-imidazole hydrochloride¹ with an appropriate secondary amine or thiourea. The intermediate 4-(hydroxymethyl)-imidazole hydrochloride was prepared by the method of Totter and Darby.² None of the compounds listed below inhibited the effect of histamine on isolated intestinal strips of guinea pigs.

Typical Procedure.—The 4-(aminomethyl)-imidazoles were prepared by heating under reflux for two hours a mixture of 0.025 mole of 4-(chloromethyl)-imidazole hydrochloride, 0.075 mole of the secondary amine and 75 cc. of absolute alcohol. The resultant solution was first made alkaline with aqueous ammonia and the ammonium chloride that formed was removed by filtration. After evaporation under reduced pressure to remove the solvent and the excess amine, the residue was dissolved in absolute alcohol, from which the hydrochloride was obtained as a sticky solid by the addition of an ether solution of anhydrous hydrogen chloride. In all cases two or three recrystallizations from alcohol-ether gave pure products in the form of hygroscopic white powders.

4-(Diethylaminomethyl)-imidazole Dihydrochloride.—Yield, 90%; m. p. 200° with softening at 172°.

Anal. Calcd. for $C_8H_{16}N_3 \cdot 2HCl$: N, 18.6; Cl, 31.3. Found: N, 18.4; Cl, 31.2.

(1) Pyman, *J. Chem. Soc.*, **99**, 674 (1911).

(2) Totter and Darby, "Organic Syntheses," **24**, 69 (1944).

4-(1-Piperidylmethyl)-imidazole Dihydrochloride.—Yield, 45.5%; m. p. 223.6–227.1°.

Anal. Calcd. for $C_9H_{15}N_3 \cdot 2HCl$: N, 17.7; Cl, 29.6. Found: N, 17.9; Cl, 29.5.

4-(4-Morpholinylmethyl)-imidazole Dihydrochloride.—Yield, 66.7%; m. p. 173°.

Anal. Calcd. for $C_8H_{13}N_3O \cdot 2HCl$: C, 39.9; H, 6.25; N, 17.5; Cl, 29.6. Found: C, 39.4; H, 6.69; N, 16.5; Cl, 29.5.

S-(4-Imidazolylmethyl)-isothiourea Dihydrochloride.—A solution of 3.5 g. (0.05 mole) of thiourea in 100 cc. of absolute alcohol was brought to a boil and then treated with 7.6 g. (0.05 mole) of 4-(chloromethyl)-imidazole hydrochloride. The resultant mixture was heated under reflux for ten minutes, and the solid which precipitated upon cooling was collected. A practically quantitative yield was obtained of milky, white prisms which, after recrystallization from dilute alcohol, melted at 227.6–229.6°.

Anal. Calcd. for $C_5H_8N_4S \cdot 2HCl$: N, 24.4; Cl, 31.0. Found: N, 24.3; Cl, 30.9.

CALCO CHEMICAL DIVISION

AMERICAN CYANAMID COMPANY

BOUND BROOK, NEW JERSEY

RICHARD J. TURNER

RECEIVED JUNE 9, 1948

Dichromate Salts of 2-Benzyl and 4-Benzylpyridine

2-Benzylpyridinium Dichromate.—2-Benzylpyridine (25 g., 0.15 mole), dissolved in hot water (600 ml.) containing sufficient sulfuric acid to yield a clear solution, was treated with a solution of chromic anhydride (30 g., 0.30 mole) in water (200 ml.). A red oil was deposited which even on seeding could not be induced to crystallize. However, upon solution in acetone (100 ml.) and reprecipitation by addition of water (600 ml.) the oil readily crystallized in bright orange prisms. By several recrystallizations from hot water (any traces of tar being removed by filtration) the melting point was raised to 95–96°, uncor., with decomposition. There was no evidence of hydration.

The salt was analyzed for chromium by ignition; the neutralization equivalent was determined in water using phenolphthalein, the color change being from orange to yellow (dichromate to chromate) and finally back to orange (red of indicator plus yellow of chromate).

Anal. Calcd. for $C_{24}H_{24}Cr_2N_2O_7$ (*i. e.*, $(C_{12}H_{11}N)_2 \cdot H_2Cr_2O_7$): Cr, 18.7; neut. equiv., 139.0. Found: Cr, 18.9, 18.9; neut. equiv., 136.7, 136.8.

4-Benzylpyridinium Dichromate.—4-Benzylpyridine (5 g., 0.03 mole) dissolved in dilute sulfuric acid (25 ml.) and treated with a solution of chromic anhydride (5 g., 0.05 mole) in water (5 ml.) precipitated a red oil. By solution in acetone (15 ml.) and pouring into water (50 ml.) the salt soon crystallized in orange flakes (7.7 g., 94% yield) melting at 110–115° uncor. with decomposition. By further recrystallization from water (as for the isomer above) the melting point of the salt was raised to 115–115.5°, uncor., with decomposition.

Anal. Calcd. for $C_{24}H_{24}Cr_2N_2O_7$ (*i. e.*, $(C_{12}H_{11}N)_2 \cdot H_2Cr_2O_7$): Cr, 18.7; neut. equiv., 139.0. Found: Cr, 18.8, 18.9; neut. equiv., 138.7, 139.1.

DEPARTMENT OF CHEMISTRY

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

CAMBRIDGE 39, MASS.

JACKSON LABORATORY

ERNEST H. HUNTRESS

E. I. DU PONT DE NEMOURS AND CO., INC.

WILMINGTON, DELAWARE

HENRY C. WALTER

RECEIVED MAY 14, 1948

1,2-Di-(4-ketocyclohexane)-ethane

1,2-Di-(4-hydroxycyclohexane)-ethane was obtained by high-pressure hydrogenation of *p,p'*-dihydroxystilbene