

Synthesis and Properties of Macroheterocyclic Compounds of ABAB-Type with *tert*-Butylpyrrolepyrazine Fragments[⊗]

Mikhail G. Stryapan, Svetlana V. Efimova, Oscar I. Koifman,
and Mikhail K. Islyaikin[@]

Ivanovo State University of Chemistry and Technology, Friedrich Engels Pr-t, 7, Ivanovo, 153000, Russia

[@]Corresponding author E-mail: islyaikin@isuct.ru

*New representatives of macroheterocyclic compounds of ABAB-type were synthesized by condensation of *m*-phenylenediamine or 2,6-diaminopyridine (A) with 5-*tert*-butyl-2,3-dicyanopyrazine (B) in *n*-BuOH. These compounds were characterized by UV-vis, IR, ¹H NMR spectroscopies, mass-spectrometry and elemental analysis.*

Keywords: Macroheterocyclic compounds, *m*-phenylenediamine, 2,6-diaminopyridine, 5-*tert*-butyl-2,3-dicyanopyrazine, hemiporphyrine.

Introduction

Macroheterocyclic compounds (Mc) of ABAB-type can be represented as structural analogues of phthalocyanine, where two opposite isoindole moieties (B) are replaced by aromatic rings (A) derived from aromatic diamines. Since their discovery,^[1] these compounds called by Campbell "hemiporphyrines"^[2] are the objects of continuous interest due to their structure particularities^[3,4] as well as attractive properties revealed recently: optical limiting, fluorescence, catalysis, devices for keeping information and so on.^[5-7]

For the moment a number of representatives of this series was synthesized,^[8,9] but macrocycles containing pyrazine fragments (B) and those of six-membered aromatic diamines (A) are studied not enough completely. The first two representatives of nonsubstituted macroheterocyclic compounds of this type were described earlier.^[10] These compounds were characterized by IR and UV-vis spectroscopies as well as elemental analysis data. Their serious imperfection is a low solubility in common organic solvents, what makes their refinement to be little tedious and constricts a number of methods which could be applied to study their structures and properties. It was established that introduction of bulky *tert*-butyl groups led to increase of solubility of Mc.^[11] Hence the subject of this work is synthesis of Mc of ABAB-type bearing *tert*-butylpyrazine rings incorporated into macrocyclic core as units (B).

Experimental

General procedure. A mixture of 5-*tert*-butyl-2,3-dicyanopyrazine (1 mmol) and *m*-phenylenediamine or 2,6-diaminopyridine (1 mmol) in butanol (10 ml) was stirred at

80 °C for 6 h and then refluxed for 23 h. The solvent was removed under vacuum, and the crude product was washed with MeOH and then extracted with CHCl₃ in Soxhlet's apparatus. After solvent evaporation, the product was purified by column chromatography on silica gel (CHCl₃:MeOH:C₆H₁₄ = 10:1:5). The solvents were removed and the solids were dried under vacuum.

2,15-Di(*tert*-butyl)-5,26:13,18-diimino-7,11:20,24-dimetheno-[*c,n*]-dipyrano-[1,6,12,17]-tetraazacyclodocosen-[1,3,5,7,9,12,14,16,21,23], 1. It was synthesized following the general procedure by interaction of 5-*tert*-butyl-2,3-dicyanopyrazine with *m*-phenylenediamine as a dark orange powder. Yield: 36 mg (26 %). UV-vis (CHCl₃) λ_{max} nm (lg ε): 361 (3.82), 383 (3.82). IR (KBr) ν cm⁻¹: 3424, 2922, 2852, 1636, 1464, 1254, 1157. MS (MALDI-TOF) *m/z*: 555 [(M+H)⁺]. Found: C 68.51, H 5.34, N 25.17 %. C₃₂H₃₀N₁₀ requires C 69.30, H 5.45, N 25.25 %.

2,15-Di(*tert*-butyl)-5,26:13,18-diimino-7,11:20,24-dinitrilo-[*c,n*]-dipyrano-[1,6,12,17]-tetraazacyclodocosen-[1,3,5,7,9,12,14,16,21,23], 2. It was synthesized following the general procedure by interaction of 5-*tert*-butyl-2,3-dicyanopyrazine with 2,6-diaminopyridine. The target product represents a dark red powder. Yield: 13 mg (9 %). UV-vis (CHCl₃) λ_{max} nm (lg ε): 379 (3.96), 425 (3.96). IR (KBr) ν cm⁻¹: 3327, 2927, 2856, 1746, 1640, 1465, 1250, 1165. ¹H NMR (CDCl₃) δ ppm: 15.09 (s, 2H), 8.9 (s, 2H), 7.6 (tr, 2H), 7.1 (d, 2H), 6.3 (d, 2H), 1.55, (18H, -CH₃). MS (MALDI-TOF) *m/z*: 556 [(M+H)⁺]. Found: C 64.41, H 4.89, N 29.63 %. C₃₀H₂₈N₁₂ requires C 64.72, H 5.03, N 30.21 %.

Results and Discussion

A general way to gain access to the Mc of ABAB-type is the interactions of alkoxy- or diiminoisoindolines with aromatic diamines. But phthalonitrile itself and substituted phthalonitriles are able to react directly with diamines without previous transformation into corresponding isoindoline derivatives.^[12,13] It is worthy to note that presence of nitrogen atoms in dicyanoconstituent makes this reaction easier to go.¹³ So we succeed in synthesizing of Mc **1**, **2** by direct condensation of 5-*tert*-butyl-2,3-dicyanopyridine with 1,3-phenylenediamine or 2,6-diaminopyridine in *n*-BuOH.

⊗ This contribution is dedicated to professor Vasilij Fedorovich Borodkin on the occasion of his 100th Anniversary.

⊗ Статья посвящена 100-летию юбилею профессора Василия Фёдоровича Бородинкина.




Figure 1. IR spectra of compounds **1** and **2**.

The reactions were monitored by TLC along with NH₃ separation. After solvent elimination, the residues were purified by extraction of topic products by CHCl₃ in Soxhlet extractor with following column chromatography using silica-gel.

The compounds **1**, **2** were characterized by UV-vis, IR, ¹H NMR spectroscopies, mass-spectrometry, and elemental analysis.

The signals at 555 and 556 *m/z* which are in good agreement with molecular ions [M+H]⁺ of **1** and **2** correspondingly were detected in MALDI-TOF. There are three groups of signals in the ¹H NMR spectra of **1** and **2**. So, in spectrum of **2**, two series of signals located at 0.8 – 1.8 and 6.0 – 9 ppm are induced by resonance of the protons of *tert*-butyl groups and those of aromatic systems correspondingly. The signals (~15 ppm) of **1** and **2** are due to the resonance of iminogroup protons and their position at a

such low field shows that these compounds have no aromatic macrocycles at the base of their skeletons.

UV-vis spectra of **1** and **2** in chloroform contain the broad bands of absorbance with maxima located at 383 and 425 nm, correspondingly. A longer wavelength shift observed on going from **1** to **2** is most probably due to more plane structure of the central core of Mc **2** if compared with **1**.

IR-spectra of compounds **1** and **2** (Figure 1) are very similar to each other. So, the strong bands at 2925 and 2855 cm⁻¹ are due to the symmetric and asymmetric valance vibrations of C-H bonds belonging to *tert*-butyl groups. The bands at 1640 and 1465 cm⁻¹ can be put in concordance with deformations of C=C and C=N bonds, correspondingly.

Conclusions

tert-Butylsubstituted azaanalogues of hemiporphyrines were obtained by direct condensation of 5-*tert*-butyl-2,3-dicyanopyridine with *m*-phenylenediamine or 2,6-diaminopyridine in boiling *n*-butanol. Sufficiently high solubility in organic solvents allows to apply the column chromatography for their purification. Their structures were established on the base of UV-vis, IR, ¹H NMR spectroscopies, mass-spectrometry and elemental analysis data.

Acknowledgements. This work was supported by Federal Principal Program “Research and research-pedagogical staffs of innovative Russia for 2009-2013” (№ 02.740.11.0106).

References

1. Elvidge J.A., Linstead R.P. *J. Chem. Soc.* **1952**, 20, 5008-5012.
2. Campbell J. B. Pat. 2765308 (USA) **1956**.
3. Rodríguez-Morgade M.S., Gema de la Torre, Torres T. Design and Synthesis of Low-Symmetry Phthalocyanines and Related Systems, in *The Porphyrin Handbook* (Kadish K.M., Smith K.M., Guillard R., Eds.) Amsterdam, Boston, London, New York, Oxford, Paris, San Diego, San Francisco, Singapore, Sydney, Tokyo: Academic Press **2003**, Vol. 15, 125-160.
4. Islyaikin M.K., Danilova E.A., Romanenko Yu.V., Khelevina O.G., Lomova T.N. Synthesis, Structure Peculiarities and Biological Properties of Macroheterocyclic Compounds, in *Chemical Processes with Participation of Biological and Related Compounds*. (Lomova T.N., Zaikov G.E., Eds.) BRILL, Leiden-Boston **2008**, 219-270.
5. Dini D., Calvete M.J.F., Hanack M., Amendolab V., Meneghetti M. *Chem. Commun.* **2006**, 22, 2394-2396.
6. Campaci F., Campestrini S. *J. Mol. Catalysis A: Chemical* **1999**, 140, 121-130.

7. Avanzino S. Pat. 7141482 (USA) **2006**.
8. Fernánrez-Lazáro F., Torres T., Hauschel B., Hanack M. *Chem. Rev.* **1998**, 98, 563-576.
9. Islyaikin M.K., Danilova E.A. *Russ. Chem. Bull., Int. Ed.* **2007**, 5, 689-706.
10. Neustroeva N.R., Kulikov M.A., Vorob'ev Yu.G., Smirnov R.P. *Zh. Obshch. Khim.* **1999**, 69, 1836-1838 (in Russ.).
11. Danilova E.A., Islyaikin M.K. Synthesis and Properties of *tert*-Butylsubstituted Macroheterocyclic Compounds and their Metal Complexes, in *Uspekhi Khimii Porfirinov (Advances in Porphyrin Chemistry)* (Golubchikov O.A., Ed.) St. Petersburg: NII Khimii SPbGU **2004**, Vol. 4, 356-375 (in Russ.).
12. Islyaikin M.K., Mayzlish V.E., Borodkin V.F. *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.* **1979**, 22, 152-154.
13. Rodriguez-Morgade M.S., Pantos G.D., Caballero E., Sessler J.L., Torres T. *Macroheterocycles* **2008**, 1, 40-43.

Received 10.11.2009

Accepted 15.03.2010

First published on the web 20.03.2010