



Figure 1. Nonradiative energy transfer in films containing a mixture of equal weights of an acceptor-labeled poly(methyl methacrylate) and donor-labeled methyl methacrylate-butyl methacrylate copolymers. The molecular weight of the polymers was estimated from solution viscosities as 380 000.

is easily observed with a difference of 5 mol % in the composition of the two polymers.

Acknowledgments. One of us (F.A.) is indebted to the American Friends of the Middle East for a research fellowship. We also wish to express our gratitude to the National Science Foundation for their support of this study through Grant DMR 77-05210.

References and Notes

- (1) This work will form part of a Ph.D. dissertation to be submitted by F.A. to the Graduate School of the Polytechnic Institute of New York in June 1978.
- (2) S. Krause, *J. Macromol. Sci., Rev. Macromol. Chem.*, **7**, 251 (1972).
- (3) G. Gee, *Q. Rev., Chem. Soc.*, **1**, 265 (1947).
- (4) T. Nishi, T. T. Wang, and T. K. Kwei, *Macromolecules*, **8**, 227 (1975).
- (5) G. L. Slonimski, *J. Polym. Sci.*, **30**, 625 (1958).
- (6) M. P. Zverev, L. A. Polovikhina, A. N. Barash, L. P. Mal'kova, and G. D. Litovchenko, *Vysokomol. Soedin.*, **16**, 2098 (1974).
- (7) S. Ichihara, A. Kamatsu, and T. Hata, *Polym. J.*, **2**, 640 (1971).
- (8) H. E. Bair, *Polym. Eng. Sci.*, **10**, 247 (1970).
- (9) J. Stoelting, F. E. Karasz, and W. J. MacKnight, *Polym. Eng. Sci.*, **10**, 133 (1970).
- (10) F. Kollinsky and G. Markert, *Adv. Chem. Ser.*, **No. 99**, 175-188 (1972).
- (11) M. Matsuo, C. Nozaki, and C. Jyo, *Polym. Eng. Sci.*, **9**, 197 (1969).
- (12) *Adv. Chem. Ser.*, **No. 99** (1972).
- (13) I. Z. Steinberg, *Annu. Rev. Biochem.*, **40**, 83 (1971).

H. Morawetz* and F. Amrani

Polytechnic Institute of New York,
Brooklyn, New York 11201

Received October 7, 1977

CORRECTIONS

"Statistical Mechanical Treatment of Protein Conformation. 2. A Three-State Model for Specific-Sequence Copolymers of Amino Acids", by Seiji Tanaka* and Harold A. Scheraga, Volume 9, Number 1, January-February 1976, pages 165 and 166.

The upper limit of the product in eq 54, p 165, should be $i + n - 2$.

Equation 65 on page 166 should read:

$$P(i|n|\{\rho\}) \neq \left[\prod_{k=i}^{i+n-1} F_{k,\eta_k} \right]_{|\rho\}}$$

or

$$P(i|n|\{\rho\}) \neq \left[\prod_{k=i}^{i+n-1} P(k|1|\eta_k) \right]_{|\rho\}}$$

"Statistical Mechanical Treatment of Protein Conformation. 3. Prediction of Protein Conformation Based on a Three-State Model", by Seiji Tanaka* and Harold A. Scheraga, Volume 9, Number 1, January-February 1976, page 170.

The upper limit of the product in eq 9 should be $i + n - 2$.

"Statistical Mechanical Treatment of Protein Conformation. 6. Elimination of Empirical Rules for Prediction by Use of a High-Order Probability. Correlation between the Amino Acid Sequences and Conformations for Homologous Neurotoxin Proteins", by Seiji Tanaka* and Harold A. Scheraga, Volume 10, Number 2, March-April 1977, page 308.

The upper limit of the product in eq 5 should be $i + n - 2$.

These were typographical errors in printing. However, these errors were *not* made in the computations.

* Deceased November 8, 1977.

"Cross Relaxation in Poly(vinylidene fluoride) from Transient Overhauser Measurements", by V. J. McBrierty and D. C. Douglass*, Volume 10, Number 4, July-August 1977, page 855.

Due to a typographical error in our manuscript eq 4 and 5 are stated incorrectly. In eq 4 τ_c should be replaced by 10^{-8} . τ_c in eq 5 should read $S_z - S_0 = S_0[e^{-t/D_1} + e^{-t/T_1}]$.

The correct equations have been used in subsequent analysis.