

CORRECTIONS

Jeffrey Pyun, Shijun Jia, Tomasz Kowalewski, Gary D. Patterson, and Krzysztof Matyjaszewski*: Synthesis and Characterization of Organic/Inorganic Hybrid Nanoparticles: Kinetics of Surface-Initiated Atom Transfer Radical Polymerization and Morphology of Hybrid Nanoparticle Ultrathin Films. Volume 36, Number 14, July 15, 2003, pp 5094–5104.

Page 5098. The values of k_p^{app} in Table 2 were from experiments using initiators attached to colloidal silica. The correct values for homogeneous ATRP using ethyl 2-bromoisobutyrate as the initiator are shown in the revised table below.

Table 2. Summary of Rate and Equilibrium Constants Determined from the Homopolymerization of Sty, BA, and MMA in the ATRP from Ethyl 2-Bromoisobutyrate

monomer	$k_p^{\text{app}}, \text{s}^{-1}$	$k_p, \text{M}^{-1} \text{s}^{-1}$	K_{eq}	$k_t, \text{M}^{-1} \text{s}^{-1}$
MMA	2×10^{-4}	1615	7×10^{-7}	3.7×10^3
Sty	6.5×10^{-6}	895	4×10^{-8}	5.5×10^3
BA	2.7×10^{-6}	56780	4×10^{-10}	1.2×10^3

MA030401+

10.1021/ma030401+

Published on Web 08/14/2003