


Corrigendum

Structures of Two Novel Triterpene Saponins from *Arenaria filicaulis* (Boiss.)

M. H. A. Elgamal, H. S. M. Soliman,
D. T. Elmunajjed, G. Tóth,* A. Simon and
H. Duddeck,*

Magn. Reson. Chem. **35**, 637 (1997).

During the continuation of our studies on the constitu-



1 R = H

2 R = SO₃H

Scheme 1. Structures of Snatzkein A (1) and B (2)

ents of *Arenaria filicaulis* we realized that the structure of glycoside **2** is wrong. A negative FAB-MS measurement proved unequivocally that the substituent at the oxygen atom of C-2' is not R = CH₂CH₂OH, but a sulfonyl group (SO₃H), similar to another saponin from the same plant which we studied later.¹ Apparently the CH₂CH₂OH group signals which we observed in the spectra have to be attributed to a 2:1 host-guest complex formation of **2** and diglycole.

REFERENCE

1. G. Tóth, A. Simon, M. H. A. Elgamal, H. S. M. Soliman, D. T. Elmunajjed, Gy. Horváth and H. Duddeck, *Magn. Reson. Chem.* submitted

* Correspondence to: G. Tóth, Technical University Budapest, Technical Analytical Research Group of the Hungarian Academy of Sciences, Institute for General and Analytical Chemistry, Szent Gellért tér 4, H-1111 Budapest, Hungary or H. Duddeck, Universität Hannover, Institut für Organische Chemie, Schneiderberg 1B, D-30167 Hannover, Germany.