

# MEMOIRS

of the  
American Mathematical Society

Volume 234 • Number 1103 (third of 5 numbers) • March 2015

## Quasi-Linear Perturbations of Hamiltonian Klein-Gordon Equations on Spheres

J.-M. Delort



ISSN 0065-9266 (print) ISSN 1947-6221 (online)

**American Mathematical Society**

[www.ebook3000.com](http://www.ebook3000.com)

# MEMOIRS

of the  
American Mathematical Society

---

Volume 234 • Number 1103 (third of 5 numbers) • March 2015

---

## Quasi-Linear Perturbations of Hamiltonian Klein-Gordon Equations on Spheres

J.-M. Delort



---

ISSN 0065-9266 (print) ISSN 1947-6221 (online)

---

**American Mathematical Society**  
Providence, Rhode Island

---

## Library of Congress Cataloging-in-Publication Data

Delort, Jean-Marc, 1961-

Quasi-linear perturbations of Hamiltonian Klein-Gordon equations on spheres / J.-M. Delort.  
pages cm. – (Memoirs of the American Mathematical Society, ISSN 0065-9266 ; volume 234,  
number 1103)

Includes bibliographical references and index.

ISBN 978-1-4704-0983-8 (alk. paper)

1. Hamiltonian systems. 2. Klein-Gordon equation. 3. Wave equation. 4. Sphere. I. Title.  
QA614.83.D45 2014

516'.156–dc23

2014041958

DOI: <http://dx.doi.org/10.1090/memo/1103>

---

## Memoirs of the American Mathematical Society

This journal is devoted entirely to research in pure and applied mathematics.

**Subscription information.** Beginning with the January 2010 issue, *Memoirs* is accessible from [www.ams.org/journals](http://www.ams.org/journals). The 2015 subscription begins with volume 233 and consists of six mailings, each containing one or more numbers. Subscription prices for 2015 are as follows: for paper delivery, US\$860 list, US\$688.00 institutional member; for electronic delivery, US\$757 list, US\$605.60 institutional member. Upon request, subscribers to paper delivery of this journal are also entitled to receive electronic delivery. If ordering the paper version, add US\$10 for delivery within the United States; US\$69 for outside the United States. Subscription renewals are subject to late fees. See [www.ams.org/help-faq](http://www.ams.org/help-faq) for more journal subscription information. Each number may be ordered separately; *please specify number* when ordering an individual number.

**Back number information.** For back issues see [www.ams.org/bookstore](http://www.ams.org/bookstore).

Subscriptions and orders should be addressed to the American Mathematical Society, P. O. Box 845904, Boston, MA 02284-5904 USA. *All orders must be accompanied by payment.* Other correspondence should be addressed to 201 Charles Street, Providence, RI 02904-2294 USA.

**Copying and reprinting.** Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy select pages for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Permissions to reuse portions of AMS publication content are handled by Copyright Clearance Center's RightsLink® service. For more information, please visit: <http://www.ams.org/rightslink>.

Send requests for translation rights and licensed reprints to [reprint-permission@ams.org](mailto:reprint-permission@ams.org).

Excluded from these provisions is material for which the author holds copyright. In such cases, requests for permission to reuse or reprint material should be addressed directly to the author(s). Copyright ownership is indicated on the copyright page, or on the lower right-hand corner of the first page of each article within proceedings volumes.

---

*Memoirs of the American Mathematical Society* (ISSN 0065-9266 (print); 1947-6221 (online)) is published bimonthly (each volume consisting usually of more than one number) by the American Mathematical Society at 201 Charles Street, Providence, RI 02904-2294 USA. Periodicals postage paid at Providence, RI. Postmaster: Send address changes to *Memoirs*, American Mathematical Society, 201 Charles Street, Providence, RI 02904-2294 USA.

© 2014 by the American Mathematical Society. All rights reserved.

Copyright of individual articles may revert to the public domain 28 years after publication. Contact the AMS for copyright status of individual articles.

This publication is indexed in *Mathematical Reviews*®, *Zentralblatt MATH*, *Science Citation Index*®, *Science Citation Index<sup>TM</sup>-Expanded*, *ISI Alerting Services<sup>SM</sup>*, *SciSearch*®, *Research Alert*®, *CompuMath Citation Index*®, *Current Contents*®/*Physical, Chemical & Earth Sciences*. This publication is archived in *Portico* and *CLOCKSS*.

Printed in the United States of America.

⊗ The paper used in this book is acid-free and falls within the guidelines established to ensure permanence and durability.

Visit the AMS home page at <http://www.ams.org/>

10 9 8 7 6 5 4 3 2 1      20 19 18 17 16 15

## Contents

Chapter 0. Introduction	1
Chapter 1. Statement of the main theorem	7
Chapter 2. Symbolic calculus	23
Chapter 3. Quasi-linear Birkhoff normal forms method	41
Chapter 4. Proof of the main theorem	53
A. Appendix	77
Bibliography	79



## Abstract

The Hamiltonian  $\int_X (|\partial_t u|^2 + |\nabla u|^2 + \mathbf{m}^2 |u|^2) dx$ , defined on functions on  $\mathbb{R} \times X$ , where  $X$  is a compact manifold, has critical points which are solutions of the linear Klein-Gordon equation. We consider perturbations of this Hamiltonian, given by polynomial expressions depending on first order derivatives of  $u$ . The associated PDE is then a *quasi-linear* Klein-Gordon equation. We show that, when  $X$  is the sphere, and when the mass parameter  $\mathbf{m}$  is outside an exceptional subset of zero measure, smooth Cauchy data of small size  $\epsilon$  give rise to almost global solutions, i.e. solutions defined on a time interval of length  $c_N \epsilon^{-N}$  for any  $N$ . Previous results were limited either to the semi-linear case (when the perturbation of the Hamiltonian depends only on  $u$ ) or to the one dimensional problem.

The proof is based on a quasi-linear version of the Birkhoff normal forms method, relying on convenient generalizations of para-differential calculus.

---

Received by the editor Received by the editor April 6, 2012 and, in revised form, April 11, 2013.

Article electronically published on July 28, 2014.

DOI: <http://dx.doi.org/10.1090/memo/1103>

2010 *Mathematics Subject Classification*. Primary 35L72, 35S50, 37K45.

*Key words and phrases*. Hamiltonian quasi-linear Klein-Gordon equations, Almost global existence, Birkhoff normal forms.



## CHAPTER 0

# Introduction

This paper is devoted to the study of solutions of small quasi-linear perturbations of an infinite dimensional Hamiltonian system. To be more specific, let  $X$  be a compact Riemannian manifold, and define on  $H^s(X, \mathbb{C})$ , for  $s$  large enough,  $G_0(U) = \int_X (\Lambda_{\mathbf{m}} u) \bar{u} d\mu$ , where  $U = (u, \bar{u})$ ,  $d\mu$  is the Riemannian volume and  $\Lambda_{\mathbf{m}} = \sqrt{-\Delta_X + \mathbf{m}^2}$ . Since  $H^s(X, \mathbb{C})$  is endowed with a symplectic form  $\omega_0(h, h') = 2\text{Im} \int_X h \bar{h}' d\mu$ , one may consider the Hamiltonian equation associated to  $G_0$ , given by

$$(1) \quad \partial_t u = i \nabla_{\bar{u}} G_0(u, \bar{u}).$$

If  $u = \frac{\sqrt{2}}{2} (\Lambda_{\mathbf{m}}^{-1/2} \partial_t v + i \Lambda_{\mathbf{m}}^{1/2} v)$ , with  $v$  in  $H^{s+1/2}(X, \mathbb{R})$  and  $\partial_t v$  in  $H^{s-1/2}(X, \mathbb{R})$ , this Hamiltonian equation is nothing but the Klein-Gordon equation

$$(2) \quad (\partial_t^2 - \Delta + \mathbf{m}^2)v = 0.$$

We want to study the solutions of equations of form (1), where  $G_0$  has been replaced by a more general Hamiltonian  $G$ , such that  $G - G_0$  is small. By rescaling, this is equivalent to the study of

$$(3) \quad \partial_t u = i \nabla_{\bar{u}} G(u, \bar{u}),$$

where  $G - G_0$  vanishes at least at order three at zero, and where the Cauchy data are small in  $H^s$  ( $s \gg 1$ ), of size  $\epsilon \rightarrow 0$ . The question is to determine whether, for  $\epsilon$  small enough, the solution exists over a long interval of time, and has Sobolev norm  $O(\epsilon)$  on that interval.

Since  $G_0$  may be written as  $\int_X (\Lambda_{\mathbf{m}}^{1/2} u) (\Lambda_{\mathbf{m}}^{1/2} \bar{u}) d\mu$ , the most natural perturbation would be of the form  $(G - G_0)(u, \bar{u}) = \int_X P(\Lambda_{\mathbf{m}}^{1/2} u, \Lambda_{\mathbf{m}}^{1/2} \bar{u}) d\mu$  for some function  $P$  vanishing at least at order three at the origin. Nevertheless, most works devoted to that question up to now concern the case when the perturbation  $G - G_0$  is given by a function of  $\Lambda_{\mathbf{m}}^{-1/2} U$ . At the level of the Hamiltonian equation, this corresponds to perturbations of (2) which are weakly semi-linear, i.e. may be written

$$(4) \quad (\partial_t^2 - \Delta + \mathbf{m}^2)v = f(v),$$

for a smooth function  $f$  vanishing at least at order 2 at zero, while, in the general case, the non-linear perturbation is quasi-linear, i.e. depends on derivatives of  $v$  up to order 2, and is linear with respect to  $(\partial^\alpha v)_{|\alpha|=2}$ .

In one dimension, i.e. when  $X$  is the circle, the weakly semi-linear problem has been solved by Bourgain [12] and Bambusi [1] (see also for related results Bambusi-Grébert [3] and the lectures of Grébert [25]). It has been proved that, if  $\mathbf{m}$  is taken outside a subset of zero measure of  $]0, +\infty[$ , for any  $N \in \mathbb{N}$ , for any  $s$  large enough, there is  $\epsilon_0 > 0$ , such that equation (4) with Cauchy data of



size  $\epsilon < \epsilon_0$  in  $H^{s+1/2}(\mathbb{S}^1, \mathbb{R}) \times H^{s-1/2}(\mathbb{S}^1, \mathbb{R})$ , has an almost global solution, i.e. a solution defined on a domain  $] -T_\epsilon, T_\epsilon[ \times \mathbb{S}^1$ , with  $T_\epsilon > c\epsilon^{-N}$ , satisfying an estimate

$$(5) \quad \sup_{|t| < T_\epsilon} \|v(t, \cdot)\|_{H^{s+1/2}} + \sup_{|t| < T_\epsilon} \|\partial_t v(t, \cdot)\|_{H^{s-1/2}} \leq C\epsilon.$$

This result has been extended to higher dimensions, i.e. to equation (4) on spheres, or more generally on Zoll manifolds, by Bambusi, Delort, Grébert and Szeftel [2], after preliminary results of Delort and Szeftel [17–19]. The method of proof relies on Birkhoff normal forms, as in one dimension, to reduce the proof of long time existence and of estimate (5) to the study of a sequence of homological equations. The resolution of these equations is possible only under a suitable small divisors property, which holds when the parameter  $\mathbf{m}$  is outside a subset of zero measure, because of the very special distribution of the eigenvalues of  $-\Delta$  on  $\mathbb{S}^d$ . Actually, the important property is that two different eigenvalues of  $\sqrt{-\Delta_{\mathbb{S}^d}}$  are separated essentially by a fixed distance. When such a property does not hold, for instance when  $X = \mathbb{T}^d$ , or when  $-\Delta$  is replaced by the harmonic oscillator  $-\Delta + |x|^2$  on  $X = \mathbb{R}^d$ , only much weaker results are known (see [15, 21, 35]).

Let us mention also that normal forms methods have been proved useful to study long time existence for solutions of equations of the form (4) on  $\mathbb{R}^d$ , for Cauchy data that are smooth and decaying at infinity. Actually, the use of normal forms in that framework has been introduced by Shatah [33] to prove global existence of solutions of non-linear Klein-Gordon equations on  $\mathbb{R}^3$  (an alternative proof has been given at the same time by Klainerman [29]). More recently, this approach has been made more systematic in a series of papers of Germain, Masmoudi and Shatah [23, 24] and Germain [22].

Our concern here is to treat the problem of general perturbations, giving rise to a quasi-linear equation instead of (4). This brings considerable new difficulties, especially in higher dimensions. Actually, in one dimension, i.e. on the circle, we could solve the problem in [16]. Our goal in this paper is to obtain a long time stability result of type (5) on any higher dimensional sphere  $\mathbb{S}^d$ . The main difference with the one dimensional case comes from the fact that one can no longer make appeal to Fourier series, as in one dimension, nor use just separation properties of the eigenvalues of the laplacian on the sphere, like in the semi-linear case. One has to rely instead on the classical property of the geodesic flow that is responsible for these behaviour of eigenvalues, namely periodicity.

Before giving some hints on the strategy we shall follow, let us comment on related – though different – results concerning existence of periodic or quasi-periodic solutions to equations of the form (4). This problem has been solved in one dimension by Kuksin [31], Wayne [34], Craig and Wayne [14], and in higher dimensions (i.e. on the torus) by Bourgain [10, 11]. More recently, existence of periodic solutions for Klein-Gordon equations on homogeneous spaces (including the sphere) has been proved by Berti and Procesi [9]. We refer also to the book of Bourgain [13] for more references, and to the paper of Berti and Bolle [8] for a complete and up to date bibliography on these matters. Let us stress upon the fact that all these results were using in an essential way the fact that the non-linearity in the equation *does not* depend on derivatives of  $v$ . The challenge of treating equations depending also on first order derivatives of  $v$  has been overcome only recently, by Berti, Biasco and Procesi [6, 7], for Hamiltonian wave equations as well as for a more general class of PDEs (reversible wave equations). Earlier results concerning

other equations with unbounded non-linear perturbations are due to Kuksin [30] and Kappeler-Pöschel [28] for KdV equations, and to Liu-Yuan [32] in a general abstract framework. In the case of quasi-linear non-linearities, like the ones we consider in this paper, very few results are known. Let us mention the construction of periodic solutions for the water waves equation by Iooss, Plotnikov and Toland [27] and very recent results of Baldi, Berti and Montalto [5], who constructed quasi-periodic solutions for quasi-linear KdV equations.

Let us describe the main new ideas of the proof of our paper in more details. As in [16], the key point is to design a Birkhoff normal forms method adapted to quasi-linear equations. This requires to express the Hamiltonian using para-differential operators. In one dimension, one uses Fourier analysis on  $\mathbb{S}^1$  to define such classes of operators globally on  $\mathbb{S}^1$ , using symbols which are functions on  $\mathbb{S}^1 \times \mathbb{Z}$  (where  $\mathbb{Z}$  should be considered as the dual group of  $\mathbb{S}^1$ ). In higher dimensions, we define para-differential operators using a characterization in terms of commutators with differential operators, similar to the Beals characterization of pseudo-differential operators [4]. The classes we need are more general than usual para-differential operators: they depend on some auxiliary functions, and have to take into account some losses relatively to small frequencies that will appear because of small divisors. Because of that, we have to rewrite the whole theory (symbolic calculus, principal symbols on compact manifolds, ...) in our framework. This is done in section 2 of the paper. Section 3 is devoted to the computation of Poisson brackets between functions defined in terms of integrals of type  $\int_X (A(U)u)\bar{u} d\mu$ ,  $\text{Re} \int_X (A(U)u)u d\mu$ , where  $A(U)$  is a para-differential operator of order one, homogeneous of some degree  $k$  in  $U$ . The proof of the main theorem of long time existence occupies section 4. Let us describe the main new idea on a toy model.

Consider equation (3) with Hamiltonian

$$(6) \quad G(u, \bar{u}) = \int_X (\Lambda_{\mathbf{m}} u) \bar{u} d\mu + \text{Re} \int_X (A(U)u) \bar{u} d\mu,$$

where  $A(U)$  is a self-adjoint para-differential operator of order one, homogeneous of degree  $k$  in  $U = (u, \bar{u})$ . Equation (3) reads

$$(7) \quad D_t u = \Lambda_{\mathbf{m}} u + A(U)u + \dots$$

where the dots represent some contributions which are of a semi-linear nature (opposed to the main quasi-linear contribution  $A(U)u$ ). To prove that (7) with  $H^s$  data of size  $\epsilon$  has a solution defined on a time interval of length  $c\epsilon^{-N}$ , one has to find a modified energy  $\Theta_s(u)$  such that  $\Theta_s(u) \sim \|u\|_{H^s}^2$  close to zero and

$$(8) \quad \frac{d}{dt} \Theta_s(u(t, \cdot)) = O(\|u(t, \cdot)\|_{H^s}^{N+2})$$

when  $u$  is a solution of (7). The usual normal forms method consists in defining  $\Theta_s = \Theta_s^0 \circ \chi$ , where  $\Theta_s^0 = \|\Lambda_{\mathbf{m}}^s u\|_{L^2}^2$  is the usual  $H^s$ -energy, and  $\chi$  is a convenient symplectic diffeomorphism on a neighborhood of zero in  $H^s$ , defined as the value at time  $t = -1$  of the flow of the Hamiltonian vector field  $X_F$  of some auxiliary function  $F$ . Then equation (3) implies

$$(9) \quad \frac{d}{dt} \Theta_s(u(t, \cdot)) = \{\Theta_s^0 \circ \chi, G\}(u(t, \cdot)) = \{\Theta_s^0, G \circ \chi^{-1}\} \circ \chi(u(t, \cdot))$$

and the definition of  $\chi$  gives an expansion

$$(10) \quad \{\Theta_s^0, G \circ \chi^{-1}\} \sim \{\Theta_s^0, \sum_p \frac{(-1)^p}{p!} \text{ad}^p F \cdot G\}$$

where  $\text{ad} F \cdot G = \{F, G\}$ . One looks for  $F$  as given by an expansion in terms of homogeneous terms  $F = \sum_{\ell \geq 1} F_\ell$ . Then the term of degree of homogeneity  $p+2$  in (10) vanishes if and only if

$$(11) \quad \{\Theta_s^0, -\{F_p, G_0\} + H_p\} = 0,$$

where  $H_p$  is computed from  $G$  and  $F_{p'}$  with  $p' < p$ . One looks for  $F_p$  as

$$(12) \quad F_p(u, \bar{u}) = \int_X (B_p(U)u) \bar{u} d\mu$$

where  $B_p(U)$  is a self-adjoint para-differential operator of order one. One proves moreover that, if  $F_{p'}$  has a similar structure for  $p' < p$ , then

$$H_p(u, \bar{u}) = \int_X (A'_p(U)u) \bar{u} d\mu + \dots$$

where  $A'_p$  is also a self-adjoint operator of first order, and the dots represent some other contributions which are of a semi-linear nature. Then  $\{F_p, G_0\} - H_p$  may be written as  $\int_X (C_p(U)u) \bar{u} d\mu$  where, if we decompose

$$B_p(U) = \sum_{\ell=0}^p B_p^\ell(u, \underbrace{\dots, u}_\ell, \bar{u}, \dots, \bar{u}),$$

and make similar decompositions of  $A'_p, C_p$ ,

$$(13) \quad \begin{aligned} C_p^\ell(u, \dots, \bar{u}) &= i[B_p^\ell, \Lambda_{\mathbf{m}}](u, \dots, \bar{u}) + i \sum_{j=1}^{\ell} B_p^\ell(u, \dots, \underbrace{\Lambda_{\mathbf{m}} u}_j, \dots, \bar{u}) \\ &\quad - i \sum_{j=\ell+1}^p B_p^\ell(u, \dots, \underbrace{\Lambda_{\mathbf{m}} \bar{u}}_j, \dots, \bar{u}) - A_p'^\ell(u, \dots, \bar{u}). \end{aligned}$$

A way to solve (11) would be to find  $B_p^\ell$  so that  $C_p^\ell = 0$ . If one replaces in (13)  $(u, \dots, \bar{u})$  by  $(\Pi_{n_1} u, \dots, \Pi_{n_p} \bar{u})$ , where  $\Pi_n$  is the spectral projector associated by the  $n$ -th eigenvalue  $\lambda_n^2$  of  $-\Delta$ , one gets to show that

$$(14) \quad [B_p^\ell, \Lambda_{\mathbf{m}}](\Pi_{n_1} u, \dots, \Pi_{n_p} \bar{u}) + G_{\mathbf{m}}^{p,\ell}(n') B_p^\ell(\Pi_{n_1} u, \dots, \Pi_{n_p} \bar{u}) = -i A_p'^\ell(\Pi_{n_1} u, \dots, \Pi_{n_p} \bar{u}),$$

where  $G_{\mathbf{m}}^{p,\ell}(n') = \sum_{j=1}^{\ell} \sqrt{\mathbf{m}^2 + \lambda_{n_j}^2} - \sum_{j=\ell+1}^p \sqrt{\mathbf{m}^2 + \lambda_{n_j}^2}$ . One may choose  $\mathbf{m}$  so that, for some  $c > 0, L_0 > 0$ ,

$$(15) \quad d(\mathbb{Z}, G_{\mathbf{m}}^{p,\ell}(n')) \geq c|n'|^{-L_0} = c(n_1 + \dots + n_p)^{-L_0}$$

as soon as the left hand side does not vanish trivially (i.e. as soon as trivial two by two cancellations in the expression of  $G_{\mathbf{m}}^{p,\ell}(n')$  are excluded). If one replaces in the bracket in (14),  $\Lambda_{\mathbf{m}}$  by  $\tilde{\Lambda} = \sqrt{-\Delta + \left(\frac{d-1}{2}\right)^2}$ , an approximate solution of (14) may be obtained defining

$$B_p^\ell(\Pi_{n_1} u, \dots, \Pi_{n_p} \bar{u}) = -i \int_0^{+\infty} e^{-it\tilde{\Lambda}} A_p'^\ell(\Pi_{n_1} u, \dots, \Pi_{n_p} \bar{u}) e^{it\tilde{\Lambda}} e^{itG_{\mathbf{m}}^{p,\ell}(n')} \theta(\epsilon t) dt$$

where  $\theta \in C_0^\infty(\mathbb{R})$ ,  $\theta \equiv 1$  close to zero. Actually, using that the eigenvalues of  $\tilde{\Lambda}$  are half integers, we see that  $t \rightarrow e^{-it\tilde{\Lambda}} A_p^\ell e^{it\tilde{\Lambda}}$  is  $4\pi$ -periodic. Decomposing this function of time in Fourier series and using the small divisor estimate (15), we check that  $B_k^\ell$  is a para-differential operator, and that (14) holds, up to some remainders. Of course, this cannot be done when estimate (15) does not hold i.e. when there is a trivial vanishing of the left hand side. But the contributions corresponding to this special case are in the kernel of  $\{\Theta_s^0, \cdot\}$ , so may be discarded since the equation we need to solve is (11) and not  $\{F_p, G_0\} - H_p = 0$ .

Using repeatedly the preceding method, we may eliminate as many terms as we want in the right hand side of (10), and deduce from (9) the wanted property (8).

Let us point out that several other difficulties have to be dealt with. First, if one really wanted to define  $\chi$  as the value at time  $-1$  of the flow of  $X_F$ , with  $F = \sum_{p \geq 1} F_p$  and  $F_p$  given by (12), one would have to solve an equation  $\dot{u} = X_F(u)$  which is *not* an ODE, at the difference with the case of semi-linear equations. This comes from the fact that in (12),  $B_p$  is of order one, so that  $X_F$  involves the loss of one derivative. We circumvent this problem defining not  $\chi$  itself, but only its action by right composition on functions, using iterated Poisson brackets as in (10).

Another complication that appears when solving the “real problem”, instead of the toy model (6), is that the Hamiltonian one has to study is of the form

$$(16) \quad G(u, \bar{u}) = \int_X (\Lambda_{\mathbf{m}} u) \bar{u} d\mu + \operatorname{Re} \int_X (A(U)u) \bar{u} d\mu + \operatorname{Re} \int_X (C(U)u) u d\mu,$$

where  $C(U)$  is also a para-differential operator of order one, vanishing at  $U = 0$ . Before performing the method we outlined above, one has first to reduce oneself to an Hamiltonian of form (6). One does that in two steps. First, one eliminates the Taylor expansion of  $U \rightarrow C(U)$  at  $U = 0$  up to a large enough degree. This is done by a Birkhoff normal form method, involving functions of type (12), but with  $B_p$  of order *zero* instead of one. Because of that, this step is purely semi-linear. When this has been achieved, one obtains still a reduced Hamiltonian of form (16), but with a  $C$  vanishing at large order at  $U = 0$ . One then makes a change of unknown  $U \rightarrow \tilde{U} = \Psi(U)$ , constructed in such a way that the new Hamiltonian is a function of  $\tilde{U}$  of form (6). This part of the reasoning is similar to the usual process of diagonalization of the principal symbol of a quasi-linear symmetrizable system, that allows one to prove energy inequalities in such a framework. Actually, the construction of the change of unknown  $\Psi$  is made from a convenient diagonalization of that type.

Let us mention that we limited ourselves to polynomial non-linearities and to spheres (instead of Zoll manifolds) to avoid some extra technicalities.



## CHAPTER 1

### Statement of the main theorem

**1.1. Notations. First statement.** We denote by  $(X, g)$  the standard  $d$ -dimensional sphere ( $d \geq 2$ ), endowed with its usual metric, by  $\Delta$  the Laplace-Beltrami operator on  $X$  and by  $d\mu$  the volume form associated to the metric, given in local coordinates by  $d\mu = (\det g)^{1/2} dx$ . The eigenvalues of  $-\Delta$  are given by  $\lambda_n^2 = (n-1)(n+d-2)$ ,  $n \in \mathbb{N}^*$ . Let  $f : \mathbb{R} \times T^*X \rightarrow \mathbb{R}$  be a smooth function,  $(z, \rho) \rightarrow f(z, \rho)$ , polynomial relatively to  $z$  and to the fiber variable of  $T^*X$ . Assume that  $f$  vanishes at least at order 3 on  $\{0\} \times X$  (where  $X$  is considered as the zero section of  $T^*X$ ). For  $v : X \rightarrow \mathbb{R}$  a smooth enough function, we define from  $f, v$  a new function  $P[v] : X \rightarrow \mathbb{R}$  in the following way: If  $x$  denotes local coordinates on an open set  $U$  of  $X$ , if  $\rho = (x, \xi)$  are the corresponding local coordinates on  $T^*U$ ,  $f$  is a function of  $(z, x, \xi)$  on  $\mathbb{R} \times T^*U$ , and we set

$$(1.1.1) \quad P[v](x) = \frac{\partial f}{\partial z} \left( v(x), x, \frac{\partial v}{\partial x}(x) \right) - \sum_{j=1}^d (\det g)^{-1/2} \frac{\partial}{\partial x_j} \left[ (\det g)^{1/2} \frac{\partial f}{\partial \xi_j} \left( v(x), x, \frac{\partial v}{\partial x}(x) \right) \right].$$

We notice that  $P[v]$  is intrinsically defined. Actually, it suffices to check that, for any smooth function  $h$  on  $X$ , compactly supported in the local chart  $U$ ,  $\int_X P[v](x) h(x) d\mu(x)$  is intrinsically defined. Denote by  $\{\cdot, \cdot\}$  the Poisson bracket on  $T^*X$ , given in local coordinates by

$$\{f, g\} = \sum_{j=1}^d \left( \frac{\partial f}{\partial \xi_j} \frac{\partial g}{\partial x_j} - \frac{\partial f}{\partial x_j} \frac{\partial g}{\partial \xi_j} \right).$$

Then it follows from (1.1.1) and the definition of  $d\mu$  that

$$(1.1.2) \quad \int_X P[v] h d\mu = \int_X \frac{\partial f}{\partial z}(v, dv) h d\mu + \int_X \{f, h\}(v, dv) d\mu$$

which is intrinsic (We denote by  $dv$  the section of  $T^*X$  given in local coordinates by  $x \rightarrow (x, \frac{\partial v}{\partial x}(x))$ ).

Our main result is the following one. For  $s \in \mathbb{R}$ , we denote by  $H^s(X, \mathbb{R})$  the Sobolev space of real valued functions on  $X$ .

**THEOREM 1.1.1.** *There is a subset  $\mathcal{N}$  of zero measure in  $]0, +\infty[$  and for any  $\mathbf{m} \in ]0, +\infty[ - \mathcal{N}$ , any  $P \in \mathbb{N}$ , there is  $s_0 > 0$  such that, for any  $s \geq s_0$ , there are  $\epsilon_0 > 0$ ,  $c > 0$ ,  $C > 0$  and, for any  $\epsilon \in ]0, \epsilon_0[$ , any  $(v_0, v_1)$  in the unit ball of  $H^{s+\frac{1}{2}}(X, \mathbb{R}) \times H^{s-\frac{1}{2}}(X, \mathbb{R})$ , there is a unique solution  $v$  in  $C^0([-T_\epsilon, T_\epsilon], H^{s+\frac{1}{2}}) \cap$*

$C^1([-T_\epsilon, T_\epsilon], H^{s-\frac{1}{2}})$  of the equation

$$(1.1.3) \quad \begin{aligned} (\partial_t^2 - \Delta + \mathbf{m}^2)v &= P[v] \\ v|_{t=0} &= \epsilon v_0 \\ \partial_t v|_{t=0} &= \epsilon v_1 \end{aligned}$$

with  $T_\epsilon \geq c\epsilon^{-P}$ . Moreover,  $v$  satisfies the uniform bound

$$(1.1.4) \quad \sup_{]-T_\epsilon, T_\epsilon[} \|v(t, \cdot)\|_{H^{s+1/2}} + \sup_{]-T_\epsilon, T_\epsilon[} \|\partial_t v(t, \cdot)\|_{H^{s-1/2}} \leq C\epsilon.$$

**Remark:** The assumption on  $f$ , together with (1.1.1), shows that the right hand side of the first equation in (1.1.3) is a quasi-linear (polynomial) non-linearity, vanishing at least at order two at zero.

Expression (1.1.1) of the non-linearity will allow us to write equation (1.1.3) as a Hamiltonian system. Let us introduce the necessary notations.

Denote by  $J$  the matrix

$$(1.1.5) \quad J = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

and for  $Z, Z'$  two functions in  $L^2(X, \mathbb{R}^2)$ , define

$$(1.1.6) \quad \omega_0(Z, Z') = \langle {}^t J Z, Z' \rangle = \langle Z, J Z' \rangle,$$

where  $\langle \cdot, \cdot \rangle$  is the  $L^2(X, \mathbb{R}^2)$  scalar product given by integration against the measure  $d\mu$ . Let  $s \in \mathbb{R}_+$ ,  $\Omega$  be an open subset of  $H^s(X, \mathbb{R}^2)$ ,  $F : \Omega \rightarrow \mathbb{R}$  a  $C^1$  function. We define the symplectic gradient  $X_F(V)$  of  $F$  at  $V \in \Omega$  as the element of  $\mathcal{D}'(X, \mathbb{R}^2)$  defined by  $\omega_0(X_F(V), Z) = dF(V) \cdot Z$  for any  $Z$  in  $C^\infty(X, \mathbb{R}^2)$ . In an equivalent way

$$(1.1.7) \quad X_F(V) = J \nabla F(V).$$

If  $G : \Omega \rightarrow \mathbb{R}$  is another  $C^1$  function, we define the Poisson bracket of  $F$  and  $G$  by

$$(1.1.8) \quad \{F, G\}(V) = dF \cdot X_G(V) = dF(V) \cdot [J \nabla G(V)]$$

as soon as the right hand side has a meaning, i.e. when  $X_G(V)$  belongs to a subspace of  $\mathcal{D}'(X, \mathbb{R})$  to which the linear form  $dF(V)$  extends. Of course, this Poisson bracket should not be confused, though we use the same notation, with the Poisson bracket on  $T^*X$  used in (1.1.2).

Let us rewrite equation (1.1.3) in the Hamiltonian framework. Let

$$(1.1.9) \quad \Lambda_{\mathbf{m}} = \sqrt{-\Delta + \mathbf{m}^2}$$

and define from the function  $v$  solving (1.1.3), an element  $V$  of  $H^s(X, \mathbb{R}^2)$  by

$$(1.1.10) \quad V(t, x) = \begin{bmatrix} \Lambda_{\mathbf{m}}^{-1/2} \partial_t v \\ \Lambda_{\mathbf{m}}^{1/2} v \end{bmatrix} = \begin{bmatrix} V^1 \\ V^2 \end{bmatrix}.$$

If we set  $F(v) = \int_X f(v, dv) d\mu$ , and if  $h \in C^\infty(X, \mathbb{R})$ ,

$$dF(v) \cdot h = \int_X P[v] h d\mu$$

as follows from (1.1.1), taking  $h$  supported in a local chart domain. We define for  $V$  in  $H^s(X, \mathbb{R}^2)$ , with  $s$  large enough

$$(1.1.11) \quad \begin{aligned} G_0(V) &= \frac{1}{2} \int_X (\Lambda_{\mathbf{m}} V) \cdot V \, d\mu \\ G(V) &= G_0(V) - F(\Lambda_{\mathbf{m}}^{-1/2} V^2). \end{aligned}$$

Then, if  $H$  is in  $C^\infty(M, \mathbb{R}^2)$ ,

$$dG(V) \cdot H = \int_X (\Lambda_{\mathbf{m}} V) \cdot H \, d\mu - \int_X P[\Lambda_{\mathbf{m}}^{-1/2} V^2](\Lambda_{\mathbf{m}}^{-1/2} H^2) \, d\mu$$

which shows that

$$\nabla G(V) = \Lambda_{\mathbf{m}} V + \begin{bmatrix} 0 \\ -\Lambda_{\mathbf{m}}^{-1/2} P[\Lambda_{\mathbf{m}}^{-1/2} V^2] \end{bmatrix}.$$

Using (1.1.10), we see that  $v$  is a solution of the first equation in (1.1.3) if and only if  $V$  satisfies

$$(1.1.12) \quad \partial_t V = X_G(V).$$

Let us write this equality using complex coordinates. We identify  $H^s(X, \mathbb{R}^2)$  to  $H^s(X) \stackrel{\text{def}}{=} H^s(X, \mathbb{C})$  through

$$(1.1.13) \quad V = \begin{bmatrix} V^1 \\ V^2 \end{bmatrix} \rightarrow u = \frac{\sqrt{2}}{2}(V^1 + iV^2).$$

More precisely, we identify  $H^s(X, \mathbb{R}^2)$  to the submanifold  $\{(U^1, U^2); U^2 = \overline{U^1}\}$  inside the product  $H^s(X, \mathbb{C}) \times H^s(X, \mathbb{C})$  through

$$(1.1.14) \quad V = \begin{bmatrix} V^1 \\ V^2 \end{bmatrix} \rightarrow U = \begin{bmatrix} u = \frac{\sqrt{2}}{2}(V^1 + iV^2) \\ \bar{u} = \frac{\sqrt{2}}{2}(V^1 - iV^2) \end{bmatrix}.$$

If  $F$  is a  $C^1$  function on an open subset of  $H^s(X, \mathbb{R}^2)$ , we define

$$(1.1.15) \quad \begin{aligned} d_u F &= \frac{\sqrt{2}}{2} [d_{V^1} F - i d_{V^2} F], \quad d_{\bar{u}} F = \frac{\sqrt{2}}{2} [d_{V^1} F + i d_{V^2} F] \\ \nabla_u F &= \frac{\sqrt{2}}{2} [\nabla_{V^1} F - i \nabla_{V^2} F], \quad \nabla_{\bar{u}} F = \frac{\sqrt{2}}{2} [\nabla_{V^1} F + i \nabla_{V^2} F]. \end{aligned}$$

If  $\begin{bmatrix} H_1 \\ H_2 \end{bmatrix}$  (resp.  $\begin{bmatrix} H'_1 \\ H'_2 \end{bmatrix}$ ), element of  $H^s(X, \mathbb{R}^2)$ , is identified through (1.1.14) to the element  $H = \begin{bmatrix} h \\ \bar{h} \end{bmatrix}$  (resp.  $H' = \begin{bmatrix} h' \\ \bar{h}' \end{bmatrix}$ ) of  $H^s(X)$ , the symplectic form in complex coordinates may be written

$$\omega_0(H, H') = i \langle H, JH' \rangle.$$

Through identification (1.1.14), the expression of the Hamiltonian vector field in complex coordinates is given by

$$(1.1.16) \quad X_F(U) = -iJ \begin{bmatrix} \nabla_u F \\ \nabla_{\bar{u}} F \end{bmatrix} = i^t J \begin{bmatrix} \nabla_u F \\ \nabla_{\bar{u}} F \end{bmatrix} = i \begin{bmatrix} \nabla_{\bar{u}} F \\ -\nabla_u F \end{bmatrix}.$$

If  $F$  and  $G$  are two  $C^1$  functions on an open set of  $H^s(X)$ , whose differentials at every point extend as bounded maps on  $L^2(X)$ , one gets the following expression for their Poisson brackets in complex coordinates

$$(1.1.17) \quad \{F, G\} = dF \cdot X_G(U) = i[d_u F \cdot \nabla_{\bar{u}} G - d_{\bar{u}} F \cdot \nabla_u G].$$



Finally, equation (1.1.12) may be written in an equivalent way

$$(1.1.18) \quad \partial_t u = i \nabla_{\bar{u}} G(u, \bar{u})$$

if we consider the function given by (1.1.11) as a function of the complex variable  $U = (u, \bar{u})$ .

**1.2. Quasi-linear Hamiltonians and general statements.** The goal of this subsection is to introduce a general class of Hamiltonian equations, containing (1.1.11), such that the associated equation (1.1.12) or (1.1.18) is a quasi-linear first order system. This class has to be wide enough to be stable under the reductions that will have to be made in the proof of Theorem 1.1.1.

Let us introduce some notation. We denote by  $\lambda_n = \sqrt{(n-1)(n+d-2)}$ ,  $n \in \mathbb{N}^*$ , the eigenvalues of  $\sqrt{-\Delta}$  on  $X = \mathbb{S}^d$ . We call  $\Pi_n$  the spectral projector associated to  $\lambda_n$ . In particular, for any  $s \in \mathbb{R}$ , there is  $C > 0$  such that, for any  $u$  in  $H^s(X)$ ,

$$(1.2.1) \quad C^{-1} n^s \|\Pi_n u\|_{L^2} \leq \|\Pi_n u\|_{H^s} \leq C n^s \|\Pi_n u\|_{L^2}.$$

Fix  $\varphi$  a non-negative real valued smooth function on  $\mathbb{R}$ , compactly supported in  $]0, +\infty[$ , such that  $\sum_{j=-\infty}^{+\infty} \varphi(2^{-j}t) \equiv 1$  for any  $t > 0$ . We define  $\psi(t) = \sum_{j=-\infty}^0 \varphi(2^{-j}t)$  and we set for a distribution  $u$  on  $X$  and a natural integer  $j$ ,

$$(1.2.2) \quad \begin{aligned} \Delta_j u &= \sum_{n=1}^{+\infty} \varphi(2^{-j} \lambda_n) \Pi_n u, \quad j \geq 1 \\ \Delta_0 u &= S_0 u = \sum_{n=1}^{+\infty} \psi(\lambda_n) \Pi_n u, \\ S_j u &= \sum_{j'=0}^j \Delta_{j'} u, \quad j \geq 0. \end{aligned}$$

In that way,  $u$  is in  $H^s(X)$  if and only if  $(2^{-js} \|\Delta_j u\|_{L^2})_j$  is in  $\ell^2(\mathbb{N})$ .

We shall study an equation of the form of (1.1.18) where the Hamiltonian  $G$  will be expressed in terms of para-differential operators acting on  $U = \begin{bmatrix} u \\ \bar{u} \end{bmatrix}$ . We introduce several classes of multi-linear operators. If  $p \in \mathbb{N}^*$ , if  $\mathcal{U} = (u_1, \dots, u_p)$  is a  $p$ -tuple of complex valued functions defined on  $X$  and if  $n' = (n_1, \dots, n_p)$  is an element of  $(\mathbb{N}^*)^p$ , we denote

$$(1.2.3) \quad \Pi_{n'} \mathcal{U} = (\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p), \quad |n'| = \max(n_1, \dots, n_p).$$

We assume given a strictly increasing function  $\nu : \mathbb{N} \rightarrow \mathbb{N}$ , such that  $\nu(0) = 0$ , that satisfies a growth condition of type

$$(1.2.4) \quad \nu(p) + \nu(q) + a \leq \nu(p+q), \quad p, q \in \mathbb{N}^*,$$

where  $a$  is some positive constant that will be adjusted later.

**DEFINITION 1.2.1.** Let  $p \in \mathbb{N}, m \in \mathbb{R}, \nu \in \mathbb{R}_+$ . One denotes by  $\tilde{\mathcal{P}}_p^{m, \nu}$  the space of  $p$ -linear maps  $\mathcal{U} = (u_1, \dots, u_p) \rightarrow A(\mathcal{U})$ , defined on  $C^\infty(X)^p$  with values in the space of linear maps from  $C^\infty(X)$  to  $\mathcal{D}'(X)$ , such that, for any  $N_1, N_2$  in  $\mathbb{N}$ , there

is  $C > 0$  so that, for any  $\mathcal{U} = (u_1, \dots, u_p)$ , any  $(n_0, \dots, n_{p+1})$  in  $(\mathbb{N}^*)^{p+2}$ ,

$$(1.2.5) \quad \begin{aligned} \|\Pi_{n_0} A(\Pi_{n'} \mathcal{U}) \Pi_{n_{p+1}}\|_{\mathcal{L}(L^2)} &\leq C \langle n_0 - n_{p+1} \rangle^{-2} \left( \min \left[ \frac{n_0}{n_{p+1}}, \frac{n_{p+1}}{n_0} \right] \right)^{N_1} n_0^m \\ &\times |n'|^{\nu(p)+\nu} \left( 1 + \frac{|n'|}{n_0} \right)^{-N_2} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}, \end{aligned}$$

where  $\langle n \rangle = (1 + n^2)^{1/2}$ .

**Remarks:** • The definition implies that for any  $j, j'$  in  $\mathbb{N}$

$$(1.2.6) \quad \|\Delta_j A(\Pi_{n'} \mathcal{U}) \Delta_{j'}\|_{\mathcal{L}(L^2)} \leq C 2^{-N_1 |j-j'|} 2^{jm} |n'|^{\nu(p)+\nu} (1 + 2^{-j} |n'|)^{-N_2} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

The characterization of Sobolev spaces in terms of dyadic decompositions implies that, for any  $s$  in  $\mathbb{R}$ , any  $N_2 \in \mathbb{N}$ ,  $|n'|^{N_2} A(\Pi_{n'} \mathcal{U})$  extends as a bounded linear map from  $H^s(X)$  to  $H^{s-m-N_2}(X)$ , with estimates

$$(1.2.7) \quad |n'|^{N_2} \|A(\Pi_{n'} \mathcal{U})\|_{\mathcal{L}(H^s, H^{s-m-N_2})} \leq C_{s, N_2} |n'|^{\nu(p)+\nu} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

In particular, if  $u_\ell \in H^\sigma(X)$ ,  $\ell = 1, \dots, p$ , with  $\sigma > \nu(p) + \nu + \frac{1}{2}$ ,  $A(\mathcal{U}) = \sum_{n'} A(\Pi_{n'} \mathcal{U})$  defines an element of  $\mathcal{L}(H^s, H^{s-m})$  for any  $s$  in  $\mathbb{R}$ .

• Estimates (1.2.6), (1.2.7) assert that elements of  $\widehat{\mathcal{P}}_p^{0, \nu}$  are bounded from  $H^s$  to  $H^s$  for any  $s$ , and that in (1.2.6), the coefficients  $\mathcal{U}$  are spectrally localized essentially for frequencies  $\lambda_{n_1}, \dots, \lambda_{n_p}$  satisfying  $|n'| = \max(n_1, \dots, n_p) \leq 2^j$ . Such properties are to be expected from para-differential operators. Nevertheless, they do not suffice to give a class of operators enjoying a symbolic calculus. To define a true class of para-differential operators, we shall not use symbols, but instead a formulation in terms of commutators with differential operators, similar to the Beals characterization of pseudo-differential operators [4]. In that way, we can give a global definition on the manifold.

If  $A, P$  are two operators, we set  $\text{Ad}_P A = [P, A]$ .

**DEFINITION 1.2.2.** We assume given a real number  $M_0 \geq 1$ . For  $p \in \mathbb{N}, m \in \mathbb{R}, \nu \in \mathbb{R}_+$ , we denote by  $\widetilde{H}_p^{m, \nu}$  the space of  $p$ -linear maps  $\mathcal{U} = (u_1, \dots, u_p) \rightarrow A(\mathcal{U})$ , defined on  $C^\infty(X)^p$  with values in the space of linear maps from  $C^\infty(X)$  to  $\mathcal{D}'(X)$ , such that, for any family  $(P_1, \dots, P_k)$  of differential operators on  $X$ , of respective orders  $d_1, \dots, d_k$ , for any  $N_1, N_2 \in \mathbb{N}$ , there is  $C > 0$  and, for any  $\mathcal{U}$  in  $C^\infty(X)^p$ , any  $n' = (n_1, \dots, n_p) \in (\mathbb{N}^*)^p$ , any  $j, j'$  in  $\mathbb{N}$

$$(1.2.8) \quad \begin{aligned} \|\Delta_j \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} A(\Pi_{n'} \mathcal{U}) \Delta_{j'}\|_{\mathcal{L}(L^2)} &\leq C 2^{-N_1 |j-j'|} 2^{j(m + \sum_{\ell=1}^k d_\ell - k)} |n'|^{\nu(p)+\nu+M_0 k} \\ &\times (1 + 2^{-j} |n'|)^{-N_2} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}. \end{aligned}$$

If we want to make explicit in the notation the constant  $M_0$  in the above estimate, we use the notation  $\widetilde{H}_p^{m, \nu}[M_0]$  for the preceding space.

**Remarks:** • In comparison with (1.2.6), we see that (1.2.8) gains  $-1$  on the order of the operator every time we make act a commutator with a  $P_{k'}$ . This gain

is traded against a loss on the smoothness of the coefficients  $\mathcal{U}$ , given by the power  $M_0 k$  of  $|n'|$ . In general,  $M_0$  will be a constant strictly larger than one.

- It follows from the definition that  $\widetilde{H\Psi}_p^{m-1,\nu+1} \subset \widetilde{H\Psi}_p^{m,\nu}$ .
- If  $A$  is in  $\widetilde{H\Psi}_p^{m,\nu}$ , it follows from the definition that for any  $P_1, \dots, P_k$  as in (1.2.8), any  $N_2 \in \mathbb{N}$ , any  $s \in \mathbb{R}$ , there is  $C > 0$ , and for any  $\mathcal{U}$  in  $C^\infty(X)^p$ , any  $n'$  in  $(\mathbb{N}^*)^p$

$$(1.2.9) \quad |n'|^{N_2} \|\text{Ad}_{P_1} \cdots \text{Ad}_{P_k} A(\Pi_{n'} \mathcal{U})\|_{\mathcal{L}(H^s, H^{s-m-N_2-\sum_1^k d_\ell+k})} \leq C_{s,N_2} |n'|^{\nu(p)+\nu+M_0 k} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

Conversely, if such an estimate holds for any  $P_1, \dots, P_k, s, N_2$ , then  $A$  satisfies (1.2.8). This equivalent characterization shows in particular that, if  $\theta$  is in  $C^\infty(X)$  and  $A$  is in  $\widetilde{H\Psi}_p^{m,\nu}$ , then  $\theta A$  and  $A\theta$  are in  $\widetilde{H\Psi}_p^{m,\nu}$ . Moreover, estimate (1.2.9) with  $N_2 = 0, k = 0$  shows that if  $u_\ell$  is in  $H^\sigma(X)$  with  $\sigma > \nu(p) + \nu + \frac{1}{2}$  for  $\ell = 1, \dots, p$ , then  $\mathcal{U} \rightarrow A(\mathcal{U})$  extends as a continuous  $p$ -linear map from  $\widetilde{H}^\sigma \times \cdots \times \widetilde{H}^\sigma$  to  $\mathcal{L}(H^s, H^{s-m})$  for any  $s$ .

- One has an inclusion

$$(1.2.10) \quad \widetilde{H\Psi}_p^{m,\nu} \subset \widetilde{\mathcal{P}}_p^{m,\nu+2M_0}.$$

Actually, we may write

$$(\lambda_{n_0}^2 - \lambda_{n_{p+1}}^2)^2 \Pi_{n_0} A(\Pi_{n'} \mathcal{U}) \Pi_{n_{p+1}} = \Pi_{n_0} \text{Ad}_\Delta \text{Ad}_\Delta A(\Pi_{n'} \mathcal{U}) \Pi_{n_{p+1}}$$

so that (1.2.8) implies estimates (1.2.9) with  $\nu$  replaced by  $\nu + 2M_0$ .

- Note that it suffices to assume that (1.2.8) or (1.2.9) holds when the orders  $d_1, \dots, d_k$  of  $P_1, \dots, P_k$  are zero or one. Actually, any differential operator on  $X$  of order  $r \geq 1$  may be written as a finite linear combination with smooth coefficients of expressions  $X_1 \cdots X_{r'}$ , with  $r' \leq r$  and  $X_j$  smooth vector field on  $X$ . This allows one to deduce (1.2.9) (and so (1.2.8)) for general  $P_\ell$ 's from the estimates corresponding to operators of order zero or one.

We define from the preceding classes operators whose coefficients are given in terms of a single function  $U = (u, \bar{u})$  instead of a  $p$ -tuple of functions.

**DEFINITION 1.2.3.** Let  $m \in \mathbb{R}, \nu \in \mathbb{R}_+, p \in \mathbb{N}$ . One denotes by  $H\Psi_p^{m,\nu}$  (resp.  $\mathcal{P}_p^{m,\nu}$ ) the space of functions  $U \rightarrow A(U)$ , defined for  $U = (u, \bar{u})$  in  $C^\infty(X)$ , with values in the vector space of linear maps from  $C^\infty(X)$  to  $\mathcal{D}'(X)$ , such that there is a family  $A_j$  of elements of  $\widetilde{H\Psi}_p^{m,\nu}$  (resp.  $\widetilde{\mathcal{P}}_p^{m,\nu}$ ),  $0 \leq j \leq p$ , so that

$$(1.2.11) \quad A(U) = \sum_{j=0}^p A_j(\underbrace{u, \dots, u}_j, \underbrace{\bar{u}, \dots, \bar{u}}_{p-j}).$$

As in the case of multi-linear operators, we see that if  $\sigma > \nu(p) + \nu + \frac{1}{2}$ , any element  $U \rightarrow A(U)$  of  $H\Psi_p^{m,\nu}$  or  $\mathcal{P}_p^{m,\nu}$  extends as a continuous map from  $\widetilde{H}^\sigma(X)$  to  $\mathcal{L}(H^s, H^{s-m})$  for any  $s$ .

We denote by  $H\Psi_p^{m,\nu}[M_0]$  the same space as above, when we want to make explicit the constant  $M_0$  used in the definition of  $\widetilde{H\Psi}_p^{m,\nu}[M_0]$ .

We may deduce from the definitions of the preceding classes composition results.

LEMMA 1.2.4. *Let  $m, m'$  be in  $\mathbb{R}$ ,  $\nu, \nu'$  be in  $\mathbb{R}_+$ ,  $p, q$  be in  $\mathbb{N}$ . Assume that (1.2.4) is satisfied for some  $a \leq \nu + \nu'$ . The following inclusions hold:*

$$(1.2.12) \quad \begin{aligned} \widetilde{H\Psi}_p^{m,\nu}[M_0] \circ \widetilde{H\Psi}_q^{m',\nu'}[M_0] &\subset \widetilde{H\Psi}_{p+q}^{m+m',\nu+\nu'-a}[M_0] \\ H\Psi_p^{m,\nu}[M_0] \circ H\Psi_q^{m',\nu'}[M_0] &\subset H\Psi_{p+q}^{m+m',\nu+\nu'-a}[M_0] \end{aligned}$$

when  $p > 0$  and  $q > 0$ ,

$$(1.2.13) \quad \begin{aligned} \widetilde{H\Psi}_p^{m,\nu}[M_0] \circ \widetilde{H\Psi}_q^{m',\nu'}[M_0] &\subset \widetilde{H\Psi}_{p+q}^{m+m',\nu+\nu'}[M_0] \\ H\Psi_p^{m,\nu}[M_0] \circ H\Psi_q^{m',\nu'}[M_0] &\subset H\Psi_{p+q}^{m+m',\nu+\nu'}[M_0] \end{aligned}$$

if  $p = 0$  or  $q = 0$ ,

$$(1.2.14) \quad \begin{aligned} \widetilde{H\Psi}_p^{m,\nu}[M_0] \circ \widetilde{\mathcal{P}}_q^{m',\nu'} &\subset \widetilde{\mathcal{P}}_{p+q}^{m+m',\nu+\nu'-a+2M_0} \\ \widetilde{\mathcal{P}}_p^{m,\nu} \circ \widetilde{H\Psi}_q^{m',\nu'}[M_0] &\subset \widetilde{\mathcal{P}}_{p+q}^{m+m',\nu+\nu'-a+2M_0} \\ \widetilde{\mathcal{P}}_p^{m,\nu} \circ \widetilde{\mathcal{P}}_q^{m',\nu'} &\subset \widetilde{\mathcal{P}}_{p+q}^{m+m',\nu+\nu'-a} \end{aligned}$$

if  $p > 0$  and  $q > 0$ . If  $p = 0$  or  $q = 0$ , one has similar inclusions with  $a = 0$  in the exponents of the right hand side. Finally, (1.2.14) holds also replacing everywhere  $\widetilde{H\Psi}$  (resp.  $\widetilde{\mathcal{P}}$ ) by  $H\Psi$  (resp.  $\mathcal{P}$ ).

*Proof:* To prove (1.2.12), one notices that if  $\mathcal{U}' = (u_1, \dots, u_p)$ ,  $\mathcal{U}'' = (u_{p+1}, \dots, u_{p+q})$ ,  $\text{Ad}_{P_1} \cdots \text{Ad}_{P_K}[A(\mathcal{U}') \circ B(\mathcal{U}'')]$  may be expressed from

$$(\text{Ad}_{P_{i_1}} \cdots \text{Ad}_{P_{i_\ell}} A(\mathcal{U}')) \circ (\text{Ad}_{P_{j_1}} \cdots \text{Ad}_{P_{j_{\ell'}}} B(\mathcal{U}''))$$

where  $\ell + \ell' = k$ ,  $\{i_1, \dots, i_\ell\} \cup \{j_1, \dots, j_{\ell'}\} = \{1, \dots, k\}$ . One has just to apply characterization (1.2.9) of the  $\widetilde{H\Psi}$ -spaces, together with (1.2.4), to obtain the first inclusion in (1.2.12), (1.2.13). The second inclusion in those formulas follows from the first one.

To prove the last inclusion (1.2.14), one writes

$$\Pi_{n_0} A(\Pi_{n'} \mathcal{U}') \circ B(\Pi_{n''} \mathcal{U}'') \Pi_{n_{p+1}} = \sum_n (\Pi_{n_0} A(\Pi_{n'} \mathcal{U}') \Pi_n) \circ (\Pi_n B(\Pi_{n''} \mathcal{U}'') \Pi_{n_{p+1}}),$$

and uses (1.2.5). The first and the second inclusions (1.2.5) follow from the third one and from (1.2.10). This concludes the proof.  $\square$

We have defined up to now operators homogeneous of order  $p$  in some function  $U$ . We shall need as well similar classes for which the  $U$ -dependence will be only  $C^1$ , with some vanishing when  $U \rightarrow 0$ . If  $\sigma$  is a real number (that will be large enough in practice), we set for  $R > 0$ ,

$$(1.2.15) \quad B_\sigma(R) = \{U \in H^\sigma(X); \|U\|_{H^\sigma} < R\}.$$

DEFINITION 1.2.5. *Let  $m \in \mathbb{R}, \nu \in \mathbb{R}_+, p \in \mathbb{N}, \sigma > 0, R > 0$ . One denotes by  $\Psi_p^{m,\nu}(B_\sigma(R))$  the space of maps  $U = (u, \bar{u}) \rightarrow A(U)$ , defined on  $B_\sigma(R)$ , with values in the space of linear maps from  $C^\infty(X)$  to  $\mathcal{D}'(X)$ , satisfying the following conditions:*

(i) *For any family  $P_1, \dots, P_k$  of differential operators of orders  $d_1, \dots, d_k$  on  $X$ , for any  $N \in \mathbb{N}$ , there is  $C > 0$  such that, for any  $U \in B_\sigma(R)$ , any  $j, j' \in \mathbb{N}$*

$$(1.2.16) \quad \|\Delta_j \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} A(U) \Delta_{j'}\|_{\mathcal{L}(L^2)} \leq C 2^{-N|j-j'|} 2^{j[m + \sum_{\ell=1}^k d_\ell - k + (M_0 k + \nu(p) + \nu - \sigma)_+]} \|U\|_{H^\sigma}^p.$$

(ii) For any  $j, j', P_1, \dots, P_k$  as above, the map  $U \rightarrow \Delta_j \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} A(\Pi_n \mathcal{U}) \Delta_{j'}$  is  $C^1$  on  $B_\sigma(R)$  with values in  $\mathcal{L}(L^2)$  and, for any  $\sigma'$  with  $|\sigma'| \leq \sigma$ , the differential in  $U$  extends as a linear map from  $H^{\sigma'}(X)$  to  $\mathcal{L}(L^2)$  such that, for any  $N \in \mathbb{N}$ , there is  $C > 0$  so that, for any  $U \in B_\sigma(R)$ , any  $j, j' \in \mathbb{N}$ , any  $H$  in  $H^{\sigma'}(X)$

$$(1.2.17) \quad \|\Delta_j \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} (\partial_U A(U) \cdot H) \Delta_{j'}\|_{\mathcal{L}(L^2)} \\ \leq C 2^{-N|j-j'|} 2^{j[m+\sum_{\ell=1}^k d_\ell - k + (M_0 k + \nu(p) + \nu - \sigma')_+]} \|U\|_{H^\sigma}^{p-1} \|H\|_{H^{\sigma'}}.$$

**Remarks:** • We may write equivalent formulations of (1.2.16), (1.2.17), asking that for any  $s \in \mathbb{R}$ , any  $P_1, \dots, P_k$  as in the statement of the definition, any  $\sigma'$  with  $|\sigma'| \leq \sigma$ , any  $U$  in  $B_\sigma(R)$ , any  $H$  in  $H^{\sigma'}$ ,

$$(1.2.18) \quad \|\text{Ad}_{P_1} \cdots \text{Ad}_{P_k} A(U)\|_{\mathcal{L}(H^s, H^{s-(m+\sum d_\ell - k) - (M_0 k + \nu(p) + \nu - \sigma')_+})} \leq C \|U\|_{H^\sigma}^p$$

and

$$(1.2.19) \quad \|\text{Ad}_{P_1} \cdots \text{Ad}_{P_k} (\partial_U A(U) \cdot H)\|_{\mathcal{L}(H^s, H^{s-(m+\sum d_\ell - k) - (M_0 k + \nu(p) + \nu - \sigma')_+})} \leq C \|U\|_{H^\sigma}^{p-1} \|H\|_{H^{\sigma'}}.$$

• In (1.2.18), (1.2.19) it would be enough to assume that these inequalities hold for all  $k$  such that  $M_0 k + \nu(p) + \nu - \sigma < 1$ . The inequalities for larger values of  $k$  will follow from the fact that  $P_\ell$  is bounded from  $H^s$  to  $H^{s-d_\ell}$  for any  $s$  and the assumption  $M_0 \geq 1$ .

• As in the case of the preceding multi-linear classes, the space  $\Psi_p^{m,\nu}(B_\sigma(R))$  is stable by multiplication at the left or at the right by functions in  $C^\infty(X)$ . Moreover, it is enough to assume conditions (1.2.16), (1.2.17) (or (1.2.18), (1.2.19)) for operators  $P_\ell$  of order zero or one.

• It follows from Definition 1.2.5 and from (1.2.8) that if  $q \geq p$  and if  $\nu'$  is any number with  $\nu' > \nu + \nu(q) - \nu(p) + \frac{1}{2}$ ,  $H\Psi_q^{m,\nu}$  is contained in  $\Psi_p^{m,\nu'}(B_\sigma(R))$  for any  $\sigma > 0$ , any  $R > 0$  (using characterization (1.2.1) of Sobolev norms).

Let us state a composition result for the operators of the class we just defined, similar to Lemma 1.2.4. We shall need later on to control some semi-norms that we introduce before the statement. If  $\mathcal{P} = (P_1, \dots, P_k)$  is a collection of differential operators of orders  $d_1, \dots, d_k$ , if  $\sigma \in \mathbb{R}$ , we set  $\text{Ad}_{\mathcal{P}} = \text{Ad}_{P_1} \cdots \text{Ad}_{P_k}$ ,

$$(1.2.20) \quad \alpha(\mathcal{P}, p, \nu, \sigma) = \sum_{\ell=1}^k d_\ell - k + (M_0 k + \nu(p) + \nu - \sigma)_+.$$

If  $A$  is in  $\Psi_p^{m,\nu}(B_\sigma(R))$  and  $\sigma'$  is in  $[-\sigma, \sigma]$ , we define

$$(1.2.21) \quad \mathfrak{N}_{(\mathcal{P}, p, \nu, \sigma)}^{0, m, s}(A) = \sum_{\mathcal{P}' \subset \mathcal{P}} \|\text{Ad}_{\mathcal{P}'} A(U)\|_{\mathcal{L}(H^s, H^{s-m-\alpha(\mathcal{P}', p, \nu, \sigma)})} \\ \mathfrak{N}_{(\mathcal{P}, p, \nu, \sigma')}^{1, m, s}(A) = \sum_{\mathcal{P}' \subset \mathcal{P}} \sup_{\|H\|_{H^{\sigma'}} \leq 1} \|\text{Ad}_{\mathcal{P}'} (\partial_u A(U) \cdot H)\|_{\mathcal{L}(H^s, H^{s-m-\alpha(\mathcal{P}', p, \nu, \sigma')})}$$

where the sums are taken over all sub-families of  $\mathcal{P}$ .

**LEMMA 1.2.6.** *Let  $m, m' \in \mathbb{R}$ ,  $\nu, \nu' \in \mathbb{R}_+$ ,  $p, p' \in \mathbb{N}$ ,  $R > 0$ . Let  $\sigma > 0$  satisfying the inequality  $\sigma \geq \min(\nu(p), \nu(p')) + \min(\nu, \nu')$  and let  $U \rightarrow A(U)$  (resp.  $U \rightarrow B(U)$ ) be in  $\Psi_p^{m,\nu}(B_\sigma(R))$  (resp.  $\Psi_{p'}^{m',\nu'}(B_\sigma(R))$ ). Then  $U \rightarrow A(U) \circ B(U)$*

is in  $\Psi_{\max(p,p')}^{m+m', \max(\nu, \nu')}(B_\sigma(R))$ . Moreover, for any  $\mathcal{P} = (P_1, \dots, P_k)$  as above, there is a constant  $C > 0$ , depending only on  $\mathcal{P}$ ,  $R, p, p', \nu, \nu', m, m', s, \sigma' \in [-\sigma, \sigma]$  such that, if  $p'' = \max(p, p')$ ,  $\nu'' = \max(\nu, \nu')$

(1.2.22)

$$\begin{aligned} \mathfrak{N}_{(\mathcal{P}, p'', \nu'', \sigma)}^{0, m+m', s}(A \circ B) &\leq C \mathfrak{N}_{(\mathcal{P}, p, \nu, \sigma)}^{0, m, s}(A) \mathfrak{N}_{(\mathcal{P}, p', \nu', \sigma)}^{0, m', s}(B) \\ \mathfrak{N}_{(\mathcal{P}, p'', \nu'', \sigma')}^{1, m+m', s}(A \circ B) &\leq C [\mathfrak{N}_{(\mathcal{P}, p, \nu, \sigma)}^{0, m, s}(A) \mathfrak{N}_{(\mathcal{P}, p', \nu', \sigma')}^{1, m', s}(B) + \mathfrak{N}_{(\mathcal{P}, p, \nu, \sigma')}^{1, m, s}(A) \mathfrak{N}_{(\mathcal{P}, p', \nu', \sigma)}^{0, m', s}(B)]. \end{aligned}$$

*Proof:* We may write  $\text{Ad}_{\mathcal{P}}(A(U) \circ B(U))$  as a linear combination of quantities of the form  $(\text{Ad}_{\mathcal{P}'} A(U)) \circ (\text{Ad}_{\mathcal{P}''} B(U))$  where  $(\mathcal{P}', \mathcal{P}'')$  is a partition of  $\mathcal{P}$ . If  $k'$  (resp.  $k''$ , resp.  $k$ ) is the cardinal of  $\mathcal{P}'$  (resp.  $\mathcal{P}''$ , resp.  $\mathcal{P}$ ), we shall deduce estimates (1.2.18), (1.2.19) for the composition from the corresponding estimates for each factor, if we have the inequality

$$\alpha(\mathcal{P}', p, \nu, \sigma) + \alpha(\mathcal{P}'', p', \nu', \sigma') \leq \alpha(\mathcal{P}, \max(p, p'), \max(\nu, \nu'), \sigma')$$

for  $|\sigma'| \leq \sigma$  (and the same inequality with  $\sigma$  and  $\sigma'$  exchanged in the left hand side). This follows from

$$(M_0 k' + \nu(p) + \nu - \sigma)_+ + (M_0 k'' + \nu(p') + \nu' - \sigma')_+ \leq (M_0 k + \max(\nu(p), \nu(p')) + \max(\nu, \nu') - \sigma')_+$$

which is a consequence of the assumptions on  $\sigma, \sigma'$ . Inequalities (1.2.22) follow from the proof.  $\square$

To state a more general version of Theorem 1.1.1, we introduce classes of functions defined on  $B_\sigma(R)$ , in terms of the preceding classes of operators.

DEFINITION 1.2.7. Let  $m \in \mathbb{R}, \nu \in \mathbb{R}_+, p \in \mathbb{N}, \sigma \in \mathbb{R}, R > 0$ . Assume

$$(1.2.23) \quad \sigma > \nu(p) + \nu + \frac{1}{2}, \quad 2\sigma \geq m.$$

(i) One denotes by  $H\mathcal{F}_{p, E}^{m, \nu}(B_\sigma(R))$  (resp.  $H\mathcal{F}_{p, H}^{m, \nu}(B_\sigma(R))$ , resp.  $H\mathcal{F}_p^{m, \nu}(B_\sigma(R))$ ) the space of functions  $F$  defined on  $B_\sigma(R)$ , real valued, that may be written, for some element  $A_p$  of  $H\Psi_p^{m, \nu}$  as

$$(1.2.24) \quad F(U) = \text{Re} \int_X [A_p(U)u]u \, d\mu$$

resp. as

$$(1.2.25) \quad F(U) = \text{Re} \int_X [A_p(U)u]\bar{u} \, d\mu,$$

resp. as the sum of an element of  $H\mathcal{F}_{p, E}^{m, \nu}(B_\sigma(R))$  and an element of  $H\mathcal{F}_{p, H}^{m, \nu}(B_\sigma(R))$ . One uses the notation  $H\mathcal{F}_p^{m, \nu}(B_\sigma(R))[M_0]$  when one wants to keep track of the constant  $M_0$  as in the definition of  $H\Psi_p^{m, \nu}[M_0]$ .

(ii) One denotes by  $\mathcal{F}_{p, E}^{m, \nu}(B_\sigma(R))$  (resp.  $\mathcal{F}_{p, H}^{m, \nu}(B_\sigma(R))$ , resp.  $\mathcal{F}_p^{m, \nu}(B_\sigma(R))$ ) the space of functions defined on  $B_\sigma(R)$  by similar expressions as above, but with  $A_p$  in  $\Psi_p^{m, \nu}(B_\sigma(R))$ .

(iii) One denotes by  $\mathcal{G}_p^{m, \nu}(B_\sigma(R))$  the space of functions defined on  $B_\sigma(R)$  as  $\mathcal{F}_p^{m, \nu}(B_\sigma(R))$ , but with  $A_p$  in  $\mathcal{P}_p^{m, \nu}$ .

**Remarks:** • We shall call elements of  $H\mathcal{F}_{p,E}^{m,\nu}(B_\sigma(R))$  (resp.  $H\mathcal{F}_{p,H}^{m,\nu}(B_\sigma(R))$ ) elliptic (resp. hyperbolic) elements.

• Since  $A_p(U)$  sends  $H^\sigma$  to  $H^{\sigma-m}$ , the second assumption (1.2.23) shows that (1.2.24), (1.2.25) make sense.

• It follows from Definition 1.2.3 and inequality (1.2.7) that if  $F$  is in  $\mathcal{G}_p^{m,\nu}(B_\sigma(R))$  and  $\sigma$  is large enough relatively to  $m, \nu$ , then  $\nabla F$  is in  $H^{\sigma-m}$ .

Let us define the set  $\mathcal{N}$  of exceptional values of the mass that have to be excluded. For  $\mathbf{m} \in ]0, +\infty[$ , define if  $p \in \mathbb{N}$ ,  $0 \leq \ell \leq p$ ,  $\xi_j \in \mathbb{R}$  for  $0 \leq j \leq p+1$ ,

$$(1.2.26) \quad \begin{aligned} F_{\mathbf{m}}^{p,\ell}(\xi_0, \dots, \xi_{p+1}) &= \sum_{j=0}^{\ell} \sqrt{\mathbf{m}^2 + \xi_j^2} - \sum_{j=\ell+1}^{p+1} \sqrt{\mathbf{m}^2 + \xi_j^2} \\ G_{\mathbf{m}}^{p,\ell}(\xi_1, \dots, \xi_p) &= \sum_{j=1}^{\ell} \sqrt{\mathbf{m}^2 + \xi_j^2} - \sum_{j=\ell+1}^p \sqrt{\mathbf{m}^2 + \xi_j^2}. \end{aligned}$$

Denote by  $Z^\ell(p)$  (resp.  $\underline{Z}^\ell(p)$ ) the set of those  $(n_0, \dots, n_{p+1}) \in (\mathbb{N}^*)^{p+2}$  (resp.  $(n_1, \dots, n_p) \in (\mathbb{N}^*)^p$ ) such that there is a bijection  $\tau : \{0, \dots, \ell\} \rightarrow \{\ell+1, \dots, p+1\}$  (resp.  $\tau : \{1, \dots, \ell\} \rightarrow \{\ell+1, \dots, p\}$ ) with  $n_{\tau(j)} = n_j$  for any  $j = 0, \dots, \ell$  (resp.  $j = 1, \dots, \ell$ ). Of course, these sets are empty if  $p$  is odd, or if  $p$  is even and  $\ell \neq p/2$ . Then, Theorem 4.7 of [17] (see also, for similar results in one dimension, Bambusi [1] and Bourgain [12]) asserts that there is  $\mathcal{N} \subset ]0, +\infty[$  of zero measure and for any  $p \in \mathbb{N}$ ,  $L_0 > 0$  such that, for any  $\mathbf{m} \in ]0, +\infty[ - \mathcal{N}$ , there is  $c > 0$  with

$$(1.2.27) \quad \begin{aligned} |F_{\mathbf{m}}^{p,\ell}(\lambda_{n_0}, \dots, \lambda_{n_{p+1}})| &\geq c(n_0 + \dots + n_{p+1})^{-L_0} \\ \inf_{\alpha \in \mathbb{Z}} \left| G_{\mathbf{m}}^{p,\ell}(\lambda_{n_1}, \dots, \lambda_{n_p}) + \frac{\alpha}{2} \right| &\geq c(n_1 + \dots + n_p)^{-L_0} \end{aligned}$$

for any  $(n_0, \dots, n_{p+1}) \in (\mathbb{N}^*)^{p+2} - Z^\ell(p)$  and any  $(n_1, \dots, n_p) \in (\mathbb{N}^*)^p - \underline{Z}^\ell(p)$  respectively.

We may state a more general version of Theorem 1.1.1.

**THEOREM 1.2.8.** *Let  $P \in \mathbb{N}^*$  be given,  $R > 0, \nu' \in \mathbb{R}_+$ . Let  $P' \in \mathbb{N}^*$ ,  $\sigma > \nu(P') + \nu + \frac{1}{2}$  and let for any  $1 \leq p \leq P'$ ,  $G_p$  be an element of  $H\mathcal{F}_p^{1,\nu'}(B_\sigma(R))$ . Denote  $G_0(U) = \int_X (\Lambda_{\mathbf{m}} u) \bar{u} d\mu$  and  $G(U) = \sum_{p=0}^{P'} G_p(U)$ . Then for any  $\mathbf{m} \in ]0, +\infty[ - \mathcal{N}$ , there is  $s_0 > 0$  such that, for any  $s > s_0$ , there are  $\epsilon_0 \in ]0, 1[$ ,  $c > 0, C > 0$  and, for any  $\epsilon \in ]0, \epsilon_0[$ , for any  $u_0$  in the unit ball of  $H^s(X)$ , there is a unique solution  $u$  in  $C^0([-T_\epsilon, T_\epsilon], H^s(X))$  of the equation*

$$(1.2.28) \quad u_t = i\nabla_{\bar{u}} G(u, \bar{u}), \quad u(0, x) = \epsilon u_0(x)$$

with  $T_\epsilon \geq c\epsilon^{-P}$ . Moreover

$$(1.2.29) \quad \sup_{|t| < T_\epsilon} \|u(t, \cdot)\|_{H^s} \leq C\epsilon.$$

In the rest of this subsection, we shall prove that Theorem 1.2.8 contains Theorem 1.1.1. We do need the more general statement of Theorem 1.2.8 because the structure of  $G = \sum_p G_p$  will be stable along the proof.

We have seen that the proof of Theorem 1.1.1 may be reduced to solving equation (1.1.18) with a Hamiltonian  $G$  given by (1.1.11), with  $V$  related to  $U = (u, \bar{u})$  by (1.1.13). Consequently, Theorem 1.1.1 follows from Theorem 1.2.8 and the following lemma:

LEMMA 1.2.9. *Set  $G(U) = G_0(U) - F(\Lambda_{\mathbf{m}}^{-1/2} \frac{\sqrt{2}}{2i}(u - \bar{u}))$ . Then  $G$  has the structure of the assumption of Theorem 1.2.8 for any fixed  $R > 0$  if  $\sigma > \nu(p) + \nu + \frac{1}{2}, \nu' \geq \frac{d}{2} + 1$  and  $\nu(p) \geq p(\frac{d}{2} + 1)$  for any  $p = 1, \dots, P'$ .*

Before giving the proof of the lemma, we shall establish some properties of  $F(v) = \int_X f(v, dv) d\mu$ , where  $(z, \rho) \rightarrow f(z, \rho)$  is a smooth function on  $\mathbb{R} \times T^*X$ , polynomial in  $z$  and in the fiber variable of  $\rho$ , and  $v$  is in  $\text{Lip}(X)$ , the space of real valued lipschitz function on  $X$ . We denote for a while by  $\{\cdot, \cdot\}$  the Poisson bracket between functions on  $T^*X$ , as at the beginning of subsection 1.1. We consider systematically any smooth enough function on  $X$  as a function on  $T^*X$ , independent of the fiber variable. For  $h$  a given element in  $\text{Lip}(X)$ ,  $f : \mathbb{R} \times T^*X \rightarrow \mathbb{R}$  a smooth function, we define a new function  $\mathcal{D}(h)f$  from  $\mathbb{R} \times T^*X$  to  $\mathbb{R}$  by the formula

$$(1.2.30) \quad \mathcal{D}(h)f = \frac{\partial f}{\partial z} h - \{h, f\}.$$

This new function is no longer smooth in  $\rho \in T^*X$ , because of the limited smoothness of  $h$ . Nevertheless, a composition  $\mathcal{D}(h_1) \cdots \mathcal{D}(h_p)f$  is still meaningful, for any  $h_1, \dots, h_p \in \text{Lip}(X)$ : for instance

$$\mathcal{D}(h_1)\mathcal{D}(h_2)f = \frac{\partial^2 f}{\partial z^2} h_1 h_2 - \left\{ h_1, \frac{\partial f}{\partial z} \right\} h_2 - \left\{ h_2, \frac{\partial f}{\partial z} \right\} h_1 + \{h_2, \{h_1, f\}\}$$

never involves more than one derivative acting on each  $h_j$ . By an immediate computation in local coordinates, one checks that  $F(U) = \int_X f(v, dv) d\mu$  is a smooth function on  $\text{Lip}(X)$ , whose differentials are given by

$$(1.2.31) \quad d^p F(v) \cdot (h_1, \dots, h_p) = \int_X \left( \prod_{j=1}^p \mathcal{D}(h_j)f \right) (v, dv) d\mu.$$

This formula, and the above expression for  $\mathcal{D}(h_j)$ , shows that when  $v$  stays in a fixed ball of  $\text{Lip}(X)$ ,

$$(1.2.32) \quad |d^p F(v) \cdot (h_1, \dots, h_p)| \leq C \prod_{j=3}^p \|h_j\|_{\text{Lip}(X)} \|h_1\|_{H^1} \|h_2\|_{H^1}.$$

We shall need bounds on the commutator between the symmetric operator associated to the bilinear form  $(h_1, h_2) \rightarrow d^2 F(v)(h_1, h_2)$  and differential operators. We fix some notation: if  $w_1, \dots, w_p$  are smooth functions on  $X$ , if  $k \in \mathbb{N}$ , we set

$$(1.2.33) \quad M_k(w_1, \dots, w_p) = \sum_{|\alpha_1| + \dots + |\alpha_p| = k} \prod_{j=1}^p \|\partial^{\alpha_j} w_j\|_{\text{Lip}(X)}.$$

DEFINITION 1.2.10. *Let  $p \in \mathbb{N}, r \in \mathbb{N}, m \in \mathbb{Z}$ . If  $m < 0$ , set  $\widetilde{H\mathcal{A}}_{p,r}^m = \{0\}$ . If  $m \geq 0$ , we denote by  $\widetilde{H\mathcal{A}}_{p,r}^m$  the space of  $p$ -linear maps  $(v_1, \dots, v_p) \rightarrow A(v_1, \dots, v_p)$ , defined on  $C^\infty(M, \mathbb{R})^p$ , with values in the space symmetric bilinear forms on  $C^\infty(X, \mathbb{R}) \times C^\infty(X, \mathbb{R})$ , satisfying the following condition: for any local chart  $\theta : V \rightarrow U$  from an open subset  $V$  of  $X$  to an open subset  $U$  of  $\mathbb{R}^d$ , for any  $\gamma_1, \gamma_2$  in  $\mathbb{N}^d$  with  $|\gamma_1| + |\gamma_2| \leq m$ , there is a map*

$$(1.2.34) \quad \begin{aligned} a_{\gamma_1, \gamma_2} : (x, v_1, \dots, v_p) &\longrightarrow a_{\gamma_1, \gamma_2}(x, v_1, \dots, v_p) \\ U \times C^\infty(M, \mathbb{R})^p &\longrightarrow \mathbb{R} \end{aligned}$$



which is  $p$ -linear in  $(v_1, \dots, v_p)$ , smooth in  $x$ , satisfying for any  $\alpha \in \mathbb{N}^d$

$$(1.2.35) \quad |\partial_x^\alpha a_{\gamma_1, \gamma_2}(x, v_1, \dots, v_p)| \leq CM_{|\alpha|+r}(v_1, \dots, v_p)$$

such that, for any  $h_1, h_2$  in  $C_0^\infty(V, \mathbb{R})$ ,

(1.2.36)

$$A(v_1, \dots, v_p)(h_1, h_2) = \sum_{|\gamma_1|+|\gamma_2| \leq m} \int_U a_{\gamma_1, \gamma_2}(x, v_1, \dots, v_p) (\partial_x^{\gamma_1} [h_1 \circ \theta^{-1}]) (\partial_x^{\gamma_2} [h_2 \circ \theta^{-1}]) dx.$$

One denotes by  $\widehat{H\mathcal{A}}_{p,r}^m$  the space of maps  $v \rightarrow A(v)$  that may be written  $\tilde{A}(v, \dots, v)$  for some  $\tilde{A}$  in  $\widehat{H\mathcal{A}}_{p,r}^m$ .

We shall use the following lemma:

LEMMA 1.2.11. *Let  $P$  be a differential operator of order  $\ell$  on  $X$ ,  $A$  be an element of  $\widehat{H\mathcal{A}}_{p,r}^m$ . Then  $(h_1, h_2) \rightarrow A(v_1, \dots, v_p)(Ph_1, h_2) - A(v_1, \dots, v_p)(h_1, {}^tPh_2)$  is in  $\sum_{\ell'=1}^\ell \widehat{H\mathcal{A}}_{p,r+\ell'}^{m+\ell-\ell'}$ .*

*Proof:* If  $\tilde{A}$  is in  $\widehat{H\mathcal{A}}_{p,r}^m$ , we may write, according to (1.2.36), the quantity under study from expressions

$$\begin{aligned} & \int_U a_{\gamma_1, \gamma_2}(x, v_1, \dots, v_p) (\partial_x^{\gamma_1} [Ph_1 \circ \theta^{-1}]) (\partial_x^{\gamma_2} [h_2 \circ \theta^{-1}]) dx \\ & - \int_U a_{\gamma_1, \gamma_2}(x, v_1, \dots, v_p) (\partial_x^{\gamma_1} [h_1 \circ \theta^{-1}]) (\partial_x^{\gamma_2} [({}^tPh_2) \circ \theta^{-1}]) dx. \end{aligned}$$

If we perform integrations by parts in the second integral, we get expressions involving at most  $\ell'$  derivatives of  $a_{\gamma_1, \gamma_2}$  and  $|\gamma_1| + |\gamma_2| - \ell' + \ell$  derivatives of  $(h_1, h_2)$  for  $1 \leq \ell' \leq \ell$ . This shows that we obtain an element of  $\sum_{\ell'=1}^\ell \widehat{H\mathcal{A}}_{p,r+\ell'}^{m+\ell-\ell'}$ .  $\square$

*Proof of Lemma 1.2.9:* We denote by  $Q$  the pseudo-differential operator of order  $-1/2$ ,  $Q = \Lambda_{\mathbf{m}}^{-1/2} \frac{\sqrt{2}}{2i} [1 - 1]$ , acting on complex valued functions  $U = \begin{bmatrix} u \\ \bar{u} \end{bmatrix}$ . The Hamiltonian  $G$  of the statement of the lemma is expressed in terms of

$$U \rightarrow F(QU) = \int_X f(QU, dQU) d\mu.$$

Using notation (1.2.2), with the convention  $S_{-1} = 0$ , we write  $\text{Id} = \sum_{j=0}^{+\infty} (S_j - S_{j-1})$  so that

$$F(QU) = \sum_{j=0}^{+\infty} (F(QS_j U) - F(QS_{j-1} U)) = \sum_{j=0}^{+\infty} \int_0^1 dF(QS_j(t)U) \cdot Q\Delta_j U dt$$

where  $S_j(t) = tS_j + (1-t)S_{j-1}$ . Applying the same decomposition to  $U \rightarrow dF(QS_j(t)U)$ , which is possible since  $dF(0) = 0$ , we get

$$(1.2.37) \quad F(U) = \sum_{j=0}^{+\infty} \sum_{j'=0}^{+\infty} \int_0^1 \int_0^1 d^2 F(QS_{jj'}(t, t')U) \cdot (Q\Delta_j U, QS_{jj'}(t, t')U) dt dt'$$

with  $S_{jj'}(t, t') = S_j(t)S_{j'}(t')$ . We use Taylor formula to decompose

$$d^2 F(W) = \sum_{p=0}^{+\infty} \frac{1}{p!} d^{p+2} F(0) \cdot W^p$$

where the sum is finite since  $F$  is polynomial. Plugging this expression in (1.2.37) and using the expression (1.2.31) of  $d^p F$ , and the fact that  $F$  vanishes at least at order three at zero, we may write

$$(1.2.38) \quad F(QU) = \sum_{p=1}^{+\infty} A_p(U) \cdot (U, U),$$

where the symmetric bilinear forms  $A_p(U)$  are given by

$$(1.2.39) \quad A_p(U) = \sum_{j_1=0}^{+\infty} \sum_{j'_1=0}^{+\infty} \int_0^1 \int_0^1 A_p^{j_1, j'_1}(U, t, t') dt dt', \quad p \geq 1,$$

denoting

$$(1.2.40) \quad A_p^{j_1, j'_1}(U, t, t') \cdot (H_1, H_2) = \frac{1}{p!} \int_X \mathcal{D}(Q\Delta_{j_1} H_1) \mathcal{D}(QS_{j_1}(t) \Delta_{j'_1} H_2) g_p(U, \cdot)|_{\{0\} \times X} d\mu$$

with

$$(1.2.41) \quad g_p(U, z, \rho) = [\mathcal{D}(QS_{j_1 j'_1}(t, t') U)]^p f(z, \rho)$$

(The restriction in the right hand side of (1.2.40) means restriction to the zero section  $\{0\} \times X$  of  $\mathbb{R} \times T^*X$ ).

To finish the proof of the lemma, we have to show that  $G_p(U) = -A_p(U) \cdot (U, U)$  is in  $H\mathcal{F}_p^{1, \nu'}(B_\sigma(R))$  with  $\sigma > \nu(p) + \nu' + \frac{1}{2}$  for any  $p \geq 1$ . This follows from next lemma.  $\square$

LEMMA 1.2.12. *For  $\mathcal{U} = (U_1, \dots, U_p)$  a family of  $C^\infty(X)$  functions, set*

$$(1.2.42) \quad \tilde{g}_p(\mathcal{U}, t, t', z, \rho) = \prod_{\ell=1}^p \mathcal{D}(QS_{j_1 j'_1}(t, t') U_\ell) \cdot f(z, \rho)$$

and define  $A_p^{j_1, j'_1}(\mathcal{U}, t, t')$  (resp.  $A_p(U)$ ) by (1.2.40) (resp. (1.2.39)) with  $g_p$  replaced by  $\tilde{g}_p$  and with  $U$  replaced by  $\mathcal{U}$ . Then, if we identify these symmetric bilinear forms on  $C^\infty(X) \times C^\infty(X)$  to the corresponding linear maps from  $C^\infty(X)$  to  $\mathcal{D}'(X)$ ,  $A_p(\mathcal{U})$  belongs to  $\widetilde{H\Psi}_p^{1, \nu'}$  if  $\nu' \geq \nu(\frac{d}{2} + 1)$  and  $\nu(p) \geq p(\frac{d}{2} + 1)$ .

*Proof:* Define for  $p \in \mathbb{N}^*$

$$(1.2.43) \quad B_p^{j_1, j'_1}(\mathcal{U}, t, t')(H_1, H_2) = \frac{1}{p!} \int_X \mathcal{D}(H_1) \mathcal{D}(H_2) \tilde{g}_p(\mathcal{U}, \cdot)|_{\{0\} \times X} d\mu.$$

Let us check that  $B_p^{j_1, j'_1}$  is in  $\widetilde{H\mathcal{A}}_{p,0}^2$ , with constants in estimates (1.2.35) uniform in  $j_1, j'_1, t, t'$ .

It follows from the definition of  $\mathcal{D}(h)$  that  $|\mathcal{D}(h_1) \cdots \mathcal{D}(h_p) f|$  is bounded from above by a constant times  $C \prod_{j=1}^p \|h_j\|_{\text{Lip}(X)}$ . Expression (1.2.42), and boundedness of pseudo-differential operators of order  $-1/2$  like  $QS_{j_1, j'_1}(t, t')$  on  $\text{Lip}(X)$ , show that  $\partial_x^\alpha$  derivatives of  $\tilde{g}_p(\mathcal{U}, t, t', z, \rho)$  are bounded, when  $(z, \rho)$  stays in a compact set, in terms of  $M_{|\alpha|}(U_1, \dots, U_p)$  defined by (1.2.33). If in (1.2.43),  $H_1, H_2$  are supported in a chart domain, it follows that  $B_p^{j_1, j'_1}(\mathcal{U}, t, t')(H_1, H_2)$  is given by an expression of form (1.2.36) with  $m = 2$ , and with  $\partial_x^\alpha$ -derivatives of the coefficients  $a_{\gamma_1, \gamma_2}$  bounded by the right hand side of (1.2.35) with  $r = 0$ . This shows that

$B_p^{j_1, j'_1}(\mathcal{U}, t, t')$  is in  $\widetilde{H\mathcal{A}}_{p,0}^2$ , uniformly in  $j_1, j'_1, t, t'$ . Moreover, because of the definition of  $\tilde{g}_p$ ,  $B_p^{j_1, j'_1}(\Pi_n \mathcal{U}, t, t') \equiv 0$  if  $|n'| \geq C2^{\min(j_1, j'_1)}$  for some large enough constant  $C$ .

We have to deduce from these informations on  $B_p^{j_1, j'_1}$  the conclusion of the lemma for  $A_p(\mathcal{U})$ . Identifying bilinear forms and linear maps, we deduce from (1.2.40), (1.2.42), (1.2.43) that  $A_p^{j_1, j'_1}$  may be written in terms of  $B_p^{j_1, j'_1}$  by

$$(1.2.44) \quad A_p^{j_1, j'_1}(\mathcal{U}, t, t') = \Delta_{j_1} {}^t Q B_p^{j_1, j'_1}(\mathcal{U}, t, t') Q S_{j_1}(t) \Delta_{j'_1}.$$

Let  $\mathcal{P} = (P_1, \dots, P_k)$  be a family of differential operators, and denote  $\text{Ad}_{\mathcal{P}} = \text{Ad}_{P_1} \cdots \text{Ad}_{P_k}$ . By definition of  $A_p(\mathcal{U})$  and (1.2.44)

$$(1.2.45) \quad \Delta_j \text{Ad}_{\mathcal{P}} A_p(\Pi_n \mathcal{U}) \Delta_{j'} = \sum_{j_1} \sum_{j'_1} \int_0^1 \int_0^1 \Delta_j \text{Ad}_{\mathcal{P}} [\Delta_{j_1} {}^t Q B_p^{j_1, j'_1}(\Pi_n \mathcal{U}, t, t') Q S_{j_1}(t) \Delta_{j'_1}] \Delta_{j'} dt dt'.$$

The integrand may be written from a sum of terms

$$(1.2.46) \quad \Delta_j \text{Ad}_{\mathcal{P}'} [\Delta_{j_1} {}^t Q] \cdot \text{Ad}_{\mathcal{P}''} [B_p^{j_1, j'_1}(\Pi_n \mathcal{U}, t, t')] \cdot \text{Ad}_{\mathcal{P}'''} [Q S_{j_1}(t) \Delta_{j'_1}] \Delta_{j'}$$

where  $\mathcal{P}' \cup \mathcal{P}'' \cup \mathcal{P}'''$  is a partition of  $\mathcal{P}$ . Let  $\tilde{\varphi} \in C_0^\infty(\mathbb{R} - \{0\})$  with  $\tilde{\varphi}\varphi = \varphi$ , and denote by  $\tilde{\Delta}_j = \tilde{\varphi}(2^{-j}\sqrt{-\Delta})$  the corresponding spectral cut-off (with the obvious modification for  $j = 0$ ). We insert  $\tilde{\Delta}_{j_2} \Delta_{j_2}$ ,  $\tilde{\Delta}_{j'_2} \Delta_{j'_2}$  in (1.2.46) and rewrite this expression as

$$\sum_{j_2} \sum_{j'_2} K(j, j_1, j_2) \Delta_{j_2} \text{Ad}_{\mathcal{P}''} [B_p^{j_1, j'_1}(\Pi_n \mathcal{U}, t, t')] \Delta_{j'_2} K'(j'_2, j'_1, j')$$

with

$$K(j, j_1, j_2) = \Delta_j \text{Ad}_{\mathcal{P}'} [\Delta_{j_1} {}^t Q] \tilde{\Delta}_{j_2}, \quad K'(j'_2, j'_1, j') = \tilde{\Delta}_{j'_2} \text{Ad}_{\mathcal{P}'''} [Q S_{j_1}(t) \Delta_{j'_1}] \Delta_{j'}.$$

By Proposition A.1 of the appendix, for any  $N \in \mathbb{N}$ , there is  $C_N > 0$  such that

$$\begin{aligned} \|K(j, j_1, j_2)\|_{\mathcal{L}(L^2)} &\leq C_N 2^{-N[|j-j_1|+|j_1-j_2|]-\frac{j}{2}+jD} \\ \|K'(j'_2, j'_1, j')\|_{\mathcal{L}(L^2)} &\leq C_N 2^{-N[|j'-j'_1|+|j'_1-j'_2|]-\frac{j'}{2}+j'D'} \end{aligned}$$

with  $D = \sum_{P \in \mathcal{P}'} \deg(P) - |\mathcal{P}'|$ ,  $D' = \sum_{P \in \mathcal{P}'''} \deg(P) - |\mathcal{P}'''|$ . These estimates, and an elementary computation, show that (1.2.45) will satisfy (1.2.8) with  $m = 1$ ,  $M_0 = 1$  if we may prove that for any  $N_1 \in \mathbb{N}$ , there is  $N'_1$  such that

$$(1.2.47) \quad \begin{aligned} \|\Delta_{j_2} \text{Ad}_{\mathcal{P}''} [B_p^{j_1, j'_1}(\Pi_n \mathcal{U}, t, t')] \Delta_{j'_2}\|_{\mathcal{L}(L^2)} &\leq C 2^{-N_1|j_2-j'_2|+N'_1(|j_1-j_2|+|j'_1-j'_2|)} 2^{j_2[2+D'']} \\ &\quad \times |n'|^{\nu(p)+|\mathcal{P}''|} (1 + 2^{-j_1}|n'|)^{-N_2} \prod \|\Pi_{n_\ell} u_\ell\|_{L^2}, \end{aligned}$$

where  $D'' = \sum_{P \in \mathcal{P}''} \deg(P) - |\mathcal{P}''|$ . Note that a commutator  $[P, B_p^{j_1, j'_1}(\Pi_n \mathcal{U}, t, t')]$  is the translation at the level of linear maps of the quantities studied in Lemma 1.2.11 at the level of bilinear forms. By this lemma,  $\text{Ad}_{\mathcal{P}''} B_p^{j_1, j'_1}(\Pi_n \mathcal{U}, t, t')$  will be the

sum indexed by  $a$  in  $\{0, \dots, D''\}$  of elements of  $\widetilde{H\mathcal{A}}_{p, |\mathcal{P}''|+a}^{D''-a+2}$ . Expressions (1.2.36) and bounds (1.2.35) imply that the left hand side of (1.2.47) is smaller than

$$CM_{|\mathcal{P}''|+a}(\Pi_{n'}\mathcal{U}) \sum_{|\gamma_2|+|\gamma'_2| \leq D''-a+2} 2^{j_2|\gamma_2|+j'_2|\gamma'_2|}.$$

By (1.2.33) and Sobolev injections, this is smaller than

$$(1.2.48) \quad C|n'|^{|\mathcal{P}''|+p\left[\frac{d}{2}+1\right]} 2^{j_2(D''+2)+(D''+2-a)|j_2-j'_2|} (|n'|2^{-j_2})^a \prod_1^p \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

We have seen that if  $B_p^{j_1, j'_1}(\Pi_{n'}\mathcal{U}, t, t') \neq 0$ , then  $|n'| \leq C \min(2^{j_1}, 2^{j'_1})$ . This shows that (1.2.48) implies (1.2.47) with  $\nu(p) = p\left[\frac{d}{2}+1\right]$  if we are able to show that we may improve (1.2.48) by a factor  $2^{N_1[|j_1-j_2|+|j'_1-j'_2|-|j_2-j'_2|]}$  for any  $N_1$ . To do so, we denote by  $\bar{\Delta}_j$  a new cut-off such that, for  $j > 0$ ,  $\bar{\Delta}_j = 2^{2j}\Delta_j$ . When  $0 < j'_2 \leq j_2$ , write

$$\begin{aligned} 2^{2j_2}\Delta_{j_2}\text{Ad}_{\mathcal{P}''}[B_p^{j_1, j'_1}(\Pi_{n'}\mathcal{U}, t, t')]\Delta_{j'_2} &= \bar{\Delta}_{j_2}\Delta\text{Ad}_{\mathcal{P}''}[B_p^{j_1, j'_1}(\Pi_{n'}\mathcal{U}, t, t')]\Delta_{j'_2} \\ &= \bar{\Delta}_{j_2}\text{Ad}_\Delta\text{Ad}_{\mathcal{P}''}[B_p^{j_1, j'_1}(\Pi_{n'}\mathcal{U}, t, t')]\Delta_{j'_2} + \bar{\Delta}_{j_2}\text{Ad}_{\mathcal{P}''}[B_p^{j_1, j'_1}(\Pi_{n'}\mathcal{U}, t, t')]\bar{\Delta}_{j'_2}2^{2j'_2}. \end{aligned}$$

If we apply estimate (1.2.48) to the right hand side of this equality, we get a bound given by the product of (1.2.48) and of

$$|n'|2^{j_2} + 2^{2j'_2} \leq C[2^{j_2+j'_1} + 2^{2j'_2}].$$

Consequently, we gained the factor

$$2^{j'_1-j_2} + 2^{2(j'_2-j_2)} \leq C2^{-|j'_2-j_2|+|j'_1-j'_2|}.$$

A symmetric reasoning applies if  $j_2 \leq j'_2$ , so that we have gained, in the special case  $N_1 = 1$ , the missing factor  $2^{N_1[|j_1-j_2|+|j'_1-j'_2|-|j_2-j'_2|]}$ . Iterating, we get the case of an arbitrary  $N_1$ .  $\square$



## CHAPTER 2

### Symbolic calculus

The quasi-linear Birkhoff normal forms method, that will be used to prove Theorem 1.2.8, will rely on properties of commutators of operators belonging to the classes defined in Definitions 1.2.3, 1.2.5. To establish such properties, we need a notion of principal symbol that will be defined through local models of our operators, acting on functions defined on the Euclidean space.

**2.1. Operators on  $\mathbb{R}^d$  and their symbols.** In the same way as we defined dyadic cut-offs acting on functions on the compact manifold  $X$  in (1.2.2), we define for  $u$  in  $\mathcal{S}'(\mathbb{R}^d)$ ,  $j$  in  $\mathbb{N}$

$$(2.1.1) \quad \begin{aligned} \Delta_j^E u &= \mathcal{F}^{-1}(\varphi(2^{-j}|\xi|)\hat{u}), \quad j \geq 1 \\ \Delta_0^E u &= S_0^E u = \mathcal{F}^{-1}(\psi(|\xi|)\hat{u}) \\ S_j^E u &= \sum_{j'=0}^j \Delta_{j'}^E u. \end{aligned}$$

where  $\hat{u}$  (resp.  $\mathcal{F}^{-1}$ ) denotes the Fourier transform (resp. the inverse Fourier transform) on  $\mathbb{R}^d$ , and where  $\varphi, \psi$  are the functions introduced at the beginning of subsection 1.2.

**DEFINITION 2.1.1.** Let  $m \in \mathbb{R}, \nu \in \mathbb{N}, p \in \mathbb{N}$ . One denotes by  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu}$  the space of  $p$ -linear maps  $\mathcal{U} = (U_1, \dots, U_p) \rightarrow A(\mathcal{U})$ , defined on  $C^\infty(X)^p$ , with values in the space of linear maps from  $C_0^\infty(\mathbb{R}^d)$  to  $\mathcal{D}'(\mathbb{R}^d)$ , such that the distribution kernel of  $A(\mathcal{U})$  is supported in a compact subset of  $\mathbb{R}^d \times \mathbb{R}^d$  (independent of  $\mathcal{U}$ ) and such that for any family of differential operators  $P_1, \dots, P_k$  of order  $d_1, \dots, d_k$ , for any  $\mathcal{U}$  in  $C^\infty(X)^p$ , any  $n' = (n_1, \dots, n_p)$  in  $(\mathbb{N}^*)^p$ , any  $j, j'$  in  $\mathbb{N}$ , any  $N_1, N_2$  in  $\mathbb{N}$ ,

$$\|\Delta_j^E \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} A(\Pi_{n'} \mathcal{U}) \Delta_{j'}^E\|_{\mathcal{L}(L^2)}$$

is bounded from above by the right hand side of (1.2.8)

As in the case of global operators on  $X$ , one can give an equivalent definition in terms of estimates of the form (1.2.9). We define from  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu}$  the space  $H\Psi_{p,\text{loc}}^{m,\nu}$  as in Definition 1.2.3

**DEFINITION 2.1.2.** Let  $m \in \mathbb{R}, \nu \in \mathbb{N}, p \in \mathbb{N}, \sigma > 0, R > 0$ . One defines  $\Psi_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$  as the space of maps  $U \rightarrow A(U)$  defined on  $B_\sigma(R)$ , with values in the vector space of linear maps from  $C_0^\infty(\mathbb{R}^d)$  to  $\mathcal{D}'(\mathbb{R}^d)$ , such that the distribution kernel of  $A(U)$  is supported in a compact subset of  $\mathbb{R}^d \times \mathbb{R}^d$  independent of  $U$ , and such that for any family of differential operators  $P_1, \dots, P_k$  of order  $d_1, \dots, d_k$ , for any  $N \in \mathbb{N}$ , for any  $\sigma' \in [-\sigma, \sigma]$ , any  $U \in B_\sigma(R)$ , any  $H \in H^{\sigma'}(X)$ , any  $j, j' \in \mathbb{N}$ ,  $\|\Delta_j^E \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} A(U) \Delta_{j'}^E\|_{\mathcal{L}(L^2)}$  (resp.  $\|\Delta_j^E \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} (\partial_U A(U) \cdot H) \Delta_{j'}^E\|_{\mathcal{L}(L^2)}$ ) is bounded from above by the right hand side of (1.2.16) (resp. (1.2.17)).

As in the case of the global definition, one may give an equivalent characterization in terms of estimates like (1.2.18), (1.2.19).

We introduce classes of symbols that will allow us to give alternative descriptions of the preceding classes of operators.

DEFINITION 2.1.3. *Let  $m \in \mathbb{R}, \nu \in \mathbb{N}, p \in \mathbb{N}$ . One denotes by  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  the space of functions*

$$a : C^\infty(X)^p \times T^*\mathbb{R}^d \rightarrow \mathbb{C} \\ (u_1, \dots, u_p, x, \xi) \rightarrow a(u_1, \dots, u_p, x, \xi)$$

which are  $p$ -linear in  $(u_1, \dots, u_p)$ , smooth in  $(x, \xi)$ , and satisfy for any  $\alpha, \beta \in \mathbb{N}^d$ , any  $\mathcal{U} = (u_1, \dots, u_p)$  in  $C^\infty(X)^p$ , any  $n' = (n_1, \dots, n_p)$  in  $(\mathbb{N}^*)^p$ , any  $(x, \xi) \in T^*\mathbb{R}^d$ , any  $N \in \mathbb{N}$

$$|\partial_x^\alpha \partial_\xi^\beta a(\Pi_{n'} \mathcal{U}, x, \xi)| \leq C_N \langle \xi \rangle^{m-|\beta|} |n'|^{\nu(p)+\nu+M_0(|\alpha|+|\beta|+d+1)} \left(1 + \frac{|n'|}{\langle \xi \rangle}\right)^{-N} \prod_1^p \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

We define also a class of symbols that are not homogeneous in  $U$ .

DEFINITION 2.1.4. *Let  $m \in \mathbb{R}, \nu \in \mathbb{N}, p \in \mathbb{N}, \sigma > 0, R > 0$ . One denotes by  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$  the space of maps*

$$a : B_\sigma(R) \times T^*\mathbb{R}^d \rightarrow \mathbb{C} \\ (U, x, \xi) \rightarrow a(U, x, \xi)$$

which are smooth in  $(x, \xi)$ ,  $C^1$  in  $U$ , and satisfy the following conditions:

(i) For any  $\alpha, \beta \in \mathbb{N}^d$ , there is  $C > 0$  and, for any  $U$  in  $B_\sigma(R)$ , any  $(x, \xi)$  in  $T^*\mathbb{R}^d$

$$(2.1.3) \quad |\partial_x^\alpha \partial_\xi^\beta a(U, x, \xi)| \leq C \langle \xi \rangle^{m-|\beta|+(M_0(|\alpha|+|\beta|+d+1)+\nu(p)+\nu-\sigma)_+} \|U\|_{H^\sigma}^p.$$

(ii) For all  $\alpha, \beta, x, \xi, U, H \rightarrow (\partial_x^\alpha \partial_\xi^\beta \partial_U a)(U, x, \xi) \cdot H$  extends as a continuous linear form on  $H^{\sigma'}(X)$  for all  $\sigma'$  with  $|\sigma'| \leq \sigma$  and satisfies

$$(2.1.4) \quad |(\partial_x^\alpha \partial_\xi^\beta \partial_U a)(U, x, \xi) \cdot H| \leq C \langle \xi \rangle^{m-|\beta|+(M_0(|\alpha|+|\beta|+d+1)+\nu(p)+\nu-\sigma')_+} \|U\|_{H^\sigma}^{p-1} \|H\|_{H^{\sigma'}}.$$

We quantize the preceding classes of symbols in the following way. If  $v$  is in  $C_0^\infty(\mathbb{R}^d)$  and  $a$  is in  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ , we set

$$(2.1.5) \quad \text{Op}(a(U; \cdot))v = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} a(U, x, \xi) \hat{v}(\xi) d\xi.$$

We use a similar formula to quantize elements of  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$ .

If  $\chi$  is in  $C_0^\infty(\mathbb{R}^d)$ ,  $a$  belongs to  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$ , we set

$$(2.1.6) \quad a_\chi(\mathcal{U}, x, \xi) = \chi(D_x / \langle \xi \rangle) [a(\mathcal{U}, x, \xi)] = \mathcal{F}_\eta^{-1} [\chi(\eta / \langle \xi \rangle) \hat{a}(\mathcal{U}, \eta, \xi)]$$

where  $\hat{a}(\mathcal{U}, \eta, \xi)$  is the  $x$ -Fourier transform of  $a(\mathcal{U}, x, \xi)$ . We use a similar notation for elements of  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ . The definition immediately implies that  $a_\chi$  belongs to the same space of symbols as  $a$ .

If  $\chi$  is supported in a small enough neighborhood of zero, on the support of  $\widehat{a_\chi}(\mathcal{U}, \eta, \xi)$ ,  $|\eta| \ll |\xi|$ , so that  $\langle \xi + \eta \rangle \sim \langle \xi \rangle$ . For technical reasons, we shall have to consider symbols for which a weaker version of this spectral localization is

satisfied, namely symbols  $a$  in  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  or  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$  such that for any  $\lambda \in \mathbb{R}$ ,  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} a$  is still in the same class. We shall use eventually the following lemma:

LEMMA 2.1.5. *Let  $a$  be an element of  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp.  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ) such that  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} a$  is in the same space for any  $\lambda \in \mathbb{R}$ . Let  $\theta$  in  $\mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$  and  $b(\mathcal{U}, x, y, \xi) = \theta(x, y) a(\mathcal{U}, x, \xi)$  (resp.  $b(U, x, y, \xi) = \theta(x, y) a(U, x, \xi)$ ). Then, for any  $n'$  in  $(\mathbb{N}^*)^p$ , any  $\lambda$  in  $\mathbb{R}$ ,*

$$|\partial_x^\alpha \partial_\xi^\beta \partial_y^\gamma [\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} b(\Pi_{n'} \mathcal{U}, x, y, \xi)]|,$$

resp.

$$|\partial_x^\alpha \partial_\xi^\beta \partial_y^\gamma [\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} b(U, x, y, \xi)]|,$$

resp.

$$|\partial_x^\alpha \partial_\xi^\beta \partial_y^\gamma [\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} (\partial_U b(U, x, y, \xi) \cdot H)]|$$

is bounded from above by the right hand side of (2.1.2) (resp. (2.1.3), resp. (2.1.4)).

*Proof:* It is enough to prove that the operator  $\langle D_x + \xi \rangle^\lambda \theta \langle D_x + \xi \rangle^{-\lambda} - \theta$  is bounded on  $L^\infty(dx)$ , uniformly in  $\xi, y$ , for any  $\lambda \in \mathbb{R}$ . The distribution kernel of this operator, at fixed  $\xi, y$ , is

$$K_{\xi,y}(x, x') = \frac{1}{(2\pi)^{2d}} e^{-i(x-x') \cdot \xi} \int e^{i(x-x') \cdot \eta + ix \cdot \zeta} \langle \eta \rangle^{-\lambda} [\langle \eta + \zeta \rangle^\lambda - \langle \eta \rangle^\lambda] \hat{\theta}(\zeta, y) d\zeta d\eta,$$

where  $\hat{\theta}(\zeta, y)$  is the Fourier transform relatively to the first variable. Integrating by parts, one checks

$$|K_{\xi,y}(x, x')| \leq C_N |x - x'|^{-d+1} \langle x - x' \rangle^{-N} \langle x \rangle^{-N}$$

for any  $N$ , from which the  $\mathcal{L}(L^\infty)$ -bound follows.  $\square$

We relate the symbols we just defined to the local operators.

PROPOSITION 2.1.6. *Let  $m \in \mathbb{R}, \nu \in \mathbb{N}, \sigma > 0, R > 0$ .*

(i) *Let  $A$  be an element of  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu}$  (resp.  $\Psi_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ). There is a (unique) symbol  $a$  in  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp.  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ) such that for any  $\lambda \in \mathbb{R}$ ,  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} a$  is still in the same space, and such that  $A(\mathcal{U}) = \text{Op}(a(\mathcal{U}; \cdot))$  (resp.  $A(U) = \text{Op}(a(U; \cdot))$ ).*

(ii) *Let  $a$  be a symbol in  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp.  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ) such that for any  $\lambda \in \mathbb{R}$ ,  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} a$  is still in the same space. Let  $\theta_1, \theta_2$  in  $C_0^\infty(\mathbb{R}^d)$ . Then  $\theta_1 \text{Op}(a) \theta_2$  belongs to  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu+2M_0(d+1)}$  (resp.  $\Psi_{p,\text{loc}}^{m,\nu+2M_0(d+1)}(B_\sigma(R))$ ).*

(iii) *Let  $a$  be a symbol in  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp.  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ). Let  $\chi$  be in  $C_0^\infty(\mathbb{R}^d)$  with small enough support,  $\chi \equiv 1$  close to zero. Then, for any  $j \geq 0$ ,  $a - a_\chi$  is in  $\widetilde{HS}_{p,\text{loc}}^{m-j,\nu+jM_0}$  (resp.  $S_{p,\text{loc}}^{m-j,\nu+jM_0}(B_\sigma(R))$ ).*

*Proof:* (i) Let  $A$  be an element of  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu}$ . Denote by  $K(\mathcal{U}, x, y)$  the distribution kernel of  $A(\mathcal{U})$  and define

$$a(\mathcal{U}, x, \xi) = \int_{\mathbb{R}^d} K(\mathcal{U}, x, y) e^{-i(x-y) \cdot \xi} dy$$



so that  $\text{Op}(a(\mathcal{U}; \cdot)) = A(\mathcal{U})$  and  $\text{Op}(\partial_x^\alpha (i\partial_\xi)^\beta a(\mathcal{U}; \cdot)) = (\text{Ad}_{\partial_x})^\alpha (\text{Ad}_x)^\beta A(\mathcal{U})$ . Since  $K$  is compactly supported, we may choose  $\theta_1, \theta_2$  in  $C_0^\infty(\mathbb{R}^d)$  such that one has  $K = \theta_1(x)K\theta_2(y)$ . We compute

$$\begin{aligned} \hat{a}(\mathcal{U}, \eta, \xi) &= \int e^{-ix \cdot (\xi + \eta)} \theta_1(x) K(\mathcal{U}, x, y) e^{iy \cdot \xi} \theta_2(y) dy d\xi dx \\ &= \int e^{-ix \cdot (\xi + \eta)} \theta_1(x) A(\mathcal{U}) [e^{i\langle \xi, \cdot \rangle} \theta_2(\cdot)] dx. \end{aligned}$$

This implies, for any  $\lambda \in \mathbb{R}$ ,

$$\left| \widehat{\partial_x^\alpha \partial_\xi^\beta a(\Pi_{n'} \mathcal{U}, \eta, \xi)} \right| \leq \left\| e^{-i\langle \cdot, \xi + \eta \rangle} \theta_1 \right\|_{H^{-\lambda}} \| (\text{Ad}_{\partial_x})^\alpha (\text{Ad}_x)^\beta A(\Pi_{n'} \mathcal{U}) [e^{i\langle \xi, \cdot \rangle} \theta_2] \|_{H^\lambda}.$$

Using that  $\left\| e^{-i\langle \cdot, \xi \rangle} \theta_j \right\|_{H^\lambda} \sim \langle \xi \rangle^\lambda$  when  $|\xi| \rightarrow +\infty$  and inequality (1.2.9), we obtain

$$\left| \widehat{\partial_x^\alpha \partial_\xi^\beta a(\Pi_{n'} \mathcal{U}, \eta, \xi)} \right| |n'|^{N_2} \leq C \langle \xi + \eta \rangle^{-\lambda} \langle \xi \rangle^{\lambda + N_2 + m - |\beta|} |n'|^{\nu(p) + \nu + M_0(|\alpha| + |\beta|)} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

By Sobolev injection, we get the same estimate for

$$\left| \partial_x^\alpha \partial_\xi^\beta (\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} a(\Pi_{n'} \mathcal{U}, x, \xi)) \right|$$

if we replace in the right hand side  $\nu$  by  $\nu + M_0(d+1)$  and discard  $\langle \xi + \eta \rangle^{-\lambda} \langle \xi \rangle^\lambda$ . This gives the wanted estimate (2.1.2), showing that for any  $\lambda$ ,  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} a$  is in  $\widetilde{HS}_{p, \text{loc}}^{m, \nu}$ .

The proof of the analogous statement for operators in  $\Psi_{p, \text{loc}}^{m, \nu}(B_\sigma(R))$  is similar, using (1.2.18) and (1.2.19).

(ii) Let  $a$  be in  $\widetilde{HS}_{p, \text{loc}}^{m, \nu}$ . Since  $\theta_1 \text{Op}(a) \theta_2$  satisfies the support property, we just need to check estimates (1.2.8) for the operator  $\text{Ad}_{P_1} \cdots \text{Ad}_{P_k} [\theta_1 \text{Op}(a(\Pi_{n'} \mathcal{U}; \cdot)) \theta_2]$ . By the last remark following Definition 1.2.2, we may assume that  $P_\ell$  is of order zero or one, so either of the form  $b(x)$  or  $b(x) \frac{\partial}{\partial x_j}$  for some function  $b$  in  $C_0^\infty(\mathbb{R}^d)$ . Let  $\alpha$  (resp.  $\beta$ ) be the number of those  $\ell$  such that  $P_\ell$  is of order 1 (resp. 0). The kernel of  $[b(x) \frac{\partial}{\partial x_j}, \theta_1 \text{Op}(a(\mathcal{U}; \cdot)) \theta_2]$  (resp.  $[b(x), \theta_1 \text{Op}(a(\mathcal{U}; \cdot)) \theta_2]$ ) is

$$\frac{1}{(2\pi)^d} \int e^{i(x-y) \cdot \xi} \tilde{a}(\mathcal{U}, x, y, \xi) d\xi$$

with

$$\begin{aligned} \tilde{a}(\mathcal{U}, x, y, \xi) &= - \sum_{k=1}^d b_k(x, y) \theta_1(x) \theta_2(y) \frac{\partial}{\partial \xi_k} [a(\mathcal{U}, x, \xi) \xi_j] + b(x) \frac{\partial}{\partial x_j} [\theta_1(x) a(\mathcal{U}, x, \xi) \theta_2(y)] \\ &\quad + \frac{\partial}{\partial y_j} [b(y) \theta_1(x) a(\mathcal{U}, x, \xi) \theta_2(y)] \end{aligned}$$

with  $b_k$  a smooth function on  $\mathbb{R}^d \times \mathbb{R}^d$  (resp. with

$$\tilde{a}(\mathcal{U}, x, y, \xi) = i \sum_{k=1}^d b_k(x, y) \theta_1(x) \frac{\partial a}{\partial \xi_k}(\mathcal{U}, x, \xi) \theta_2(y).$$

Computing in the same way iterated commutators, and using Lemma 2.1.5, we conclude that  $\text{Ad}_{P_1} \cdots \text{Ad}_{P_k} [\theta_1 \text{Op}(a(\mathcal{U}; \cdot)) \theta_2] = \text{Op}(b(\mathcal{U}; \cdot))$ , where, for any  $\lambda \in \mathbb{R}$ ,

any  $N \in \mathbb{N}$ ,

(2.1.7)

$$|\partial_x^{\alpha'} \partial_\xi^{\beta'} \partial_y^{\gamma'} [(D_x + \xi)^\lambda \langle \xi \rangle^{-\lambda} b(\Pi_{n'} \mathcal{U}, x, y, \xi)]| \leq C \langle \xi \rangle^{m-\beta-|\beta'|} |n'|^{\nu(p)+\nu+M_0(\alpha+\beta+|\alpha'|+|\beta'|+d+1)} \\ \times \left(1 + \frac{|n'|}{\langle \xi \rangle}\right)^{-N} \prod_1^p \|\Pi_{n_\ell} u_\ell\|_{L^2},$$

and where  $\text{Op}(b)$  is the operator with kernel  $(2\pi)^{-d} \int e^{i(x-y)\cdot\xi} b(\mathcal{U}, x, y, \xi) d\xi$ . Let us estimate  $\|\Delta_j \text{Op}(b(\Pi_{n'} \mathcal{U}; \cdot)) \Delta_{j'}\|_{\mathcal{L}(L^2)}$ . We decompose  $b(\mathcal{U}, x, y, \xi) = \sum_{j''} b_{j''}(\mathcal{U}, x, y, \xi)$  with the notation  $b_{j''} = b\varphi(2^{-j''}|\xi|)$  for  $j \geq 1$ ,  $b_0 = b\psi(|\xi|)$ . The kernel  $K_{j''}(\mathcal{U}, x, y)$  of  $\Delta_j \text{Op}(b_{j''}) \Delta_{j'}$  may be written by a direct computation

$$K_{j''}(\mathcal{U}, x, y) = \frac{1}{(2\pi)^d} \int e^{i(x-y)\cdot\xi} c_{j''}(\mathcal{U}, x, y, \xi) d\xi$$

with

$$c_{j''}(\mathcal{U}, x, y, \xi) = \frac{1}{(2\pi)^{2d}} \int e^{i[z\zeta+z'\zeta']} \varphi(2^{-j}|\xi-\zeta|) \varphi(2^{-j'}|\xi-\zeta'|) \varphi(2^{-j''}|\xi|) \\ \times b(\mathcal{U}, x-z, y-z', \xi) dz dz' d\zeta d\zeta'$$

when  $j'' > 0, j > 0, j' > 0$ , and similar expressions when  $j, j'$  or  $j''$  is zero. If we write

$$b(\mathcal{U}, x-z, y-z', \xi) = \langle D_z - \xi \rangle^{-\lambda} \langle D_x + \xi \rangle^\lambda b(\mathcal{U}, x-z, y-z', \xi)$$

and integrate by parts in  $z$ , we get the expression

$$(2.1.8) \quad c_{j''}(\Pi_{n'} \mathcal{U}, x, y, \xi) = \frac{1}{(2\pi)^{2d}} \langle \xi \rangle^\lambda 2^{-j\lambda} \int e^{i[z\zeta+z'\zeta']} \phi_j(2^{-j}|\xi-\zeta|) \varphi(2^{-j'}|\xi-\zeta'|) \varphi(2^{-j''}|\xi|) \\ \times \tilde{b}(\mathcal{U}, x-z, y-z', \xi) dz dz' d\zeta d\zeta'$$

where  $\tilde{b}$  is the symbol in the left hand side of (2.1.7), and where  $\phi_j$  has the same support properties as  $\varphi$ , with estimates of its derivatives uniform in  $j$ . Let us show that, for any  $N, N'$

(2.1.9)

$$|\partial_\xi^{\beta'} c_{j''}(\Pi_{n'} \mathcal{U}, x, y, \xi)| \leq C 2^{j''(m-\beta)} [\max(2^{-j}, 2^{-j'}, 2^{-j''})]^{|\beta'|} 2^{-N'(|j-j'|+|j'-j''|)} \\ \times |n'|^{\nu(p)+\nu+M_0(\alpha+\beta+d+1+|\beta'|)} (1+2^{-j''}|n'|)^{-N} \prod_1^p \|u_\ell\|_{L^2}.$$

To do so, we perform  $N_1$  integrations by parts in (2.1.8) using the vector field  $L = (1+|\zeta'|^2)^{-1}(1+\zeta' \cdot D_{z'})$ . The estimate (2.1.7) show that we gain in that way a  $\langle \zeta' \rangle^{-N_1}$  factor. Next, perform  $N_2$  integrations by parts using  $L_2 = (1+|z'|^2)^{-1}(1+z' \cdot D_{\zeta'})$ . We gain a factor  $\langle z' \rangle^{-N_2}$ . Finally, we integrate by parts using  $L_3 = (1+2^{2j}|z|^2)^{-1}(1+2^{2j}z \cdot D_\zeta)$  to gain a factor  $\langle 2^j z \rangle^{-N_3}$ . We obtain for the modulus of  $\partial_\xi^{\beta'} c_{j''}$  a bound given by the product of

$$C 2^{j''(m-\beta)} [\max((2^{-j}, 2^{-j'}, 2^{-j''}))^{|\beta'|} |n'|^{\nu(p)+\nu+M_0(\alpha+\beta+d+1+|\beta'|)} (1+2^{-j''}|n'|)^{-N} \prod_1^p \|u_\ell\|_{L^2}$$

and of

$$2^{(j''-j)\lambda} \int \mathbb{1}_{|\xi-\zeta|\sim 2^j} \mathbb{1}_{|\xi-\zeta'|\sim 2^{j'}} \mathbb{1}_{|\xi|\sim 2^{j''}} \langle \zeta' \rangle^{-N_1} \langle z' \rangle^{-N_2} \langle 2^j z \rangle^{-N_3} dz dz' d\zeta d\zeta'.$$

If we integrate for  $|\zeta'| \geq c2^{|j'-j''|}$  for some  $c > 0$ , we get, choosing  $\lambda$  conveniently and  $N_1$  large enough, a factor  $O(2^{-N'(|j-j'|+|j'-j''|)})$ , whence the upper bound (2.1.9). If we integrate for  $|\zeta'| \ll 2^{|j'-j''|}$ , the cut-offs show that  $|j' - j''|$  has to stay smaller than a fixed constant, so that we get also the wanted estimate (2.1.9). If we go back to the kernel  $K_{j''}$ , perform  $d+1$  integrations by parts in  $\xi$  and use (2.1.9), we get a bound

$$|K_{j''}(\Pi_n \mathcal{U}, x, y)| \leq C 2^{j''d} [1 + \min(2^j, 2^{j'}, 2^{j''}) |x - y|]^{-d-1} 2^{-N'(|j-j'|+|j'-j''|)} \\ \times 2^{j''(m-\beta)} |n'|^{\nu(p)+\nu+M_0(\alpha+\beta+2(d+1))} (1 + 2^{-j''} |n'|)^{-N} \prod_{\ell=1}^p \|u_\ell\|_{L^2}.$$

The  $\mathcal{L}(L^2)$ -norm of  $\Delta_j \text{Op}(b) \Delta_{j'}$  is bounded from above by the sum in  $j''$  of the  $L^1(dx)$  (or  $L^1(dy)$ ) norm of  $K_{j''}(\Pi_n \mathcal{U}, x, y)$ , so by

$$C 2^{j(m-\beta)} |n'|^{\nu(p)+\nu+M_0(\alpha+\beta+2(d+1))} (1 + 2^{-j} |n'|)^{-N} 2^{-N'|j-j'|} \prod_{\ell=1}^p \|u_\ell\|_{L^2}$$

for some new value of  $N'$ . This shows that  $\theta_1 \text{Op}(a) \theta_2$  is in  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu+2M_0(d+1)}$  as claimed.

The proof of the corresponding statement in the framework of the  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$  class of symbols is identical.

(iii) This statement follows from the formula

$$(a - a_\chi)(\mathcal{U}, x, \xi) = \left[ \left( \frac{D_x}{\langle \xi \rangle} \right)^{-j} (1 - \chi) \left( \frac{D_x}{\langle \xi \rangle} \right) \right] \left( \frac{D_x}{\langle \xi \rangle} \right)^j a$$

for any  $j > 0$ .  $\square$

We have already obtained, in the global framework, results of composition of operators (see Lemma 1.2.4 and Lemma 1.2.6). Nevertheless, to see that a commutator between two operators is one order less than the sum of the orders, we need to study, for local operators, the symbol of the composition.

**THEOREM 2.1.7.** *Let  $m, m'$  be real numbers,  $\nu, \nu'$  in  $\mathbb{R}_+$ ,  $\sigma > \min(\nu(p), \nu(q)) + \min(\nu, \nu') + M_0(2d+3)$ . Assume that*

$$(2.1.10) \quad \nu(p) + \nu(q) + \nu' + M_0(2d+3) \leq \nu(p+q), \quad p > 0, q > 0.$$

*Let  $A$  be in  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu}$  (resp.  $\Psi_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ) and  $B$  be in  $\widetilde{H\Psi}_{q,\text{loc}}^{m',\nu'}$  (resp.  $\Psi_{q,\text{loc}}^{m',\nu'}(B_\sigma(R))$ ).*

*Denote by  $a$  and  $b$  the element of  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp.  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ) and  $\widetilde{HS}_{q,\text{loc}}^{m',\nu'}$  (resp.  $S_{q,\text{loc}}^{m',\nu'}(B_\sigma(R))$ ) given by (i) of Proposition 2.1.6. Let  $\chi \in C_0^\infty(\mathbb{R}^d)$  be equal to one close to zero, with small enough support. Then, if  $\theta \in C_0^\infty(\mathbb{R}^d)$  is equal to one on a large enough compact subset of  $\mathbb{R}^d$ , we may write for any  $\mathcal{U}'$  in  $C^\infty(X)^p$ ,  $\mathcal{U}''$  in  $C^\infty(X)^q$*

$$(2.1.11) \quad A(\mathcal{U}') \circ B(\mathcal{U}'') = \theta \text{Op}[(a(\mathcal{U}', \cdot) b(\mathcal{U}'', \cdot))_\chi] \theta + C(\mathcal{U}', \mathcal{U}''),$$

resp. for  $U$  in  $B_\sigma(R)$ ,

$$(2.1.12) \quad A(U) \circ B(U) = \theta \text{Op}[(a(U, \cdot) b(U, \cdot))_\chi] \theta + C(U),$$

where  $C$  is an element of

$$\widetilde{H\Psi}_{p+q, \text{loc}}^{m+m'-1, \nu+M_0(2d+3)} \quad (\text{resp. } \Psi_{\max(p,q), \text{loc}}^{m+m'-1, \max(\nu, \nu')+M_0(2d+3)}(B_\sigma(R))) \text{ if } p > 0, q > 0 \text{ and of}$$

$$\widetilde{H\Psi}_{p+q, \text{loc}}^{m+m'-1, \nu+\nu'+2M_0(2d+3)} \quad (\text{resp. } \Psi_{\max(p,q), \text{loc}}^{m+m'-1, \max(\nu, \nu')+M_0(2d+3)}(B_\sigma(R))) \text{ if } p = 0 \text{ or } q = 0.$$

**Remark:** Of course, if one assumes instead of (2.1.10),  $\nu(p) + \nu(q) + \nu + M_0(2d+3) \leq \nu(p+q)$  for  $p > 0, q > 0$ , one gets the same conclusion with  $\nu$  and  $\nu'$  exchanged.

*Proof:* Let  $A$  be in  $\widetilde{H\Psi}_{p, \text{loc}}^{m, \nu}$  and  $B$  in  $\widetilde{H\Psi}_{q, \text{loc}}^{m', \nu'}$ . Let  $\theta, \theta_1$  be in  $C_0^\infty(\mathbb{R}^d)$ , with  $\theta_1 \theta = \theta$  and  $\theta A \theta = A, \theta B \theta = B$ . By (iii) of Proposition 2.1.6, for any  $\lambda \in \mathbb{R}$ ,  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} (a - a_\chi)$  is in  $\widetilde{HS}_{p, \text{loc}}^{m-1, \nu+M_0}$  and we have a similar statement with  $b$ . Applying (ii) of Proposition 2.1.6, we may write

$$A = \theta_1 \text{Op}(a_\chi) \theta_1 + A_1, \quad B = \theta_1 \text{Op}(b_\chi) \theta_1 + B_1$$

with  $A_1 \in \widetilde{H\Psi}_{p, \text{loc}}^{m-1, \nu+M_0(2d+3)}$  and  $B_1 \in \widetilde{H\Psi}_{q, \text{loc}}^{m'-1, \nu'+M_0(2d+3)}$ . Using (2.1.10) and Lemma 1.2.4, we write

$$(2.1.13) \quad AB = \theta_1 \text{Op}(a_\chi) \theta_1^2 \text{Op}(b_\chi) \theta_1 + C_1$$

where  $C_1$  is a contribution to  $C$  in (2.1.11). The first term in the right hand side may be written  $\theta_1 \text{Op}(c(\mathcal{U}, \cdot)) \theta_1$  where

$$c(\mathcal{U}, x, \xi) = \frac{1}{(2\pi)^d} \int e^{-iz \cdot \zeta} a_\chi(\mathcal{U}', x, \xi - \zeta) \theta_1^2(x - z) b_\chi(\mathcal{U}'', x - z, \xi) dz d\zeta$$

if  $\mathcal{U} \in C^\infty(X)^{p+q}$  has been split in  $\mathcal{U} = (\mathcal{U}', \mathcal{U}'') \in C^\infty(X)^p \times C^\infty(X)^q$ . Then

$$(2.1.14) \quad c(\mathcal{U}, x, \xi) - a_\chi b_\chi(\mathcal{U}, x, \xi) \theta_1(x)^2 = \frac{1}{(2\pi)^d} \int e^{-iz \cdot \zeta} [a_\chi(\mathcal{U}', x, \xi - \zeta) - a_\chi(\mathcal{U}', x, \xi)] \theta_1^2(x - z) \\ \times b_\chi(\mathcal{U}'', x - z, \xi) dz d\zeta.$$

The right hand side may be written, performing an integration by parts, as

$$(2.1.15) \quad \frac{1}{(2\pi)^d} \int_0^1 \int e^{-iz \cdot \zeta} (\partial_\xi a_\chi)(\mathcal{U}', x, \xi - t\zeta) D_x [\theta_1^2(x - z) b_\chi(\mathcal{U}'', x - z, \xi)] dz d\zeta dt.$$

We perform  $d+1$  integrations by parts using the vector field  $L = (1 + \zeta^2)^{-1} (1 - \zeta \cdot D_z)$  to get an absolutely convergent integral in  $\zeta$ . If we replace in (2.1.15)  $\mathcal{U}'$  by  $\Pi_n \mathcal{U}'$ ,  $\mathcal{U}''$  by  $\Pi_{n'} \mathcal{U}''$ , each  $D_z$  derivative acting on  $b_\chi$  makes lose a power  $|n''|^{M_0}$ . We decompose  $\theta_1^2$  as

$$(2.1.16) \quad \theta_1^2(w) = (\chi(D/\langle \xi \rangle) \theta_1^2)(w) + \tilde{\theta}(w, \xi)$$

where  $|\partial_w^\gamma \tilde{\theta}(w, \xi)| = O(\langle \xi \rangle^{-N} \langle w \rangle^{-N})$  for any  $N, \gamma$ . Let us study first the contribution to (2.1.15) coming from the first term in the right hand side of (2.1.16). If  $\text{Supp } \chi$  is small enough, the support properties of  $\tilde{b}_\chi(\mathcal{U}'', \eta, \xi)$  imply that, in

(2.1.15), we may assume  $|\zeta| \ll |\xi|$  on the domain of integration. This integral, computed at  $\Pi_n \mathcal{U}$  instead of  $\mathcal{U}$ , is thus bounded from above in modulus by

$$C|n'|^{\nu(p)+\nu+M_0(d+2)}|n''|^{\nu(q)+\nu'+M_0(2d+3)}\langle \xi \rangle^{m+m'-1} \left(1 + \frac{|n'|}{\langle \xi \rangle}\right)^{-N} \left(1 + \frac{|n''|}{\langle \xi \rangle}\right)^{-N} \prod_1^{p+q} \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

Using (2.1.10), we get estimates (2.1.2) of an element of  $\widetilde{HS}_{p+q, \text{loc}}^{m+m'-1, \nu+M_0}$  if  $p, q > 0$  and of  $\widetilde{HS}_{p+q, \text{loc}}^{m+m'-1, \nu+M_0(d+2)}$  if  $p$  or  $q$  is zero. The derivatives are bounded in the same way, and the symbol obtained making act  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda}$  on (2.1.15) satisfies similar estimates, again because the  $x$ -Fourier transform of (2.1.15) is supported for  $|\eta| \ll |\xi|$  if  $\text{Supp } \chi$  is small enough. If we consider the contribution of  $\tilde{\theta}$  in (2.1.16) to (2.1.15), and use that  $\partial_w^\gamma \tilde{\theta} = O(\langle \xi \rangle^{-N} \langle w \rangle^{-N})$  for any  $N$ , and the fact that  $L$ -integrations by parts allow one to gain any  $\langle \xi \rangle^{-N} \langle \xi \rangle^N$  factor, one sees that the corresponding contribution to (2.1.15) is in  $\widetilde{HS}_{p+q, \text{loc}}^{-N, 0}$  for any  $N$ , as well as the action of  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda}$  on that function for any  $\lambda$ . It follows from (2.1.13) and from (ii) of Proposition 2.1.6 that

$$(2.1.17) \quad AB = \theta_1 \text{Op}[\theta_1^2 a_\chi b_\chi] \theta_1$$

modulo an operator  $C$  satisfying the conclusion of the theorem. Since  $AB = \theta AB \theta$ , we may write as well, modulo such a  $C$ ,  $AB = \theta \text{Op}[a_\chi b_\chi] \theta$ . Write  $a_\chi b_\chi = (ab)_\chi - r$  with

$$r = [(a_\chi b_\chi)_\chi - (a_\chi b_\chi)] + ((a - a_\chi)b)_\chi + (a_\chi(b - b_\chi))_\chi.$$

By (iii) of proposition 2.1.6 and (2.1.10),  $r$  belongs to  $\widetilde{HS}_{p+q, \text{loc}}^{m+m'-1, \nu+M_0}$  if  $p > 0, q > 0$  and to  $\widetilde{HS}_{p+q, \text{loc}}^{m+m'-1, \nu+M_0(d+2)}$  if  $p$  or  $q$  is zero. The associated operator contributes to  $C$  in (2.1.12). This concludes the proof in the case of homogeneous operators. The case of operators in  $\Psi_{p, \text{loc}}^{m, \nu}(B_\sigma(R))$ ,  $\Psi_{q, \text{loc}}^{m', \nu'}(B_\sigma(R))$  is treated similarly.  $\square$

Let us finish this subsection with a similar result for transpose and adjoint.

**THEOREM 2.1.8.** *Let  $m \in \mathbb{R}, \nu \in \mathbb{R}_+, p \in \mathbb{N}, R > 0, \sigma > 0$ . Let  $A$  be an element of the space  $\widetilde{H\Psi}_{p, \text{loc}}^{m, \nu}$  (resp.  $\Psi_{p, \text{loc}}^{m, \nu}(B_\sigma(R))$ ) and let  $a$  be a symbol in  $\widetilde{HS}_{p, \text{loc}}^{m, \nu}$  (resp.  $S_{p, \text{loc}}^{m, \nu}(B_\sigma(R))$ ) given by (i) of Proposition 2.1.6. If  $\chi$  is in  $C_0^\infty(\mathbb{R})$ , has small enough support, and is equal to one close to zero, if  $\theta$  is in  $C_0^\infty(\mathbb{R}^d)$  and is equal to one on a large enough compact set, then, denoting by  ${}^t A$  and  $A^*$  the transpose and the adjoint of  $A$  for the  $L^2(\mathbb{R}^d, dx)$ -scalar product,*

$$(2.1.18) \quad {}^t A = \theta \text{Op}(a_\chi^\vee) \theta + C_1, \quad A^* = \theta \text{Op}(\bar{a}_\chi) \theta + C_2$$

where  $C_1, C_2$  are in  $\widetilde{H\Psi}_{p, \text{loc}}^{m-1, \nu+M_0(3d+5)}$  (resp.  $\Psi_{p, \text{loc}}^{m-1, \nu+M_0(3d+5)}(B_\sigma(R))$ ) and where  $a^\vee(x, \xi) = a(x, -\xi)$ .

*Proof:* Because of the support assumption on the kernel of  $A$  and of (iii) of Proposition 2.1.6, we know that we may write  $A(\mathcal{U}) = \theta \text{Op}(a_\chi) \theta$  modulo a remainder of

form  $C_1, C_2$ . Moreover,  ${}^t\text{Op}(a_\chi) = \text{Op}(b_1)$ ,  $\text{Op}(a_\chi)^* = \text{Op}(b_2)$  with

$$(2.1.19) \quad \begin{aligned} b_1(x, \xi) &= \frac{1}{(2\pi)^d} \int e^{-iz \cdot \zeta} a_\chi(x - z, -\xi + \zeta) dz d\zeta \\ b_2(x, \xi) &= \frac{1}{(2\pi)^d} \int e^{-iz \cdot \zeta} \bar{a}_\chi(x - z, \xi - \zeta) dz d\zeta. \end{aligned}$$

Because of the support of  $\widehat{a}_\chi(\zeta, -\xi + \zeta)$ ,  $\widehat{\bar{a}}_\chi(\zeta, \xi - \zeta)$ , we may insert a cut-off  $\chi_1(\zeta/\langle \xi \rangle)$  inside the integrals in (2.1.19) if  $\chi_1 \equiv 1$  close to zero,  $\text{Supp } \chi_1$  small enough. Then

$$b_1(x, \xi) - a_\chi(x, -\xi) = \frac{1}{(2\pi)^d} \int e^{-iz \cdot \zeta} \chi_1(\zeta/\langle \xi \rangle) [a_\chi(x - z, -\xi + \zeta) - a_\chi(x, -\xi)] dz d\zeta.$$

Since the term between brackets vanishes at  $(z = 0, \zeta = 0)$ , we see, performing  $d+1$  integrations by parts using  $L = \frac{1-i\langle \xi \rangle^2 z \cdot \nabla_\zeta}{1+\langle \xi \rangle^2 |z|^2}$  that  $b_1(x, \xi) - a_\chi(x, -\xi)$  may be written as  $c_{\chi_1}(x, \xi)$  for some symbol  $c$  in  $\widetilde{HS}_{p, \text{loc}}^{m-1, \nu+M_0(d+3)}$  (resp.  $\Psi_{p, \text{loc}}^{m-1, \nu+M_0(d+3)}(B_\sigma(R))$ ). Using (ii) of Proposition 2.1.6, we get that  $C_1 = \text{Op}(c_{\chi_1})$  satisfies the conclusion of the theorem. One treats in the same way the case of the adjoint.  $\square$

**2.2. Principal symbols.** In this subsection, we shall define principal symbols for para-differential operators on the manifold  $X$ , and use them to study commutators between such operators. We first study the action of a change of variables on local operators. If  $X, X'$  are two manifolds and  $\kappa : X \rightarrow X'$  is a diffeomorphism, we denote by  $\tau_\kappa : T^*X \rightarrow T^*X'$  the canonical transformation induced by  $\kappa$ . In local coordinates

$$(2.2.1) \quad \tau_\kappa : (x, \xi) \rightarrow \tau_\kappa(x, \xi) = (\kappa(x), {}^t d\kappa(x)^{-1} \xi).$$

We denote by  $\kappa^* : C_0^\infty(X') \rightarrow C_0^\infty(X)$  the map defined by  $\kappa^* u = u \circ \kappa$ .

**PROPOSITION 2.2.1.** *Let  $m \in \mathbb{R}, \nu \in \mathbb{R}_+, R > 0, \sigma > 0, V, \tilde{V}$  be two open sets of  $\mathbb{R}^d$ ,  $\kappa : V \rightarrow \tilde{V}$  be a  $C^\infty$ -diffeomorphism, let  $\theta, \theta', \tilde{\theta}, \tilde{\theta}'$  be in  $C_0^\infty(V)$  with  $\theta\tilde{\theta} = \theta, \theta'\tilde{\theta}' = \theta'$ . Let  $A(\mathcal{U})$  (resp.  $A(U)$ ) be an element of  $\widetilde{H\Psi}_{p, \text{loc}}^{m, \nu}$  (resp.  $\Psi_{p, \text{loc}}^{m, \nu}(B_\sigma(R))$ ). Then  $B = (\kappa^{-1})^* \theta A \theta' \kappa^*$  is in  $\widetilde{H\Psi}_{p, \text{loc}}^{m, \nu}$  (resp.  $\Psi_{p, \text{loc}}^{m, \nu}(B_\sigma(R))$ ). Moreover, if  $\chi_1 \in C_0^\infty(\mathbb{R})$  equals one close to zero and has small enough support, if  $a$  is the symbol in  $\widetilde{HS}_{p, \text{loc}}^{m, \nu}$  (resp.  $S_{p, \text{loc}}^{m, \nu}(B_\sigma(R))$ ) associated to  $A$  by (i) of Proposition 2.1.6, we may write*

$$(2.2.2) \quad B = (\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta\theta' a \circ \tau_\kappa^{-1})_{\chi_1}] (\tilde{\theta}' \circ \kappa^{-1}) + C$$

where  $C$  is in  $\widetilde{H\Psi}_{p, \text{loc}}^{m-1, \nu+M_0(3d+4)}$  (resp.  $\Psi_{p, \text{loc}}^{m-1, \nu+M_0(3d+4)}(B_\sigma(R))$ ).

Since  $\theta' \kappa^* = \theta' \kappa^* (\tilde{\theta}' \circ \kappa^{-1})$ , this operator preserves Sobolev spaces. The characterization of  $\widetilde{H\Psi}_p^{m, \nu}$  and  $\Psi_p^{m, \nu}(B_\sigma(R))$  in terms of estimates of type (1.2.9), (1.2.18), (1.2.19) thus shows that  $B$  is in the indicated space of operators. We just have to show the statements concerning symbols.

**LEMMA 2.2.2.** (i) *Let  $W$  be a neighborhood of the diagonal in  $X \times X$ . Let  $(\theta_i)_{i \in I}$  be a smooth partition of unity on  $X$  such that  $\theta_i(x)\theta_j(y) \neq 0$  implies  $(x, y) \in W$ . Let  $J = \{(i, j) \in I \times I; \text{Supp } \theta_i \cap \text{Supp } \theta_j \neq \emptyset\}$ . For any  $A$  in  $\widetilde{H\Psi}_p^{m, \nu}$  (resp.  $\Psi_p^{m, \nu}(B_\sigma(R))$ ),  $A - \sum_{(i, j) \in J} \theta_i A \theta_j$  is in  $\widetilde{H\Psi}_p^{m-r, \nu+rM_0}$  (resp.  $\Psi_p^{m-r, \nu+rM_0}(B_\sigma(R))$ ) for any  $r \geq 0$ .*

(ii) Let  $W$  be a neighborhood of the diagonal in  $\mathbb{R}^d \times \mathbb{R}^d$ ,  $K$  a compact subset of  $\mathbb{R}^d$ ,  $\chi \in C_0^\infty(\mathbb{R}^d)$ ,  $\chi \equiv 1$  close to zero, with small enough support. Let  $(\theta_i)_{i \in I}$  be a smooth partition of unity on a neighborhood of  $K$ . For any  $A$  in  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu}$  (resp.  $\Psi_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ) with support in  $K \times K$ , there is a symbol  $a$  in  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp.  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ), compactly supported in  $x$ , such that  $A - \sum_{(i,j) \in J} \theta_i \text{Op}(a_\chi) \theta_j$  is in  $\widetilde{H\Psi}_{p,\text{loc}}^{m-r,\nu+M_0(r+2d+2)}$  (resp.  $\Psi_p^{m-r,\nu+M_0(r+2d+2)}(B_\sigma(R))$ ) for any  $r \geq 0$ .

*Proof:* (i) We notice that if  $\theta_1, \theta_2$  are smooth functions with disjoint support,  $\theta_1 A \theta_2$  is in  $\widetilde{H\Psi}_p^{m-r,\nu+rM_0}$  (resp.  $\Psi_p^{m-r,\nu+rM_0}(B_\sigma(R))$ ) for any  $r \geq 0$ : actually, if  $\tilde{\theta}$  is in  $C^\infty(X)$  and  $\tilde{\theta}\theta_1 = \theta_1$ ,  $\tilde{\theta}\theta_2 = 0$ , we may write for any  $r \in \mathbb{N}$

$$\theta_1 A \theta_2 = \underbrace{\text{Ad}_{\tilde{\theta}} \cdots \text{Ad}_{\tilde{\theta}}}_r (\theta_1 A \theta_2).$$

We just have to apply the characterization of  $\widetilde{H\Psi}_p^{m,\nu}$  (resp.  $\Psi_p^{m,\nu}(B_\sigma(R))$ ) to get the conclusion.

(ii) The same proof as above, together with the fact that  $A$  is supported in  $K \times K$ , shows that  $A - \sum_{(i,j) \in J} \theta_i A \theta_j$  is in the wanted space. By (i), (ii), (iii) of Proposition 2.1.6, there is a symbol  $a$  in  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu}$  (resp.  $\Psi_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ) such that  $\theta_i A \theta_j - \theta_i \text{Op}(a_\chi) \theta_j$  belongs to  $\widetilde{H\Psi}_{p,\text{loc}}^{m-r,\nu+M_0(r+2d+2)}$  (resp.  $\Psi_{p,\text{loc}}^{m-r,\nu+M_0(r+2d+2)}(B_\sigma(R))$ ). We may assume moreover that  $a$  is compactly supported in a neighborhood of  $K$ . This concludes the proof.  $\square$

**LEMMA 2.2.3.** *Let  $\kappa$  be as in the statement of proposition 2.2.1. Let  $a$  be in  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp.  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ), compactly supported in  $x$  in a compact subset of  $V$ . Let  $\chi \in C_0^\infty(\mathbb{R}^d)$ ,  $\chi \equiv 1$  close to zero, with small enough support. Let  $(x, z) \rightarrow \gamma(x, z)$  be a smooth function, such that  $(x, y) \rightarrow \gamma(x, x - y)$  is compactly supported in  $\tilde{V} \times \tilde{V}$ , and such that  $(x, z) \rightarrow \gamma(x, z)$  is supported for  $|z| \leq \rho \ll 1$ . Define for  $\mathcal{U} = (U_1, \dots, U_p) \in C^\infty(X)^p$  (resp. for  $U \in B_\sigma(R)$ )*

$$(2.2.3) \quad b(\mathcal{U}, x, \xi) = \frac{1}{(2\pi)^d} \int e^{i[\kappa^{-1}(x) - \kappa^{-1}(x-z)]\zeta - iz \cdot \xi} \gamma(x, z) a_\chi(\mathcal{U}, \kappa^{-1}(x), \zeta) dz d\zeta$$

(resp.  $b(U, x, \xi)$  given by the same formula with  $\mathcal{U}$  replaced by  $U$ ). Let  $\Gamma \in C_0^\infty(\mathbb{R} - \{0\})$  be equal to one on a large enough compact subset of  $\mathbb{R} - \{0\}$ . Then one may write

$$(2.2.4) \quad b(\mathcal{U}, x, \xi) = \frac{1}{(2\pi)^d} \int e^{i[(x-x') \cdot \eta' + z \cdot (\zeta' - \xi) + (\kappa^{-1}(x') - x'') \cdot \eta'']} \\ \times c(\mathcal{U}, x', x'', z, \eta', \eta'', \zeta', \xi) dx' dx'' dz d\eta' d\eta'' d\zeta' + b_1(\mathcal{U}, x, \xi)$$

where, for any  $\lambda \in \mathbb{R}$ ,  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} b_1$  is in  $H\Psi_{p,\text{loc}}^{m-r,\nu}$  for all  $r \geq 0$  (resp.  $b(U, x, \xi)$  may be written as (2.2.4) with  $\mathcal{U}$  replaced by  $U$  and  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda} b_1$  belonging to  $\Psi_{p,\text{loc}}^{m-r,\nu}(B_\sigma(R))$  for any  $r \geq 0$ ,  $\lambda \in \mathbb{R}$ ), and where  $c$  is a function supported for

$$(2.2.5) \quad |\eta''| \ll \langle \zeta' \rangle \sim \langle \xi \rangle \sim \langle \xi + \eta' \rangle$$

and satisfies estimates

$$(2.2.6) \quad |\partial_{x'}^{\alpha'} \partial_{x''}^{\alpha''} \partial_z^\delta \partial_\xi^\beta \partial_{\eta'}^{\beta'} \partial_{\eta''}^{\beta''} \partial_{\zeta'}^\gamma c(\Pi_n \mathcal{U}, x', x'', z, \eta', \eta'', \zeta', \xi)| \\ \leq C \langle \xi \rangle^{m-|\beta|-|\beta'|-|\beta''|-|\gamma|} |n'|^{\nu(p)+\nu+M_0[|\alpha'|+|\alpha''|+|\delta|+|\gamma|+d+1]} \left(1 + \frac{|n'|}{\langle \xi \rangle}\right)^{-N} \prod_1^p \|u_\ell\|_{L^2},$$

resp.  $c$  satisfies the estimates

$$(2.2.7) \quad |\partial_{x'}^{\alpha'} \partial_{x''}^{\alpha''} \partial_z^\delta \partial_\xi^\beta \partial_{\eta'}^{\beta'} \partial_{\eta''}^{\beta''} \partial_{\zeta'}^\gamma c(U, x', x'', z, \eta', \eta'', \zeta', \xi)| \\ \leq C \langle \xi \rangle^{m-|\beta|-|\beta'|-|\beta''|-|\gamma|+(\nu(p)+\nu+M_0[|\alpha'|+|\alpha''|+|\delta|+|\gamma|+d+1]-\sigma)_+} \|U\|_{H^\sigma}^p,$$

and the estimate

$$(2.2.8) \quad |\partial_{x'}^{\alpha'} \partial_{x''}^{\alpha''} \partial_z^\delta \partial_\xi^\beta \partial_{\eta'}^{\beta'} \partial_{\eta''}^{\beta''} \partial_{\zeta'}^\gamma [\partial_U c(U, x', x'', z, \eta', \eta'', \zeta', \xi) \cdot H]| \\ \leq C \langle \xi \rangle^{m-|\beta|-|\beta'|-|\beta''|-|\gamma|+(\nu(p)+\nu+M_0[|\alpha'|+|\alpha''|+|\delta|+|\gamma|+d+1]-\sigma')_+} \|U\|_{H^{\sigma'}}^{p-1} \|H\|_{H^{\sigma'}},$$

for any  $\sigma' \in [-\sigma, \sigma]$ , any  $H$  in  $H^{\sigma'}$ .

Moreover, the restriction of  $c$  at  $z = 0$  is given by

$$(2.2.9) \quad c|_{z=0} = \frac{1}{(2\pi)^d} \gamma(x', 0) |\det \kappa(\kappa^{-1}(x'))| \chi(\eta'' / \langle {}^t d\kappa(\kappa^{-1}(x')) \cdot \zeta' \rangle) \\ \times \Gamma\left(\frac{\langle \zeta' \rangle}{\langle \xi \rangle}\right) \Gamma\left(\frac{\langle \zeta' \rangle}{\langle \xi + \eta' \rangle}\right) a(\mathcal{U}, x'', {}^t d\kappa(\kappa^{-1}(x')) \cdot \zeta')$$

(resp. by the same formula with  $\mathcal{U}$  replaced by  $U$ ).

*Proof:* Let us prove the lemma in the case  $a \in \widetilde{HS}_{p,\text{loc}}^{m,\nu}$ . We may write the definition (2.2.3) of  $b$

$$(2.2.10) \quad b(\mathcal{U}, x, \xi) = \frac{1}{(2\pi)^{3d}} \int e^{i[(x-x')\eta' + (\kappa^{-1}(x') - \kappa^{-1}(x'-z))\zeta + (\kappa^{-1}(x') - x'')\eta'' - z\xi]} \gamma(x', z) \chi(\eta'' / \langle \zeta \rangle) \\ \times a(\mathcal{U}, x'', \zeta) dx' dx'' dz d\eta' d\eta'' d\zeta.$$

On the support of  $\gamma(x', z) \Gamma(\langle \nabla_{x'}(\kappa^{-1}(x' - z) \cdot \zeta) \rangle / \langle \xi \rangle)$ , the quotient  $\langle \zeta \rangle / \langle \xi \rangle$  stays between two uniform constants (since  $x' - z$  stays in a compact subset of  $\tilde{V}$ ), and the function  $\Gamma_1(x' - z, \zeta, \xi) = (1 - \Gamma)(\langle \nabla_{x'}(\kappa^{-1}(x' - z) \cdot \zeta) \rangle / \langle \xi \rangle)$  satisfies

$$(2.2.11) \quad \partial_{x'}^{\alpha'} \partial_z^\delta \Gamma_1(x' - z, \zeta, \xi) = O(1).$$

Define  $b'_1$  as the function given by formula (2.2.10) with  $\Gamma_1$  inserted inside the integral, and define

$$L_1 = \frac{1 - i(\nabla_{x'}(\kappa^{-1}(x' - z) \cdot \zeta) - \xi) \cdot \nabla_z}{1 + |\nabla_{x'}(\kappa^{-1}(x' - z) \cdot \zeta) - \xi|^2}.$$

Performing integrations by parts using  $L_1$ , and denoting by  $\Phi$  the phase function in (2.2.10), we write

$$(2.2.12) \quad b'_1(\mathcal{U}, x, \xi) = \frac{1}{(2\pi)^{3d}} \int e^{i\Phi(x, x', x'', z, \eta', \eta'', \zeta, \xi)} \\ \times ({}^t L_1)^N [\gamma(x', z) \chi(\eta'' / \langle \zeta \rangle) \Gamma_1(x' - z, \zeta, \xi) a(\mathcal{U}, x'', \zeta)] dx' dx'' dz d\eta' d\eta'' d\zeta.$$



We note that, if the domain over which  $\Gamma \equiv 1$  is large enough, on the support of  $\Gamma_1 = 1 - \Gamma$ , either  $\langle \zeta \rangle \gg \langle \xi \rangle$  or  $\langle \zeta \rangle \ll \langle \xi \rangle$ , so that every  $L_1$ -integration by parts will provide a gain of  $(1 + |\zeta| + |\xi|)^{-1}$ . Consequently, the symbol in the integral in (2.2.12) satisfies the same estimates as  $a$ , as well as its derivatives, with a gain in  $(1 + |\zeta| + |\xi|)^{-N}$ . If we insert inside the integral another cut-off  $\omega(\eta'/(1 + \xi^2 + \zeta^2)^{1/2})$ , with  $\omega$  in  $C_0^\infty(\mathbb{R}^d)$ , and use that the integrand is compactly supported in  $(x', x'', z)$ , we obtain for the corresponding contribution to  $b'_1$  a bound in

$$(2.2.13) \quad C_N \langle \xi \rangle^{-N'} |n'|^{\nu(p)+\nu} \left(1 + \frac{|n'|}{\langle \xi \rangle}\right)^{-N'}$$

if  $N \gg N'$ . Similar estimates hold for derivatives. Moreover, if we make act on  $b'_1$  the operator  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda}$ , and use that  $\langle \eta' + \xi \rangle$  is controlled by the gain coming from the integrations by parts, we still get a bound of type (2.2.13).

Consider now the contribution to  $b'_1$  given by (2.2.12) in which we insert  $(1 - \omega)(\eta'/(1 + \xi^2 + \zeta^2)^{1/2})$ , with  $\omega \equiv 1$  on a large enough compact set. We consider

$$L_2 = \frac{1 - i(\nabla_{x'}(\kappa^{-1}(x') \cdot (\eta'' + \zeta)) - (\eta' + \xi)) \cdot (\nabla_{x'} + \nabla_z)}{1 + |\nabla_{x'}(\kappa^{-1}(x') \cdot (\eta'' + \zeta)) - (\eta' + \xi)|^2}$$

and perform  $N$  integrations by parts using  $L_2$ . We get

$$\begin{aligned} & \frac{1}{(2\pi)^{3d}} \int e^{i\Phi(x, x', x'', z, \eta', \eta'', \zeta, \xi)} ({}^t L_2)^N \left[ (1 - \omega) \left( \frac{\eta'}{(1 + \xi^2 + \zeta^2)^{1/2}} \right) \right. \\ & \quad \times \left. ({}^t L_1)^N [\gamma(x', z) \chi(\eta''/\langle \zeta \rangle) \Gamma_1(x' - z, \zeta, \xi) a(\mathcal{U}, x'', \zeta)] \right] dx' dx'' dz d\eta' d\eta'' d\zeta. \end{aligned}$$

Since on the support of the integrand,  $|\eta''| \ll \langle \zeta \rangle$ ,  $|\eta'| \gg (1 + |\xi| + |\zeta|)$ , the integrations by parts gain a  $(1 + |\eta'| + |\eta''| + |\zeta| + |\xi|)^{-N}$  factor. We obtain again a quantity bounded by (2.2.13), with similar estimates for the derivatives or for the action of  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda}$  for any  $\lambda$ . Consequently,  $b'_1$  contributes to  $b_1$  in (2.2.4). Let us study next  $b''_1$ , which is given by (2.2.10) where we insert the cut-off  $\Gamma(\langle \nabla_{x'}(\kappa^{-1}(x' - z) \cdot \zeta) \rangle / \langle \xi \rangle)$ . In particular,  $\langle \zeta \rangle \sim \langle \xi \rangle$  on the domain of integration. We decompose  $b''_1 = b_0 + b'''_1$  with

$$(2.2.14) \quad \begin{aligned} b_0(\mathcal{U}, x, \xi) &= \frac{1}{(2\pi)^{3d}} \int e^{i\Phi(x, x', x'', z, \eta', \eta'', \zeta, \xi)} \Gamma \left( \frac{\langle \nabla_{x'}(\kappa^{-1}(x') \cdot \zeta) \rangle}{\langle \xi + \eta' \rangle} \right) \\ & \quad \times \gamma(x', z) \chi(\eta''/\langle \zeta \rangle) \Gamma \left( \frac{\langle \nabla_{x'}(\kappa^{-1}(x' - z) \cdot \zeta) \rangle}{\langle \xi \rangle} \right) a(\mathcal{U}, x'', \zeta) dx' dx'' dz d\eta' d\eta'' d\zeta \end{aligned}$$

and  $b'''_1$  being given by the same integral, with the first  $\Gamma$  replaced by  $1 - \Gamma$ . We study first  $b'''_1$ , performing  $N$  integrations by parts using  $L_2$ . We get

$$(2.2.15) \quad \begin{aligned} & \frac{1}{(2\pi)^{3d}} \int e^{i\Phi(x, x', x'', z, \eta', \eta'', \zeta, \xi)} ({}^t L_2)^N \left[ (1 - \Gamma) \left( \frac{\langle \nabla_{x'}(\kappa^{-1}(x') \cdot \zeta) \rangle}{\langle \xi + \eta' \rangle} \right) \right. \\ & \quad \times \gamma(x', z) \chi(\eta''/\langle \zeta \rangle) \Gamma \left( \frac{\langle \nabla_{x'}(\kappa^{-1}(x' - z) \cdot \zeta) \rangle}{\langle \xi \rangle} \right) a(\mathcal{U}, x'', \zeta) \left. \right] dx' dx'' dz d\eta' d\eta'' d\zeta. \end{aligned}$$

On the support of the domain of integration,  $|\eta''| \ll \langle \zeta \rangle$ ,  $\langle \zeta \rangle \sim \langle \xi \rangle$  and either  $\langle \zeta \rangle \ll \langle \xi + \eta' \rangle$  or  $\langle \zeta \rangle \gg \langle \xi + \eta' \rangle$ . This implies that the integrations by parts gain a factor  $(1 + |\zeta| + |\xi + \eta'|)^{-N}$ , and that (2.2.15) has a bound of form (2.2.13), as well as the action of  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda}$  on this integral. Since derivatives are estimated in the same way, we conclude that  $b'''_1$  contributes to  $b_1$  in (2.2.4).

Finally, we are left with showing that  $b_0$  may be written as the first term in the right hand side of (2.2.4). Since the cut-off  $\gamma$  is supported for  $|z'| \leq \rho \ll 1$ , we may write, if  $\rho$  is small enough,  $\kappa^{-1}(x') - \kappa^{-1}(x' - z) = {}^t B(x', z)^{-1} z$ , where  $B(x', 0) = {}^t d\kappa(\kappa^{-1}(x'))$  and  $B(x', z)$  is an invertible matrix for any  $z$  with  $|z| \ll \rho$ . Setting  $\zeta = B(x', z) \cdot \zeta'$  in (2.2.14), we rewrite

$$b_0(\mathcal{U}, x, \xi) = \frac{1}{(2\pi)^{2d}} \int e^{i[(x-x') \cdot \eta' + z \cdot (\zeta' - \xi) + (\kappa^{-1}(x') - x'') \cdot \eta'']} \\ \times c(\mathcal{U}, x', x'', z, \eta', \eta'', \zeta', \xi) dx' dx'' dz d\eta' d\eta'' d\zeta'$$

with

$$c(\mathcal{U}, x', x'', z, \eta', \eta'', \zeta', \xi) = \frac{\tilde{\gamma}(x', z)}{(2\pi)^d} \chi\left(\frac{\eta''}{\langle B(x', z) \cdot \zeta' \rangle}\right) \Gamma\left(\frac{\langle (\nabla_{x'}(\kappa^{-1}(x')) \cdot B(x', z) \cdot \zeta') \rangle}{\langle \xi + \eta' \rangle}\right) \\ \times \Gamma\left(\frac{\langle \nabla_{x'}(\kappa^{-1}(x' - z)) \cdot (B(x', z) \cdot \zeta') \rangle}{\langle \xi \rangle}\right) a(\mathcal{U}, x'', B(x', z) \cdot \zeta')$$

for another smooth function  $\tilde{\gamma}$ , compactly supported in  $(x', z)$  in  $|z| \leq \rho \ll 1$ , such that

$$\tilde{\gamma}(x', 0) = \gamma(x', 0) |\det d\kappa(\kappa^{-1}(x'))|.$$

We note that (2.2.5) and estimates (2.2.6) hold true, and that  $c|_{z=0}$  is given by (2.2.9). This concludes the proof of Lemma 2.2.3, as the case of symbols in  $S_{p, \text{loc}}^{m, \nu}(B_\sigma(R))$  is identical.  $\square$

*Proof of proposition 2.2.1:* We consider  $A$  in  $\widetilde{H\Psi}_{p, \text{loc}}^{m, \nu}$ . According to Lemma 2.2.2, we may find a  $C_0^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ -function,  $(x, z) \rightarrow \gamma_1(x, z)$ , supported for  $|z| \leq \rho \ll 1$ , satisfying  $\theta\theta'(x)\gamma_1(x, 0) = \theta\theta'(x)$  and such that  $A(\mathcal{U})$  is given, up to an operator in  $\widetilde{H\Psi}_{p, \text{loc}}^{m-r, \nu+M_0(r+2d+2)}$  for any  $r$ , by the operator  $\tilde{A}(\mathcal{U})$  with kernel

$$K(\mathcal{U}, x, y) = \frac{1}{(2\pi)^d} \int e^{i(x-y) \cdot \xi} \gamma_1(x, x-y) a_\chi(\mathcal{U}, x, \xi) d\xi$$

where  $a$  is an element of  $\widetilde{H\Psi}_{p, \text{loc}}^{m, \nu}$ , compactly supported in  $x$ . The distribution kernel of the operator  $(\kappa^{-1})^* \theta \tilde{A} \theta' \kappa^*$  will be

$$\tilde{\theta} \circ \kappa^{-1}(x) \frac{1}{(2\pi)^d} \int e^{i(\kappa^{-1}(x) - \kappa^{-1}(y)) \cdot \xi} \gamma(x, x-y) a_\chi(\mathcal{U}, \kappa^{-1}(x), \xi) d\xi \tilde{\theta}' \circ \kappa^{-1}(y)$$

where

$$\gamma(x, z) = \theta \circ \kappa^{-1}(x) \gamma_1(\kappa^{-1}(x), \kappa^{-1}(x) - \kappa^{-1}(x-z)) \theta' \circ \kappa^{-1}(x-z) |\det d\kappa^{-1}(x-z)|.$$

Consequently,  $(\kappa^{-1})^* \theta \tilde{A}(\mathcal{U}) \theta' \kappa^* = (\tilde{\theta} \circ \kappa^{-1}) \text{Op}(b(\mathcal{U}, \cdot)) (\tilde{\theta}' \circ \kappa^{-1})$  where  $b$  is given by (2.2.3). We apply Lemma 2.2.3. By (2.2.4) and (ii) of proposition 2.1.6, we may replace  $b$  by the first term in the right hand side of (2.2.4), up to an operator in  $\widetilde{H\Psi}_{p, \text{loc}}^{m-r, \nu+2M_0(d+1)}$  for any  $r \geq 0$ . Write  $c = c_0 + c_1$  where  $c_0 = c|_{z=0}$  is given by (2.2.9). The contribution corresponding to  $c_0$  to the first term in the right hand

side of (2.2.4) is

$$(2.2.16) \quad \Gamma\left(\frac{\langle \xi \rangle}{\langle D_x + \xi \rangle}\right) \left[ \gamma(x, 0) |\det d\kappa(\kappa^{-1}(x))| a_\chi(\mathcal{U}, \kappa^{-1}(x), {}^t d\kappa(\kappa^{-1}(x)) \cdot \xi) \right] \\ = \Gamma\left(\frac{\langle \xi \rangle}{\langle D_x + \xi \rangle}\right) \left[ (\theta\theta') \circ \kappa^{-1} a_\chi(\mathcal{U}, \kappa^{-1}(x), {}^t d\kappa(\kappa^{-1}(x)) \cdot \xi) \right].$$

We write

$(\theta\theta' a_\chi) \circ \tau_\kappa = (\theta\theta'(a_\chi - a)) \circ \tau_\kappa + [(\theta\theta' a) \circ \tau_\kappa - ((\theta\theta' a) \circ \tau_\kappa)_{\chi_1}] + ((\theta\theta' a) \circ \tau_\kappa)_{\chi_1}$  with  $\chi_1 \in C_0^\infty(\mathbb{R}^d)$ ,  $\chi_1 \equiv 1$  close to zero. By (iii) of Proposition 2.1.6, the first two terms in the right hand side belong to  $\widetilde{HS}_{p, \text{loc}}^{m-r, \nu+rM_0}$  for any  $r \geq 0$ . If one makes act on them the spectral cut-off  $\Gamma(\frac{\langle \xi \rangle}{\langle D_x + \xi \rangle})$  of (2.2.16), one obtains elements of the same space, that stay in it if one applies  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda}$  for any  $\lambda \in \mathbb{R}$ . Consequently, since  $\Gamma(\frac{\langle \xi \rangle}{\langle D_x + \xi \rangle}) \chi_1(D/\langle \xi \rangle) = \chi_1(D/\langle \xi \rangle)$  if  $\text{Supp } \chi_1$  is small enough, we may write

$$(\kappa^{-1})^* \theta \tilde{A} \theta' \kappa^* = (\tilde{\theta} \circ \kappa^{-1}) \text{Op}(((\theta\theta' a) \circ \tau_\kappa)_{\chi_1}(\tilde{\theta}' \circ \kappa^{-1}) + (\tilde{\theta} \circ \kappa^{-1}) \text{Op}(\tilde{b})(\tilde{\theta}' \circ \kappa^{-1}) + C_1$$

where  $C_1$  contributes to  $C$  in (2.2.2) and  $\tilde{b}$  is given by the first term in the right hand side of (2.2.4), with  $c$  replaced by  $c_1 = c - c|_{z=0}$ . We may write  $c_1 = z c'_1$ , where  $c'_1$  satisfies estimates (2.2.6), with  $\nu$  replaced by  $\nu + M_0$ .

Then

$$(2.2.17) \quad \tilde{b}(\mathcal{U}, x, \xi) = \frac{1}{(2\pi)^{2d}} \int e^{i[(x-x') \cdot \eta' + z \cdot (\zeta' - \xi) + (\kappa^{-1}(x') - x'') \cdot \eta'']} \\ \times i(\partial_{\zeta'} c'_1)(\mathcal{U}, x', x'', z, \eta', \eta'', \zeta', \xi) dx' dx'' dz d\eta' d\eta'' d\zeta'.$$

Consider the vector fields

$$L_3 = \frac{1 - i\langle \xi \rangle^2 (\kappa^{-1}(x') - x'') \cdot \nabla_{\eta''}}{1 + \langle \xi \rangle^2 |\kappa^{-1}(x') - x''|^2} \\ L_4 = \frac{1 - i\langle \xi \rangle^2 (x - x') \cdot \nabla_{\eta'}}{1 + \langle \xi \rangle^2 |x - x'|^2} \\ L_5 = \frac{1 - i\langle \xi \rangle^2 |n'|^{-2M_0} z \cdot \nabla_{\zeta'}}{1 + |z|^2 \langle \xi \rangle^2 |n'|^{-2M_0}}.$$

If we perform  $N$  integrations by parts using  $L_3, L_4, L_5$ , use (2.2.6) and the support condition (2.2.5), we estimate (2.2.17) by

$$\int (1 + \langle \xi \rangle |\kappa^{-1}(x') - x''|)^{-N} (1 + \langle \xi \rangle |x - x'|)^{-N} (1 + |z| \langle \xi \rangle |n'|^{-M_0})^{-N} \\ \times \mathbb{1}_{|\eta''| \ll \langle \zeta' \rangle \sim \langle \xi \rangle \sim \langle \xi + \eta' \rangle} dx' dx'' dz d\eta' d\eta'' d\zeta' \\ \times \langle \xi \rangle^{m-1} |n'|^{\nu(p)+\nu+M_0(d+3)} \left(1 + \frac{|n'|}{\langle \xi \rangle}\right)^{-N} \\ \leq C \langle \xi \rangle^{m-1} |n'|^{\nu(p)+\nu+M_0(2d+3)} \left(1 + \frac{|n'|}{\langle \xi \rangle}\right)^{-N}.$$

Since the derivatives are estimated in the same way, we get an element of  $\widetilde{HS}_{p, \text{loc}}^{m-1, \nu+M_0(d+2)}$ , which stays in that space if we apply any  $\langle D_x + \xi \rangle^\lambda \langle \xi \rangle^{-\lambda}$

because of the support condition (2.2.5). The corresponding operator is in  $\widetilde{H\Psi}_{p,\text{loc}}^{m-1,\nu+M_0(3d+4)}$  by (ii) of proposition 2.1.6. This concludes the proof, as the case of operators in  $\Psi_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$  is similar.  $\square$

Let us define a principal symbol.

**DEFINITION 2.2.4.** *Let  $m \in \mathbb{R}, \nu \in \mathbb{R}_+, p \in \mathbb{N}, \sigma > 0, R > 0$ . One denotes by  $\widetilde{HS}_p^{m,\nu}(T^*X)$  (resp.  $S_p^{m,\nu}(B_\sigma(R), T^*X)$ ) the space of maps  $(\mathcal{U}, \rho) = (U_1, \dots, U_p, \rho) \rightarrow a(\mathcal{U}, \rho)$  (resp.  $(U, \rho) \rightarrow a(U, \rho)$ ) which are defined on  $C^\infty(X)^p \times T^*X$  (resp.  $B_\sigma(R) \times T^*X$ ), with values in  $\mathbb{C}$ , which are  $p$ -linear in  $\mathcal{U}$  (resp.  $C^1$  in  $U$ ), smooth in  $\rho$ , and such that for any local coordinate system  $\kappa$  defined on an open set  $V$  of  $X$ , with values in  $\mathbb{R}^d$ , for any  $\theta$  in  $C_0^\infty(V)$ ,  $(\theta a) \circ \tau_\kappa^{-1}$  is an element of  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp.  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ).*

**DEFINITION 2.2.5.** *One says that an element  $A$  of  $\widetilde{H\Psi}_p^{m,\nu}$  (resp.  $\Psi_p^{m,\nu}(B_\sigma(R))$ ) admits an element  $a$  of  $\widetilde{HS}_p^{m,\nu}(T^*X)$  (resp.  $S_p^{m,\nu}(B_\sigma(R), T^*X)$ ) as a principal symbol modulo  $\widetilde{H\Psi}_p^{m-1,\nu+\ell}$  (resp.  $\Psi_p^{m-1,\nu+\ell}(B_\sigma(R))$ ) if and only if, for any  $\chi$  in  $C_0^\infty(\mathbb{R}^d)$ ,  $\chi \equiv 1$  close to zero, with small enough support, for any local chart  $\kappa : V \subset X \rightarrow \mathbb{R}^d$ , for any functions  $\theta, \theta', \tilde{\theta}, \tilde{\theta}'$  in  $C_0^\infty(V)$ , with  $\theta\tilde{\theta} = \theta$ ,  $\theta'\tilde{\theta}' = \theta'$ ,*

$$(2.2.18) \quad (\kappa^{-1})^* \theta A \theta' \kappa^* - (\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta \theta' a) \circ \tau_\kappa^{-1}]_\chi (\tilde{\theta}' \circ \kappa^{-1})$$

*is in  $\widetilde{H\Psi}_{p,\text{loc}}^{m-1,\nu+\ell}$  (resp.  $\Psi_{p,\text{loc}}^{m-1,\nu+\ell}(B_\sigma(R))$ ).*

**Remark:** If  $\ell \geq M_0(3d+4)$ , the above definition is coherent. Actually, consider two local coordinate systems  $\kappa : V \xrightarrow{\sim} \tilde{V}$ ,  $\kappa_1 : W \xrightarrow{\sim} \tilde{W}$ , where  $V, W$  are non disjoint open subsets of  $X$ , and  $\tilde{V}, \tilde{W}$  are open subsets of  $\mathbb{R}^d$ . Let  $\theta, \theta', \tilde{\theta}, \tilde{\theta}'$  be in  $C_0^\infty(V \cap W)$ , with  $\theta\tilde{\theta} = \theta$ ,  $\theta'\tilde{\theta}' = \theta'$  and set  $\mathcal{O} = \kappa(V \cap W)$ ,  $\mathcal{O}_1 = \kappa_1(V \cap W)$ . Then

$$\begin{aligned} (\kappa_1^{-1})^* \theta A \theta' \kappa_1^* &= (\kappa \circ \kappa_1^{-1}|_{\mathcal{O}_1})^* (\kappa^{-1})^* \theta A \theta' \kappa^* (\kappa_1 \circ \kappa^{-1}|_{\mathcal{O}})^* \\ &= (\kappa \circ \kappa_1^{-1}|_{\mathcal{O}_1})^* (\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta \theta' a) \circ \tau_\kappa^{-1}]_\chi (\tilde{\theta}' \circ \kappa^{-1}) (\kappa_1 \circ \kappa^{-1}|_{\mathcal{O}})^* \end{aligned}$$

modulo  $\widetilde{H\Psi}_{p,\text{loc}}^{m-1,\nu+\ell}$  (resp.  $\Psi_{p,\text{loc}}^{m-1,\nu+\ell}(B_\sigma(R))$ ). By proposition 2.2.1, this is equal to

$$(\tilde{\theta} \circ \kappa_1^{-1}) \text{Op}[(\theta \theta' a) \circ \tau_{\kappa_1}^{-1}]_{\chi_1} (\tilde{\theta}' \circ \kappa_1^{-1})$$

modulo  $\widetilde{H\Psi}_{p,\text{loc}}^{m-1,\nu+M_0(3d+4)}$  (resp.  $\Psi_{p,\text{loc}}^{m-1,\nu+M_0(3d+4)}(B_\sigma(R))$ ). Of course, using (iii) of Proposition 2.1.6, we may replace  $\chi_1$  by any other function  $\chi$  in  $C_0^\infty(\mathbb{R}^d)$ ,  $\chi \equiv 1$  close to zero, with small enough support.

**PROPOSITION 2.2.6.** *Let  $m \in \mathbb{R}, \nu \in \mathbb{R}_+, \nu' \in \mathbb{R}_+, p, q \in \mathbb{N}, \sigma > 0$ .*

(i) *Every element of  $\widetilde{H\Psi}_p^{m,\nu}$  (resp.  $\Psi_p^{m,\nu}(B_\sigma(R))$ ) has a principal symbol  $a$  in  $\widetilde{HS}_p^{m,\nu}(T^*X)$  (resp.  $S_p^{m,\nu}(B_\sigma(R), T^*X)$ ) modulo  $\widetilde{H\Psi}_p^{m-1,\nu+M_0(3d+4)}$  (resp.  $\Psi_p^{m-1,\nu+M_0(3d+4)}(B_\sigma(R))$ ).*

(ii) *Let  $a$  be an element of  $\widetilde{HS}_p^{m,\nu}(T^*X)$  (resp.  $S_p^{m,\nu}(B_\sigma(R), T^*X)$ ). There is an operator  $A$  in  $\widetilde{H\Psi}_p^{m,\nu}$  (resp.  $\Psi_p^{m,\nu}(B_\sigma(R))$ ) whose principal symbol modulo  $\widetilde{H\Psi}_p^{m-1,\nu+M_0(3d+4)}$  (resp.  $\Psi_p^{m-1,\nu+M_0(3d+4)}(B_\sigma(R))$ ) is  $a$ .*

(iii) Assume  $\sigma > \min(\nu(p), \nu(q)) + \min(\nu, \nu') + M_0(3d+4)$ , and, when  $p > 0$  and  $q > 0$ , assume moreover that (2.1.10) holds.

Let  $A$  be in  $\widetilde{H\Psi}_p^{m,\nu}$  (resp.  $\Psi_p^{m,\nu}(B_\sigma(R))$ ) and  $B$  be in  $\widetilde{H\Psi}_q^{m',\nu'}$  (resp.  $\Psi_q^{m',\nu'}(B_\sigma(R))$ ). Denote by  $a$  a principal symbol of  $A$  modulo  $\widetilde{H\Psi}_p^{m-1,\nu+M_0(3d+4)}$  (resp.  $\Psi_p^{m-1,\nu+M_0(3d+4)}(B_\sigma(R))$ ) and by  $b$  a principal symbol of  $B$  modulo  $\widetilde{H\Psi}_p^{m'-1,\nu'+M_0(3d+4)}$  (resp.  $\Psi_q^{m'-1,\nu'+M_0(3d+4)}(B_\sigma(R))$ ). Then  $ab$  is a principal symbol for  $A \circ B$  modulo  $\widetilde{H\Psi}_{p+q}^{m+m'-1,\nu+M_0(3d+4)}$  when  $p > 0, q > 0$ , modulo  $\widetilde{H\Psi}_{p+q}^{m+m'-1,\nu+\nu'+2M_0(2d+3)}$  if  $p$  or  $q$  is zero (resp. modulo the space  $\Psi_{\max(p,q)}^{m+m'-1,\max(\nu,\nu')+M_0(3d+4)}(B_\sigma(R))$ ).

(iv) With the notations of (iii),  $a^\vee$  (resp.  $\bar{a}$ ) is a principal symbols of  ${}^tA$  and  $A^*$  modulo  $\widetilde{H\Psi}_p^{m-1,\nu+M_0(3d+5)}$  (resp.  $\Psi_p^{m,\nu+M_0(3d+5)}(B_\sigma(R))$ ).

*Proof:* (i) We consider  $(\theta_i)_{i \in I}$  a finite partition of unity on  $X$ , such that, for each couple  $(i, j)$  of  $J = \{(i, j) \in I \times I; \text{Supp } \theta_i \cap \text{Supp } \theta_j \neq \emptyset\}$ , there is a chart  $\kappa_{ij} : V_{ij} \xrightarrow{\sim} \tilde{V}_{ij}$  from the open set  $V_{ij}$  of  $X$  to the open set  $\tilde{V}_{ij}$  of  $\mathbb{R}^d$  such that  $\text{Supp } \theta_i \cup \text{Supp } \theta_j \subset V_{ij}$ . We have seen already in the proof of Lemma 2.2.2, that  $\sum_{(i,j) \notin J} \theta_i A \theta_j$  is in  $\widetilde{H\Psi}_p^{m-1,\nu+M_0}$  (resp.  $\Psi_p^{m-1,\nu+M_0}(B_\sigma(R))$ ). For each  $i$  in  $I$ , choose  $\tilde{\theta}_i$  a smooth function such that  $\tilde{\theta}_i \theta_i = \theta_i$  and that  $\text{Supp } \tilde{\theta}_i$  is a compact subset of  $V_{ij}$  for any  $j$  such that  $(i, j) \in J$ . We consider

$$(\kappa_{ij}^{-1})^* \theta_i A \theta_j (\kappa_{ij})^* = (\theta_i \circ \kappa_{ij}^{-1}) [(\kappa_{ij}^{-1})^* \tilde{\theta}_i A \tilde{\theta}_j (\kappa_{ij})^*] (\theta_j \circ \kappa_{ij}^{-1}).$$

The characterization of  $\widetilde{H\Psi}_p^{m,\nu}$  (resp.  $\Psi_p^{m,\nu}(B_\sigma(R))$ ) in terms of Sobolev spaces given in (1.2.7), (1.2.18), (1.2.19) and Definition 2.1.1 imply that the term between brackets is in  $\widetilde{H\Psi}_{p,\text{loc}}^{m,\nu}$  (resp.  $\Psi_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ). By Proposition 2.1.6, we may find a symbol  $a_{ij}$  in  $\widetilde{HS}_{p,\text{loc}}^{m,\nu}$  (resp. in  $S_{p,\text{loc}}^{m,\nu}(B_\sigma(R))$ ) such that

$$(2.2.19) \quad (\kappa_{ij}^{-1})^* \theta_i A \theta_j (\kappa_{ij})^* = (\theta_i \circ \kappa_{ij}^{-1}) \text{Op}(a_{ij,\chi}) (\theta_j \circ \kappa_{ij}^{-1})$$

modulo  $\widetilde{H\Psi}_{p,\text{loc}}^{m-1,\nu+M_0(2d+3)}$  (resp.  $\Psi_{p,\text{loc}}^{m-1,\nu+M_0(2d+3)}(B_\sigma(R))$ ). We define a symbol on  $C^\infty(X)^p \times T^*X$  (resp. on  $B_\sigma(R) \times T^*X$ ) by

$$(2.2.20) \quad a = \sum_{(i,j) \in I \times I} (\theta_i \theta_j a_{ij}) \circ \tau_{\kappa_{ij}}.$$

Let  $\kappa : V \xrightarrow{\sim} \tilde{V}$  be a local coordinate system,  $\theta, \tilde{\theta}, \theta', \tilde{\theta}'$  be in  $C_0^\infty(V)$  with  $\tilde{\theta}\theta = \theta$ ,  $\tilde{\theta}'\theta' = \theta'$ . Let us show that (2.2.18) holds with the symbol we just defined. We write, modulo  $\widetilde{H\Psi}_{p,\text{loc}}^{m-1,\nu+M_0(2d+3)}$  (resp.  $\Psi_p^{m-1,\nu+M_0(2d+3)}(B_\sigma(R))$ ),

$$\begin{aligned} (\kappa^{-1})^* \theta A \theta' \kappa^* &= \sum_{(i,j) \in J} (\kappa^{-1})^* \theta \theta_i A \theta_j \theta' \kappa^* \\ &= \sum_{(i,j) \in J} (\kappa^{-1})^* \theta (\kappa_{ij})^* (\kappa_{ij}^{-1})^* (\theta_i A \theta_j) (\kappa_{ij})^* (\kappa_{ij}^{-1})^* \theta' \kappa^* \\ &= \sum_{(i,j) \in J} ((\kappa \circ \kappa_{ij}^{-1}|_{\kappa_{ij}(V \cap V_{ij})})^{-1})^* (\theta \theta_i) \circ \kappa_{ij}^{-1} \text{Op}(a_{ij,\chi}) (\theta' \theta_j) \circ \kappa_{ij}^{-1} (\kappa \circ \kappa_{ij}^{-1}|_{\kappa_{ij}(V \cap V_{ij})})^* \end{aligned}$$

where the last equality comes from (2.2.19). By formula (2.2.2) of Proposition 2.2.1, the last term may be written, modulo  $\widetilde{H\Psi}_{p,\text{loc}}^{m-1,\nu+M_0(3d+4)}$  (resp. modulo the space  $\Psi_{p,\text{loc}}^{m-1,\nu+M_0(3d+4)}(B_\sigma(R))$ ) as

$$\sum_{(i,j) \in J} (\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta\theta'\theta_i\theta_j a_{ij}) \circ \tau_{\kappa_{ij}}^{-1}]_{\chi_1} (\tilde{\theta}' \circ \kappa^{-1}).$$

Using (2.2.20) (and that  $\theta_i\theta_j = 0$  if  $(i,j) \notin J$ ), we obtain  $(\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(a \circ \tau_\kappa^{-1})_{\chi_1}] (\tilde{\theta}' \circ \kappa^{-1})$  i.e. the right hand side of (2.2.18).

(ii) Let  $(\theta_i)_{i \in I}$  be a partition of unity on  $X$  such that  $\text{Supp } \theta_i$  is contained in an open set  $V_i$  so that there is a local chart  $\kappa_i : V_i \xrightarrow{\sim} \tilde{V}_i \subset \mathbb{R}^d$ . Take  $\tilde{\theta}_i$  in  $C_0^\infty(V_i)$  with  $\tilde{\theta}_i\theta_i = \theta_i$  and define  $A$  by

$$A = \sum_{i \in I} \kappa_i^* (\tilde{\theta}_i \circ \kappa_i^{-1}) \text{Op}[(\theta_i a) \circ \tau_{\kappa_i}^{-1}]_\chi (\tilde{\theta}_i \circ \kappa_i^{-1}) (\kappa_i^{-1})^*$$

where  $\chi \in C_0^\infty(\mathbb{R}^d)$ ,  $\chi \equiv 1$  close to zero,  $\text{Supp } \chi$  small enough. By (ii) of Proposition 2.1.6,  $A$  is an element of  $\widetilde{H\Psi}_p^{m,\nu+2M_0(d+1)}$  (resp.  $\Psi_p^{m,\nu+2M_0(d+1)}(B_\sigma(R))$ ). Let us check that  $a$  satisfies (2.2.18). If  $\kappa, \theta, \theta'$  are as in the statement of Definition 2.1.5,

$$(2.2.21) \quad (\kappa^{-1})^* \theta A \theta' \kappa^* \\ = \sum_{i \in I} (\tilde{\theta} \circ \kappa^{-1}) [(\kappa_i \circ \kappa^{-1})^* (\theta \tilde{\theta}_i \circ \kappa_i^{-1}) \text{Op}[(\theta_i a) \circ \tau_{\kappa_i}^{-1}]_\chi (\theta' \tilde{\theta}_i \circ \kappa_i^{-1}) (\kappa \circ \kappa_i^{-1})^*] (\tilde{\theta}' \circ \kappa^{-1})$$

if  $\tilde{\theta}, \tilde{\theta}'$  are in  $C_0^\infty(V)$  with  $\tilde{\theta} = \theta$ ,  $\theta' \tilde{\theta}' = \theta'$ . We apply Proposition 2.2.1, to get modulo  $\widetilde{H\Psi}_{p,\text{loc}}^{m-1,\nu+M_0(3d+4)}$  (resp.  $\Psi_{p,\text{loc}}^{m-1,\nu+M_0(3d+4)}(B_\sigma(R))$ ) the expression

$$\sum_{i \in I} (\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta \theta' \theta_i a) \circ \tau_\kappa^{-1}]_{\chi_1} (\tilde{\theta}' \circ \kappa^{-1}) = (\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta \theta' a) \circ \tau_\kappa^{-1}]_{\chi_1} (\tilde{\theta}' \circ \kappa^{-1}).$$

(iii) We just need to check the property locally. We may always assume that the distribution kernels of  $A$  and  $B$  are supported in a small neighborhood  $W$  of the diagonal. Let  $\kappa : V \xrightarrow{\sim} \tilde{V}$  be a chart and  $\theta, \theta'$  in  $C_0^\infty(V)$ . If  $W$  is small enough, we may find  $\theta_1$  in  $C_0^\infty(V)$  with  $\theta\theta_1 = \theta$ ,  $\theta'\theta_1 = \theta'$  so that

$$(\kappa^{-1})^* \theta A B \theta' \kappa^* = [(\kappa^{-1})^* \theta A \theta_1 \kappa^*] [(\kappa^{-1})^* \theta_1 B \theta' \kappa^*].$$

By Definition 2.2.5, Lemma 1.2.4 and Lemma 1.2.6, the last expression may be written, modulo  $\widetilde{H\Psi}_{p+q,\text{loc}}^{m+m'-1,\nu+\nu'+M_0(3d+4)}$  (resp.  $\Psi_{\max(p,q),\text{loc}}^{m+m'-1,\max(\nu,\nu')+M_0(3d+4)}(B_\sigma(R))$ ) as

$$(2.2.22) \quad (\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta a) \circ \tau_\kappa^{-1}]_\chi (\tilde{\theta}_1 \circ \kappa^{-1})^2 \text{Op}[(\theta' b) \circ \tau_\kappa^{-1}]_\chi (\tilde{\theta}' \circ \kappa^{-1})$$

where  $\tilde{\theta}_1\theta_1 = \theta_1$ ,  $\tilde{\theta}\theta = \theta$ ,  $\tilde{\theta}'\theta' = \theta'$ , when  $p > 0$  and  $q > 0$ . When  $p = 0$  or  $q = 0$ , the equality is valid modulo  $\widetilde{H\Psi}_{p+q,\text{loc}}^{m+m'-1,\nu+\nu'+M_0(3d+4)}$  (resp. modulo the space  $\Psi_{\max(p,q),\text{loc}}^{m+m'-1,\max(\nu,\nu')+M_0(3d+4)}(B_\sigma(R))$ ). By Theorem 2.1.7 (see (2.1.13) and (2.1.17)), we may write (2.2.22) as

$$(\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta \theta' a b) \circ \tau_\kappa^{-1}]_\chi (\tilde{\theta}_1 \circ \kappa^{-1})$$

modulo an element of  $\widetilde{H\Psi}_{p+q,\text{loc}}^{m+m'-1,\nu+\nu'+M_0(2d+3)}$  when  $p > 0, q > 0$  and modulo an element of  $\widetilde{H\Psi}_{p+q,\text{loc}}^{m+m'-1,\nu+\nu'+2M_0(2d+3)}$  if  $p = 0$  or  $q = 0$  (resp. modulo

$\Psi_{\max(p,q),\text{loc}}^{m+m'-1,\max(\nu,\nu')+M_0(2d+3)}(B_\sigma(R))$ ). This shows that  $ab$  is a principal symbol of the composition.

(iv) With the same notations as above, we need to study  ${}^t[(\kappa^{-1})^* \theta A \theta' \kappa^*]$  (where we denote by the same notation transpose of maps from  $L^2(X, d\mu)$  to  $L^2(X, d\mu)$ , from  $L^2(\mathbb{R}^d, dx)$  to  $L^2(\mathbb{R}^d, dx)$ , from  $L^2(X, d\mu)$  to  $L^2(\mathbb{R}^d, dx)$  and from  $L^2(\mathbb{R}^d, dx)$  to  $L^2(X, d\mu)$ ). Denote by  $\alpha$  the density of the measure  $d\mu$  relatively to Lebesgue measure in the coordinate patch i.e.  $d\mu(y) = \alpha(x)dx$  if  $y = \kappa^{-1}(x)$ . Then  ${}^t(\kappa^*) = (\kappa^{-1})^*(\alpha \circ \kappa)$ ,  ${}^t(\kappa^{-1})^* = (\alpha \circ \kappa)^{-1} \kappa^*$ , so that

$$[(\kappa^{-1})^* \theta^t A \theta' \kappa^*] = {}^t[(\kappa^{-1})^* \theta' (\alpha \circ \kappa) A (\alpha \circ \kappa)^{-1} \theta \kappa^*].$$

We apply (2.2.18) and (2.1.18) to write this quantity as

$$(\tilde{\theta} \circ \kappa^{-1}) \text{Op}[(\theta \theta' a^\vee)_\chi] (\tilde{\theta}' \circ \kappa^{-1})$$

modulo  $\widehat{H\Psi}_{p,\text{loc}}^{m-1,\nu+M_0(3d+5)}$  (resp.  $\Psi_{p,\text{loc}}^{m-1,\nu+M_0(3d+5)}(B_\sigma(R))$ ). This concludes the proof, the case of the adjoint being similar.  $\square$

## CHAPTER 3

### Quasi-linear Birkhoff normal forms method

**3.1. Description of the method.** Our goal is to prove Theorem 1.2.8. We fix some integer  $P \in \mathbb{N}^*$  and take  $G$  a function as in the statement of the theorem. We fix also  $\mathbf{m} \in ]0, +\infty[-\mathcal{N}$ , where  $\mathcal{N}$  is the zero measure subset of  $]0, +\infty[$  introduced before the statement of Theorem 1.2.8. Then inequalities (1.2.27) hold.

To prove the theorem, i.e. to show that the solution of (1.2.28) extends to a time interval  $] -T_\epsilon, T_\epsilon[$  with  $T_\epsilon \geq c\epsilon^{-P}$ , for all  $u_0$  in the unit ball of  $H^s(X)$  ( $s > s_0$ ) and any small enough  $\epsilon$ , it is enough to construct a function  $\Theta_s$ , defined on a neighborhood of zero in  $H^s(X)$ , real valued, such that there is  $C > 0$ ,  $R > 0$  with

$$(3.1.1) \quad C^{-1}\|u\|_{H^s}^2 \leq \Theta_s(u) \leq C\|u\|_{H^s}^2 \text{ for all } u \text{ in } B_s(R)$$

and for any solution  $u$  of  $u_t = i\nabla_{\bar{u}}G(u, \bar{u})$ , defined and smooth on  $] -T, T[ \times X$  for some  $T$ , the estimate

$$(3.1.2) \quad \Theta_s(u(t, \cdot)) \leq \Theta_s(u(0, \cdot)) + C \left| \int_0^t \|u(\tau, \cdot)\|_{H^s}^{P+2} d\tau \right| + C\epsilon^P \left| \int_0^t \|u(\tau, \cdot)\|_{H^s}^2 d\tau \right|$$

holds for all  $t$  in  $] -T, T[$  such that  $u(t, \cdot) \in B_s(R)$ . Actually, (3.1.1) and (3.1.2) imply an estimate

$$\|u(t, \cdot)\|_{H^s}^2 \leq C \left[ \|u(0, \cdot)\|_{H^s}^2 + \left| \int_0^t \|u(\tau, \cdot)\|_{H^s}^{P+2} d\tau \right| + \epsilon^P \left| \int_0^t \|u(\tau, \cdot)\|_{H^s}^2 d\tau \right| \right]$$

as long as  $u(t, \cdot)$  stays in  $B_s(R) \cap C^\infty$ . This shows that for  $u(0, \cdot)$  smooth and  $\|u(0, \cdot)\|_{H^s} \leq \epsilon$  small enough, the solution extends up to a time  $T_\epsilon \geq c\epsilon^{-P}$  for some small  $c > 0$ . When  $u(0, \cdot)$  is only in  $H^s(X)$ , with small enough  $H^s$ -norm, we may always approximate the Cauchy data by  $C^\infty(X)$  functions, solve the equation for these approximations on  $] -T_\epsilon, T_\epsilon[$  and pass to the limit to get the solution of (1.2.28) for any  $u_0$  in the unit ball of  $H^s(X)$  and any  $\epsilon > 0$  small enough.

The construction of  $\Theta_s$  will be made in three steps, given by the following three theorems.

**THEOREM 3.1.1.** *There is  $\nu \in \mathbb{R}_+$ ,  $s_0 > 0$ ,  $R > 0$  and for any  $s > s_0$ , a canonical transformation  $\chi$  defined on  $B_s(R)$ , with  $\chi(0) = 0, \chi'(0) = \text{Id}$ , and there are*

- Elements  $A_p$  of  $H\Psi_p^{1,\nu}$ ,  $1 \leq p \leq P-1$ , self-adjoint,
- Elements  $A_P, B_P$  of  $\Psi_P^{1,\nu}(B_s(R))$ , with  $A_P$  self-adjoint,
- Functions  $G_{L,p}$  in  $\mathcal{G}_p^{1-\infty, \nu+\infty}(B_s(R))$ ,  $1 \leq p \leq P-1$ ,

*such that for any  $C^1$  function  $\Theta_s^1$ , defined on  $B_s(R)$ , vanishing at order 2 at the origin, one has, for any  $U$  in  $B_s(R) \cap C^\infty$*

$$(3.1.3) \quad \{\Theta_s^1 \circ \chi, G\}(U) = \{\Theta_s^1, G_L + G_H + \tilde{G}_P\} \circ \chi(U) + O(\|u\|_{H^s}^{P+2}), \quad U \rightarrow 0$$



with

$$(3.1.4) \quad G_L(U) = \sum_{p=1}^{P-1} G_{L,p}(U)$$

$$(3.1.5) \quad G_H(U) = G_0(u) + \sum_{p=1}^{P-1} \int_X (A_p(U)u)\bar{u} d\mu$$

and

$$(3.1.6) \quad \tilde{G}_P(U) = \int_X (A_P(U)u)\bar{u} d\mu + \operatorname{Re} \int_X (B_P(U)u)u d\mu.$$

**Remarks •** We write the preceding statement for functions  $U = (u, \bar{u})$  in  $B_s(R) \cap C^\infty$  so that the Poisson brackets in (3.1.3) are well defined. Actually, since (3.1.5), (3.1.6) involve operators of order one,  $X_{G_H}$ ,  $X_{\tilde{G}_P}$  are in  $H^{s-1}(X)$  if  $U$  is in  $H^s$ . On the other hand,  $d\Theta_s^1(U)$  is in  $\mathcal{L}(H^s, \mathbb{R})$ , so that  $d\Theta_s^1(U) \cdot X_{G_L+G_H+\tilde{G}_P}$  is not necessarily well defined if  $U$  is assumed to be only in  $H^s$ . Of course, at the end of the proof,  $\Theta_s^1$  will be defined in such a way that the right hand side of (3.1.3) will extend even to function  $U$  that belong only to  $H^s$ .

• The conclusion of the theorem asserts that we may choose  $\chi$  in such a way that the new Hamiltonian in the right hand side of (3.1.3) contains contributions homogeneous of lower degree  $0 \leq p \leq P-1$  that are *hyperbolic* (in the sense of Definition 1.2.7), which form  $G_H$ , “lower order” contributions giving  $G_L$ , and a term  $\tilde{G}_P$ , which has no special structure, but vanishes at high order when  $U$  goes to zero.

The second step in the proof of theorem 1.2.8 will eliminate the elliptic part of  $\tilde{G}_P$ .

**THEOREM 3.1.2.** *Let  $\nu \in \mathbb{R}_+$  be given. There is  $s_0 > 0$  such that if  $s \geq s_0$ , there is a local diffeomorphism  $\psi$  from a neighborhood of 0 in  $H^s(X)$  to a neighborhood of zero in  $H^s(X)$ , with  $\psi(0) = 0$ , such that, for any function  $\Theta_s^2$  which is in  $\bigoplus_{p=0}^{P-1} (H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R)) + \mathcal{G}_p^{2s-1,\nu}(B_s(R)))$ , for any  $U$  in  $C^\infty(X)$  belonging to a small enough neighborhood of zero in  $H^s(X)$ ,*

$$(3.1.7) \quad \{\Theta_s^2 \circ \psi, G_L + G_H + \tilde{G}_P\}(U) = \{\Theta_s^2, G_L + G_H\} \circ \psi(U) + O(\|U\|_{H^s}^{P+2}).$$

The last step of the proof will be to choose  $\Theta_s^2$  so that the first term in the right hand side of (3.1.7) vanishes essentially at order  $P+2$  at zero.

**THEOREM 3.1.3.** *There are  $\nu \in \mathbb{R}_+$ ,  $s_0 > 0$  and, for any  $s \geq s_0$ , a constant  $C_0 > 0$  and, for any  $\epsilon \in ]0, 1[$ , any  $R > 0$ , an element  $\Theta_s^2 = \Theta_s^2(\epsilon)$  in  $\bigoplus_{p \geq 0} H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R)) + \mathcal{G}_p^{2s-1,\nu}(B_s(R))$ , with estimates uniform in  $\epsilon \in ]0, 1[$ , satisfying (3.1.1) uniformly in  $\epsilon \in ]0, 1[$ , such that*

$$(3.1.8) \quad |\{\Theta_s^2, G_L + G_H\}(U)| \leq C_0[\|U\|_{H^s}^{P+2} + \epsilon^P \|U\|_{H^s}^2]$$

for any  $U$  in  $C^\infty(X) \cap B_s(R)$ .

*Proof of theorem 1.2.8:* As already seen, we just need to construct a function  $\Theta_s$  such that (3.1.1) and (3.1.2) hold (actually,  $\Theta_s$  will be a family of functions, depending on  $\epsilon$ , and for which (3.1.1), (3.1.2) hold with constants independent of  $\epsilon \in ]0, 1[$ ). We take  $\nu$  large enough so that Theorems 3.1.1 and 3.1.3 hold, and  $s_0$  large enough so that Theorems 3.1.1, 3.1.2 and 3.1.3 apply. We use Theorem 3.1.1

to determine  $\chi$ ,  $G_H, \tilde{G}_P$ . Then, Theorem 3.1.3 determines  $\Theta_s^2$  and Theorem 3.1.2 determines  $\psi$ . We set  $\Theta_s^1 = \Theta_s^2 \circ \psi$  and  $\Theta_s = \Theta_s^1 \circ \chi$ . Then, (3.1.3), (3.1.7) and (3.1.8) show that  $|\{\Theta_s, G\}|$  is bounded from above by the right hand side of (3.1.8). Since it follows from equation (1.2.18) that  $\frac{d}{dt}\Theta_s(u(t, \cdot)) = \{\Theta_s, G\}(u(t, \cdot))$ , this gives (3.1.2) by integration. Inequality (3.1.1) holds by construction. This concludes the proof of Theorem 1.2.8 from the three results in this subsection.  $\square$

The proofs of Theorems 3.1.1 to 3.1.3 will depend on the study of Poisson brackets between functions of the classes of Definition 1.2.7. We shall study such Poisson brackets in the following subsection.

**3.2. Computation of Poisson brackets.** Let  $U \rightarrow A(U)$  be a  $C^1$  function defined on an open set  $\Omega$  of  $H^\sigma(X)$ , with  $\sigma$  large enough, with values in  $\mathcal{L}(H^s, H^{s-m})$  for some  $m$  and any  $s$  with  $s \geq \sigma$  and  $m \leq 2s$ . Consider the functions

$$(3.2.1) \quad \begin{aligned} H_1^A(W, U) &= \int_X (A(W)u)u \, d\mu \\ H_0^A(W, U) &= \int_X (A(W)u)\bar{u} \, d\mu \\ H_{-1}^A(W, U) &= \int_X (A(W)\bar{u})\bar{u} \, d\mu \end{aligned}$$

defined for  $W \in \Omega$ ,  $U = (u, \bar{u}) \in H^s(X)$ . If  $U$  is in  $\Omega \cap H^s(X)$ , set  $F_a^A(U) = H_a^A(U, U)$ ,  $a = -1, 0, 1$ . Let  $B$  be another  $C^1$  map from  $\Omega$  to  $\mathcal{L}(H^s, H^{s-m'})$  for some  $m'$  and any  $s \geq \sigma$ ,  $m' \leq 2s$ . Assume moreover  $m + m' \leq 2s$ . By formulas (1.1.17), (1.1.16)

$$(3.2.2) \quad \begin{aligned} \{F_a^A, F_b^B\} &= \{H_a^A(W, \cdot), H_b^B(W, \cdot)\}|_{W=U} + i(d_W H_a^A)(U, U)^t J(\nabla_U H_b^B)(U, U) \\ &\quad - i(d_W H_b^B)(U, U)^t J(\nabla_U H_a^A)(U, U) + \{H_a^A(\cdot, U), H_b^B(\cdot, U)\}|_{W=U} \end{aligned}$$

where the first (resp. the last) term in the right hand side is the Poisson bracket of the functions  $U \rightarrow H_a^A(W, U)$ ,  $U \rightarrow H_b^B(W, U)$  (resp.  $W \rightarrow H_a^A(W, U)$ ,  $W \rightarrow H_b^B(W, U)$ ) at fixed  $W$  (resp.  $U$ ). Let us compute explicitly the first Poisson bracket. According to formula (1.1.17)

$$(3.2.3) \quad \begin{aligned} \{H_1^A(W, \cdot), H_0^B(W, \cdot)\} &= i \int_X ((A(W) + {}^t A(W))u)(B(W)u) \, d\mu \\ \{H_1^A(W, \cdot), H_{-1}^B(W, \cdot)\} &= i \int_X ((A(W) + {}^t A(W))u)((B(W) + {}^t B(W))\bar{u}) \, d\mu \\ \{H_0^A(W, \cdot), H_{-1}^B(W, \cdot)\} &= i \int_X ({}^t A(W)\bar{u})((B(W) + {}^t B(W))\bar{u}) \, d\mu \end{aligned}$$

and

$$(3.2.4) \quad \begin{aligned} \{H_1^A(W, \cdot), H_1^B(W, \cdot)\} &= 0, \quad \{H_{-1}^A(W, \cdot), H_{-1}^B(W, \cdot)\} = 0 \\ \{H_0^A(W, \cdot), H_0^B(W, \cdot)\} &= i \int_X ([A, B](W)u)\bar{u} \, d\mu. \end{aligned}$$

Consequently, the first term in the right hand side of (3.2.2) may be written as  $H_c^C(U, U)$  for some  $C$  that may be computed from compositions of  $A, B, {}^t A, {}^t B$  or

from a commutator  $[A, B]$ . On the other hand, the second term in the right hand side of (3.2.2), has the following structure for  $a = 1, 0, -1$

$$(3.2.5) \quad \begin{aligned} & i \int_X [d_W A(W) \cdot ({}^t J \nabla_U H_b^B) u] u \, d\mu \\ & i \int_X [d_W A(W) \cdot ({}^t J \nabla_U H_b^B) u] \bar{u} \, d\mu \\ & i \int_X [d_W A(W) \cdot ({}^t J \nabla_U H_b^B) \bar{u}] \bar{u} \, d\mu. \end{aligned}$$

By definition of  $H_b^B$ ,

$$(3.2.6) \quad \begin{aligned} \nabla_U H_1^B(W, U) &= \begin{bmatrix} (B(W) + {}^t B(W))u \\ 0 \end{bmatrix}, \quad \nabla_U H_0^B(W, U) = \begin{bmatrix} {}^t B(W)\bar{u} \\ B(W)u \end{bmatrix} \\ \nabla_U H_{-1}^B(W, U) &= \begin{bmatrix} 0 \\ (B(W) + {}^t B(W))\bar{u} \end{bmatrix} \end{aligned}$$

so that (3.2.5) will be computed from quantities of the form

$$(3.2.7) \quad d_w A(W) \cdot (C(W)u), \quad d_w A(W) \cdot (C(W)\bar{u}), \quad d_{\bar{w}} A(W) \cdot (C(W)u), \quad d_{\bar{w}} A(W) \cdot (C(W)\bar{u})$$

where  $C(W)$  will be an operator of the same form as  $A(W), B(W)$ .

We introduce an auxiliary class of multi-linear operators.

**DEFINITION 3.2.1.** *Let  $m \in \mathbb{R}, \nu \in \mathbb{R}_+, p \in \mathbb{N}^*$ . We denote by  $\widetilde{HM}_p^{m, \nu}$  the space of  $p$ -linear maps  $(u_1, \dots, u_p) \rightarrow M(u_1, \dots, u_p)$ , defined on  $C^\infty(X)^p$ , with values in the space of linear maps from  $C^\infty(X)$  to  $\mathcal{D}'(X)$ , that may be written as a sum  $\sum_{i=1}^p M_i(u_1, \dots, u_p)$ , where  $M_i$  satisfies the following estimates: for any  $N_1, N_2$  in  $\mathbb{N}$ , there is  $C > 0$  and for any  $n_0, \dots, n_{p+1}$  in  $\mathbb{N}^*$*

$$(3.2.8) \quad \begin{aligned} \|\Pi_{n_0} M_i(\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}\|_{\mathcal{L}(L^2)} &\leq C \left[ \min\left(\frac{n_i}{n_{p+1}}, \frac{n_{p+1}}{n_i}\right) \right]^{N_1} \max(n_0, \dots, \widehat{n_i}, \dots, n_p)^{\nu(p)+\nu} \\ &\quad \times (n_i + n_{p+1})^m \left( 1 + \frac{\max(n_0, \dots, \widehat{n_i}, \dots, n_p)}{n_i + n_{p+1}} \right)^{-N_2} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}. \end{aligned}$$

We shall denote by  $HM_p^{m, \nu}$  the space of operator valued maps  $U \rightarrow M(U)$  that may be written

$$(3.2.9) \quad M(U) = \sum_{\ell=0}^p M_\ell(\underbrace{u, \dots, u}_\ell, \underbrace{\bar{u}, \dots, \bar{u}}_{p-\ell})$$

with  $M_\ell$  in  $\widetilde{HM}_p^{m, \nu}$ .

**LEMMA 3.2.2.** *Let  $A$  be an element of  $\mathcal{P}_p^{m, \nu}$  and set for  $j \in \mathbb{N}$*

$$H^{A, \leq j}(W, U) = \int_X (A(W) S_j u) u \, d\mu \quad \text{or} \quad \int_X (A(W) S_j u) \bar{u} \, d\mu \quad \text{or} \quad \int_X (A(W) S_j \bar{u}) \bar{u} \, d\mu.$$

*Then  $\nabla_w H^{A, \leq j}(W, U)|_{W=U}$  and  $\nabla_{\bar{w}} H^{A, \leq j}(W, U)|_{W=U}$  may be written as a linear combination of  $M_\alpha(U) S_j u$ ,  $M_\alpha(U) S_j \bar{u}$  for elements  $M_\alpha$  of  $HM_p^{m, \nu}$ .*

*Proof:* We consider for instance the first expression of  $H^{A, \leq j}$ . We decompose  $A(W) = \sum_{\ell=0}^p A_\ell(\underbrace{w, \dots, w}_\ell, \bar{w}, \dots, \bar{w})$  with  $A_\ell$  in  $\tilde{\mathcal{P}}_p^{m, \nu}$ . Then  $d_w H^{A, \leq j}(W, U) \cdot h$  may be written as a sum

$$\sum_{\ell=0}^p \sum_{i=1}^\ell \int_X (A_\ell(w, \dots, \underbrace{h}_i, \dots, w, \bar{w}, \dots, \bar{w}) S_j u) u \, d\mu.$$

Consequently,  $\nabla_w H^{A, \leq j}(W, U)|_{W=U}$  may be written as a sum

$$\sum_{\ell=0}^p \sum_{i=1}^\ell M_{\ell i}(\underbrace{u, \dots, u}_\ell, \bar{u}, \dots, \bar{u}) S_j u$$

with

$$\int_X (M_{\ell i}(u_1, \dots, u_p) u_{p+1}) h \, d\mu = \int_X (A_\ell(u_1, \dots, u_{i-1}, h, u_{i+1}, \dots, u_p) u_{p+1}) u_i \, d\mu.$$

The fact that  $\|\Pi_{n_0} M_{\ell i}(\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}\|_{\mathcal{L}(L^2)}$  is bounded by the right hand side of (3.2.8) follows from (1.2.5).  $\square$

The goal of this subsection is to prove the following:

**PROPOSITION 3.2.3.** *Let  $m, m'$  in  $\mathbb{R}$ ,  $\nu, \nu'$  in  $\mathbb{R}_+$ ,  $p, q$  in  $\mathbb{N}$ ,  $\sigma \in \mathbb{R}$ . Assume*

$$(3.2.10) \quad \nu(p) + \nu(q) + \max(\nu, \nu') + M_0(3d+4) \leq \nu(p+q) \text{ if } p > 0, q > 0$$

*and that  $\sigma$  is large enough so that condition (1.2.23) holds for all the spaces of multi-linear forms below. Then, with the notations introduced in definition 1.2.7, one has the following inclusions*

$$(3.2.11) \quad \begin{aligned} & \left\{ H\mathcal{F}_{p,E}^{m, \nu}(B_\sigma(R)), H\mathcal{F}_{q,E}^{m', \nu'}(B_\sigma(R)) \right\} \cup \left\{ H\mathcal{F}_{p,H}^{m, \nu}(B_\sigma(R)), H\mathcal{F}_{q,E}^{m', \nu'}(B_\sigma(R)) \right\} \\ & \cup \left\{ H\mathcal{F}_{p,H}^{m, \nu}(B_\sigma(R)), H\mathcal{F}_{q,H}^{m', \nu'}(B_\sigma(R)) \right\} \\ & \subset H\mathcal{F}_{p+q}^{m+m', \min(\nu, \nu')}(B_\sigma(R)) + H\mathcal{F}_{p+q}^{m, m'_+ + \min(\nu, \nu')}(B_\sigma(R)) + H\mathcal{F}_{p+q}^{m', m_+ + \min(\nu, \nu')}(B_\sigma(R)) \end{aligned}$$

*if  $p > 0$  and  $q > 0$ . If  $p$  or  $q$  is zero, one gets the same inclusion with  $\min(\nu, \nu')$  replaced by  $\nu + \nu'$ ;*

$$(3.2.12) \quad \begin{aligned} & \left\{ H\mathcal{F}_{p,H}^{m, \nu}(B_\sigma(R)), H\mathcal{F}_{q,H}^{m', \nu'}(B_\sigma(R)) \right\} \subset H\mathcal{F}_{p+q,H}^{m+m'-1, \min(\nu, \nu') + M_0(3d+4)}(B_\sigma(R)) \\ & + H\mathcal{F}_{p+q,H}^{m, m'_+ + \min(\nu, \nu')}(B_\sigma(R)) + H\mathcal{F}_{p+q,H}^{m', m_+ + \min(\nu, \nu')}(B_\sigma(R)) \end{aligned}$$

*if  $p > 0$  and  $q > 0$ . If  $p$  or  $q$  is zero, one gets the same inclusion with  $\min(\nu, \nu')$  replaced by  $\nu + \nu'$  and  $M_0(3d+4)$  replaced by  $2M_0(2d+3)$ . Moreover, one has also the inclusions*

$$(3.2.13) \quad \begin{aligned} & \left\{ \mathcal{G}_p^{m, \nu}(B_\sigma(R)), \mathcal{G}_q^{m', \nu'}(B_\sigma(R)) \right\} \\ & \subset \mathcal{G}_{p+q}^{m+m', \min(\nu, \nu')}(B_\sigma(R)) + \mathcal{G}_{p+q}^{m, m'_+ + \min(\nu, \nu')}(B_\sigma(R)) + \mathcal{G}_{p+q}^{m', m_+ + \min(\nu, \nu')}(B_\sigma(R)) \end{aligned}$$

if  $p > 0$  and  $q > 0$ . If  $p$  or  $q$  is zero, one gets the same inclusion with  $\min(\nu, \nu')$  replaced by  $\nu + \nu'$ . One has also

$$(3.2.14) \quad \left\{ H\mathcal{F}_{p,\mathbb{H}}^{m,\nu}(B_\sigma(R)), \mathcal{G}_q^{m',\nu'}(B_\sigma(R)) \right\} \\ \subset H\mathcal{F}_{p+q,\mathbb{H}}^{m,m'_+ + \min(\nu,\nu')}(B_\sigma(R)) + \mathcal{G}_{p+q}^{m+m', \min(\nu,\nu') + 2M_0}(B_\sigma(R)) + \mathcal{G}_{p+q}^{m', m_+ + \min(\nu,\nu') + 2M_0}(B_\sigma(R))$$

if  $p > 0$  and  $q > 0$ . If  $p$  or  $q$  is zero, one gets the same inclusion with  $\min(\nu, \nu')$  replaced by  $\nu + \nu'$  (resp.  $\nu + \nu' + 2M_0$ ) in the first (resp. last) term in the right hand side.

Finally, one has the inclusion

$$(3.2.15) \quad \left\{ H\mathcal{F}_p^{m,\nu}(B_\sigma(R)), \mathcal{G}_q^{m',\nu'}(B_\sigma(R)) \right\} \\ \subset \mathcal{G}_{p+q}^{m+m', \min(\nu,\nu') + 2M_0}(B_\sigma(R)) + \mathcal{G}_{p+q}^{m, m'_+ + \min(\nu,\nu') + 2M_0}(B_\sigma(R)) + \mathcal{G}_{p+q}^{m', m_+ + \min(\nu,\nu') + 2M_0}(B_\sigma(R))$$

if  $p > 0$  and  $q > 0$ . If  $p$  or  $q$  is zero, one gets the same inclusion with  $\min(\nu, \nu')$  replaced by  $\nu + \nu'$ .

The main step to prove the proposition will be to study expressions of type (3.2.7) where  $A$  will be an element of  $H\Psi_p^{m,\nu}$ . We consider the different possible situations in the following lemma.

LEMMA 3.2.4. Assume, for some  $a \geq 0$ ,

$$(3.2.16) \quad \nu(p) + \nu(q) + \max(\nu, \nu') + a \leq \nu(p+q) \text{ if } p > 0 \text{ and } q > 0.$$

Let  $A$  be an element of  $\widetilde{H\Psi}_p^{m,\nu}$  (resp.  $\widetilde{\mathcal{P}}_p^{m,\nu}$ ) and  $B$  be an element of  $\widetilde{\mathcal{P}}_q^{m',\nu'+a}$ . Define

$$(3.2.17) \quad C(u_1, \dots, u_{p+q}) = A(B(u_1, \dots, u_q)u_{q+1}, u_{q+2}, \dots, u_{p+q}).$$

Then  $C$  is in  $\widetilde{H\Psi}_{p+q}^{m, \min(\nu,\nu') + m'_+}$  if  $p > 0, q > 0$  and in  $\widetilde{H\Psi}_{p+q}^{m, \nu + \nu' + m'_+ + a}$  if  $p = 0$  or  $q = 0$ . (resp. in  $\widetilde{\mathcal{P}}_{p+q}^{m, \min(\nu,\nu') + m'_+}$  if  $p > 0, q > 0$  and in  $\widetilde{\mathcal{P}}_{p+q}^{m, \nu + \nu' + m'_+ + a}$  if  $p = 0$  or  $q = 0$ ).

*Proof:* Let  $P_1, \dots, P_k$  be differential operators as in Definition 1.2.1. When  $A$  is in  $\widetilde{H\Psi}_p^{m,\nu}$ , denote  $\tilde{A} = \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} A$  and  $\tilde{C} = \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} C$ . Compute

$$(3.2.18) \quad \Delta_j \tilde{C}(\Pi_{n_1} u_1, \dots, \Pi_{n_{p+q}} u_{p+q}) \Delta_{j'} \\ = \sum_n \Delta_j \tilde{A}(\Pi_n B(\Pi_{n_1} u_1, \dots, \Pi_{n_q} u_q) \Pi_{n_{q+1}} u_{q+1}, \Pi_{n_{q+2}} u_{q+2}, \dots, \Pi_{n_{p+q}} u_{p+q}) \Delta_{j'}.$$

By symmetry, we may assume  $n_1 \geq \dots \geq n_q, n_{q+2} \geq \dots \geq n_{p+q}$ . By (1.2.8) and (1.2.5), the  $\mathcal{L}(L^2)$ -norm of the general term of the sum is bounded from above by a constant times  $\prod_1^{p+q} \|\Pi_{n_\ell} u_\ell\|_{L^2}$  times

$$(3.2.19) \quad 2^{-|j-j'|N'_1} (n + n_{q+2})^{\nu(p) + \nu + M_0 k} 2^{j\tilde{m}} (1 + 2^{-j}(n + n_{q+2}))^{-N'_2} \\ \times \left[ \min\left(\frac{n}{n_{q+1}}, \frac{n_{q+1}}{n}\right) \right]^{N''_1} n^{m'} n_1^{\nu(q) + \nu' + a} \left(1 + \frac{n_1}{n}\right)^{-N''_2} \langle n - n_{q+1} \rangle^{-2}$$

where  $\tilde{m} = m + \sum_1^k d_\ell - k$ . We write

$$(3.2.20) \quad \left(1 + \frac{n + n_{q+2}}{2^j}\right) \left(1 + \frac{n_1}{n}\right) \geq 1 + \frac{n + n_{q+2} + n_1}{2^j} \\ \geq \left(1 + \frac{n_{q+1} + n_{q+2} + n_1}{2^j}\right) \min\left(\frac{n}{n_{q+1}}, \frac{n_{q+1}}{n}\right),$$

so that (3.2.19) is bounded from above by

$$(3.2.21) \quad 2^{j\tilde{m}} 2^{-|j-j'|N'_1} (n_{q+1} + n_{q+2} + n_1)^{\nu(p)+\nu(q)+\nu+\nu'+a+m'_++M_0k} \left(1 + \frac{n_{q+1} + n_{q+2} + n_1}{2^j}\right)^{-N'_2} \\ \times \langle n - n_{q+1} \rangle^{-2} \left[ \min\left(\frac{n}{n_{q+1}}, \frac{n_{q+1}}{n}\right) \right]^{N'_1 - N'_2 - \nu(p) - \nu - m'_+ - M_0k}$$

if  $N'_1, N'_2$  are large enough relatively to  $N'_2, \nu, m'_+, k$ . If we sum this in  $n$ , we get estimate (1.2.8) of an element of  $\widetilde{H\Psi}_{p+q}^{m, \min(\nu, \nu') + m'_+}$  when  $p > 0, q > 0$  and (3.2.16) holds. If  $p = 0$  or  $q = 0$ , we get an element of  $\widetilde{H\Psi}_{p+q}^{m, \nu + \nu' + m'_+ + a}$ . When  $A$  is in  $\widetilde{\mathcal{P}}_p^{m, \nu}$ , we bound

$$\|\Pi_{n_0} C(\Pi_{n_1} u_1, \dots, \Pi_{n_{p+q}} u_{p+q}) \Pi_{n_{p+q+1}}\|_{\mathcal{L}(L^2)}$$

using (1.2.5), by the product of a constant times  $\prod_1^{p+q} \|\Pi_{n_\ell} u_\ell\|_{L^2}$  times

$$(3.2.22) \quad \langle n_0 - n_{p+q+1} \rangle^{-2} \left[ \min\left(\frac{n_0}{n_{p+q+1}}, \frac{n_{p+q+1}}{n_0}\right) \right]^{N'_1} (n + n_{q+2})^{\nu + \nu(p)} n_0^m \left(1 + \frac{n + n_{q+2}}{n_0}\right)^{-N'_2} \\ \times \langle n - n_{q+1} \rangle^{-2} \left[ \min\left(\frac{n}{n_{q+1}}, \frac{n_{q+1}}{n}\right) \right]^{N'_1} n^{m'} n_1^{\nu(q) + \nu'} \left(1 + \frac{n_1}{n}\right)^{-N'_2}.$$

The rest of the computation is similar as above.  $\square$

LEMMA 3.2.5. *Let  $p, q \in \mathbb{N}^*$ . Assume for some  $a \geq 0$ ,*

$$(3.2.23) \quad \nu(p) + \nu(q) + \max(\nu, \nu') + a + 1 \leq \nu(p + q).$$

*Let  $A$  be in  $\widetilde{H\Psi}_p^{m, \nu}$  (resp.  $\widetilde{\mathcal{P}}_p^{m, \nu}$ ) and  $M = \sum_{i=1}^q M_i$  be in  $\widetilde{H\mathcal{M}}_q^{m', \nu' + a}$ . Consider*

$$(3.2.24) \quad C(u_1, \dots, u_{p+q}) = \sum_{j''=0}^{+\infty} \sum_{i=1}^q A(M_i(u_1, \dots, u_q) S_{j''} u_{q+1}, u_{q+2}, \dots, u_{p+q}) \Delta_{j''}.$$

*Then  $C$  is in  $\widetilde{H\Psi}_{p+q}^{m, \min(\nu, \nu') + m'_+}$  (resp.  $\widetilde{\mathcal{P}}_{p+q}^{m, \min(\nu, \nu') + m'_+}$ ).*

*Proof:* We consider the contribution to (3.2.24) of the term corresponding to  $i = 1$ . We write  $M$  instead of  $M_1$ . Let  $P_1, \dots, P_k$  be differential operators on  $X$  of orders  $d_1, \dots, d_k$ . We may express  $\text{Ad}_{P_1} \cdots \text{Ad}_{P_k} C$  from

$$\sum_{j''} \text{Ad}_{\mathcal{P}'} A(M(u_1, \dots, u_q) S_{j''} u_{q+1}, u_{q+2}, \dots, u_{p+q}) \text{Ad}_{\mathcal{P}''} \Delta_{j''}$$

where  $\mathcal{P} = \mathcal{P}' \cup \mathcal{P}''$  is any partition of  $\mathcal{P} = (P_1, \dots, P_k)$ . Denote by  $k'$  (resp.  $k''$ ) the cardinal of  $\mathcal{P}'$  (resp.  $\mathcal{P}''$ ) and set  $d' = \sum_{P_\ell \in \mathcal{P}'} d_\ell$ ,  $d'' = \sum_{P_\ell \in \mathcal{P}''} d_\ell$ . Then  $\|\Delta_j \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} C(\Pi_n \mathcal{U}) \Delta_{j'}\|_{\mathcal{L}(L^2)}$  will be bounded from above by the sum for  $\mathcal{P} = \mathcal{P}' \cup \mathcal{P}''$  of

$$(3.2.25) \quad \sum_n \sum_{j''} \sum_{\tilde{j}''} \|\Delta_j \text{Ad}_{\mathcal{P}'} A(\Pi_n M(\Pi_{n_1} u_1, \dots, \Pi_{n_q} u_q) S_{j''} \Pi_{n_{q+1}} u_{q+1}, \\ \Pi_{n_{q+2}} u_{q+2}, \dots, \Pi_{n_{p+q}} u_{p+q}) \Delta_{\tilde{j}''}\|_{\mathcal{L}(L^2)} \|\tilde{\Delta}_{\tilde{j}''}(\text{Ad}_{\mathcal{P}''} \Delta_{j''}) \Delta_{j'}\|_{\mathcal{L}(L^2)}$$

where  $\tilde{\Delta}_{\tilde{j}''}$  is a cut-off such that  $\tilde{\Delta}_{\tilde{j}''} \Delta_{\tilde{j}''} = \Delta_{\tilde{j}''}$ . By symmetry, we may assume that  $n_2 \geq n_3 \geq \dots \geq n_q$ ,  $n_{q+2} \geq \dots \geq n_{p+q}$ . Moreover, in the sum,  $n_{q+1} \leq C2^{j''}$  for some  $C > 0$ . By (1.2.8) and (3.2.8) with  $i = 1$ , the first factor in the general term of (3.2.25) is bounded from above by the product of  $\prod_1^{p+q} \|\Pi_{n_\ell} u_\ell\|_{L^2}$  and

$$(3.2.26) \quad 2^{-|j-j''|N'_1} (n + n_{q+2})^{\nu(p)+\nu+M_0k'} 2^{j(m+d'-k')} (1 + 2^{-j}(n + n_{q+2}))^{-N'_2} \\ \times \left( \min\left(\frac{n_1}{n_{q+1}}, \frac{n_{q+1}}{n_1}\right) \right)^{N''_1} (n+n_2)^{\nu(q)+\nu'} (n_1+n_{q+1})^{m'} \left(1 + \frac{n+n_2}{n_1+n_{q+1}}\right)^{-N''_2} \mathbb{1}_{n_{q+1} \leq C2^{j''}}.$$

We notice that

$$(3.2.27) \quad (1 + 2^{-j}(n + n_{q+2})) \left(1 + \frac{n+n_2}{n_1+n_{q+2}}\right) \max\left[\frac{n_1}{n_{q+1}}, \frac{n_{q+1}}{n_1}\right] 2^{|j-j''|} \\ \geq c(1 + 2^{-j}(n_{q+2} + n_{q+1} + n_2 + n_1))$$

since  $n_{q+1} \leq C2^{j''}$ . Moreover, for  $\ell = 2$  or  $\ell = q+2$

$$(n + n_\ell) \left(1 + \frac{n+n_2}{n_1+n_{q+2}}\right)^{-1} \leq C(n_{q+2} + n_{q+1} + n_2 + n_1).$$

If  $N''_1, N''_2$  are large enough, we bound the sum in  $n$  of (3.2.26) by

$$(3.2.28) \quad C2^{-N'_1|j-j''|} 2^{j(m+d'-k')} (n_1 + \dots + n_{p+q})^{\nu(p)+\nu(q)+\nu+\nu'+1+m'_++M_0k'} \\ \times (1 + 2^{-j}(n_1 + \dots + n_{p+q}))^{-N'_2} 2^{N'_2|j-j''|}.$$

We use Proposition A.1 of the appendix to bound the last factor in (3.2.25) by the expression  $C_N 2^{-N[|j''-j''|+|j''-j'|]2^{j(d''-k'')}}$ . If we take  $N \gg N'_1 \gg N'_2 \gg N_1, N_2$  and use (3.2.23), we bound (3.2.25) by

$$(3.2.29) \quad C2^{-N_1|j-j'|} 2^{j(m+\sum d_\ell-k)} (n_1 + \dots + n_{p+q})^{\nu(p+q)+\min(\nu,\nu')+m'_++M_0k} \\ \times (1 + 2^{-j}(n_1 + \dots + n_{p+q}))^{-N_2} \prod_1^{p+q} \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

This is the wanted estimate.

The case when  $A$  is in  $\tilde{\mathcal{P}}_p^{m,\nu}$  is treated in the same way.  $\square$

*Proof of Proposition 3.2.3:* Let  $A$  (resp.  $B$ ) be an element of  $H\Psi_p^{m,\nu}$  or  $\mathcal{P}_p^{m,\nu}$  (resp.  $H\Psi_q^{m',\nu'}$  or  $\mathcal{P}_q^{m',\nu'}$ ). Denote by  $H_a^1, H_b^b$ ,  $-1 \leq a, b \leq 1$ , the functions given by any of the formulas (3.2.1). We have to compute  $\{H_a^A(U, U), H_b^B(U, U)\}$ . According to

(3.2.2), the first quantity to study is given by (3.2.3) or (3.2.4) with  $W = U$ . They may be written as expressions of type

$$(3.2.30) \quad \int_X (C(U)u)\bar{u} d\mu, \quad \int_X (C(U)u)u d\mu, \quad \int_X (C(U)\bar{u})\bar{u} d\mu.$$

Consider first the contribution to the left hand side of (3.2.11) given by (3.2.3), (3.2.4). Then  $C$  may be expressed from the composition of an element of  $H\Psi_p^{m,\nu}$  and of  $H\Psi_q^{m',\nu'}$ . By lemma 1.2.4 with  $a = \max(\nu, \nu')$  and (3.2.10), we obtain that  $C$  is in  $H\Psi_{p+q}^{m+m', \min(\nu, \nu')}$  if  $p > 0, q > 0$  and  $H\Psi_{p+q}^{m+m', \nu+\nu'}$  if  $p = 0$  or  $q = 0$ . This shows that (3.2.30) belongs to the right hand side of (3.2.11). If we consider the contribution of (3.2.4) to (3.2.12), we get expressions of form (3.2.30) with  $C = [A, B]$ . We apply Proposition 2.2.6 (iii), which is possible under assumption (3.2.10), and see that  $C$  is in  $H\Psi_{p+q}^{m+m'-1, \min(\nu, \nu')+M_0(3d+4)}$  if  $p > 0, q > 0$  and  $H\Psi_{p+q}^{m+m'-1, \nu+\nu'+2M_0(2d+3)}$  if  $p = 0$  or  $q = 0$ .

Let us study as well the contributions of (3.2.3), (3.2.4) to the left hand side of (3.2.13), (3.2.14). For the first of these inclusions,  $C$  may be expressed from the composition of an element of  $\mathcal{P}_p^{m,\nu}$  and  $\mathcal{P}_q^{m',\nu'}$ , so belongs to  $\mathcal{P}_{p+q}^{m+m', \min(\nu, \nu')}$  (if  $p > 0, q > 0$ ) and  $\mathcal{P}_{p+q}^{m+m', \nu+\nu'}$  if  $p = 0$  or  $q = 0$  by (1.2.14), so that (3.2.30) contributes to the right hand side of inclusion (3.2.13). The inclusion (1.2.10) shows that the contributions of (3.2.3), (3.2.4) to the left hand side of (3.2.14) belong to  $\mathcal{G}_{p+q}^{m+m', \min(\nu, \nu')+2M_0}(B_\sigma(R))$  when if  $p > 0, q > 0$  and to  $\mathcal{G}_{p+q}^{m+m', \nu+\nu'+2M_0}(B_\sigma(R))$  if  $p = 0$  or  $q = 0$ .

We consider now the contributions of type (3.2.5) to the inclusions (3.2.11)–(3.2.14) (as well as the symmetric ones obtained exchanging  $A$  and  $B$ ). According to (3.2.7), we must study

$$(3.2.31) \quad \int_X (d_{(w, \bar{w})}A(W) \cdot C(W)u_1)u_2 d\mu|_{W=U},$$

where  $u_1, u_2$  stand for  $u$  or  $\bar{u}$ ,  $C(W)$  stands for  $B(W)$  or  ${}^tB(W)$  or  $B(W) + {}^tB(W)$ , and the symmetric expressions obtained exchanging  $A$  and  $B$ .

Let us study first contributions of type (3.2.31) to the left hand side of (3.2.11), (3.2.12). Then  $A$  is in  $H\Psi_p^{m,\nu}$  and  $C$  is in  $H\Psi_q^{m',\nu'} \subset \mathcal{P}_q^{m',\nu'+2M_0}$ . It follows from Lemma 3.2.4 that (3.2.31) is in  $H\mathcal{F}_{p+q}^{m, m'_+ + \min(\nu, \nu')}(B_\sigma(R))$  if  $p > 0, q > 0$  and to  $H\mathcal{F}_{p+q}^{m, m'_+ + \nu + \nu' + 2M_0}(B_\sigma(R))$  if  $p = 0$  or  $q = 0$ . Consider now (3.2.13). Then  $A$  (resp.  $C$ ) in (3.2.31) is in  $\mathcal{P}_p^{m,\nu}$  (resp.  $\mathcal{P}_q^{m',\nu'}$ ) so that Lemma 3.2.4 shows that (3.2.31) is in  $\mathcal{G}_{p+q}^{m, m'_+ + \min(\nu, \nu')}(B_\sigma(R))$  if  $p > 0, q > 0$  and in  $\mathcal{G}_{p+q}^{m, m'_+ + \nu + \nu'}(B_\sigma(R))$  when  $p = 0$  or  $q = 0$ . For inclusion (3.2.14), we use that in (3.2.31),  $A$  is in  $H\Psi_p^{m,\nu}$  and  $B$  in  $\mathcal{P}_q^{m',\nu'}$ , so that Lemma 3.2.4 shows that (3.2.31) is in  $H\mathcal{F}_{p+q}^{m, m'_+ + \min(\nu, \nu')}(B_\sigma(R))$  when  $p > 0, q > 0$  and in  $H\mathcal{F}_{p+q}^{m, m'_+ + \nu + \nu'}(B_\sigma(R))$  when  $p = 0$  or  $q = 0$ .

Those contributions coming from the symmetric version of (3.2.31) with  $A$  and  $B$  exchanged belong to the space of functions obtained exchanging  $(m, \nu)$  and  $(m', \nu')$ , except for inclusion (3.2.14). In this case, we have to consider (3.2.31) with  $A$  replaced by an element of  $\mathcal{P}_q^{m',\nu'}$  and  $C$  by an element of  $H\Psi_p^{m,\nu} \subset \mathcal{P}_p^{m,\nu+2M_0}$ .



Again, by Lemma 3.2.4, we get a contribution belonging to  $\mathcal{G}_{p+q}^{m', m_+ + \min(\nu, \nu')}(B_\sigma(R))$  if  $p > 0, q > 0$  and to  $\mathcal{G}_{p+q}^{m', m_+ + \nu + \nu' + 2M_0}(B_\sigma(R))$  if  $p$  or  $q$  is zero.

Finally, we have to study the last contribution to (3.2.2), namely

$$\{H_a^A(\cdot, U), H_b^B(\cdot, U)\}.$$

We decompose

$$H_a^A(W, U) = \sum_{j=0}^{+\infty} H_a^{A,j}(W, U)$$

where

$$H_a^{A,j}(W, U) = \int_X (A(W) \Delta_j u_1) u_2 d\mu$$

with  $(u_1, u_2) = (u, u)$  if  $a = 1$ ,  $(u_1, u_2) = (u, \bar{u})$  if  $a = 0$ ,  $(u_1, u_2) = (\bar{u}, \bar{u})$  if  $a = -1$ . We use a similar decomposition for  $H_b^B$  and set  $H_a^{A, < j} = \sum_{j' < j} H_a^{A, j'}$ ,  $H_a^{A, \leq j} = \sum_{j' \leq j} H_a^{A, j'}$ . Then

$$\{H_a^A(\cdot, U), H_b^B(\cdot, U)\} = \sum_j \{H_a^{A, \leq j}(\cdot, U), H_b^{B, j}(\cdot, U)\} + \sum_j \{H_a^{A, j}(\cdot, U), H_b^{B, < j}(\cdot, U)\}.$$

We study the first sum in the right hand side. The general term in that sum may be written  $-i(d_W H_b^{B, j}) \cdot ({}^t J \nabla_W H_a^{A, j})$  so is equal to

$$(3.2.33) \quad -i \int_X ([d_W B(W) \cdot ({}^t J \nabla_W H_a^{A, j}(W, U))] \Delta_j u_1) u_2 d\mu|_{W=U}.$$

We apply Lemma 3.2.2 considering  $A$  as an element of  $\mathcal{P}_p^{m, \nu + 2M_0}$ . This allows us to write (3.2.33) as a linear combination of quantities

$$\int_X [(d_U B(U) \cdot M(U) S_j u_3) \Delta_j u_1] u_2 d\mu$$

where  $u_3 = u$  or  $\bar{u}$  and  $M$  is in  $H\mathcal{M}_p^{m, \nu + 2M_0}$ . The sum in  $j$  of these quantities may be computed from expressions

$$\sum_j \int_X (B(u_1, \dots, u_\ell, M(u_{\ell+1}, \dots, u_{\ell+p}) S_j u_{\ell+p+1}, u_{\ell+p+2}, \dots, u_{p+q}) \Delta_j u_{p+q+1}) u_0 d\mu$$

where  $u_i$  has to be replaced by  $u$  or  $\bar{u}$ . This is an expression of the form

$$\int_X (C(u_1, \dots, u_{p+q}) u_{p+q+1}) u_0 d\mu$$

with  $C$  given by (3.2.24) (up to a change of indices). By Lemma 3.2.5,  $C$  is in  $\widetilde{H\Psi}_{p+q}^{m', m_+ + \min(\nu, \nu')}$  (resp.  $\widetilde{\mathcal{P}}_{p+q}^{m', m_+ + \min(\nu, \nu')}$ ) if  $B$  is in  $H\Psi_q^{m', \nu'}$  (resp.  $\mathcal{P}_q^{m', \nu'}$ ). This gives to (3.2.32) a contribution belonging to  $H\mathcal{F}_{p+q}^{m', m_+ + \min(\nu, \nu')}(B_\sigma(R))$ , when we consider (3.2.11), (3.2.12) and to  $\mathcal{G}_{p+q}^{m', m_+ + \min(\nu, \nu')}(B_\sigma(R))$  for (3.2.13), (3.2.14).

The second contribution to (3.2.32) belongs to the same classes with  $(m, \nu)$  and  $(m', \nu')$  exchanged in the case of (3.2.11), (3.2.12), (3.2.13). For (3.2.14), we get a contribution in  $H\mathcal{F}_{p+q}^{m, m'_+ + \min(\nu, \nu')}(B_\sigma(R))$  (Note that when studying (3.2.32) we may assume  $p > 0$  and  $q > 0$ , as otherwise the corresponding contribution is zero).

Inclusion (3.2.15) follows from (3.2.13) and the inclusion  $H\mathcal{F}_p^{m, \nu}(B_\sigma(R)) \subset \mathcal{G}_p^{m, \nu + 2M_0}(B_\sigma(R))$ .  $\square$

We shall need a version of inclusion (3.2.12), when one of the functions involved in the bracket is not homogeneous, and with a weaker conclusion.

LEMMA 3.2.6. *Let  $\nu, \nu'$  in  $\mathbb{R}_+$ ,  $p$  in  $\mathbb{N}$ ,  $q$  in  $\mathbb{N}^*$ ,  $R > 0$ . Assume (2.1.10). There is  $s_0$ , depending only on  $\nu, \nu', p, q$  such that if  $s \geq s_0$ , if  $\Theta$  is in  $H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R)) + \mathcal{G}_p^{2s-1,\nu}(B_s(R))$  and  $F$  is in  $\mathcal{F}_{q,H}^{1,\nu'}(B_s(R))$ , then  $\{\Theta, F\}(U) = O(\|U\|_{H^s}^{q+2})$ ,  $U \rightarrow 0$ .*

*Proof:* If  $\Theta$  is in  $\mathcal{G}_p^{2s-1,\nu}(B_s(R))$ ,  $\{\Theta, F\}(U) = d\Theta \cdot X_F(U)$  is well defined and is  $O(\|U\|_{H^s}^{q+2})$  since, by the last remark following Definition 1.2.7,  $X_F(U)$  is in  $H^{s-1}$  while  $d\Theta$  belongs to  $\mathcal{L}(H^{s-1}, \mathbb{R})$  because of the definition of  $\mathcal{G}_p^{2s-1,\nu}(B_s(R))$  (if  $s$  is large enough).

Assume from now on that  $\Theta$  is in  $H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R))$ . Using notation (3.2.1), we may write  $\Theta(U) = H_0^A(U, U)$ ,  $F(U) = H_0^B(U, U)$  with  $A$  in  $H\Psi_p^{2s,\nu}$  and  $B$  in  $\Psi_q^{1,\nu'}(B_s(R))$ . We have seen, in the last remark following Definition 1.2.5, that we may consider  $A$  as an element of  $\Psi_0^{2s,\bar{\nu}}(B_s(R))$  with  $\bar{\nu} > \nu + \nu(p) + \frac{1}{2}$ . In particular, by (iii) of Proposition 2.2.6,  $[A, B]$  is an element of  $\Psi_q^{2s, \max(\bar{\nu}, \nu') + M_0(3d+4)}(B_s(R))$ . We use (3.2.2) to express  $\{\Theta, F\}$ . The first term in the right hand side of this equality is given by the last formula in (3.2.4). Since  $[A, B]$  is of order  $2s$ , we do get, for  $s \geq s_0$  large enough, a  $O(\|U\|_{H^s}^{q+2})$  contribution.

We consider next the second term in the right hand side of (3.2.2). By (3.2.6),  ${}^t J \nabla_U H_0^B(U, U)$  is in  $H^{s-1}$ , and is  $O(\|U\|_{H^s}^{q+1})$  (if  $s \geq s_0$  large enough) using (1.2.18). The term under study, given by the second formula in (3.2.5), is thus  $O(\|U\|_{H^s}^{q+1})$ , since  $d_W A(W) \cdot ({}^t J \nabla_U H_0^B(U, U))$  is  $\mathcal{L}(H^s, H^{-s})$  by (1.2.18).

Take now the third term in the right hand side of (3.2.2). By (3.2.6), the expression  ${}^t J \nabla_U H_0^A(U, U)$  is in  $H^{-s}$ , with norm in that space  $O(\|U\|_{H^s})$ . We apply (1.2.19) to  $B$ , with  $m = 1$ ,  $\sigma' = -s$ . We get, if  $s$  is large enough relatively to  $\nu(q)$ ,  $\nu'$ , that  $dB(U) \cdot ({}^t J \nabla_U H_0^A(U, U))$  is in  $H^{-s}$ , with norm in that space  $O(\|U\|_{H^s}^{q+1})$ . The corresponding contribution to  $\{\Theta, F\}$  is again  $O(\|U\|_{H^s}^{q+1})$ .

Finally, the last contribution to (3.2.2) will be expressed from

$$\int_X (d_W A(W) \cdot ({}^t J \nabla_W H_0^B(W, U))u) \bar{u} d\mu|_{W=U}$$

and from the similar expression exchanging  $A$  and  $B$ . We need to check that

$$d_W A(W) \cdot ({}^t J \nabla_W H_0^B(W, U)), d_W B(W) \cdot ({}^t J \nabla_W H_0^A(W, U))$$

are in  $\mathcal{L}(H^s, H^{-s})$ . This will follow from (1.2.19) with  $m = 2s$  (resp.  $m = 1$ ) if we show that  $\nabla_W H_0^B(W, U)$  (resp.  $\nabla_W H_0^A(W, U)$ ) belongs to  $H^{\sigma'}$  for some  $\sigma' \geq \bar{\nu}$  (resp. to  $H^{-s}$ ) and if  $s$  is large enough. In other words, we have to check that

$$H \rightarrow \int_X ((d_W B(W) \cdot H)u) \bar{u} d\mu|_{W=U}, H \rightarrow \int_X ((d_W A(W) \cdot H)u) \bar{u} d\mu|_{W=U}$$

are linear continuous on  $H^{-\sigma'}$  (resp.  $H^s$ ). This follows again from (1.2.19).  $\square$



## CHAPTER 4

### Proof of the main theorem

We have seen that Theorem 1.2.8 follows from Theorem 3.1.1, Theorem 3.1.2 and Theorem 3.1.3. This section is devoted to the proof of these results.

**4.1. Elimination of elliptic terms of lower degree.** This subsection will be devoted to the proof of Theorem 3.1.1: we shall construct a canonical transformation  $\chi$  such that (3.1.3) holds, i.e. such that we may reduce the Hamiltonian  $G$  to another Hamiltonian  $G_L + G_H + \tilde{G}_P$ , where all elliptic terms of degree strictly smaller than  $P$  have been eliminated. Let us introduce a class of smoothing operators.

**DEFINITION 4.1.1.** *Let  $m \in \mathbb{R}$ ,  $p \in \mathbb{N}$ ,  $p \geq 2$ ,  $\gamma \in \mathbb{R}_+$ ,  $s \geq \gamma$ ,  $R > 0$ . One denotes by  $\mathcal{R}_{\gamma,p}^m(B_s(R))$  the space of maps  $U \rightarrow \mathfrak{R}(U)$ , defined on  $B_s(R)$ , with values in  $H^{2s-m-\gamma}$ , satisfying the following:*

(i) *The map  $U \rightarrow \mathfrak{R}(U)$  is  $C^1$  from  $B_s(R)$  to  $H^{2s-m-\gamma}$  and there is  $C > 0$  such that, for any  $U$  in  $B_s(R)$*

$$(4.1.1) \quad \|\mathfrak{R}(U)\|_{H^{2s-m-\gamma}} \leq C \|U\|_{H^s}^p.$$

(ii) *For any  $\theta, \theta'$  in  $\mathbb{R}$  satisfying*

$$(4.1.2) \quad s - m \geq |\theta|, \quad s - m \geq |\theta'|, \quad s + \theta + \theta' > m + \gamma,$$

*for any  $U$  in  $B_s(R)$ ,  $\partial_U \mathfrak{R}(U)$  extends as an element of  $\mathcal{L}(H^{\theta'}, H^{-\theta})$  with estimates*

$$(4.1.3) \quad \|\partial_U \mathfrak{R}(U)\|_{\mathcal{L}(H^{\theta'}, H^{-\theta})} \leq C \|U\|_{H^s}^{p-1}.$$

The main example of remainders satisfying the conditions of the preceding definition is given by the following lemma:

**LEMMA 4.1.2.** *Let  $m \in \mathbb{R}$ ,  $\nu \in \mathbb{R}_+$ ,  $p \leq q$  in  $\mathbb{N}^*$ . Let  $s, \gamma$  such that*

$$(4.1.4) \quad s \geq \gamma > \nu(q) + \nu + \frac{1}{2}, \quad 2s \geq m + \gamma.$$

*Let  $A$  be in  $H\Psi_q^{m,\nu}$  and define  $H(W, U) = \int_X (A(W)w_1)w_2 d\mu$  where  $w_1, w_2$  stand for  $u$  or  $\bar{u}$ . Then  $\mathfrak{R}(U) = X_{H(\cdot, U)}|_{W=U} = i^t J \nabla_W H(W, U)|_{W=U}$  is in  $\mathcal{R}_{\gamma,p+1}^m(B_s(R))$  for any  $R > 0$ .*

*Proof:* Decomposing  $A(W) = \sum_{\ell=0}^q A_\ell(w, \dots, w, \bar{w}, \dots, \bar{w})$  with  $A_\ell$  in  $\widetilde{H\Psi_q^{m,\nu}}$ , we may assume that  $A$  is one of the terms in that sum. If  $K = (k, \bar{k})$  is a smooth function on  $X$ , we may write by duality  $\int_X \mathfrak{R}(U) \cdot K d\mu$  from expressions of type

$$(4.1.5) \quad \int_X (A(Jh, u_2, \dots, u_q)u_{q+1})u_0 d\mu$$

where  $h$  stands for  $k$  or  $\bar{k}$  and  $u_\ell$  is  $u$  or  $\bar{u}$ , and from similar expressions where the  $Jh$  term replaces any other argument of  $A$ . In the same way,  $\int_X (\partial_U \Re(U) \cdot K') K d\mu$  may be written from expressions of type

$$(4.1.6) \quad \int_X (A(Jh, h', u_3, \dots, u_q) u_{q+1}) u_0 d\mu$$

$$(4.1.7) \quad \int_X (A(Jh, u_2, \dots, u_q) h') u_0 d\mu$$

and from similar expressions, where  $Jh$  replaces any other argument inside  $A$ , or  $h' = k'$  or  $\bar{k}'$  replaces any other argument of  $A$ , or where  $h'$  and  $u_0$  are exchanged in (4.1.7). To prove (4.1.1), we must bound the modulus of (4.1.5) by

$$(4.1.8) \quad C \|h\|_{H^{-2s+\gamma+m}} \|u_0\|_{H^s} \prod_{\ell=2}^{q+1} \|u_\ell\|_{H^s}.$$

In the same way, to show (4.1.3), we have to bound the modulus of (4.1.6) and (4.1.7) respectively by

$$(4.1.9) \quad C \|h\|_{H^\theta} \|h'\|_{H^{\theta'}} \prod_{\ell=3}^{q+1} \|u_\ell\|_{H^s} \|u_0\|_{H^s}, \quad C \|h\|_{H^\theta} \|h'\|_{H^{\theta'}} \prod_{\ell=2}^q \|u_\ell\|_{H^s} \|u_0\|_{H^s}.$$

To obtain (4.1.8), we apply (1.2.9) with  $N_2 = 2s - m$ . We obtain a bound of (4.1.5) by the product of  $C \|u_0\|_{H^s} \|u_{q+1}\|_{H^s}$  and of

$$(4.1.10) \quad \sum_{n_1} \cdots \sum_{n_q} (n_1 + \cdots + n_q)^{\nu(q)+\nu-2s+m} \|\Pi_{n_1} h\|_{L^2} \prod_{\ell=2}^q \|\Pi_{n_\ell} u_\ell\|_{L^2} \\ \leq C \sum_{n_1} \cdots \sum_{n_q} n_1^{\nu(q)+\nu-\gamma} \prod_{\ell=2}^q n_\ell^{-s} \prod_{\ell=1}^q c_{n_\ell}^\ell \|h\|_{H^{-2s+\gamma+m}} \prod_{\ell=2}^q \|u_\ell\|_{H^s},$$

where  $(c_{n_\ell}^\ell)_{n_\ell}$  are  $\ell^2$ -sequences, and where we used that the exponent in the left hand side of (4.1.10) is negative by (4.1.4). The assumptions on  $\gamma, s$  show that the series converges, which gives (4.1.8).

To estimate (4.1.6) by the first term in (4.1.9), we bound the modulus of (4.1.6) by

$$\sum_n \sum_{n'} \sum_{n_3} \cdots \sum_{n_q} \|A(J\Pi_n h, \Pi_{n'} h', \Pi_{n_3} u_3, \dots, \Pi_{n_q} u_q) u_{q+1}\|_{H^{-s}} \|u_0\|_{H^s}.$$

We apply (1.2.9) with  $N_2 = 2s - m$  to get a bound given by the product of the factor  $C \|h\|_{H^\theta} \|h'\|_{H^{\theta'}} \prod_{\ell=3}^{q+1} \|u_\ell\|_{H^s}$  and of

$$\sum_n \sum_{n'} \sum_{n_3} \cdots \sum_{n_q} (n + n' + n_3 + \cdots + n_q)^{\nu(q)+\nu-2s+m} n^{-\theta} n'^{-\theta'} c_n c_{n'} \prod_{\ell=3}^q n_\ell^{-s} c_{n_\ell}^\ell$$

with  $\ell^2$ -sequences  $(c_n)_n$ ,  $(c_{n'})_{n'}$ ,  $(c_{n_\ell}^\ell)_{n_\ell}$ . Since the first exponent is negative and  $s > 1/2$ , this sum is finite as soon as

$$(4.1.11) \quad \int_{\mathbb{R}_+^2} (1 + X + X')^{-\alpha} (1 + X)^{-\theta} (1 + X')^{-\theta'} c(X) c'(X') dX dX' < +\infty$$

where  $c(\cdot)$ ,  $c'(\cdot)$  are  $L^2$  functions,  $\alpha = 2s - m - \nu(q) - \nu$ . One checks that (4.1.11) holds as soon as

$$(4.1.12) \quad \alpha > \max \left[ \left( \frac{1}{2} - \theta \right)_+ + \left( \frac{1}{2} - \theta' \right), \left( \frac{1}{2} - \theta \right) + \left( \frac{1}{2} - \theta' \right)_+ \right]$$

which follows from assumptions (4.1.2), (4.1.4).

Let us study (4.1.7). We bound the modulus of this quantity by

$$\sum_n \sum_{n_2} \cdots \sum_{n_q} \|A(J\Pi_n h, \Pi_{n_2} u_2, \dots, \Pi_{n_q} u_q) h'\|_{H^{-s}} \|u_0\|_{H^s}.$$

We apply estimate (1.2.9) with  $N_2 = s + \theta' - m \geq 0$  and  $s$  replaced by  $\theta'$ , and get a bound given by the product of  $C\|h\|_{H^\theta}\|h'\|_{H^{\theta'}} \prod_2^q \|u_\ell\|_{H^s} \|u_0\|_{H^s}$  and of

$$(4.1.13) \quad \sum_n \sum_{n_2} \cdots \sum_{n_q} (n + n_2 + \cdots + n_q)^{\nu(q) + \nu - s - \theta' + m} n^{-\theta} c_n \prod_2^q n_\ell^{-s} c_{n_\ell}^\ell$$

for  $\ell^2$ -sequence  $(c_n)_n$ ,  $(c_{n_\ell}^\ell)_{n_\ell}$ . By symmetry, we may assume  $n_2 \geq \cdots \geq n_q$ . This reduces the verification of the finiteness of (4.1.13) to the study of an integral of form (4.1.11) with  $(\alpha, \theta, \theta')$  replaced by  $(s + \theta' - m - \nu(q) - \nu, \theta, s)$ . One checks that condition (4.1.12) follows from (4.1.2), (4.1.4). This concludes the proof.  $\square$

To prepare the proof of Theorem 3.1.1, we have to study the Hamiltonian flow of some auxiliary functions. Let  $A_p$  be an element of  $H\Psi_p^{0,\nu}$ ,  $1 \leq p \leq P-1$ , for some  $\nu \in \mathbb{R}_+$ . Define  $A = \sum_{p=1}^{P-1} A_p$ , and set

$$F(U) = \operatorname{Re} \int_X (A(U)u)u \, d\mu.$$

By (1.1.16) and Lemma 4.1.2

$$(4.1.14) \quad X_F(U) = \frac{i}{2} \left[ \overline{(A(U) + {}^t A(U))\bar{u}} \right] + \Re(U) = \mathcal{A}(U)U + \Re(U)$$

where  $\mathcal{A}(U) = \frac{i}{2} \begin{bmatrix} 0 & \bar{A} + {}^t \bar{A} \\ -(A + {}^t A) & 0 \end{bmatrix}$  and  $\Re$  is an element of  $\mathcal{R}_{\gamma,2}^0(B_\sigma(R))$  if  $\gamma > \nu(P-1) + \nu + \frac{1}{2}$ ,  $\sigma > \gamma$ . By the Cauchy-Lipschitz theorem, the equation

$$(4.1.15) \quad \begin{aligned} \dot{\Phi}(t, U) &= X_F(\Phi(t, U)) \\ \Phi(0, U) &= U, \end{aligned}$$

where  $\Phi(t, U) = \left[ \frac{\phi(t, U)}{\phi(t, U)} \right]$ , has a unique solution defined for  $t \in [-1, 1]$ , for  $U$  in  $B_\sigma(R)$ , with  $R$  small enough. Moreover,  $\Phi(t, U)$  stays in a bounded subset of  $H^s$  if  $U$  stays in  $B_s(R)$  for some  $s \geq \sigma$  and  $t \in [-1, 1]$ .

Let us notice that if  $s \geq \sigma$  and  $U$  is in  $B_s(R)$ ,  $D\Phi(t, U)$  extends as an element of  $\mathcal{L}(H^{s'}, H^{s'})$  for any  $s'$  with  $|s'| \leq s$ , any  $t$  in  $[-1, 1]$ . Actually, consider the solution  $W(t, U)$  of the linear ODE

$$(4.1.16) \quad \begin{aligned} \dot{W}(t, U) &= \mathcal{A}(\Phi(t, U))W(t, U) + [D\mathcal{A}(\Phi(t, U)) \cdot W(t, U)]\Phi(t, U) + D\Re(\Phi(t, U)) \cdot W(t, U) \\ W(0, U) &= \operatorname{Id}. \end{aligned}$$

We just need to check that the right hand side of the first equation in (4.1.16) is, as a function of  $W$ , a bounded linear map from  $\mathcal{L}(H^{s'}, H^{s'})$  to itself, for any  $s'$  with  $|s'| \leq s$ , when  $U$  is taken in  $B_s(R)$ . The boundedness of  $W \rightarrow D\Re(\Phi(t, U)) \cdot W$

follows from (4.1.3), since  $\Phi(t, U)$  stays in a bounded subset of  $H^s$  and (4.1.2) holds by the conditions imposed to  $\sigma, \gamma$ . The continuity of  $W \rightarrow \mathcal{A}(\Phi(t, U))W$  is a consequence of (1.2.9) and the fact that  $\sigma > \nu(P-1) + \nu + \frac{1}{2}$ . We are left with examining the middle term in the right hand side of the first equation (4.1.16). Let  $\bar{\nu} = \nu + \nu(P-1) - \nu(1) + \frac{1}{2} + 0$ . By the last remark following Definition 1.2.5,  $\mathcal{A}$  is in  $\Psi_1^{0, \bar{\nu}}(B_\sigma(R)) \otimes \mathcal{M}_2(\mathbb{R})$ . We write for  $H$  in  $H^{s'}$

$$\begin{aligned} \|D\mathcal{A}(\Phi(t, U)) \cdot (W \cdot H)\Phi(t, U)\|_{H^{s'}} &\leq C\|D\mathcal{A}(\Phi(t, U)) \cdot (W \cdot H)\|_{\mathcal{L}(H^s, H^{s'})}\|\Phi(t, U)\|_{H^s} \\ &\leq C\|W \cdot H\|_{H^{s'}}\|\Phi(t, U)\|_{H^s}, \end{aligned}$$

where the last inequality comes from (1.2.19) applied with  $\sigma = s, \sigma' = s'$ , and noticing that  $s - (\nu(1) + \bar{\nu} - s')_+ \geq s'$  because if the assumptions on  $s, s'$ . We get the wanted inequality

$$\|(D\mathcal{A}(\Phi(t, U)) \cdot W)\Phi(t, U)\|_{\mathcal{L}(H^{s'}, H^{s'})} \leq C\|W\|_{\mathcal{L}(H^{s'}, H^{s'})}.$$

This gives the fact that  $D\Phi(t, U)$  is bounded in  $\mathcal{L}(H^{s'}, H^{s'})$  uniformly for  $t \in [-1, 1]$ , if  $|s'| \leq s$ .

We deduce from this and from (1.2.18), (1.2.19):

**COROLLARY 4.1.3.** *If  $s > \gamma > \nu(P-1) + \nu + \frac{1}{2}$  and  $\bar{\nu} = \nu + \nu(P-1) - \nu(1) + \frac{1}{2} + 0$ ,  $(\mathcal{A}(\Phi(t, \cdot)))_{-1 \leq t \leq 1}$  is a bounded family of matrices of elements of  $\Psi_1^{0, \bar{\nu}}(B_s(R))$ .*

Let us describe the structure of  $\Phi(t, U)$ .

**PROPOSITION 4.1.4.** *Let  $P \in \mathbb{N}^*$ ,  $s \geq \sigma > \gamma > \nu(P-1) + \nu + \frac{1}{2}$ . Let  $A$  be in  $\sum_{p=1}^{P-1} H\Psi_p^{0, \nu} \subset \Psi_1^{0, \bar{\nu}}(B_\sigma(R))$  with  $\bar{\nu} = \nu + \nu(P-1) - \nu(1) + \frac{1}{2} + 0$ . Let  $\Phi$  be the solution of (4.1.15), defined for  $U$  in  $B_s(R)$  for some  $R > 0$ . If  $R$  is small enough, there is a bounded family  $(B(t, \cdot))_{-1 \leq t \leq 1}$  (resp.  $(\tilde{\mathfrak{R}}(t, \cdot))_{-1 \leq t \leq 1}$ ) of  $2 \times 2$ -matrices of elements of  $\Psi_1^{0, \bar{\nu}}(B_\sigma(R))$  (resp. of elements of  $\mathcal{R}_{\gamma, 2}^0(B_\sigma(R))$ ) such that, for any  $t \in [0, 1]$*

$$(4.1.17) \quad \Phi(t, U) = U + B(t, U)U + \tilde{\mathfrak{R}}(t, U).$$

*Proof:* By (4.1.15), (4.1.14),

$$(4.1.18) \quad \Phi(t, U) = U + t \int_0^1 \mathcal{A}(\Phi(\alpha_1 t, U))\Phi(\alpha_1 t, U) d\alpha_1 + t \int_0^1 \mathfrak{R}(\Phi(\alpha_1 t, U)) d\alpha_1.$$

If  $R > 0$  is small enough, there is  $C > 0$  such that  $\|\Phi(t, U)\|_{H^s} \leq C\|U\|_{H^s}$  for any  $U$  in  $B_s(R)$ , any  $t \in [-1, 1]$ . Iterating (4.1.18), we get

$$(4.1.19) \quad \Phi(t, U) = U + \sum_{k=1}^N t^k \mathcal{A}^{[k]}(t, U)U + t^{N+1} C_N(t, U) + \sum_{k=1}^{N+1} t^k \mathfrak{R}^{[k]}(t, U)$$

where

(4.1.20)

$$\begin{aligned}\mathcal{A}^{[k]}(t, U) &= \int_0^1 \dots \int_0^1 \prod_{j=1}^k \mathcal{A}(\Phi(t\alpha_1 \dots \alpha_j, U)) \prod_{j=1}^{k-1} \alpha_j^{k-j} d\alpha_1 \dots d\alpha_k \\ C_N(t, U) &= \int_0^1 \dots \int_0^1 \prod_{j=1}^{N+1} \mathcal{A}(\Phi(t\alpha_1 \dots \alpha_j, U)) \prod_{j=1}^N \alpha_j^{N+1-j} \Phi(t\alpha_1 \dots \alpha_{N+1}, U) d\alpha_1 \dots d\alpha_{N+1} \\ \mathfrak{R}^{[k]}(t, U) &= \int_0^1 \dots \int_0^1 \prod_{j=1}^{k-1} \mathcal{A}(\Phi(t\alpha_1 \dots \alpha_j, U)) \prod_{j=1}^k \alpha_j^{k-j} \mathfrak{R}(\Phi(t\alpha_1 \dots \alpha_k, U)) d\alpha_1 \dots d\alpha_k.\end{aligned}$$

We apply Lemma 1.2.6 with  $m = m' = 0$ ,  $p = p' = 1$ ,  $\nu = \nu' = \bar{\nu}$ . For some  $\mathcal{P} = (P_1, \dots, P_k)$  as in the statement of this lemma, we set for short  $\mathfrak{R}_\sigma^{\ell, s} = \mathfrak{R}_{(\mathcal{P}, 1, \bar{\nu}, \sigma)}^{\ell, s}$ . Then, if  $|\sigma'| \leq \sigma$ ,  $s \in \mathbb{R}$ , there is  $C > 0$  such that for any  $U$  in  $B_\sigma(R)$ , any  $t \in [0, 1]$

$$\begin{aligned}\mathfrak{R}_\sigma^{0, s}(\mathcal{A}^{[k]}(t, U)) &\leq \frac{C^k}{k!} [\mathfrak{R}_\sigma^{0, s}(A)]^k \\ \mathfrak{R}_{\sigma'}^{1, s}(\mathcal{A}^{[k]}(t, U)) &\leq \frac{C^k}{k!} [\mathfrak{R}_\sigma^{0, s}(A)]^{k-1} \mathfrak{R}_{\sigma'}^{1, s}(A)\end{aligned}$$

so that  $\sum_{k=1}^{+\infty} t^k \mathcal{A}^{[k]}(t, U)$  converges to an element  $B(t, \cdot)$  of  $\Psi_1^{0, \bar{\nu}}(B_\sigma(R))$ . Since  $\mathfrak{R}_\sigma^{0, s}(C_N(t, U)) = O(\frac{C^{N+1}}{(N+1)!} \mathfrak{R}_\sigma^{0, s}(A)^{N+1})$ , the last but one term in (4.1.19) goes to zero.

Let us check that the last series in (4.1.19) converges in the space  $\mathcal{R}_{\gamma, 2}^0(B_s(R))$ .

Note first that by (4.1.1) and (1.2.18)  $\|\mathfrak{R}^{[k]}(t, U)\|_{H^{2s-\gamma}} \leq \frac{C^k}{k!} \|U\|_{H^s}^2$  for  $U$  in  $B_s(R)$ , whence an estimate of type (4.1.1) for  $\sum_k t^k \mathfrak{R}^{[k]}(t, U)$ , uniformly for  $t \in [-1, 1]$ . Let us study the norm  $\|\partial_U \mathfrak{R}^{[k]}(t, U)\|_{\mathcal{L}(H^{\theta'}, H^{-\theta})}$  for  $\theta, \theta'$  satisfying (4.1.2). If, when computing  $\partial_U \mathfrak{R}^{[k]}$  from its expression (4.1.20), the derivative  $\partial_U$  falls on  $\mathfrak{R}(\Phi(\alpha_1 \dots \alpha_k t, U))$ , it follows from (4.1.3), the boundedness of  $\partial_U \Phi$  on  $H^{\theta'}$  for  $|\theta'| \leq s$ , and (1.2.18) that the corresponding contribution to  $\|\partial_U \mathfrak{R}^{[k]}(t, U)\|_{\mathcal{L}(H^{\theta'}, H^{-\theta})}$  is bounded from above by  $\frac{C^k}{k!} \|U\|_{H^s}$ . On the other hand, if the derivative falls on  $\mathcal{A}(\Phi(\alpha_1 \dots \alpha_j t, U))$ , we have to estimate for  $H'$  in  $H^{\theta'}$

$$(4.1.21) \quad \|(\partial_U A)(\Phi(t\alpha_1 \dots \alpha_j, U)) \cdot (\partial_U \Phi(t\alpha_1 \dots \alpha_j, U) \cdot H')\|_{\mathcal{L}(H^{2s-\gamma}, H^{-\theta})}$$

since  $\mathfrak{R}(\Phi(t, \cdot))$  is in  $H^{2s-\gamma}$  by (4.1.1). We apply (1.2.19) with  $\sigma = s$ ,  $\sigma' = \theta'$ . This allows us to estimate (4.1.21) for  $H' \in H^{\theta'}$  if  $2s - \gamma - (\nu(1) + \bar{\nu} - \theta')_+ \geq -\theta$ , which follows from (4.1.2), the definition of  $\bar{\nu}$  and the assumptions of the proposition. Again, we get convergence of the corresponding series. This concludes the proof.  $\square$

*Proof of Theorem 3.1.1:* We consider a function  $G(U) = \sum_{p=0}^{P'} G_p(U)$  as in the statement of Theorem 1.2.8. For  $p = 1, \dots, P-1$ ,  $G_p$  is an element of  $H\mathcal{F}_p^{1, \nu'}(B_\sigma(R))$  for some  $\nu'$ . We shall construct the canonical transformation  $\chi$  through a standard Birkhoff normal forms method. We use an auxiliary function  $F(U) = \sum_{p=1}^{P-1} F_p(U)$  with

$$(4.1.22) \quad F_p(U) = \operatorname{Re} \int_X (A_p(U)u)u d\mu$$



with  $A_p$  element of  $H\Psi_p^{0,\nu}$  for some  $\nu \geq \nu'$  to be chosen. Let us assume that the increasing function  $p \rightarrow \nu(p)$  satisfies

$$(4.1.23) \quad \nu(p) + \nu(q) + \nu + M_0(3d+4) \leq \nu(p+q) \text{ if } p > 0, q > 0.$$

We denote by  $\Phi$  the flow of (4.1.15) defined from  $F$  on  $B_s(R)$  where  $s > \nu(P-1) + \nu + \frac{1}{2}$ . We set  $\chi(U) = \Phi(-1, U)$ : this is a canonical transformation close to zero, with  $\chi(0) = 0, \chi'(0) = \text{Id}$ . For  $U$  in  $B_s(R) \cap C^\infty$ , we compute

$$(4.1.24) \quad \begin{aligned} G \circ \chi^{-1}(U) &= G(\Phi(1, U)) \\ &= \sum_{k=0}^{P-1} \frac{1}{k!} \frac{d^k}{dt^k} (G \circ \Phi(t, U))|_{t=0} + \int_0^1 \frac{(1-t)^{P-1}}{(P-1)!} \frac{d^P}{dt^P} (G \circ \Phi(t, U)) dt \\ &= \sum_{k=0}^{P-1} \frac{(-1)^k}{k!} \text{ad}^k F \cdot G + \frac{(-1)^P}{(P-1)!} \int_0^1 (1-t)^{P-1} (\text{ad}^P F \cdot G)(\Phi(t, U)) dt, \end{aligned}$$

using the notation  $\text{ad} F \cdot G = \{F, G\}$  and  $\text{ad}^p F \cdot G = \{F, \text{ad}^{p-1} F \cdot G\}$ . We decompose

$$(4.1.25) \quad \sum_{k=0}^{P-1} \frac{(-1)^k}{k!} \text{ad}^k F \cdot G = G_0 - \sum_{\ell=1}^{P-1} \{F_\ell, G_0\} + \sum_{\ell \geq 1} H_\ell$$

where the last sum is finite, and where  $H_\ell$  is a linear combination of quantities of the form

$$(4.1.26) \quad \text{ad} F_{i_1} \cdots \text{ad} F_{i_r} G_p$$

with  $1 \leq i_j < \ell, p \geq 0, i_1 + \cdots + i_r + p = \ell$  and  $r \geq 2$  if  $p = 0$ . Assume

$$(4.1.27) \quad F_p \in H\mathcal{F}_p^{0,\nu}(B_\sigma(R)) \text{ and } \{F_p, G_0\} \in H\mathcal{F}_p^{1,\nu'}(B_\sigma(R)) \quad 1 \leq p \leq P-1.$$

Then, by inclusion (3.2.11) and assumption (4.1.23), (4.1.26) belongs to  $H\mathcal{F}_\ell^{1,\nu'}(B_\sigma(R)) + H\mathcal{F}_\ell^{0,\nu'+1}(B_\sigma(R)) \subset H\mathcal{F}_\ell^{1,\nu'}(B_\sigma(R))$  if  $\sigma$  is large enough so that (1.2.23) with  $p = \ell$  is satisfied. Consequently, for any  $\ell \geq 1$ ,  $H_\ell$  is in  $H\mathcal{F}_\ell^{1,\nu'}(B_\sigma(R))$ , and depends only on  $F_{\ell'}$  for  $\ell' < \ell$ .

LEMMA 4.1.5. *Let  $\nu \geq \nu' + 2M_0$ ,  $\sigma$  satisfying (1.2.23), and denote*

$$\mathcal{G}_p^{m-\infty, \nu+\infty}(B_\sigma(R)) = \bigcap_{N \in \mathbb{N}} \mathcal{G}_p^{m-N, \nu+N}(B_\sigma(R))$$

(with the notations introduced in definition 1.2.7). For each  $p = 1, \dots, P-1$  there is  $F_p$  in  $H\mathcal{F}_{p,E}^{0,\nu}(B_\sigma(R))$  such that  $\{F_p, G_0\} - H_p$  is in  $H\mathcal{F}_{p,H}^{1,\nu'}(B_\sigma(R)) + \mathcal{G}_p^{1-\infty, \nu'+2M_0+\infty}(B_\sigma(R))$  and that (4.1.27) holds.

*Proof:* We decompose  $H_p = H_{p,E} + H_{p,H}$  with  $H_{p,E}$  in  $H\mathcal{F}_{p,E}^{1,\nu'}(B_\sigma(R))$ ,  $H_{p,H}$  in  $H\mathcal{F}_{p,H}^{1,\nu'}(B_\sigma(R))$ . We may write  $H_{p,E}(U) = \text{Re} \int_X (B_p(U)u)u d\mu$  with  $B_p$  in  $H\Psi_p^{1,\nu'}$ . By definition of this space

$$B_p(U) = \sum_{\ell=1}^p B_{p,\ell}(\underbrace{u, \dots, u}_\ell, \bar{u}, \dots, \bar{u})$$

where  $B_{p,\ell}$  is in  $\widetilde{H\Psi}_p^{1,\nu'}$ . We decompose  $B_{p,\ell} = B_{p,\ell}^{(1)} + B_{p,\ell}^{(2)}$  where for any  $n' \in (\mathbb{N}^*)^p$ ,  $\mathcal{U} = (u_1, \dots, u_p)$ ,

$$B_{p,\ell}^{(1)}(\Pi_{n'}\mathcal{U}) = \sum_{j; |n'| \leq c2^j} B_{p,\ell}(\Pi_{n'}\mathcal{U})\Delta_j$$

with  $c > 0$  a small enough constant. By formula (A.1) of the appendix,  $B_{p,\ell}^{(1)}$  is in  $\widetilde{H\Psi}_p^{1,\nu'}$ . Moreover, if  $c$  is small enough,

$$(4.1.28) \quad \Pi_{n_0} B_{p,\ell}^{(1)}(\Pi_{n'}\mathcal{U})\Pi_{n_{p+1}} \neq 0 \Rightarrow \sqrt{\mathbf{m}^2 + \lambda_{n_{p+1}}^2} \geq 2 \left( \sum_1^p \sqrt{\mathbf{m}^2 + \lambda_{n_\ell}^2} \right)$$

and by construction (and the inclusion (1.2.10)),  $B_{p,\ell}^{(2)}$  is in  $\mathcal{G}_p^{1-\infty, \nu'+2M_0+\infty}(B_\sigma(R))$ . To prove the lemma, we just need to find  $F_p$  in  $H\mathcal{F}_{p,E}^{0,\nu}(B_\sigma(R))$  such that

$$(4.1.29) \quad \{F_p, G_0\} = \text{Re} \sum_\ell \int_X (B_{p,\ell}^{(1)}(\underbrace{u, \dots, u}_\ell, \bar{u}, \dots, \bar{u})u)u \, d\mu.$$

We look for  $F_p$  under the form

$$F_p(U) = \sum_{\ell=0}^p \text{Re} \int_X (C_{p,\ell}(u, \dots, u, \bar{u}, \dots, \bar{u})u)u \, d\mu$$

with  $C_{p,\ell}$  in  $\widetilde{H\Psi}_p^{0,\nu}$ . To solve (4.1.29), it is enough to find  $C_{p,\ell}$  such that

$$(4.1.30) \quad \sum_{\ell'=1}^\ell C_{p,\ell}(u_1, \dots, \Lambda_{\mathbf{m}} u_{\ell'}, \dots, u_p) - \sum_{\ell'=\ell+1}^p C_{p,\ell}(u_1, \dots, \Lambda_{\mathbf{m}} u_{\ell'}, \dots, u_p) \\ + \Lambda_{\mathbf{m}} C_{p,\ell}(u_1, \dots, u_p) + C_{p,\ell}(u_1, \dots, u_p) \Lambda_{\mathbf{m}} = -i B_{p,\ell}^{(1)}(u_1, \dots, u_p).$$

We replace in the formula  $u_j$  by  $\Pi_{n_j} u_j$ , compose at the left (resp. at the right) with  $\Pi_{n_0}$  (resp.  $\Pi_{n_{p+1}}$ ). We get

$$(4.1.31) \quad \tilde{F}_{\mathbf{m}}^{p,\ell}(\lambda_{n_0}, \dots, \lambda_{n_{p+1}}) \Pi_{n_0} C_{p,\ell}(\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}} \\ = -i \Pi_{n_0} B_{p,\ell}^{(1)}(\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}$$

with

$$\tilde{F}_{\mathbf{m}}^{p,\ell}(\xi_0, \dots, \xi_{p+1}) = \sqrt{\mathbf{m}^2 + \xi_0^2} + \sqrt{\mathbf{m}^2 + \xi_{p+1}^2} + \sum_{\ell'=1}^\ell \sqrt{\mathbf{m}^2 + \xi_{\ell'}^2} - \sum_{\ell'=\ell+1}^p \sqrt{\mathbf{m}^2 + \xi_{\ell'}^2}.$$

We define

$$C_{p,\ell}(u_1, \dots, u_p) = -i \sum_{n_0} \dots \sum_{n_{p+1}} \tilde{F}_{\mathbf{m}}^{p,\ell}(\lambda_{n_0}, \dots, \lambda_{n_{p+1}})^{-1} \Pi_{n_0} B_{p,\ell}^{(1)}(\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}.$$

By inclusion (1.2.10),  $\Pi_{n_0} B_{p,\ell}^{(1)}(\Pi_{n'}\mathcal{U})\Pi_{n_{p+1}}$  satisfies estimate (1.2.5) with  $\nu = \nu' + 2M_0, m = 1$ . Since, by (4.1.28),  $|\tilde{F}_{\mathbf{m}}^{p,\ell}(\lambda_{n_0}, \dots, \lambda_{n_{p+1}})| \geq c(n_0 + n_{p+1})$  when

the right hand side of (4.1.31) is not zero, we get

(4.1.32)

$$\begin{aligned} \|\Pi_{n_0} C_{p,\ell}(\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}\|_{\mathcal{L}(L^2)} &\leq C \langle n_0 - n_{p+1} \rangle^{-2} \left( \min \left[ \frac{n_0}{n_{p+1}}, \frac{n_{p+1}}{n_0} \right] \right)^{N_1} \\ &\quad \times |n'|^{\nu(p)+\nu} \left( 1 + \frac{|n'|}{n_0} \right)^{-N_2} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2} \end{aligned}$$

for any  $N_1, N_2$ . To prove similar estimates for commutators with differential operators, we consider  $P_1$  a differential operator of order  $d_1$  and deduce from (4.1.30)

(4.1.33)

$$\begin{aligned} \sum_{\ell'=1}^{\ell} [P_1, C_{p,\ell'}](u_1, \dots, \Lambda_{\mathbf{m}} u_{\ell'}, \dots, u_p) - \sum_{\ell'=\ell+1}^p [P_1, C_{p,\ell'}](u_1, \dots, \Lambda_{\mathbf{m}} u_{\ell'}, \dots, u_p) \\ + \Lambda_{\mathbf{m}} [P_1, C_{p,\ell}](u_1, \dots, u_p) + [P_1, C_{p,\ell}](u_1, \dots, u_p) \Lambda_{\mathbf{m}} = -i \tilde{B}_{p,\ell}^{(1)}(u_1, \dots, u_p) \end{aligned}$$

where

$$\begin{aligned} \tilde{B}_{p,\ell}^{(1)}(u_1, \dots, u_p) &= -i [P_1, \Lambda_{\mathbf{m}}] C_{p,\ell}(u_1, \dots, u_p) - i C_{p,\ell}(u_1, \dots, u_p) [P_1, \Lambda_{\mathbf{m}}] \\ &\quad + [P_1, B_{p,\ell}^{(1)}](u_1, \dots, u_p). \end{aligned}$$

We need to prove

$$\begin{aligned} (4.1.34) \quad \|\Pi_{n_0} [P_1, C_{p,\ell}](\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}\|_{\mathcal{L}(L^2)} \\ \leq C \langle n_0 - n_{p+1} \rangle^{-2} n_0^{d_1-1} \left( \min \left[ \frac{n_0}{n_{p+1}}, \frac{n_{p+1}}{n_0} \right] \right)^{N_1} |n'|^{\nu(p)+\nu+M_0} \\ \quad \times \left( 1 + \frac{|n'|}{n_0} \right)^{-N_2} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2} \end{aligned}$$

for any  $N_1, N_2$ . Since  $B_{p,\ell}^{(1)}$  is in  $\widetilde{H\Psi}_p^{1,\nu'}$ ,  $[P_1, B_{p,\ell}^{(1)}]$  is in  $\widetilde{H\Psi}_p^{d_1,\nu'+M_0}$  by Definition 1.2.2, so in  $\widetilde{\mathcal{P}}_p^{d_1,\nu'+M_0}$  by inclusion (1.2.10). Consequently

$$\begin{aligned} (4.1.35) \quad \|\Pi_{n_0} [P_1, B_{p,\ell}^{(1)}](\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}\|_{\mathcal{L}(L^2)} \\ \leq C \langle n_0 - n_{p+1} \rangle^{-2} n_0^{d_1} \left( \min \left[ \frac{n_0}{n_{p+1}}, \frac{n_{p+1}}{n_0} \right] \right)^{N_1} |n'|^{\nu(p)+\nu+M_0} \\ \quad \times \left( 1 + \frac{|n'|}{n_0} \right)^{-N_2} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}. \end{aligned}$$

Since  $[P_1, \Lambda_{\mathbf{m}}]$  is a pseudo-differential operator of order  $d_1$ , we have estimates

$$(4.1.36) \quad \|\Pi_{n_1} [P_1, \Lambda_{\mathbf{m}}] \Pi_{n_2}\|_{\mathcal{L}(L^2)} \leq C_N \langle n_1 - n_2 \rangle^{-N} n_1^{d_1}$$

for any  $N$ . Combining this with (4.1.32), we conclude that the right hand side of (4.1.33) is such that  $\|\Pi_{n_0} \tilde{B}_{p,\ell}^{(1)}(\Pi_{n'}(\mathcal{U})) \Pi_{n_{p+1}}\|_{\mathcal{L}(L^2)}$  is bounded from above by the right hand side of (4.1.35). If we use (4.1.36) for  $P_1$  instead of  $[P_1, \Lambda_{\mathbf{m}}]$  and (4.1.32), we conclude that  $P_1 C_{p,\ell}$ ,  $C_{p,\ell} P_1$  and so  $[P_1, C_{p,\ell}]$  satisfy estimate (4.1.32)

in which we replace, in the right hand side,  $|n'|^{\nu(p)+\nu}$  by  $|n'|^{\nu(p)+\nu} n_0^{d_1}$ . We decompose  $[P_1, C_{p,\ell}] = C_{p,\ell}^{(1)} + C_{p,\ell}^{(2)}$  where

$$C_{p,\ell}^{(1)} = \sum_{|n'| \leq cn_{p+1}} \Pi_{n_0}[P_1, C_{p,\ell}](\Pi_{n'}\mathcal{U})\Pi_{n_{p+1}}.$$

Then  $C_{p,\ell}^{(2)}$  satisfies (4.1.35) with  $n_0^{d_1}$  replaced by  $n_0^{d_1-1}$  in the right hand side, because of  $|n'| \geq cn_{p+1}$ , so that

$$\begin{aligned} & \Pi_{n_0}\Lambda_{\mathbf{m}}C_{p,\ell}^{(2)}(\Pi_{n'}\mathcal{U})\Pi_{n_{p+1}}, \quad \Pi_{n_0}C_{p,\ell}^{(2)}(\Pi_{n'}\mathcal{U})\Lambda_{\mathbf{m}}\Pi_{n_{p+1}}, \\ & \Pi_{n_0}C_{p,\ell}^{(2)}(\Pi_{n_1}u_1, \dots, \Pi_{n_{\ell'}}\Lambda_{\mathbf{m}}u_{\ell'}, \dots, \Pi_{n_p}u_p)\Pi_{n_{p+1}} \end{aligned}$$

satisfy (4.1.35). We obtain in that way from (4.1.33) an equation

$$(4.1.37) \quad \sum_{\ell'=1}^{\ell} C_{p,\ell}^{(1)}(u_1, \dots, \Lambda_{\mathbf{m}}u_{\ell'}, \dots, u_p) - \sum_{\ell'=\ell+1}^{p+1} C_{p,\ell}^{(1)}(u_1, \dots, \Lambda_{\mathbf{m}}u_{\ell'}, \dots, u_p) \\ + \Lambda_{\mathbf{m}}C_{p,\ell}^{(1)}(u_1, \dots, u_p) + C_{p,\ell}^{(1)}(u_1, \dots, u_p)\Lambda_{\mathbf{m}} = -i\tilde{B}_{p,\ell}^{(2)}(u_1, \dots, u_p),$$

where  $\|\Pi_{n_0}\tilde{B}_{p,\ell}^{(2)}(\Pi_{n'}\mathcal{U})\Pi_{n_{p+1}}\|_{\mathcal{L}(L^2)}$  is bounded by the right hand side of (4.1.35) and where, if the constant  $c$  in the definition of  $C_{p,\ell}^{(1)}$  is small enough, this operator satisfies a condition of type (4.1.28). We may thus solve (4.1.37) in the same way as (4.1.30) and find  $C_{p,\ell}^{(1)}$  and  $C_{p,\ell}$  such that (4.1.34) holds. It follows from this inequality and from (4.1.32) that  $C_{p,\ell}$  satisfies the bounds (1.2.8) (or (1.2.9)) for  $m = 0, k = 0, 1$ . Iterating the reasoning, we conclude that  $C_{p,\ell}$  is in  $\tilde{H}\Psi_p^{0,\nu}$  as wanted.  $\square$

*End of proof of Theorem 3.1.1.* We consider the expression (4.1.24) of  $G \circ \chi^{-1}$ . The first term in the right hand side is given by (4.1.25). If  $\sigma$  is large enough, the contributions  $H_\ell$  with  $\ell \geq P$  belong to  $H\mathcal{F}_\ell^{1,\nu'}(B_\sigma(R))$  which is contained by the last remark following Definition 1.2.5 in  $\mathcal{F}_P^{1,\nu}(B_\sigma(R))$  if  $\nu > \nu' + \nu(\ell) - \nu(P) + \frac{1}{2}$ . Since there are only finitely many  $H_\ell$ 's, this gives a contribution of the form of  $\tilde{G}_P$  given by (3.1.6) to  $G \circ \chi^{-1}$ . For  $1 \leq \ell \leq P-1$ , we have constructed  $F_\ell$  in Lemma 4.1.5 so that  $\{F_\ell, G_0\} - H_\ell$  belongs to  $H\mathcal{F}_{\ell,\mathbf{H}}^{1,\nu}(B_\sigma(R)) + \mathcal{G}_\ell^{1-\infty, \nu+\infty}(B_\sigma(R))$ , if  $\nu$  is taken large enough relatively to  $\nu'$ . These terms give a contribution to  $G_L + G_H$  defined by (3.1.5), (3.1.4). To prove (3.1.3), we write, since  $\chi$  is canonical,

$$\{\Theta_s^1 \circ \chi, G\} = \{\Theta_s^1, G \circ \chi^{-1}\} \circ \chi$$

and since the sum in the right hand side of (4.1.24) gives to  $G \circ \chi^{-1}$  a contribution of the form  $G_L + G_H + \tilde{G}_P$ , we are left with showing that, if

$$S = \frac{(-1)^P}{(P-1)!} \int_0^1 (1-t)^{P-1} ((\text{ad}^P F) \cdot G)(\Phi(t, U)) dt,$$

then

$$(4.1.38) \quad \{\Theta_s^1, S\} \circ \chi = \{\Theta_s^1, \tilde{G}_P\} \circ \chi + O(\|u\|_{H^s}^{P+2})$$

for some (new) contribution  $\tilde{G}_P$  of form (3.1.6). The term  $S$  may be written as a sum of expressions

$$(4.1.39) \quad \int_0^1 (1-t)^{P-1} H_p(\Phi(t, U)) dt$$

where  $p \geq P$  and  $H_p$  is an expression of form (4.1.26), so belongs to  $H\mathcal{F}_p^{1,\nu'}(B_\sigma(R))$ , i.e. (4.1.39) may be written from

$$(4.1.40) \quad \begin{aligned} & \operatorname{Re} \int_0^1 (1-t)^{P-1} \int_X (A_p(\Phi(t, U))\phi(t, U))\phi(t, U) d\mu dt \\ & \operatorname{Re} \int_0^1 (1-t)^{P-1} \int_X (A_p(\Phi(t, U))\phi(t, U))\overline{\phi(t, U)} d\mu dt \end{aligned}$$

with  $A_p$  in  $H\Psi_p^{1,\nu'}$ . The first component  $\phi$  of  $\Phi$  is given, according to (4.1.17), by

$$\phi(t, U) = u + B_1(t, U)u + B_2(t, U)\bar{u} + R(t, U)$$

where  $(B_j(t, \cdot))_{t \in [0,1]}$  are bounded families of elements of  $\Psi_1^{0,\bar{\nu}}(B_s(R))$  for  $j = 1, 2$  and where  $(R(t, \cdot))_{t \in [0,1]}$  is a bounded family of elements of  $\mathcal{R}_{\gamma,2}^0(B_s(R))$  (for  $s$  large enough). Plugging this expression inside (4.1.40), and using Corollary 4.1.3 and Lemma 1.2.6, we see that (4.1.40) may be written as a linear combination of elements of  $\mathcal{F}_P^{1,\nu}(B_s(R))$  (for some large enough  $\nu$  and some large enough  $s$ ) and of quantities of type

$$(4.1.41) \quad \int_0^1 \int_X (C(t, U)u)R(t, U) d\mu dt, \quad \int_0^1 \int_X (C(t, U)R(t, U))R(t, U) d\mu dt$$

with  $(C(t, \cdot))_{t \in [0,1]}$  in  $\Psi_P^{1,\nu}(B_s(R))$ ,  $(R(t, \cdot))_{t \in [0,1]}$  in  $\mathcal{R}_{\gamma,2}^0(B_s(R))$ , as well as similar expressions with  $u, R$  replaced by  $\bar{u}, \bar{R}$ . We just need to check that the Poisson bracket between  $\Theta_s^1$  and (4.1.41) is  $O(\|U\|_{H^s}^{P+2})$  to get (4.1.38). By assumption,  $\|D\Theta_s^1(U)\|_{\mathcal{L}(H^s, \mathbb{R})} = O(\|U\|_{H^s})$ , so that it is enough to check that if  $H(U)$  is any of the expressions (4.1.41),  $\nabla H$  belongs to  $H^s$ , with  $H^s$ -norm  $O(\|U\|_{H^s}^{P+1})$ . In other words, we have to study, for  $K$  in  $H^{-s}$ ,

$$\begin{aligned} & \int_X (C(t, U)K)R(t, U) d\mu, \quad \int_X ((d_U C(t, U) \cdot K)u)R(t, U) d\mu \\ & \int_X (C(t, U)u)(d_U R(t, U) \cdot K) d\mu, \quad \int_X (C(t, U)R(t, U))(d_U R(t, U) \cdot K) d\mu \\ & \int_X ((d_U C(t, U) \cdot K)R(t, U))R(t, U) d\mu \end{aligned}$$

(and similar expressions involving transposes). By (4.1.1),  $R(t, U)$  is in  $H^{2s-\gamma} \subset H^{s+1}$  (if  $s \geq s_0$  large enough), by (1.2.19),  $(d_U C(t, U) \cdot K)u \in H^{-\nu(p)-\nu} \subset H^{-s-1}$ ,  $(d_U C(t, U) \cdot K)R(t, U) \in H^{-s}$ , by (4.1.3),  $d_U R(t, U) \cdot K \in H^{-s+1}$ . This shows that all the above integrals are meaningful and vanish at least at order  $P+2$  at zero. This concludes the proof.  $\square$

**4.2. Diagonalization of the remainder term.** The goal of this subsection is to prove Theorem 3.1.2. Theorem 3.1.1 has reduced ourselves to the study of a new Hamiltonian  $G^1 = G_L + G_H + \tilde{G}_P$ , that may be written

$$(4.2.1) \quad G^1 = G_L(U) + \int_X ((\Lambda_{\mathbf{m}} + A(U))u)\bar{u} d\mu + \frac{1}{2} \int_X (B_P(U)u)u d\mu + \frac{1}{2} \int_X (\overline{B_P(U)}\bar{u})\bar{u} d\mu$$

where  $A(U) = \sum_{j=1}^P A_j(U)$ . By the last remark following Definition 1.2.5, we may assume that  $A$  is in  $\Psi_1^{1,\nu}(B_s(R))$  (increasing the value of  $\nu$  given by Theorem 3.1.1). Moreover,  $B_P$  is in  $\Psi_P^{1,\nu}(B_s(R))$ , and we may assume  $A(U)^* = A(U)$ ,  ${}^t B_P(U) = B_P(U)$ . We denote by  $A_H(U) = \sum_{p=1}^{P-1} A_p(U)$  the part of  $A$  made of homogeneous terms. Consider the matrices of para-differential operators

$$(4.2.2) \quad M(U) = \begin{bmatrix} B_P(U) & \Lambda_{\mathbf{m}} + {}^t A(U) \\ \Lambda_{\mathbf{m}} + A(U) & \overline{B_P(U)} \end{bmatrix}, \quad M_H(U) = \begin{bmatrix} 0 & \Lambda_{\mathbf{m}} + {}^t A_H(U) \\ \Lambda_{\mathbf{m}} + A_H(U) & 0 \end{bmatrix},$$

so that

$$(4.2.3) \quad (G_H + \tilde{G}_P)(U) = \frac{1}{2} \int_X (M(U) \cdot U)U d\mu, \quad G_H(U) = \frac{1}{2} \int_X (M_H(U) \cdot U)U d\mu.$$

Let us start with a diagonalization lemma.

**LEMMA 4.2.1.** *There are  $s_0$  large enough,  $\tilde{\nu} \geq \nu$  large enough,  $R > 0$  and for  $s \geq s_0$ , operators  $Q$  in  $\Psi_P^{0,\tilde{\nu}}(B_s(R)) \otimes \mathcal{M}_2(\mathbb{R})$ ,  $C$  in  $\Psi_P^{1,\tilde{\nu}}(B_s(R))$ , with  $C$  self-adjoint and symmetric, such that, if we set*

$$(4.2.4) \quad \Sigma(U) = \begin{bmatrix} \Lambda_{\mathbf{m}} + A(U) + C(U) & 0 \\ 0 & -\Lambda_{\mathbf{m}} - {}^t A(U) - C(U) \end{bmatrix},$$

one has the relations

$$(4.2.5) \quad \begin{aligned} & {}^t(I + Q(U)) {}^t J(I + Q(U)) - {}^t J \in \Psi_P^{-1,\tilde{\nu}}(B_s(R)) \otimes \mathcal{M}_2(\mathbb{R}) \\ & (I + Q(U))({}^t J M(U)) J^t(I + Q(U)) {}^t J - \Sigma(U) \in \Psi_P^{0,\tilde{\nu}}(B_s(R)) \otimes \mathcal{M}_2(\mathbb{R}). \end{aligned}$$

*Proof:* Define.

$$A^1 = \frac{1}{2}(A + \bar{A}), \quad A^2 = \frac{1}{2}(A - \bar{A}), \quad B^1 = \frac{1}{2}(B_P + \bar{B}_P), \quad B^2 = \frac{1}{2i}(B_P - \bar{B}_P).$$

The assumptions on  $A, B$  imply

$$(A^1)^* = {}^t A^1 = A^1, \quad (A^2)^* = A^2 = -{}^t A^2, \quad (B^1)^* = B^1, \quad (B^2)^* = B^2$$

so that

$$(4.2.6) \quad {}^t J M(U) = \begin{bmatrix} \Lambda_{\mathbf{m}} + A^1 + A^2 & B^1 - iB^2 \\ -B^1 - iB^2 & -\Lambda_{\mathbf{m}} - A^1 + A^2 \end{bmatrix}.$$

By (i) of Proposition 2.2.6, there are principal symbols  $a^1, a^2$  in  $S_1^{1,\nu}(B_s(R), T^*X)$ ,  $b^1, b^2$  in  $S_P^{1,\nu}(B_s(R), T^*X)$  of  $A^1, A^2, B^1, B^2$  respectively, modulo  $\Psi_1^{0,\tilde{\nu}}(B_s(R))$  and  $\Psi_P^{0,\tilde{\nu}}(B_s(R))$  for  $\tilde{\nu} \geq \nu + M_0(3d + 4)$ . Since  $A^1, A^2, B^1, B^2$  (resp.  $A^1, B^1, B^2$ ) are self-adjoint (resp. symmetric) we may assume by (iv) of Proposition 2.2.6 that  $a^1, a^2, b^1, b^2$  are real valued and that  $a^1 = (a^1)^\vee$ ,  $b^1 = (b^1)^\vee$ ,  $b^2 = (b^2)^\vee$  (increasing

eventually  $\tilde{\nu}$ ). We denote by  $\lambda \in S_0^{1,0}(T^*M)$  the principal symbol of  $\Lambda_{\mathbf{m}}$ , given in local coordinates by  $(\mathbf{m}^2 + g_x(\xi, \xi))^{1/2}$  where  $g$  is the metric, and set

$$b = b^1 + ib^2, \quad c = \frac{b}{\lambda + a^1 + \sqrt{(\lambda + a^1)^2 + |b|^2}}.$$

If  $R$  is small enough, the definition is meaningful since  $a^1, b$  vanish at  $U = 0$ , and we get that  $c$  is in  $S_P^{0,\nu}(B_s(R), T^*X)$ . We define

$$(4.2.7) \quad \text{Id} + q = (1 - |c|^2)^{-1/2} \begin{bmatrix} 1 & \bar{c} \\ c & 1 \end{bmatrix}.$$

By a direct computation using that  $c^\vee = c$ , we get

$$(4.2.8) \quad {}^t(\text{Id} + q^\vee) {}^t J (\text{Id} + q) = {}^t J$$

and, if we define the matrix of symbols  $\mathbf{m}$  by

$${}^t J \mathbf{m} = \begin{bmatrix} \lambda + a^1 + a^2 & b^1 - ib^2 \\ -(b^1 + ib^2) & -\lambda - a^1 + a^2 \end{bmatrix},$$

we obtain

$$(4.2.9) \quad (\text{Id} + q)({}^t J \mathbf{m}) J {}^t (\text{Id} + q^\vee) {}^t J = \begin{bmatrix} \sqrt{(\lambda + a^1)^2 - |b|^2} + a^2 & 0 \\ 0 & -\sqrt{(\lambda + a^1)^2 - |b|^2} + a^2 \end{bmatrix}.$$

Actually, the eigenvalues of  $({}^t J \mathbf{m})$  are  $a^2 \pm \sqrt{(\lambda + a^1)^2 - |b|^2}$  and  $\begin{bmatrix} 1 \\ -c \end{bmatrix}$  (resp.  $\begin{bmatrix} -\bar{c} \\ 1 \end{bmatrix}$ ) is an eigenvector associated to the first (resp. second) eigenvalue. If we set

$$(\text{Id} + p) = (1 - |c|^2)^{-1/2} \begin{bmatrix} 1 & -\bar{c} \\ -c & 1 \end{bmatrix},$$

$(\text{Id} + p)^{-1}({}^t J \mathbf{m})(\text{Id} + p)$  equals the right hand side of (4.2.9). One has just to define  $(\text{Id} + q) = (\text{Id} + p)^{-1}$  and to use (4.2.8) to get (4.2.9).

By (ii) of proposition 2.2.6, we may find operators  $E, F$  in  $\Psi_P^{0,\tilde{\nu}}(B_s(R))$  whose principal symbols modulo  $\Psi_P^{-1,\tilde{\nu}}(B_s(R))$  are  $(1 - |c|^2)^{-1/2}$  and  $\bar{c}(1 - |c|^2)^{-1/2}$  respectively (taking again  $\tilde{\nu}$  large enough). Since  $c^\vee = c$ , we may assume, using (iv) of Proposition 2.2.6, that  $E = {}^t E$ ,  $F = {}^t F$ . We define  $Q$  by

$$(4.2.10) \quad I + Q(U) = \begin{bmatrix} E & F \\ \bar{F} & \bar{E} \end{bmatrix}.$$

Then (4.2.8) and the properties of symbolic calculus of (iii) of Proposition 2.2.6 imply that

$${}^t(I + Q(U)) {}^t J (I + Q(U)) - {}^t J \in \Psi_P^{-1,\tilde{\nu}}(B_s(R)).$$

Let  $C$  be an element of  $\Psi_P^{1,\tilde{\nu}}(B_s(R))$ , self-adjoint and symmetric, whose principal symbol modulo  $\Psi_P^{0,\tilde{\nu}}(B_s(R))$  is given by  $\sqrt{(\lambda + a^1)^2 - |b|^2} - (\lambda + a^1)$ . Then (4.2.6), the definition (4.2.4) of  $\Sigma$ , the equality (4.2.9) and (iii) of proposition 2.2.6 imply that the second equality (4.2.5) holds (for  $\tilde{\nu}$  large enough).  $\square$

Before continuing the proof of Theorem 3.1.2, we write some properties that will be used repeatedly below. Consider  $P$  an element of  $\Psi_P^{m,\tilde{\nu}}(B_s(R))$  for  $m = 0, 1$ . We apply (1.2.19) with  $\sigma = s$ ,  $\sigma' = -s$  and the smoothness index denoted by  $s$  in

(1.2.19) taken to be  $s$  or  $s - 1$ . We get that if  $H \in H^{-s}$ ,  $d_U P(U) \cdot H$  belongs to  $\mathcal{L}(H^s, H^{-s+1})$  and to  $\mathcal{L}(H^{s-1}, H^{-s})$ , if  $s$  is taken large enough relatively to  $\nu(p)$ ,  $\tilde{\nu}$ . Moreover, the estimate

$$(4.2.11) \quad \|d_U P(U) \cdot H\|_{\mathcal{L}(H^s, H^{-s+1})} + \|d_U P(U) \cdot H\|_{\mathcal{L}(H^{s-1}, H^{-s})} \leq C \|H\|_{H^{-s}} \|U\|_{H^s}^{p-1}$$

holds. In the same way, again for  $s$  large enough, if  $H$  is in  $H^{s-1}$ ,  $d_U P(U) \cdot H$  is in  $\mathcal{L}(H^s, H^{s-m})$ , with

$$(4.2.12) \quad \|d_U P(U) \cdot H\|_{\mathcal{L}(H^s, H^{s-m})} \leq C \|H\|_{H^{s-1}} \|U\|_{H^s}^{p-1}.$$

If  $P(U) = \tilde{P}(u, \dots, u, \bar{u}, \dots, \bar{u})$  with  $\tilde{P}$  in  $\tilde{\mathcal{P}}_p^{m, \tilde{\nu}}$  for some  $p \in \mathbb{N}^*$ ,  $p \leq P - 1$ , estimates (1.2.7) show that (4.2.11), (4.2.12) hold. Actually, we shall also use similar estimates for the difference  $(d_U P(U) - d_U P(U')) \cdot H$ , replacing in the right hand side of (4.2.11), (4.2.12)  $\|U\|_{H^s}^{p-1}$  by the quantity  $(\|U\|_{H^s} + \|U'\|_{H^s})^{p-2} \|U - U'\|_{H^s}$ .

LEMMA 4.2.2. *There is  $s_0 > 0$  and for any  $s \geq s_0$ , the map  $U \rightarrow \psi(U) = (I + Q(U))U$  defines a local diffeomorphism from a neighborhood of zero in  $H^s$  to a neighborhood of zero in  $H^s$ , coming from a real diffeomorphism in real coordinates. If we define, for  $V$  in a small enough neighborhood of zero in  $H^s$ ,*

$$(4.2.13) \quad \hat{\Sigma}(V) = \Sigma(\psi^{-1}(V)), \quad \hat{G}(V) = \frac{1}{2} \int_X (J\hat{\Sigma}(V)V)V d\mu,$$

*then for any  $U$  in a small enough neighborhood of zero in  $H^s$*

$$(4.2.14) \quad \|d\psi(U) \cdot X_{[G_H + \tilde{G}_P]}(U) - X_{\hat{G}(\psi(U))}\|_{H^s} = O(\|U\|_{H^s}^{P+1}), \quad U \rightarrow 0.$$

*Proof:* We shall call a “good term” any function  $U \rightarrow S(U)$  defined on a neighborhood of zero in  $H^s$ , such that  $\|S(U)\|_{H^s} = O(\|U\|_{H^s}^{P+1})$ ,  $U \rightarrow 0$ . It follows from (4.2.3) that  $X_{[G_H + \tilde{G}_P]} = i^t J\nabla(G_H + \tilde{G}_P)$  is given by

$$(4.2.15) \quad \begin{aligned} \int_X X_{[G_H + \tilde{G}_P]}(U) \cdot H d\mu &= id(G_H + \tilde{G}_P)(U) \cdot (JH) \\ &= i \int_X ({}^t JM(U) \cdot U)H d\mu + \frac{i}{2} \int_X (dM(U) \cdot (JH)U)U d\mu \end{aligned}$$

for any  $H$  in  $C^\infty(X)$ . We may write

$$(4.2.16) \quad X_{[G_H + \tilde{G}_P]}(U) = i^t JM(U)U + Z(U)$$

where  $Z(U)$  is defined by the equality

$$(4.2.17) \quad \int_X Z(U)H d\mu = \frac{i}{2} \int_X (dM(U) \cdot (JH)U)U d\mu \text{ for any } H.$$

By (4.2.11), the right hand side is defined and continuous if  $H$  is in  $H^{-s}$ , so that  $Z$  is in  $H^s$  and  $\|Z(U)\|_{H^s} = O(\|U\|_{H^s}^2)$ . By the inverse function theorem,  $\psi$  is a local diffeomorphism at 0, and the form (4.2.10) of  $Q$  shows that  $\psi$  sends a vector  $\begin{bmatrix} u \\ u \end{bmatrix}$

on a vector  $\begin{bmatrix} v \\ v \end{bmatrix}$  i.e. that it is induced by a local diffeomorphism in real coordinates.

Compute

$$(4.2.18) \quad \begin{aligned} d\psi(U) \cdot X_{[G_H + \tilde{G}_P]}(U) &= (I + Q(U))i^t JM(U)U + (I + Q(U))Z(U) \\ &\quad + (dQ(U) \cdot X_{[G_H + \tilde{G}_P]}(U))U. \end{aligned}$$



The fact that (4.2.16) belongs to  $H^{s-1}$ , and (4.2.12) with  $m = 0$ , show that the last term in (4.2.18) is a good term. Since  $Z(U)$  is in  $H^s$ ,  $Q(U)Z(U)$  is also a good term, so that

$$(4.2.19) \quad d\psi(U) \cdot X_{[G_H + \tilde{G}_P]}(U) = (I + Q(U))i^t J M(U)U + Z(U) + S(U)$$

where  $S(U)$  is a good term. By the first equality (4.2.5), we may write

$$U = J^t(I + Q(U))^t J(I + Q(U)U + S_1(U))$$

where  $\|S_1(U)\|_{H^{s+1}} \leq C\|U\|_{H^s}^{P+1}$ . Inserting this into the right hand side of (4.2.19) and using the second formula (4.2.5), we get

$$(4.2.20) \quad d\psi(U) \cdot X_{[G_H + \tilde{G}_P]}(U) = i\Sigma(U)\psi(U) + Z(U) + S(U)$$

for a new good term  $S(U)$ . We define  $\hat{\Sigma}$  by (4.2.13). By (1.2.19),  $dQ(U) \cdot H$  is in  $\mathcal{L}(H^s, H^{s'})$  for any  $s'$  with  $|s'| \leq s$ , any  $H$  in  $H^{s'}$ , if  $s$  is large enough. Consequently,  $d\psi(U)$  and  $d\psi(U)^{-1}$  are bounded linear maps on  $H^{s'}$ , for any  $s'$  with  $|s'| \leq s$ , if  $U$  stays in a small enough neighborhood of zero. Moreover, the operator norm of  $d\psi(U) - \text{Id}$  in these spaces is  $O(\|U\|_{H^s}^P)$ . A consequence of this and the definition of  $\hat{\Sigma}$  is that  $\hat{\Sigma}(V) - \begin{bmatrix} \Lambda_m & 0 \\ 0 & -\Lambda_m \end{bmatrix}$  belongs to  $\Psi_1^{1, \tilde{\nu}}(B_s(R)) \otimes \mathcal{M}_2(\mathbb{R})$  for  $R > 0$  small enough. Let us define  $\widehat{M}(V) = J\hat{\Sigma}(V)$ . By (4.2.13), we may write

$$(4.2.21) \quad X_{\hat{G}}(V) = i\hat{\Sigma}(V)V + \widehat{Z}(V)$$

where  $\widehat{Z}(V)$  is defined through the equality

$$(4.2.22) \quad \int_X \widehat{Z}(V) \cdot H \, d\mu = \frac{i}{2} \int_X (d\widehat{M}(V) \cdot (JH)V)V \, d\mu \text{ for any } H.$$

To obtain (4.2.14), we must check that  $d\psi(U) \cdot X_{[G_H + \tilde{G}_P]}(U) - X_{\hat{G}}(\psi(U))$  is a good term. By (4.2.20), (4.2.21), this difference is given by  $Z(U) - \widehat{Z}(\psi(U))$ , modulo a good term. To conclude the proof, we have to show, taking (4.2.17) and (4.2.22) into account, that, for any  $H$  in  $H^{-s}$ , the modulus of

$$(4.2.23) \quad \int_X (Z(U) - \widehat{Z}(\psi(U)))H \, d\mu = \frac{i}{2} \int_X [((dM(U) \cdot (JH))U)U - ((d\widehat{M}(\psi(U))(JH))\psi(U))\psi(U)] \, d\mu$$

is bounded from above by  $C\|H\|_{H^{-s}}\|U\|_{H^s}^{P+1}$ . By (4.2.11),  $d\widehat{M}(\psi(U))(JH)$  is in  $\mathcal{L}(H^s, H^{-s})$  and  $\|\psi(U) - U\|_{H^s} = O(\|U\|_{H^s}^P)$ . Consequently, we are reduced to the study of

$$(4.2.24) \quad \int_X (((dM(U) - d\widehat{M}(\psi(U))) \cdot (JH))U)U \, d\mu.$$

By definition of  $\hat{\Sigma}$  and the second relation (4.2.5)

$$\widehat{M}(\psi(U)) = J\Sigma(U) = J(I + Q(U))^t J M(U) J^t (I + Q(U))^t J + \mathfrak{R}(U)$$

with  $\mathfrak{R}(U)$  in  $\Psi_P^{0, \tilde{\nu}}(B_s(R)) \otimes \mathcal{M}_2(\mathbb{R})$ . We decompose

$$(d\widehat{M}(\psi(U)) - dM(U)) \cdot (JH) = I + II + III$$

where

$$I = ((d\widehat{M}(\psi(U)) - d[\widehat{M}(\psi(U))]) \cdot (JH) = (d\widehat{M})(\psi(U))[\text{Id} - d\psi(U)] \cdot JH$$

$$\begin{aligned}
II &= J(dQ(U) \cdot (JH))^t JM(U) J^t(I + Q(U))^t J \\
&\quad + J(I + Q(U))^t JM(U) J(d^t Q(U) \cdot (JH))^t J + d\mathfrak{R}(U) \cdot JH \\
III &= J(I + Q(U))^t J(dM(U) \cdot (JH)) J^t(I + Q(U))^t J - dM(U) \cdot (JH).
\end{aligned}$$

To finish to prove that (4.2.24) is  $O(\|H\|_{H^{-s}}\|U\|_{H^s}^{P+1})$ , it remains to show that the  $\mathcal{L}(H^s, H^{-s})$  norm of I, II and III is  $O(\|H\|_{H^{-s}}\|U\|_{H^s}^{P-1})$ .

Since  $\|d\psi(U) - \text{Id}\|_{\mathcal{L}(H^{-s}, H^{-s})} = O(\|U\|_{H^s}^P)$ , the estimate of I follows from (4.2.11). The bound of II follows from (4.2.11) applied to  $P = Q$ , and from the fact that  $d\mathfrak{R}(U) \cdot (JH)$  is in  $\mathcal{L}(H^s, H^{-s})$  with norm  $O(\|H\|_{H^{-s}}\|U\|_{H^s}^{P-1})$ . Finally, the estimate of III follows again from (4.2.11), and from the fact that  $\|Q(U)\|_{\mathcal{L}(H^{s'}, H^{s'})} = O(\|U\|_{H^s}^P)$  for any  $s'$ . This concludes the proof of the lemma.  $\square$

*End of the proof of Theorem 3.1.2:* The left hand side of (3.1.7) is

$$(4.2.25) \quad \{\Theta_s^2 \circ \psi, G_L + G_H + \tilde{G}_P\} = d\Theta_s^2(\psi(U)) \cdot d\psi(U) \cdot [X_{G_L}(U) + X_{[G_H + \tilde{G}_P]}(U)].$$

The contribution of  $G_L$  to the right hand side may be written

$$(4.2.26) \quad \{\Theta_s^2, G_L\}(\psi(U)) + d\Theta_s^2(\psi(U)) [d\psi(U) X_{G_L}(U) - X_{G_L}(\psi(U))].$$

Since  $d\Theta_s^2 \in \mathcal{L}(H^s, \mathbb{R})$ , it is enough, to show that the last term is  $O(\|U\|_{H^s}^{P+2})$ , to prove that

$$(4.2.27) \quad d\psi(U) X_{G_L}(U) - X_{G_L}(\psi(U)) = X_{G_L}(U) - X_{G_L}(\psi(U)) + (d\psi(U) - \text{Id}) \cdot X_{G_L}(U)$$

is in  $H^s$  and has  $H^s$ -norm  $O(\|U\|_{H^s}^{P+1})$ . Since  $G_L$  may be written  $\sum_{p=1}^{P-1} G_{L,p}$ , with  $G_{L,p}$  in  $\bigcap_N \mathcal{G}_p^{1-N, \nu+N}(B_s(R)) \subset \mathcal{G}_p^{0, \nu+1}(B_s(R))$ , we may write

$$G_{L,p} = \text{Re} \int_X (A_{L,p}(U)u) \bar{u} d\mu + \text{Re} \int_X (B_{L,p}u)u d\mu$$

with  $A_{L,p}, B_{L,p}$  in  $\mathcal{P}_p^{0, \nu+1}$ . If we use (4.2.11), (4.2.12) applied to such operators (see the statement following these formulas), and the fact that  $\|\psi(U) - U\|_{H^s} = O(\|U\|_{H^s}^{P+1})$ ,  $\|d\psi(U) - \text{Id}\|_{\mathcal{L}(H^{\pm s}, H^{\pm s})} = O(\|U\|_{H^s}^P)$ , we get the wanted conclusion for (4.2.26).

Let us study the contribution of  $G_H + \tilde{G}_P$  to (4.2.25). We must prove that

$$d\Theta_s^2(\psi(U)) \cdot d\psi(U) \cdot X_{[G_H + \tilde{G}_P]}(U) - d\Theta_s^2(\psi(U)) \cdot X_{G_H}(\psi(U)) = O(\|U\|_{H^s}^{P+2}).$$

According to Lemma 4.2.2, and to the boundedness of  $d\Theta_s^2$  (resp.  $d\psi$ ) from  $H^s$  to  $\mathbb{R}$  (resp. to  $H^s$ ), this will follow from the estimate

$$(4.2.28) \quad d\Theta_s^2(\psi(U)) [X_{\tilde{G}}(\psi(U)) - X_{G_H}(\psi(U))] = O(\|U\|_{H^s}^{P+2}), \quad U \rightarrow 0.$$

Using expressions (4.2.3), (4.2.13) of  $G_H, \tilde{G}$ , and setting  $\Sigma_H = {}^t JM_H$  we write

$$(4.2.29) \quad \tilde{G}(V) - G_H(V) = \frac{1}{2} \int_X (J[\Sigma(\psi^{-1}(V)) - \Sigma_H(V)]V) V d\mu$$

and

$$\begin{aligned}
(4.2.30) \quad &\Sigma(\psi^{-1}(V)) - \Sigma_H(V) \\
&= \begin{bmatrix} A(\psi^{-1}(V)) - A_H(V) + C(\psi^{-1}(V)) & 0 \\ 0 & -({}^t A(\psi^{-1}(V)) - {}^t A_H(V) + C(\psi^{-1}(V))) \end{bmatrix}
\end{aligned}$$

By definition of  $A_H$  (which is the sum of contributions to  $A$  homogeneous of order up to  $P-1$ ),  $A - A_H$  is in  $\Psi_P^{1, \nu}(B_s(R))$ . Moreover,  $C$  is in  $\Psi_P^{1, \nu}(B_s(R))$ . Since

we have seen that  $\|d\psi^{-1}(V) - \text{Id}\|_{\mathcal{L}(H^{s'}, H^{s'})}$  is  $O(\|V\|_{H^s}^P)$  for any  $s'$  with  $|s'| \leq s$ , we deduce that  $A_H(V) - A_H(\psi^{-1}(V))$  is in  $\Psi_P^{1,\nu}(B_s(R))$  and that  $C(\psi^{-1}(V))$  is in  $\Psi_P^{1,\nu}(B_s(R))$ . Consequently, (4.2.29), (4.2.30) show that  $\widehat{G} - G_H$  belongs to  $\mathcal{F}_{P,H}^{1,\nu}(B_s(R))$ . The left hand side of (4.2.28) may be written  $\{\Theta_s^2, \widehat{G} - G_H\}(\psi(U))$ . It follows from Lemma 3.2.6, and the assumption that  $\Theta_s^2$  is a sum of elements of  $H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R)) + \mathcal{G}_p^{2s-1,\nu}(B_s(R))$  for some  $\nu$  and  $0 \leq p \leq P-1$ , that this last term is  $O(\|U\|_{H^s}^{P+2})$  as wanted.  $\square$

**4.3. Elimination of hyperbolic terms.** This section is devoted to the proof of Theorem 3.1.3. We prove first several helpful lemmas.

Remember that we denoted by  $(\lambda_n^2)_{n \in \mathbb{N}^*}$  the eigenvalues of  $-\Delta$  on  $X = \mathbb{S}^d$  and that the mass parameter  $\mathbf{m}$  has been chosen so that (1.2.27) holds. We set  $\tilde{\Lambda} = \sqrt{-\Delta + (\frac{d-1}{2})^2}$  so that  $\tilde{\Lambda}\Pi_n = (n-1 + \frac{d-1}{2})\Pi_n$  for  $n \in \mathbb{N}^*$ . In particular,  $t \rightarrow e^{it\tilde{\Lambda}}$  is a  $4\pi$ -periodic function. Moreover,  $\Lambda_{\mathbf{m}} - \tilde{\Lambda}$  is a pseudo-differential operator of order  $-1$ .

**LEMMA 4.3.1.** *Let  $\nu \in \mathbb{R}_+$ ,  $p \in \mathbb{N}^*$ ,  $\ell \in \{0, \dots, p\}$  and  $A'$  be an element of the space  $\widetilde{H\Psi}_p^{1,\nu}[M_0]$  introduced in Definition 1.2.2. Assume that  $A'$  satisfies the following condition (using the notations introduced before the statement of Theorem 1.2.8):*

$$(4.3.1) \quad (n_0, \dots, n_{p+1}) \in Z^\ell(p) \Rightarrow \Pi_{n_0} A' (\Pi_{n'} \mathcal{U}) \Pi_{n_{p+1}} = 0.$$

Denote  $M_1 = 2M_0(2d+3)$ ,  $\nu_1 = 2M_1$ . There is a family  $(C_\alpha(\mathcal{U}))_{\alpha \in \mathbb{Z}}$  of linear operators from  $C^\infty(X)$  to  $\mathcal{D}'(X)$ , such that, for any  $t \in [0, 4\pi]$

$$(4.3.2) \quad e^{-it\tilde{\Lambda}} A'(\mathcal{U}) e^{it\tilde{\Lambda}} = \sum_{\alpha \in \mathbb{Z}} e^{i\alpha t/2} C_\alpha(\mathcal{U})$$

and, for any family of differential operators  $P_1, \dots, P_k$  of orders  $d_1, \dots, d_k$ , and  $N_1, N_2$  in  $\mathbb{N}$ , there is a constant  $C > 0$  such that, for any  $\alpha \in \mathbb{Z}$ , any  $j, j'$  in  $\mathbb{N}$ , any  $n'$  in  $(\mathbb{N}^*)^p$ , any  $u_1, \dots, u_p$  in  $C^\infty(X)$

$$(4.3.3) \quad \|\Delta_j \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} C_\alpha(\Pi_{n'} \mathcal{U}) \Delta_{j'}\|_{\mathcal{L}(L^2)} \leq \frac{C}{1+\alpha^2} 2^{-N_1|j-j'|} 2^{j(1+\sum d_\ell - k)} |n'|^{\nu(p)+\nu+\nu_1+M_1k} \\ \times (1+2^{-j}|n'|)^{-N_2} \prod_{\ell=1}^p \|\Pi_{n_\ell} u_\ell\|_{L^2}.$$

Moreover  $C_0(\Pi_{n'} \mathcal{U}) \equiv 0$  if  $n'$  is in  $\underline{Z}^\ell(p) \subset (\mathbb{N}^*)^p$ . Equation (4.3.3) implies in particular that  $\sum_\alpha C_\alpha$  defines an element of  $\widetilde{H\Psi}_p^{1,\nu+\nu_1}[M_1]$ .

*Proof:* Using the  $4\pi$ -time periodicity of the left hand side of (4.3.2), we write its Fourier series decomposition, with coefficients

$$C_\alpha(\mathcal{U}) = \frac{1}{4\pi} \int_0^{4\pi} e^{-it\tilde{\Lambda}} A'(\mathcal{U}) e^{it\tilde{\Lambda}} e^{-it\alpha/2} dt.$$

Denote by  $P_{k'}(t) = e^{it\tilde{\Lambda}} P_{k'} e^{-it\tilde{\Lambda}}$ ,  $1 \leq k' \leq k$ . By the Egorov theorem, this is a family of pseudo-differential operators of order  $d_k$ , with uniform estimates for

$t \in [0, 4\pi]$ . We may write, using two integrations by parts in  $t$ ,

$$(4.3.4) \quad \alpha^2 \text{Ad}_{P_1} \cdots \text{Ad}_{P_k} C_\alpha(\mathcal{U}) = \frac{1}{\pi} \int_0^{4\pi} e^{-it\tilde{\Lambda}} \text{Ad}_{P_1(t)} \cdots \text{Ad}_{P_k(t)} \text{Ad}_{\tilde{\Lambda}}^2 A'(\mathcal{U}) e^{it\tilde{\Lambda}} e^{-it\alpha/2} dt.$$

By (iii) of Proposition 2.2.6, each commutator with  $P_{k'}(t)$  (resp.  $\tilde{\Lambda}$ ) makes gain  $d_{k'} - 1$  (resp. 0) units on the order of the operator, and makes lose  $M_1 = 2M_0(2d+3)$  on the second index of this order. It follows that (4.3.3) holds, with the preceding value of  $M_1$  and with  $\nu_1 = 2M_1$ .

To get the statement concerning  $C_0$ , we compute

$$\Pi_{n_0} C_0(\Pi_{n'} \mathcal{U}) \Pi_{n_{p+1}} = \frac{1}{4\pi} \int_0^{4\pi} e^{-it(n_0 - n_{p+1})} \Pi_{n_0} A'(\Pi_{n'} \mathcal{U}) \Pi_{n_{p+1}} dt.$$

This quantity vanishes if  $n_0 \neq n_{p+1}$ . If  $n_0 = n_{p+1}$ , the condition  $n' \in \underline{Z}^\ell(p)$  is equivalent to  $(n_0, n', n_{p+1}) \in Z^\ell(p)$ , so that assumption (4.3.1) implies that the integrand vanishes. This concludes the proof.  $\square$

LEMMA 4.3.2. *Let  $\mu \in \mathbb{R}$ ,  $\nu \in \mathbb{R}_+$ ,  $P \in \mathbb{N}$ ,  $p \in \mathbb{N}$ ,  $1 \leq p \leq P-1$ ,  $\ell \in \{0, \dots, p\}$ ,  $N \in \mathbb{N}$ . There are positive numbers  $\bar{\nu}(N)$ ,  $M_0(N) = M_0(4d+6)^N$  such that the following holds:*

(i) *For any  $A'$  in  $\widetilde{H\Psi}_p^{1,\nu}[M_0]$  satisfying (4.3.1), one may find operators  $B$  (resp.  $S$ , resp.  $R$ ) belonging to  $\widetilde{H\Psi}_p^{1,\nu+\bar{\nu}(N)}[M_0(N)]$  (resp.  $\widetilde{H\Psi}_p^{1-N,\nu+\bar{\nu}(N)}[M_0(N)]$ ,  $\widetilde{H\Psi}_p^{1,\nu+\bar{\nu}(N)}[M_0(N)]$ ), such that*

$$(4.3.5) \quad [B(\mathcal{U}), \Lambda_{\mathbf{m}}] + \sum_{j=1}^{\ell} B(u_1, \dots, \Lambda_{\mathbf{m}} u_j, \dots, u_p) - \sum_{j=\ell+1}^p B(u_1, \dots, \Lambda_{\mathbf{m}} u_j, \dots, u_p) = A'(\mathcal{U}) + S(\mathcal{U}) + \epsilon^P R(\mathcal{U}).$$

Moreover,  $S$  satisfies (4.3.1).

(ii) *Let  $A'$  be an element of  $\widetilde{\mathcal{P}}_p^{\mu,\nu}$ , satisfying (4.3.1). There is  $B'$  in  $\widetilde{\mathcal{P}}_p^{\mu+L_0,\nu}$  such that*

$$(4.3.6) \quad [B(\mathcal{U}), \Lambda_{\mathbf{m}}] + \sum_{j=1}^{\ell} B(u_1, \dots, \Lambda_{\mathbf{m}} u_j, \dots, u_p) - \sum_{j=\ell+1}^p B(u_1, \dots, \Lambda_{\mathbf{m}} u_j, \dots, u_p) = A'(\mathcal{U}).$$

The same conclusion holds for the equations

$$(4.3.7) \quad \pm [\Lambda_{\mathbf{m}} B(\mathcal{U}) + B(\mathcal{U}) \Lambda_{\mathbf{m}}] + \sum_{j=1}^{\ell} B(u_1, \dots, \Lambda_{\mathbf{m}} u_j, \dots, u_p) - \sum_{j=\ell+1}^p B(u_1, \dots, \Lambda_{\mathbf{m}} u_j, \dots, u_p) = A'(\mathcal{U}).$$

Proof: (i) Take  $\theta \in C_0^\infty(\mathbb{R})$ ,  $\theta \equiv 1$  close to zero, and define

$$(4.3.8) \quad B_1(\Pi_{n'} \mathcal{U}) = -i \int_0^{+\infty} e^{-it\tilde{\Lambda}} A'(\Pi_{n'} \mathcal{U}) e^{it\tilde{\Lambda}} e^{itG_{\mathbf{m}}^{p,\ell}(n')} \theta(\epsilon t) dt$$

where we denoted for short by  $G_{\mathbf{m}}^{p,\ell}(n')$  the function  $G_{\mathbf{m}}^{p,\ell}(\lambda_{n_1}, \dots, \lambda_{n_p})$  defined in (1.2.26). Using expansion (4.3.2), we write

$$B_1(\Pi_n \mathcal{U}) = -i \sum_{\alpha \in \mathbb{Z}} a_\alpha(n', \epsilon) C_\alpha(\Pi_n \mathcal{U})$$

with

$$a_\alpha(n', \epsilon) = \int_0^{+\infty} e^{it\left(\frac{\alpha}{2} + G_{\mathbf{m}}^{p,\ell}(n')\right)} \theta(\epsilon t) dt.$$

By (1.2.27), we know that  $\left|\frac{\alpha}{2} + G_{\mathbf{m}}^{p,\ell}(n')\right| \geq c|n'|^{-L_0}$  unless  $\alpha = 0$  and  $n' \in \underline{Z}^\ell(p)$ . This last case is excluded since, by assumption and Lemma 4.3.1,  $C_0(\Pi_n \mathcal{U}) = 0$  when  $n' \in \underline{Z}^\ell(p)$ . Consequently, an integration by parts shows that  $|a_\alpha(n', \epsilon)| \leq C|n'|^{L_0}$ . Combining this with (4.3.3) and summing in  $\alpha$ , we see that  $B_1$  satisfies the estimates of elements of  $\widetilde{H\Psi}_p^{1, \nu + \nu_1 + L_0}[M_1]$ , where  $M_1 = 2M_0(2d + 3)$ . Let us compute

$$\begin{aligned} (4.3.9) \quad & -[\tilde{\Lambda}, B_1(\Pi_n \mathcal{U})] + G_{\mathbf{m}}^{p,\ell}(n') B_1(\Pi_n \mathcal{U}) \\ &= - \int_0^{+\infty} \frac{d}{dt} \left[ e^{-it\tilde{\Lambda}} A'(\Pi_n \mathcal{U}) e^{it\tilde{\Lambda}} e^{itG_{\mathbf{m}}^{p,\ell}(n')} \right] \theta(\epsilon t) dt \\ &= A'(\Pi_n \mathcal{U}) + \epsilon^P R_1(\Pi_n \mathcal{U}) \end{aligned}$$

with

$$R_1(\Pi_n \mathcal{U}) = \int_0^{+\infty} e^{-it\tilde{\Lambda}} A'(\Pi_n \mathcal{U}) e^{it\tilde{\Lambda}} e^{itG_{\mathbf{m}}^{p,\ell}(n')} \epsilon^{-P+1} \theta'(\epsilon t) dt.$$

Let  $\theta_1 \in C_0^\infty([0, +\infty[)$  such that  $\theta_1 \equiv \theta'$  on  $[0, +\infty[$ . Using again decomposition (4.3.2), we write

$$R_1(\Pi_n \mathcal{U}) = \sum_{\alpha \in \mathbb{Z}} C_\alpha(\Pi_n \mathcal{U}) \epsilon^{-P} \hat{\theta}_1 \left( -\epsilon^{-1} \left( G_{\mathbf{m}}^{p,\ell}(n') + \frac{\alpha}{2} \right) \right).$$

By (1.2.27),  $\epsilon^{-P} \left| \hat{\theta}_1 \left( -\epsilon^{-1} \left( G_{\mathbf{m}}^{p,\ell}(n') + \frac{\alpha}{2} \right) \right) \right| \leq C|n'|^{PL_0}$  so that (4.3.3) shows that  $R_1$  belongs to the space  $\widetilde{H\Psi}_p^{1, \nu + \nu_1 + PL_0}[M_1]$ . We deduce from (4.3.9) that

$$\begin{aligned} & -[\Lambda_{\mathbf{m}}, B_1(\mathcal{U})] + \sum_{j=1}^{\ell} B_1(u_1, \dots, \Lambda_{\mathbf{m}} u_j, \dots, u_p) - \sum_{j=\ell+1}^{p+1} B_1(u_1, \dots, \Lambda_{\mathbf{m}} u_j, \dots, u_p) \\ &= A'(\mathcal{U}) + \epsilon^P R_1(\mathcal{U}) - [\Lambda_{\mathbf{m}} - \tilde{\Lambda}, B_1(\mathcal{U})]. \end{aligned}$$

Since  $\Lambda_{\mathbf{m}} - \tilde{\Lambda}$  is of order  $-1$ ,  $[\Lambda_{\mathbf{m}} - \tilde{\Lambda}, B_1(\mathcal{U})]$  is in  $\widetilde{H\Psi}_p^{0, \nu + \nu_1 + L_0}[M_1]$ , by Lemma 1.2.4, and satisfies condition (4.3.1), because  $\Pi_n$  commutes to  $\Lambda_{\mathbf{m}} - \tilde{\Lambda}$ , and because  $B_1$  satisfies (4.3.1), as follows from its expression (4.3.8) and the assumption on  $A'$ . We see that we have solved an approximate version of (4.3.5), where  $S$  is replaced by an operator  $[\tilde{\Lambda} - \Lambda_{\mathbf{m}}, B_1(\mathcal{U})]$  of order 0 instead of  $1 - N$ . Since condition (4.3.1) is satisfied by this operator, we may repeat the construction of  $B_1$ , defining successively  $B_2, B_3, \dots$  until  $B = B_1 + B_2 + \dots$  satisfies (4.3.5) with an error  $S$  of the wanted order  $1 - N$ . At each step, we have to increase the number of derivative losses on small frequencies  $\nu$ , and multiply the constant  $M_1$  of estimate (4.3.3) by  $2(2d + 3)$ . We call  $\bar{\nu}(N)$  the total loss at the end of the process, and get the wanted conclusion.

(ii) We compute (4.3.6) at  $\Pi_n \mathcal{U}$ , and compose at the left with  $\Pi_{n_{p+1}}$  and at the right with  $\Pi_{n_0}$ . We get

$$(4.3.10) \quad F_{\mathbf{m}}^{p,\ell}(\lambda_{n_0}, \dots, \lambda_{n_{p+1}}) \Pi_{n_{p+1}} B(\Pi_n \mathcal{U}) \Pi_{n_0} = \Pi_{n_{p+1}} A'(\Pi_n \mathcal{U}) \Pi_{n_0}.$$

Since  $A'$  satisfies condition (4.3.1),  $|F_{\mathbf{m}}^{p,\ell}(\lambda_{n_0}, \dots, \lambda_{n_{p+1}})|$  is bounded from below by the right hand side of the first estimate (1.2.27). Going back to the estimate (1.2.5) defining  $\mathcal{P}_p^{\mu,\nu}$ , we see that we may find a solution  $B$  to (4.3.10) in  $\mathcal{P}_p^{\mu+L_0,\nu}$ . This concludes the proof since the same reasoning applies to (4.3.7).  $\square$

Before proving Theorem 3.1.3, we recall some notations and results of [16]. Let  $F = \sum_{p=1}^{P-1} F_p$ ,  $H = \sum_{p=0}^{P-1} H_p$  be two functions defined on a neighborhood of zero in  $H^s(X)$ , which are sums of components homogeneous of order  $p+2$ . Assume also that  $F, H$  have enough smoothness so that all Poisson brackets below are meaningful. We define a truncated Poisson bracket at order  $P$  by

$$(4.3.11) \quad \{F, H\}_P = \sum_{\substack{p+p' \leq P-1 \\ p \geq 1, p' \geq 0}} \{F_p, H_{p'}\}.$$

The iterates of  $\{\cdot, \cdot\}_P$  are defined by

$$(4.3.12) \quad \begin{aligned} \text{ad}_P F \cdot H &= \{F, H\}_P \\ \text{ad}_P^j F \cdot H &= \text{ad}_P F \cdot (\text{ad}_P^{j-1} F \cdot H). \end{aligned}$$

If  $T$  is some indeterminate, we define

$$(4.3.13) \quad \exp(T \text{ad}_P F) \cdot H = \sum_{j=0}^{+\infty} \frac{T^j}{j!} \text{ad}_P^j F \cdot H.$$

Notice that the sum in the last series is finite, as all terms with  $j \geq P$  vanish. Let  $\Theta$  be another multi-linear expression in  $U = (u, \bar{u})$ . It is proved in Lemma 5.1.3 of [16] that

$$(4.3.14) \quad \{\exp(T \text{ad}_P F) \cdot \Theta, H\}_P = \exp(T \text{ad}_P F) \cdot \{\Theta, \exp(-T \text{ad}_P F) \cdot H\}_P.$$

We define from  $\Theta$  (resp.  $H$ ) and from  $F$  new functions, denoted formally by  $\Theta \circ \chi_F^P$  (resp.  $H \circ (\chi_F^P)^{-1}$ ), given by

$$(4.3.15) \quad \begin{aligned} \Theta \circ \chi_F^P &= (\exp(\text{ad}_P F)) \cdot \Theta \\ H \circ (\chi_F^P)^{-1} &= (\exp(-\text{ad}_P F)) \cdot H \end{aligned}$$

so that (4.3.14) may be written, taking  $T = 1$ ,

$$(4.3.16) \quad \{\Theta \circ \chi_F^P, H\}_P = \{\Theta, H \circ (\chi_F^P)^{-1}\}_P \circ \chi_F^P.$$

The main remaining steps in the proof of Theorem 3.1.3 will be to apply (4.3.16) with  $\Theta$  replaced by  $\Theta_s^0(u, \bar{u}) = \int_X (\Lambda_{\mathbf{m}}^{2s} u) \bar{u} d\mu$ ,  $H$  replaced by the Hamiltonian  $G_L + G_H$  found in Theorem 3.1.2, and to construct a function  $F$  so that  $\{\Theta_s^0, (G_L + G_H) \circ (\chi_F^P)^{-1}\}_P \circ \chi_F^P = O(\epsilon^P \|u\|_{H^s}^2)$  (for  $s$  large enough).

Let us fix some notations. We fix  $P \in \mathbb{N}^*$  and  $N$  large enough relatively to  $P$ . We shall need several constants indexed by  $p = 1, \dots, P-1$ , growing quickly enough. We take elements of  $[1, +\infty[$ ,  $(M_i(p))_{1 \leq p \leq P-1}$ ,  $(\tilde{\nu}_i(p))_{1 \leq p \leq P-1}$ ,  $i = 1, \dots, 4$ . We

assume for  $p, q \in \{1, \dots, P-1\}$  such that  $p+q \leq P-1$ , and for some large  $C_0$ , depending on  $P, N$ ,

$$(4.3.17) \quad \begin{aligned} M_1(p) &\geq C_0 M_2(p) \geq C_0^2 M_3(p) \geq C_0^3 M_4(p) \geq C_0^3 M_0 \\ M_4(p+q) &\geq \max(M_1(p), M_2(q)) \\ \min(\tilde{\nu}_1(p), \tilde{\nu}_2(q)) + C_0[M_1(p) + M_1(q)] &\leq \tilde{\nu}_4(p+q) \\ \tilde{\nu}_j(p) &\geq \tilde{\nu}_{j+1}(p) + C_0 M_{j+1}(p), \quad j = 1, 2, 3, \end{aligned}$$

where  $M_0$  is the constant introduced in Definition 1.2.2. We assume moreover that, if  $p \rightarrow \nu(p)$  is the function of the definition of classes  $\widehat{H\Psi}$ , it satisfies, for  $p, q$  in  $\{1, \dots, P-1\}$ ,

$$(4.3.18) \quad \nu(p) + \nu(q) + \nu + \max(\tilde{\nu}_1(p), \tilde{\nu}_2(q)) + C_0 \max(M_1(p), M_2(q)) \leq \tilde{\nu}_4(p+q).$$

Finally, we take two other functions  $p \rightarrow N_i(p)$ ,  $i = 1, 2$ , such that  $N_1(1) + M_0 = N_2(1) = N$  and that, for any  $p, q \in \{1, \dots, P-1\}$  with  $p+q \leq P-1$ ,

$$(4.3.19) \quad \begin{aligned} N_2(p+q) &\leq N_2(q) - M_0 - 1, \quad N_2(p+q) \leq N_1(p) - M_0 - 1, \quad N_1(p) \leq N_2(p) - L_0 \end{aligned}$$

(where  $L_0$  is the exponent in (1.2.27)). If  $N = N_2(1) \gg P$ , we may find such  $N_i(p)$ 's, which are moreover larger or equal to  $P$  for  $p = 1, \dots, P-1$ . We may assume as well that  $\tilde{\nu}_4(p) \geq N_2(p)$  for any  $p = 1, \dots, P-1$ , since it is always possible to increase the  $\tilde{\nu}_j(p)$  in such a way they will continue to satisfy (4.3.17).

We use inclusions (3.2.12), (3.2.14) and (3.2.13) (with the constant  $M_0$  of (3.2.10) replaced by  $M_1(p)$  or  $M_2(q)$ ) to get, taking into account assumptions (4.3.17), (4.3.18), (4.3.19), the inclusion

$$(4.3.20) \quad \begin{aligned} &\{H\mathcal{F}_{p,H}^{1,\nu+\tilde{\nu}_1(p)}(B_s(R))[M_1(p)] + \mathcal{G}_p^{1-N_1(p),\nu+\tilde{\nu}_1(p)}(B_s(R)), \\ &\quad H\mathcal{F}_{q,H}^{1,\nu+\tilde{\nu}_2(q)}(B_s(R))[M_2(q)] + \mathcal{G}_q^{1-N_2(q),\nu+\tilde{\nu}_2(q)}(B_s(R))\} \\ &\subset H\mathcal{F}_{p+q,H}^{1,\nu+\tilde{\nu}_4(p+q)}(B_s(R))[M_4(p+q)] + \mathcal{G}_{p+q}^{1-N_2(p+q)-M_0,\nu+\tilde{\nu}_4(p+q)}(B_s(R)) \end{aligned}$$

for any  $p > 0, q > 0$  such that  $p, q, p+q$  are smaller than  $P-1$ . Let us prove:

LEMMA 4.3.3. *For any  $p = 1, \dots, P-1$ , there is an element*

$$(4.3.21) \quad F_p \in H\mathcal{F}_{p,H}^{1,\nu+\tilde{\nu}_1(p)}(B_s(R))[M_1(p)] + \mathcal{G}_p^{1-N_1(p),\nu+\tilde{\nu}_1(p)}(B_s(R))$$

satisfying

$$(4.3.22) \quad \{F_p, G_0\} \in H\mathcal{F}_{p,H}^{1,\nu+\tilde{\nu}_2(p)}(B_s(R))[M_2(p)] + \mathcal{G}_p^{1-N_2(p),\nu+\tilde{\nu}_2(p)}(B_s(R))$$

such that, if  $F = \sum_{p=1}^{P-1} F_p$

$$(4.3.23) \quad \{\Theta_s^0, (G_L + G_H) \circ (\chi_F^P)^{-1}\}_P = \epsilon^P \{\Theta_s^0, R\},$$

with  $R$  in  $\bigoplus_{p=1}^{P-1} H\mathcal{F}_{p,H}^{1,\nu+\tilde{\nu}_2(p)}(B_s(R))[M_2(p)]$ .

*Proof:* By the definition (3.1.5), (3.1.4) of  $G_L, G_H$ , we may write  $G_L + G_H = G_0 + \sum_{p=1}^{P-1} (G_{L,p} + G_{H,p})$  and  $G_{L,p} + G_{H,p} \in H\mathcal{F}_{p,H}^{1,\nu}(B_s(R))[M_0] + \bigcap_N \mathcal{G}_p^{1-N,\nu+N}(B_s(R))$  for some  $\nu$ . In particular,  $G_{L,p} + G_{H,p}$  belongs to

$$(4.3.24) \quad H\mathcal{F}_{p,H}^{1,\nu+\tilde{\nu}_4(p)}(B_s(R))[M_4(p)] + \mathcal{G}_p^{1-N_2(p),\nu+\tilde{\nu}_4(p)}(B_s(R)).$$

We compute  $(G_L + G_H) \circ (\chi_F^P)^{-1}$ . By (4.3.15), (4.3.13), this may be written

$$G_0 - \sum_{p=1}^{P-1} \{F_p, G_0\} + \sum_{p=1}^{P-1} H_p$$

where  $H_p$  is a linear combination of quantities

$$(4.3.25) \quad \{F_{p_1}, \{F_{p_2}, \dots, \{F_{p_\gamma}, G_{L,p_{\gamma+1}} + G_{H,p_{\gamma+1}}\} \dots\}\}$$

where  $p_1 + \dots + p_{\gamma+1} = p$ ,  $p_\beta \geq 1$  if  $1 \leq \beta \leq \gamma$ ,  $p_{\gamma+1} \geq 0$  (with by convention the notation  $G_{L,0} + G_{H,0} = G_0$ ) and where  $\gamma \geq 2$  if  $p_{\gamma+1} = 0$ . Since  $G_{L,p_{\gamma+1}} + G_{H,p_{\gamma+1}}$  is in (4.3.24) with  $p = p_{\gamma+1}$  and  $F_{p_\gamma}$  satisfies (4.3.21) with  $p = p_\gamma$ , successive applications of inclusion (4.3.20) show that (4.3.25) belongs to (4.3.24) when  $p_{\gamma+1} > 0$ . If  $p_{\gamma+1} = 0$ , we use that  $\gamma \geq 2$ , so that we make use of (4.3.22) to reach the same conclusion. Consequently,  $H_p$  belongs to (4.3.24), and the lemma will be proved if we construct  $F_p$ , satisfying (4.3.21) and (4.3.22) and such that

$$(4.3.26) \quad \{\Theta_s^0, H_p - \{F_p, G_0\}\} = \epsilon^P \{\Theta_s^0, R_p\}$$

with  $R_p$  in  $H\mathcal{F}_{p,H}^{1,\nu+\tilde{\nu}_2(p)}(B_s(R))[M_2(p)]$ . We decompose  $H_p = H_{p,\mathcal{F}} + H_{p,\mathcal{G}}$  according to (4.3.24) and write

$$(4.3.27) \quad \begin{aligned} H_{p,\mathcal{F}}(U) &= \operatorname{Re} \int_X (C_p(U)u)\bar{u} d\mu \\ H_{p,\mathcal{G}} &= \operatorname{Re} \int_X (E_{p,0}(U)u)\bar{u} d\mu + \operatorname{Re} \int_X (E_{p,1}(U)u)u d\mu, \end{aligned}$$

with  $C_p$  in  $H\Psi_p^{1,\nu+\tilde{\nu}_4(p)}[M_4(p)]$ ,  $E_{p,0}, E_{p,1}$  in  $\mathcal{P}_p^{1-N_2(p),\nu+\tilde{\nu}_4(p)} \subset \mathcal{P}_p^{1-N_2(p),\nu+\tilde{\nu}_3(p)}$ . Coming back to the definition of these spaces, we decompose

$$(4.3.28) \quad \begin{aligned} C_p(U) &= \sum_{\ell=0}^p C_p^\ell(u, \dots, u, \bar{u}, \dots, \bar{u}) \\ E_{p,i}(U) &= \sum_{\ell=0}^p E_{p,i}^\ell(u, \dots, u, \bar{u}, \dots, \bar{u}), i = 0, 1 \end{aligned}$$

with  $C_p^\ell$  in  $\widetilde{H\Psi}_p^{1,\nu+\tilde{\nu}_4(p)}[M_4(p)]$ ,  $E_{p,i}^\ell$  in  $\widetilde{\mathcal{P}}_p^{1-N_2(p),\nu+\tilde{\nu}_3(p)}$ . We may assume that, for a fixed large enough  $C$ ,

$$(4.3.29) \quad \Pi_{n_0} C_p^\ell(\Pi_{n'} \mathcal{U}) \Pi_{n_{p+1}} \neq 0 \Rightarrow n_0, n_{p+1} \geq C|n'|.$$

Actually, to ensure that condition, we may replace  $C_p^\ell$  by

$$(4.3.30) \quad \sum_{n'} \sum_j \sum_{j'} \mathbb{1}_{\{2^j \geq C|n'|, 2^{j'} \geq C|n'|\}} \Delta_j C_p^\ell(\Pi_{n'} \mathcal{U}) \Delta_{j'}$$

since, using formula (A.1) of the appendix, one checks that (4.3.30) is an element of the space  $\widetilde{H\Psi}_p^{1,\nu+\tilde{\nu}_4(p)}[M_4(p)]$  satisfying (4.3.29). The difference between  $C_p^\ell$  and (4.3.30) may be incorporated to  $E_{p,0}$ : taking into account that this difference belongs to

$$\widetilde{H\Psi}_p^{1-N',\nu+\tilde{\nu}_4(p)+N'}[M_4(p)] \subset \widetilde{\mathcal{P}}_p^{1-N',\nu+\tilde{\nu}_4(p)+N'+2M_4(p)}$$



for any  $N'$ , we take  $N' = N_2(p)$ , and obtain that the perturbation of  $E_{p,0}$  will be in the class  $\widetilde{\mathcal{P}}_p^{1-N_2(p), \nu + \tilde{\nu}_3(p)}$  (using the last inequality (4.3.17) with a large enough  $C_0$ ).

Let us refine the decomposition of  $C_p^\ell, E_{p,i}^\ell$ . When  $\ell \neq p/2$ , we set  $E_{p,0}^{\ell,Z} = 0$ . When  $\ell = p/2$ , we define

$$(4.3.31) \quad E_{p,0}^{\ell,Z} = \sum_{\substack{\{n_0, \dots, n_\ell\} = \\ \{n_{\ell+1}, \dots, n_{p+1}\}}} \Pi_{n_0} E_{p,0}^\ell (\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}.$$

In the same way, if  $\ell + 1 \neq p/2$ , we set  $E_{p,1}^{\ell,Z} = 0$ . If  $\ell + 1 = p/2$ , we set

$$E_{p,1}^{\ell,Z} = \sum_{\substack{\{n_0, \dots, n_\ell, n_{p+1}\} = \\ \{n_{\ell+1}, \dots, n_p\}}} \Pi_{n_0} E_{p,1}^\ell (\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}}.$$

Then  $E_{p,i}^{\ell,Z}$  are elements of  $\widetilde{\mathcal{P}}_p^{1-N_2(p), \nu + \tilde{\nu}_3(p)}$ . We define  $E_{p,i}^{\ell,B} = E_{p,i}^\ell - E_{p,i}^{\ell,Z}$ . In that way,  $E_{p,0}^{\ell,B}$  satisfies (4.3.1) and  $E_{p,1}^{\ell,B}$  satisfies

$$(4.3.32) \quad \{n_0, \dots, n_\ell, n_{p+1}\} = \{n_{\ell+1}, \dots, n_p\} \Rightarrow \Pi_{n_0} E_{p,1}^{\ell,B} (\Pi_{n_1} u_1, \dots, \Pi_{n_p} u_p) \Pi_{n_{p+1}} = 0.$$

We perform a similar decomposition of  $C_p^\ell = C_p^{\ell,Z} + C_p^{\ell,B}$  using formula (4.3.31).

Let us show that  $C_p^{\ell,Z}$  belongs to  $\widetilde{H\Psi}_p^{1, \nu + \tilde{\nu}_3(p)}[M_3(p)]$ . Actually, because of condition (4.3.29), the property  $\{n_0, \dots, n_\ell\} = \{n_{\ell+1}, \dots, n_{p+1}\}$  in the sum of type (4.3.31) defining  $C_p^{\ell,Z}$  from  $C_p^\ell$ , is equivalent to  $(n_0 = n_{p+1})$  and  $\{n_1, \dots, n_\ell\} = \{n_{\ell+1}, \dots, n_p\}$ . The definition of  $C_p^{\ell,Z}$  may be written equivalently as

$$C_p^{\ell,Z}(u_1, \dots, u_p) = \sum_{\substack{\{n_1, \dots, n_\ell\} = \\ \{n_{\ell+1}, \dots, n_p\}}} \frac{1}{4\pi} \int_0^{4\pi} e^{-it\tilde{\Lambda}} C_p^\ell(\Pi_n \mathcal{U}) e^{it\tilde{\Lambda}} dt.$$

We have seen in the proof of Lemma 4.3.1 that this time average belongs to  $\widetilde{H\Psi}_p^{1, \nu + \tilde{\nu}_3(p)}[M_3(p)]$  (This follows from the fact that by (4.3.17),  $M_3(p) \geq C_0 M_4(p)$ ,  $\tilde{\nu}_3(p) \geq \tilde{\nu}_4(p) + C_0 M_4(p)$ , which are the requirement that the new constants  $(\nu_1, M_1)$  in the right hand side of (4.3.3) have to satisfy). Set

$$\begin{aligned} C_p^Z(U) &= \sum_{\ell=0}^p C_p^{\ell,Z}(\underbrace{u, \dots, u}_\ell, \bar{u}, \dots, \bar{u}), \quad E_{p,i}^Z(U) = \sum_{\ell=0}^p E_{p,i}^{\ell,Z}(\underbrace{u, \dots, u}_\ell, \bar{u}, \dots, \bar{u}) \\ C_p^B(U) &= (C_p - C_p^Z)(U), \quad E_{p,i}^B(U) = E_{p,i}(U) - E_{p,i}^Z(U) \\ H_{p,\mathcal{F}}^Z(U) &= \text{Re} \int_X (C_p^Z(U)u)\bar{u} d\mu, \quad H_{p,\mathcal{G}}^Z(U) = \text{Re} \int_X (E_{p,0}^Z(U)u)\bar{u} d\mu + \text{Re} \int_X (E_{p,1}^Z(U)u)u d\mu \end{aligned}$$

and define similarly  $H_{p,\mathcal{F}}^B(U), H_{p,\mathcal{G}}^B(U)$ . By construction and the definition of the Sobolev energy  $\Theta_s^0(u, \bar{u}) = \int_X (\Lambda_{\mathbf{m}}^{2s} u)\bar{u} d\mu$ ,  $\{\Theta_s^0, H_{p,\mathcal{F}}^Z + H_{p,\mathcal{G}}^Z\} = 0$ , so that equation (4.3.26) is equivalent to

$$(4.3.33) \quad \{\Theta_s^0, H_{p,\mathcal{F}}^B + H_{p,\mathcal{G}}^B - \{F_p, G_0\}\} = \epsilon^P \{\Theta_s^0, R_p\}.$$

Let us show that there is an element  $R_p$  in  $H\mathcal{F}_{p,H}^{1, \nu + \tilde{\nu}_2(p)}(B_s(R))[M_2(p)]$  and  $F_p$  satisfying (4.3.21), (4.3.22) such that

$$(4.3.34) \quad \{F_p, G_0\} = H_{p,\mathcal{F}}^B + H_{p,\mathcal{G}}^B - \epsilon^P R_p.$$

We look for  $F_p$  as

(4.3.35)

$$F_p(U) = \operatorname{Re} \int_X (B_p(U)u)\bar{u} d\mu + \operatorname{Re} \int_X (D_{p,0}(U)u)\bar{u} d\mu + \operatorname{Re} \int_X (D_{p,1}(U)u)u d\mu$$

where

$$B_p(U) = \sum_{\ell=0}^p B_p^\ell(u, \dots, u, \bar{u}, \dots, \bar{u}), D_{p,i}(U) = \sum_{\ell=0}^p D_{p,i}^\ell(u, \dots, u, \bar{u}, \dots, \bar{u}),$$

with  $B_p^\ell$  in  $\widetilde{H\Psi}_p^{1,\nu+\bar{\nu}_1(p)}[M_1(p)]$ ,  $D_{p,i}^\ell$  in  $\widetilde{\mathcal{P}}_p^{1-N_1(p),\nu+\bar{\nu}_1(p)}$ . Plugging (4.3.35) inside (4.3.34), we see that  $B_p^\ell$  may be computed from  $C_p^{\ell,B}$  solving an equation of type (4.3.5). Applying (i) of Lemma 4.3.2, we obtain  $B_p^\ell$  in  $\widetilde{H\Psi}_p^{1,\nu+\bar{\nu}_1(p)}[M_1(p)]$ , using (4.3.17), and assuming that  $C_0$  is larger than the loss  $\bar{\nu}(N)$  of Lemma 4.3.2. We get the wanted property for  $B_p^\ell$ .

The remainder  $S$  in the right hand side of (4.3.5) belongs to

$$\widetilde{H\Psi}_p^{1-N,\nu+\bar{\nu}_3(p)+\bar{\nu}(N)}[M_3(p)] \subset \widetilde{\mathcal{P}}_p^{1-N,\nu+\bar{\nu}_3(p)+\bar{\nu}(N)+2M_3(p)}.$$

Since  $N \geq N_2(p)$ , we get an element of  $\widetilde{\mathcal{P}}_p^{1-N_2(p),\nu+\bar{\nu}_2(p)}$ , if the constant  $C_0$  in (4.3.17) has been taken large enough with respect to  $N$ . The contributions  $E_{p,i}^{\ell,B}$  belong to the same space. By (ii) of Lemma 4.3.2, we may construct  $D_{p,0}$  (resp.  $D_{p,1}$ ) in  $\widetilde{\mathcal{P}}_p^{1-N_2(p)+L_0,\nu+\bar{\nu}_2(p)}$  solving (4.3.6) (resp. (4.3.7)) when the right hand side  $A'$  is given by  $E_{p,0}^{\ell,B} + S$  (resp  $E_{p,1}^{\ell,B}$ ). If we assume  $N_1(p) \leq N_2(p) - L_0$  as in (4.3.19), we get the wanted conclusion.

Finally, the term  $\epsilon^P R$  in the right hand side of (4.3.5) belongs to  $\epsilon^P \widetilde{H\Psi}_p^{1,\nu+\bar{\nu}_3(p)+\bar{\nu}(N)}[M_2(p)] \subset \epsilon^P \widetilde{H\Psi}_p^{1,\nu+\bar{\nu}_2(p)}[M_2(p)]$ . The contribution of this term to an integral of the form of the first one in (4.3.27) gives the last term in (4.3.34).  $\square$

*Proof of Theorem 3.1.3:* We set  $\Theta_s^2 = \Theta_s^0 \circ \chi_F^P$ . This is linear combination of quantities

$$(4.3.36) \quad \{F_{p_1}, \{F_{p_2}, \dots, \{F_{p_\ell}, \Theta_s^0\} \dots\}\}, \ell \leq P-1,$$

with  $F_p$  satisfying (4.3.21),  $1 \leq p \leq P-1$ . Since we assume  $N_1(p) \geq P$ , we have

$$(4.3.37) \quad F_p \in H\mathcal{F}_{p,H}^{1,\nu}(B_s(R)) + \mathcal{G}_p^{1-P,\nu}(B_s(R))$$

for some large enough value of  $\nu$ . Moreover,  $\Theta_s^0$  is in  $H\mathcal{F}_{0,H}^{2s,0}(B_s(R))$ . If  $s \gg P$ , it follows from inclusions (3.2.12), (3.2.13) and (3.2.14) that (4.3.36) is in  $H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R)) + \mathcal{G}_p^{2s+\ell-P,\nu}(B_s(R))$  for  $p = p_1 + \dots + p_\ell$  and a new value of  $\nu$ , independent of  $s$ . Consequently  $\Theta_s^2$  is in

$$(4.3.38) \quad \bigoplus_{p=0}^{P-1} (H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R)) + \mathcal{G}_p^{2s-1,\nu}(B_s(R)))$$

for some  $\nu$ . By (4.3.16)

(4.3.39)

$$\begin{aligned} \{\Theta_s^2, G_L + G_H\} &= \{\Theta_s^2, G_L + G_H\}_P + (\{\Theta_s^2, G_L + G_H\} - \{\Theta_s^2, G_L + G_H\}_P) \\ &= \{\Theta_s^0, (G_L + G_H) \circ (\chi_F^P)^{-1}\}_P \circ \chi_F^P + (\{\Theta_s^2, G_L + G_H\} - \{\Theta_s^2, G_L + G_H\}_P). \end{aligned}$$

The first term in the right hand side may be written according to (4.3.23) as  $\epsilon^P \{\Theta_s^0, R\} \circ \chi_F^P$  with  $R$  in  $\bigoplus_{p=1}^{P-1} H\mathcal{F}_{p,H}^{1,\nu+\bar{\nu}_2(p)}(B_s(R))$ . Since  $\Theta_s^0$  is in  $H\mathcal{F}_{0,H}^{2s,0}(B_s(R))$ , it follows from (3.2.12) that  $\{\Theta_s^0, R\}$  is in  $\bigoplus_{p=1}^{P-1} H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R))$  for some  $\nu$  independent of  $s$ . Using (4.3.37) and again inclusions (3.2.12), (3.2.14), (3.2.13), we conclude that  $\{\Theta_s^0, R\} \circ \chi_F^P$  belongs to  $\bigoplus_{p=0}^{P-1} (H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R)) + \mathcal{G}_p^{2s,\nu}(B_s(R)))$ . This implies that  $\{\Theta_s^0, R\} \circ \chi_F^P(U) = O(\|U\|_{H^s}^2)$  if  $s$  is large enough.

The last term in (4.3.39) may be written as a sum of expressions

$$(4.3.40) \quad \{\theta_p, G_{L,q} + G_{H,q}\}$$

where  $\theta_p$  is in  $H\mathcal{F}_{p,H}^{2s,\nu}(B_s(R)) + \mathcal{G}_p^{2s-1,\nu}(B_s(R))$ , and where  $G_{L,q} + G_{H,q}$  belongs to the space  $H\mathcal{F}_{q,H}^{1,\nu'}(B_s(R)) + \mathcal{G}_q^{0,\nu'}(B_s(R))$  for some  $\nu, \nu'$  depending on  $P$  but not on  $s$ , and with  $p+q \geq P$ . By (3.2.12), (3.2.13), (3.2.14), we obtain elements of  $H\mathcal{F}_{p+q,H}^{2s,\nu''}(B_s(R)) + \mathcal{G}_{p+q}^{2s,\nu''}(B_s(R))$  for some new  $\nu''$  independent of  $s$ . This implies that (4.3.40) is  $O(\|U\|_{H^s}^{P+2})$ , which is the wanted conclusion.  $\square$

## A. Appendix

This appendix is devoted to an estimate that is used several times in the paper.

PROPOSITION A.1. *Let  $P_1, \dots, P_k$  be differential operators of order  $d_1, \dots, d_k$  on the manifold  $X$ . Let  $Q, Q'$  be two pseudo-differential operators of order  $m, m'$ . For any  $N$  in  $\mathbb{N}$ , there is  $C > 0$  and for any  $j, j', j''$  in  $\mathbb{N}$*

$$(A.1) \quad \|\Delta_{j'}(\text{Ad}_{P_1} \cdots \text{Ad}_{P_k}(Q\Delta_j Q'))\Delta_{j''}\|_{\mathcal{L}(L^2)} \leq C 2^{-N[|j-j'|+|j-j''|]} 2^j (m+m'+\sum_{\ell=1}^k d_\ell - k).$$

If we prove

$$(A.2) \quad \|\Delta_{j'} \text{Ad}_{P_1} \cdots \text{Ad}_{P_k}(Q\Delta_j Q')\|_{\mathcal{L}(L^2)} \leq C 2^{-N|j-j'|} 2^j (m+m'+\sum_{\ell=1}^k d_\ell - k)$$

we may deduce (A.1) from (A.2) by duality. To prove (A.2), we introduce, with the notations (1.2.2),  $\varphi_1(t) = \varphi(\sqrt{t})$ ,  $\psi_1(t) = \psi(\sqrt{t})$ , so that  $\Delta_j = \varphi_1(-2^{-2j}\Delta)$ , for  $j \geq 1$ ,  $\Delta_0 = \psi_1(-\Delta)$ . We use the same letters  $\varphi_1, \psi_1$  to denote almost holomorphic extensions of this functions to  $\mathbb{C}$  (see Dimassi-Sjöstrand [20], Chapter 8). Then  $\varphi_1, \psi_1$  are smooth functions on  $\mathbb{C}$ ,  $\varphi_1$  (resp.  $\psi_1$ ) is supported in a domain  $[a, b] + i[-c, c]$  with  $0 < a < b$ ,  $c > 0$  (resp.  $a < 0 < b$ ,  $c > 0$ ), and  $\bar{\partial}\varphi_1, \bar{\partial}\psi_1$  vanish at infinite order on  $\mathbb{R}$ . The Helffer-Sjöstrand formula [26], [20] Theorem 8.1 implies that

$$(A.3) \quad \Delta_j = \frac{-1}{\pi} 2^{-2j} \int_{\mathbb{C}} (\bar{\partial}\varphi_1)(2^{-2j}\zeta)(\zeta + \Delta)^{-1} d\zeta d\bar{\zeta}$$

when  $j \geq 1$ , and a similar expression for  $\Delta_0$ , with  $\varphi_1$  replaced by  $\psi_1$ . We shall prove (A.2) when, more generally,  $P_1, \dots, P_k$  are pseudo-differential operators of order  $d_1, \dots, d_k$ . Writing  $[P_j, (\zeta + \Delta)^{-1}] = -(\zeta + \Delta)^{-1}[P_j, \Delta](\zeta + \Delta)^{-1}$ , we see that  $\text{Ad}_{P_1} \cdots \text{Ad}_{P_k} \Delta_j$  may be written as a linear combination of expressions

$$(A.4) \quad 2^{-2j} \int_{\mathbb{C}} (\bar{\partial}\varphi_1)(2^{-2j}\zeta) \left[ \prod_{\ell'=1}^{k'} (\zeta + \Delta)^{-1} Q_{\ell'} \right] (\zeta + \Delta)^{-1} d\zeta d\bar{\zeta}$$

where  $0 \leq k' \leq k$ ,  $Q_{\ell'}$ ,  $\ell' = 1, \dots, k'$  are pseudo-differential operators of order  $D_{\ell'} + 2$  and  $\sum_1^{k'+1} D_{\ell'} = \sum_1^k d_\ell - k$ .

LEMMA A.2. *For any  $s \in \mathbb{R}$*

$$(A.5) \quad \sup_j \|\text{Ad}_{P_1} \cdots \text{Ad}_{P_k} \Delta_j\|_{\mathcal{L}(H^s, H^{s-\sum_1^k d_\ell + k})} < +\infty.$$

*Proof:* We note that for any  $\sigma \in \mathbb{R}$

$$\|(\zeta + \Delta)^{-1}\|_{\mathcal{L}(H^\sigma, H^{\sigma+2})} \leq \frac{C(1+|\zeta|)}{|\text{Im } \zeta|}, \quad \|(\zeta + \Delta)^{-1}\|_{\mathcal{L}(H^\sigma, H^\sigma)} \leq C|\text{Im } \zeta|^{-1}.$$

We deduce from (A.4), the above inequalities, the fact that on the support of  $\varphi_1(2^{-2j}\zeta)$ ,  $|\zeta| \leq C2^{2j}$  and the vanishing of  $\bar{\partial}\varphi_1$  at infinite order on  $\mathbb{R}$ , that (A.5) is bounded from above by the supremum in  $j$  of quantities of the form

$$C2^{-2j} \int_{\mathbb{C}} |\bar{\partial}\varphi_1(2^{-2j}\zeta)| \frac{2^{2jk'}}{|\operatorname{Im} \zeta|^{k'+1}} d\zeta d\bar{\zeta} \leq C2^{2j(k'-1-N)} \int_{|\zeta| \leq C2^{2j}} |\operatorname{Im} \zeta|^{N-k'-1} d\zeta d\bar{\zeta} \leq C$$

for  $0 \leq k' \leq k$ . This gives the lemma.  $\square$

*Proof of Proposition A.1:* For  $k \in \mathbb{N}$ ,  $a \in \mathbb{R}$ , denote by  $\mathcal{E}_k(a)$  the family of operators which may be written as a finite sum of terms  $Q_0(\operatorname{Ad}_{Q_1} \cdots \operatorname{Ad}_{Q_{k'}} \Delta_j) Q_{k'+1}$  where  $k' \leq k$  and  $Q_\ell$ ,  $\ell = 0, \dots, k' + 1$  are pseudo-differential operators such that  $\sum_{\ell=0}^{k'+1} \deg Q_\ell - k' = a$ . Let  $Q, Q'$  be a pseudo-differential operator of order  $m, m'$ . Then

(A.6)

$$\operatorname{Ad}_{P_1} \cdots \operatorname{Ad}_{P_k} [Q \Delta_j Q'] = Q(\operatorname{Ad}_{P_1} \cdots \operatorname{Ad}_{P_k} \Delta_j) Q' \text{ modulo } \mathcal{E}_{k-1}(m + \sum_1^k d_\ell - k + m').$$

Actually, if (A.6) holds at order  $k$ , and if we consider another operator  $P_0$  of order  $d_0$

$$\begin{aligned} \operatorname{Ad}_{P_0} \operatorname{Ad}_{P_1} \cdots \operatorname{Ad}_{P_k} [Q \Delta_j Q'] &= Q(\operatorname{Ad}_{P_0} \cdots \operatorname{Ad}_{P_k} \Delta_j) Q' + [P_0, Q](\operatorname{Ad}_{P_1} \cdots \operatorname{Ad}_{P_k} \Delta_j) Q' \\ &\quad + Q(\operatorname{Ad}_{P_1} \cdots \operatorname{Ad}_{P_k} \Delta_j) [P_0, Q'] \text{ modulo } [P_0, \mathcal{E}_{k-1}(m + \sum_1^k d_\ell - k + m')]. \end{aligned}$$

We have just to notice that the second and third term in the right hand side are in  $\mathcal{E}_k(m + \sum_0^k d_\ell - (k+1) + m')$ , as well as the elements of  $[P_0, \mathcal{E}_{k-1}(m + \sum_1^k d_\ell - k + m')]$ .

Let us prove estimate (A.2). Assume first  $j' > j > 0$  and write

$$(A.7) \quad \Delta_{j'} \operatorname{Ad}_{P_1} \cdots \operatorname{Ad}_{P_k} (Q \Delta_j Q') = 2^{2jN} \Delta_{j'} \operatorname{Ad}_{P_1} \cdots \operatorname{Ad}_{P_k} (\tilde{Q} \tilde{\Delta}_j Q')$$

where  $\tilde{Q} = Q \Delta^{-N}$  and  $\tilde{\Delta}_j = \tilde{\varphi}(-2^{-2j} \Delta)$  for a new  $\tilde{\varphi} \in C_0^\infty([0, +\infty[)$ . By (A.6), we may write (A.7) as the product of  $2^{2jN}$  times operators of the form  $\Delta_{j'} Q_0(\operatorname{Ad}_{Q_1} \cdots \operatorname{Ad}_{Q_{k'}} \tilde{\Delta}_j) Q_{k'+1}$ , where  $k' \leq k$  and  $\sum_0^{k'+1} \deg Q_\ell - k' = -2N + m + \sum_1^k d_\ell + m' - k$ . By Lemma A.2

$$\begin{aligned} \|\Delta_{j'} Q_0(\operatorname{Ad}_{Q_1} \cdots \operatorname{Ad}_{Q_{k'}} \tilde{\Delta}_j) Q_{k'+1} u\|_{L^2} &\leq C2^{-j'\sigma} \|Q_0(\operatorname{Ad}_{Q_1} \cdots \operatorname{Ad}_{Q_{k'}} \tilde{\Delta}_j) Q_{k'+1} u\|_{H^\sigma} \\ &\leq C2^{-j'\sigma} \|u\|_{H^{\sigma-2N+m+\sum_1^k d_\ell+m'-k}}. \end{aligned}$$

We take  $\sigma = -m - \sum_1^k d_\ell - m' + 2N + k$  to conclude that

$$\|\Delta_{j'} \operatorname{Ad}_{P_1} \cdots \operatorname{Ad}_{P_k} (Q \Delta_j Q')\|_{\mathcal{L}(L^2)} \leq C_N 2^{2(j-j')N+j'(m+\sum_1^k d_\ell+m'-k)}$$

for any  $N$ . When  $j < j'$ , we perform the same computation taking  $\tilde{Q} = Q \Delta^N$ . This concludes the proof.  $\square$

## Bibliography

- [1] D. Bambusi, *Birkhoff normal form for some nonlinear PDEs*, Comm. Math. Phys. **234** (2003), no. 2, 253–285, DOI 10.1007/s00220-002-0774-4. MR1962462 (2003k:37121)
- [2] D. Bambusi, J.-M. Delort, B. Grébert, and J. Szeftel, *Almost global existence for Hamiltonian semilinear Klein-Gordon equations with small Cauchy data on Zoll manifolds*, Comm. Pure Appl. Math. **60** (2007), no. 11, 1665–1690, DOI 10.1002/cpa.20181. MR2349351 (2009b:35279)
- [3] D. Bambusi and B. Grébert, *Birkhoff normal form for partial differential equations with tame modulus*, Duke Math. J. **135** (2006), no. 3, 507–567, DOI 10.1215/S0012-7094-06-13534-2. MR2272975 (2007j:37124)
- [4] R. Beals, *Characterization of pseudodifferential operators and applications*, Duke Math. J. **44** (1977), no. 1, 45–57. MR0435933 (55 #8884)
- [5] P. Baldi, M. Berti and R. Montalto: *KAM for quasi-linear and fully nonlinear forced perturbations of Airy equation*, Math. Ann. **359** (2014), no. 1-2, 471–536. MR3201904
- [6] M. Berti, L. Biasco and L. Procesi: *KAM theory for the Hamiltonian wave equation*, Ann. Sci. Éc. Norm. Supér. (4) **46** (2013), no. 2, 301–373 (2013). MR3112201
- [7] M. Berti, L. Biasco and L. Procesi: *KAM for reversible derivative wave equation*, Arch. Ration. Mech. Anal. **212** (2014), no. 3, 905–955. MR3187681
- [8] M. Berti and P. Bolle, *Sobolev quasi-periodic solutions of multidimensional wave equations with a multiplicative potential*, Nonlinearity **25** (2012), no. 9, 2579–2613, DOI 10.1088/0951-7715/25/9/2579. MR2967117
- [9] M. Berti and M. Procesi, *Nonlinear wave and Schrödinger equations on compact Lie groups and homogeneous spaces*, Duke Math. J. **159** (2011), no. 3, 479–538, DOI 10.1215/00127094-1433403. MR2831876 (2012j:37109)
- [10] J. Bourgain, *Construction of quasi-periodic solutions for Hamiltonian perturbations of linear equations and applications to nonlinear PDE*, Internat. Math. Res. Notices **11** (1994), 475ff., approx. 21 pp. (electronic), DOI 10.1155/S1073792894000516. MR1316975 (96f:58170)
- [11] J. Bourgain, *Construction of periodic solutions of nonlinear wave equations in higher dimension*, Geom. Funct. Anal. **5** (1995), no. 4, 629–639, DOI 10.1007/BF01902055. MR1345016 (96h:35011)
- [12] J. Bourgain, *Construction of approximative and almost periodic solutions of perturbed linear Schrödinger and wave equations*, Geom. Funct. Anal. **6** (1996), no. 2, 201–230, DOI 10.1007/BF02247885. MR1384610 (97f:35013)
- [13] J. Bourgain, *Green’s function estimates for lattice Schrödinger operators and applications*, Annals of Mathematics Studies, vol. 158, Princeton University Press, Princeton, NJ, 2005. MR2100420 (2005j:35184)
- [14] W. Craig and E. Wayne, *Newton’s method and periodic solutions of nonlinear wave equations*, Comm. Pure Appl. Math. **46** (1993), no. 11, 1409–1498, DOI 10.1002/cpa.3160461102. MR1239318 (94m:35023)
- [15] J.-M. Delort, *On long time existence for small solutions of semi-linear Klein-Gordon equations on the torus*, J. Anal. Math. **107** (2009), 161–194, DOI 10.1007/s11854-009-0007-2. MR2496403 (2010i:35247)
- [16] J.-M. Delort, *A quasi-linear Birkhoff normal forms method. Application to the quasi-linear Klein-Gordon equation on  $\mathbb{S}^1$*  (English, with English and French summaries), Astérisque **341** (2012), vi+113. MR2952065
- [17] J.-M. Delort and J. Szeftel, *Long-time existence for small data nonlinear Klein-Gordon equations on tori and spheres*, Int. Math. Res. Not. **37** (2004), 1897–1966, DOI 10.1155/S1073792804133321. MR2056326 (2005k:35281)

- [18] J.-M. Delort and J. Szeftel, *Long-time existence for semi-linear Klein-Gordon equations with small Cauchy data on Zoll manifolds*, Amer. J. Math. **128** (2006), no. 5, 1187–1218. MR2262173 (2007i:58039)
- [19] J.-M. Delort and J. Szeftel, *Bounded almost global solutions for non Hamiltonian semi-linear Klein-Gordon equations with radial data on compact revolution hypersurfaces* (English, with English and French summaries), Ann. Inst. Fourier (Grenoble) **56** (2006), no. 5, 1419–1456. MR2273861 (2008c:35191)
- [20] M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series, vol. 268, Cambridge University Press, Cambridge, 1999. MR1735654 (2001b:35237)
- [21] D. Fang and Q. Zhang, *Long-time existence for semi-linear Klein-Gordon equations on tori*, J. Differential Equations **249** (2010), no. 1, 151–179, DOI 10.1016/j.jde.2010.03.025. MR2644131 (2011f:35224)
- [22] P. Germain, *Global existence for coupled Klein-Gordon equations with different speeds* (English, with English and French summaries), Ann. Inst. Fourier (Grenoble) **61** (2011), no. 6, 2463–2506 (2012), DOI 10.5802/aif.2680. MR2976318
- [23] P. Germain, N. Masmoudi, and J. Shatah, *Global solutions for 3D quadratic Schrödinger equations*, Int. Math. Res. Not. IMRN **3** (2009), 414–432, DOI 10.1093/imrn/rnn135. MR2482120 (2010b:35431)
- [24] P. Germain, N. Masmoudi, and J. Shatah, *Global solutions for 2D quadratic Schrödinger equations* (English, with English and French summaries), J. Math. Pures Appl. (9) **97** (2012), no. 5, 505–543, DOI 10.1016/j.matpur.2011.09.008. MR2914945
- [25] B. Grébert, *Birkhoff normal form and Hamiltonian PDEs* (English, with English and French summaries), Partial differential equations and applications, Sémin. Congr., vol. 15, Soc. Math. France, Paris, 2007, pp. 1–46. MR2352816 (2009d:37130)
- [26] B. Helffer and J. Sjöstrand, *Équation de Schrödinger avec champ magnétique et équation de Harper* (French), Schrödinger operators (Sønderborg, 1988), Lecture Notes in Phys., vol. 345, Springer, Berlin, 1989, pp. 118–197, DOI 10.1007/3-540-51783-9\_19. MR1037319 (91g:35078)
- [27] G. Iooss, P. I. Plotnikov, and J. F. Toland, *Standing waves on an infinitely deep perfect fluid under gravity*, Arch. Ration. Mech. Anal. **177** (2005), no. 3, 367–478, DOI 10.1007/s00205-005-0381-6. MR2187619 (2007a:76017)
- [28] T. Kappeler and J. Pöschel, *KdV & KAM*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 45, Springer-Verlag, Berlin, 2003. MR1997070 (2004g:37099)
- [29] S. Klainerman, *Global existence of small amplitude solutions to nonlinear Klein-Gordon equations in four space-time dimensions*, Comm. Pure Appl. Math. **38** (1985), no. 5, 631–641, DOI 10.1002/cpa.3160380512. MR803252 (87e:35080)
- [30] S. Kuksin, *A KAM-theorem for equations of the Korteweg-de Vries type*, Rev. Math. Math. Phys. **10** (1998), no. 3, ii+64. MR1754991 (2001g:37140)
- [31] S. Kuksin, *Nearly integrable infinite-dimensional Hamiltonian systems*, Lecture Notes in Mathematics, vol. 1556, Springer-Verlag, Berlin, 1993. MR1290785 (95k:58145)
- [32] J. Liu and X. Yuan, *A KAM theorem for Hamiltonian partial differential equations with unbounded perturbations*, Comm. Math. Phys. **307** (2011), no. 3, 629–673, DOI 10.1007/s00220-011-1353-3. MR2842962 (2012i:37124)
- [33] J. Shatah, *Normal forms and quadratic nonlinear Klein-Gordon equations*, Comm. Pure Appl. Math. **38** (1985), no. 5, 685–696, DOI 10.1002/cpa.3160380516. MR803256 (87b:35160)
- [34] E. Wayne, *Periodic and quasi-periodic solutions of nonlinear wave equations via KAM theory*, Comm. Math. Phys. **127** (1990), no. 3, 479–528. MR1040892 (91b:58236)
- [35] Q. Zhang, *Long-time existence for semi-linear Klein-Gordon equations with quadratic potential*, Comm. Partial Differential Equations **35** (2010), no. 4, 630–668, DOI 10.1080/03605300903509112. MR2753615 (2012c:35288)

### Editorial Information

To be published in the *Memoirs*, a paper must be correct, new, nontrivial, and significant. Further, it must be well written and of interest to a substantial number of mathematicians. Piecemeal results, such as an inconclusive step toward an unproved major theorem or a minor variation on a known result, are in general not acceptable for publication.

Papers appearing in *Memoirs* are generally at least 80 and not more than 200 published pages in length. Papers less than 80 or more than 200 published pages require the approval of the Managing Editor of the Transactions/Memoirs Editorial Board. Published pages are the same size as those generated in the style files provided for  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$  or  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$ .

Information on the backlog for this journal can be found on the AMS website starting from <http://www.ams.org/memo>.

A Consent to Publish is required before we can begin processing your paper. After a paper is accepted for publication, the Providence office will send a Consent to Publish and Copyright Agreement to all authors of the paper. By submitting a paper to the *Memoirs*, authors certify that the results have not been submitted to nor are they under consideration for publication by another journal, conference proceedings, or similar publication.

### Information for Authors

*Memoirs* is an author-prepared publication. Once formatted for print and on-line publication, articles will be published as is with the addition of AMS-prepared frontmatter and backmatter. Articles are not copyedited; however, confirmation copy will be sent to the authors.

**Initial submission.** The AMS uses Centralized Manuscript Processing for initial submissions. Authors should submit a PDF file using the Initial Manuscript Submission form found at [www.ams.org/submission/memo](http://www.ams.org/submission/memo), or send one copy of the manuscript to the following address: Centralized Manuscript Processing, MEMOIRS OF THE AMS, 201 Charles Street, Providence, RI 02904-2294 USA. If a paper copy is being forwarded to the AMS, indicate that it is for *Memoirs* and include the name of the corresponding author, contact information such as email address or mailing address, and the name of an appropriate Editor to review the paper (see the list of Editors below).

The paper must contain a *descriptive title* and an *abstract* that summarizes the article in language suitable for workers in the general field (algebra, analysis, etc.). The *descriptive title* should be short, but informative; useless or vague phrases such as “some remarks about” or “concerning” should be avoided. The *abstract* should be at least one complete sentence, and at most 300 words. Included with the footnotes to the paper should be the 2010 *Mathematics Subject Classification* representing the primary and secondary subjects of the article. The classifications are accessible from [www.ams.org/msc/](http://www.ams.org/msc/). The Mathematics Subject Classification footnote may be followed by a list of *key words and phrases* describing the subject matter of the article and taken from it. Journal abbreviations used in bibliographies are listed in the latest *Mathematical Reviews* annual index. The series abbreviations are also accessible from [www.ams.org/msnhtml/serials.pdf](http://www.ams.org/msnhtml/serials.pdf). To help in preparing and verifying references, the AMS offers MR Lookup, a Reference Tool for Linking, at [www.ams.org/mrlookup/](http://www.ams.org/mrlookup/).

**Electronically prepared manuscripts.** The AMS encourages electronically prepared manuscripts, with a strong preference for  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$ . To this end, the Society has prepared  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$  author packages for each AMS publication. Author packages include instructions for preparing electronic manuscripts, samples, and a style file that generates the particular design specifications of that publication series. Though  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$  is the highly preferred format of  $\mathcal{T}\mathcal{E}\mathcal{X}$ , author packages are also available in  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$ .

Authors may retrieve an author package for *Memoirs of the AMS* from [www.ams.org/journals/memo/memoauthorpac.html](http://www.ams.org/journals/memo/memoauthorpac.html) or via FTP to [ftp.ams.org](ftp://ftp.ams.org) (login as `anonymous`, enter your complete email address as password, and type `cd pub/author-info`). The



*AMS Author Handbook* and the *Instruction Manual* are available in PDF format from the author package link. The author package can also be obtained free of charge by sending email to [tech-support@ams.org](mailto:tech-support@ams.org) or from the Publication Division, American Mathematical Society, 201 Charles St., Providence, RI 02904-2294, USA. When requesting an author package, please specify  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$  or  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$  and the publication in which your paper will appear. Please be sure to include your complete mailing address.

**After acceptance.** The source files for the final version of the electronic manuscript should be sent to the Providence office immediately after the paper has been accepted for publication. The author should also submit a PDF of the final version of the paper to the editor, who will forward a copy to the Providence office.

Accepted electronically prepared files can be submitted via the web at [www.ams.org/submit-book-journal/](http://www.ams.org/submit-book-journal/), sent via FTP, or sent on CD to the Electronic Prepress Department, American Mathematical Society, 201 Charles Street, Providence, RI 02904-2294 USA.  $\mathcal{T}\mathcal{E}\mathcal{X}$  source files and graphic files can be transferred over the Internet by FTP to the Internet node [ftp.ams.org](ftp://ftp.ams.org) (130.44.1.100). When sending a manuscript electronically via CD, please be sure to include a message indicating that the paper is for the *Memoirs*.

**Electronic graphics.** Comprehensive instructions on preparing graphics are available at [www.ams.org/authors/journals.html](http://www.ams.org/authors/journals.html). A few of the major requirements are given here.

Submit files for graphics as EPS (Encapsulated PostScript) files. This includes graphics originated via a graphics application as well as scanned photographs or other computer-generated images. If this is not possible, TIFF files are acceptable as long as they can be opened in Adobe Photoshop or Illustrator.

Authors using graphics packages for the creation of electronic art should also avoid the use of any lines thinner than 0.5 points in width. Many graphics packages allow the user to specify a “hairline” for a very thin line. Hairlines often look acceptable when proofed on a typical laser printer. However, when produced on a high-resolution laser imagesetter, hairlines become nearly invisible and will be lost entirely in the final printing process.

Screens should be set to values between 15% and 85%. Screens which fall outside of this range are too light or too dark to print correctly. Variations of screens within a graphic should be no less than 10%.

Any graphics created in color will be rendered in grayscale for the printed version unless color printing is authorized by the Managing Editor and the Publisher. In general, color graphics will appear in color in the online version.

**Inquiries.** Any inquiries concerning a paper that has been accepted for publication should be sent to [memo-query@ams.org](mailto:memo-query@ams.org) or directly to the Electronic Prepress Department, American Mathematical Society, 201 Charles St., Providence, RI 02904-2294 USA.

## Editors

This journal is designed particularly for long research papers, normally at least 80 pages in length, and groups of cognate papers in pure and applied mathematics. Papers intended for publication in the *Memoirs* should be addressed to one of the following editors. The AMS uses Centralized Manuscript Processing for initial submissions to AMS journals. Authors should follow instructions listed on the Initial Submission page found at [www.ams.org/memo/memosubmit.html](http://www.ams.org/memo/memosubmit.html).

**Algebra**, to MICHAEL LARSEN, Department of Mathematics, Rawles Hall, Indiana University, 831 E 3rd Street, Bloomington, IN 47405, USA; e-mail: [mjlarsen@indiana.edu](mailto:mjlarsen@indiana.edu)

**Algebraic geometry**, to LUCIA CAPORASO, Department of Mathematics and Physics, Roma Tre University, Largo San Leonardo Murialdo, I-00146 Roma, Italy; e-mail: [caporaso@mat.uniroma3.it](mailto:caporaso@mat.uniroma3.it)

**Algebraic topology**, to SOREN GALATIUS, Department of Mathematics, Stanford University, Stanford, CA 94305 USA; e-mail: [transactions@lists.stanford.edu](mailto:transactions@lists.stanford.edu)

**Arithmetic geometry**, to TED CHINBURG, Department of Mathematics, University of Pennsylvania, Philadelphia, PA 19104-6395; e-mail: [math-tams@math.upenn.edu](mailto:math-tams@math.upenn.edu)

**Automorphic forms, representation theory and combinatorics**, to DANIEL BUMP, Department of Mathematics, Stanford University, Building 380, Sloan Hall, Stanford, California 94305; e-mail: [bump@math.stanford.edu](mailto:bump@math.stanford.edu)

**Combinatorics and discrete geometry**, to IGOR PAK, Department of Mathematics, University of California, Los Angeles, California 90095; e-mail: [pak@math.ucla.edu](mailto:pak@math.ucla.edu)

**Commutative and homological algebra**, to LUCHEZAR L. AVRAMOV, Department of Mathematics, University of Nebraska, Lincoln, NE 68588-0130; e-mail: [avramov@math.unl.edu](mailto:avramov@math.unl.edu)

**Differential geometry and global analysis**, to CHRIS WOODWARD, Department of Mathematics, Rutgers University, 110 Frelinghuysen Road, Piscataway, NJ 08854; e-mail: [ctw@math.rutgers.edu](mailto:ctw@math.rutgers.edu)

**Dynamical systems and ergodic theory and complex analysis**, to YUNPING JIANG, Department of Mathematics, CUNY Queens College and Graduate Center, 65-30 Kissena Blvd., Flushing, NY 11367; e-mail: [Yunping.Jiang@qc.cuny.edu](mailto:Yunping.Jiang@qc.cuny.edu)

**Ergodic theory and combinatorics**, to VITALY BERGELSON, Ohio State University, Department of Mathematics, 231 W. 18th Ave, Columbus, OH 43210; e-mail: [vitaly@math.ohio-state.edu](mailto:vitaly@math.ohio-state.edu)

**Functional analysis and operator algebras**, to NATHANIEL BROWN, Department of Mathematics, 320 McAllister Building, Penn State University, University Park, PA 16802; e-mail: [nbrown@math.psu.edu](mailto:nbrown@math.psu.edu)

**Geometric analysis**, to TATIANA TORO, Department of Mathematics, University of Washington, Box 354350; e-mail: [toro@uw.edu](mailto:toro@uw.edu)

**Geometric topology**, to MARK FEIGN, Math Department, Rutgers University, Newark, NJ 07102; e-mail: [feighn@andromeda.rutgers.edu](mailto:feighn@andromeda.rutgers.edu)

**Harmonic analysis, complex analysis**, to MALABIKA PRAMANIK, Department of Mathematics, 1984 Mathematics Road, University of British Columbia, Vancouver, BC, Canada V6T 1Z2; e-mail: [malabika@math.ubc.ca](mailto:malabika@math.ubc.ca)

**Harmonic analysis, representation theory, and Lie theory**, to E. P. VAN DEN BAN, Department of Mathematics, Utrecht University, P.O. Box 80 010, 3508 TA Utrecht, The Netherlands; e-mail: [E.P.vandenBan@uu.nl](mailto:E.P.vandenBan@uu.nl)

**Logic**, to ANTONIO MONTALBAN, Department of Mathematics, The University of California, Berkeley, Evans Hall #3840, Berkeley, California, CA 94720; e-mail: [antonio@math.berkeley.edu](mailto:antonio@math.berkeley.edu)

**Number theory**, to SHANKAR SEN, Department of Mathematics, 505 Malott Hall, Cornell University, Ithaca, NY 14853; e-mail: [ss70@cornell.edu](mailto:ss70@cornell.edu)

**Partial differential equations**, to MARKUS KEEL, School of Mathematics, University of Minnesota, Minneapolis, MN 55455; e-mail: [keel@math.umn.edu](mailto:keel@math.umn.edu)

**Partial differential equations and functional analysis**, to ALEXANDER KISELEV, Department of Mathematics, MS-136, Rice University, 6100 Main Street, Houston, TX 77005; e-mail: [kisilev@rice.edu](mailto:kisilev@rice.edu)

**Probability and statistics**, to PATRICK FITZSIMMONS, Department of Mathematics, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0112; e-mail: [pfitzsim@math.ucsd.edu](mailto:pfitzsim@math.ucsd.edu)

**Real analysis and partial differential equations**, to WILHELM SCHLAG, Department of Mathematics, The University of Chicago, 5734 South University Avenue, Chicago, IL 60615; e-mail: [schlag@math.uchicago.edu](mailto:schlag@math.uchicago.edu)

**All other communications to the editors**, should be addressed to the Managing Editor, ALEJANDRO ADEM, Department of Mathematics, The University of British Columbia, Room 121, 1984 Mathematics Road, Vancouver, B.C., Canada V6T 1Z2; e-mail: [adem@math.ubc.ca](mailto:adem@math.ubc.ca)

# SELECTED PUBLISHED TITLES IN THIS SERIES

- 1094 **Ian F. Putnam**, A Homology Theory for Smale Spaces, 2014
- 1093 **Ron Blei**, The Grothendieck Inequality Revisited, 2014
- 1092 **Yun Long, Asaf Nachmias, Weiyang Ning, and Yuval Peres**, A Power Law of Order  $1/4$  for Critical Mean Field Swendsen-Wang Dynamics, 2014
- 1091 **Vilmos Totik**, Polynomial Approximation on Polytopes, 2014
- 1090 **Ameya Pitale, Abhishek Saha, and Ralf Schmidt**, Transfer of Siegel Cusp Forms of Degree 2, 2014
- 1089 **Peter Šemrl**, The Optimal Version of Hua's Fundamental Theorem of Geometry of Rectangular Matrices, 2014
- 1088 **Mark Green, Phillip Griffiths, and Matt Kerr**, Special Values of Automorphic Cohomology Classes, 2014
- 1087 **Colin J. Bushnell and Guy Henniart**, To an Effective Local Langlands Correspondence, 2014
- 1086 **Stefan Ivanov, Ivan Minchev, and Dimitar Vassilev**, Quaternionic Contact Einstein Structures and the Quaternionic Contact Yamabe Problem, 2014
- 1085 **A. L. Carey, V. Gayral, A. Rennie, and F. A. Sukochev**, Index Theory for Locally Compact Noncommutative Geometries, 2014
- 1084 **Michael S. Weiss and Bruce E. Williams**, Automorphisms of Manifolds and Algebraic  $K$ -Theory: Part III, 2014
- 1083 **Jakob Wachsmuth and Stefan Teufel**, Effective Hamiltonians for Constrained Quantum Systems, 2014
- 1082 **Fabian Ziltener**, A Quantum Kirwan Map: Bubbling and Fredholm Theory for Symplectic Vortices over the Plane, 2014
- 1081 **Sy-David Friedman, Tapani Hyttinen, and Vadim Kulikov**, Generalized Descriptive Set Theory and Classification Theory, 2014
- 1080 **Vin de Silva, Joel W. Robbin, and Dietmar A. Salamon**, Combinatorial Floer Homology, 2014
- 1079 **Pascal Lambrechts and Ismar Volić**, Formality of the Little  $N$ -disks Operad, 2013
- 1078 **Milen Yakimov**, On the Spectra of Quantum Groups, 2013
- 1077 **Christopher P. Bendel, Daniel K. Nakano, Brian J. Parshall, and Cornelius Pillen**, Cohomology for Quantum Groups via the Geometry of the Nullcone, 2013
- 1076 **Jaeyoung Byeon and Kazunaga Tanaka**, Semiclassical Standing Waves with Clustering Peaks for Nonlinear Schrödinger Equations, 2013
- 1075 **Deguang Han, David R. Larson, Bei Liu, and Rui Liu**, Operator-Valued Measures, Dilations, and the Theory of Frames, 2013
- 1074 **David Dos Santos Ferreira and Wolfgang Staubach**, Global and Local Regularity of Fourier Integral Operators on Weighted and Unweighted Spaces, 2013
- 1073 **Hajime Koba**, Nonlinear Stability of Ekman Boundary Layers in Rotating Stratified Fluids, 2014
- 1072 **Victor Reiner, Franco Saliola, and Volkmar Welker**, Spectra of Symmetrized Shuffling Operators, 2014
- 1071 **Florin Diacu**, Relative Equilibria in the 3-Dimensional Curved  $n$ -Body Problem, 2014
- 1070 **Alejandro D. de Acosta and Peter Ney**, Large Deviations for Additive Functionals of Markov Chains, 2014
- 1069 **Ioan Bejenaru and Daniel Tataru**, Near Soliton Evolution for Equivariant Schrödinger Maps in Two Spatial Dimensions, 2014
- 1068 **Florica C. Cîrstea**, A Complete Classification of the Isolated Singularities for Nonlinear Elliptic Equations with Inverse Square Potentials, 2014

For a complete list of titles in this series, visit the  
AMS Bookstore at [www.ams.org/bookstore/memoseries/](http://www.ams.org/bookstore/memoseries/).

ISBN 978-1-4704-0983-8



9 781470 409838

MEMO/234/1103

AMS *on the Web*  
[www.ams.org](http://www.ams.org)