

Synthesis of 2,4,6-Tris(trinitromethyl)-1,3,5-triazine

Alexey V. Shastin,^a Tamara I. Godovikova,^b Svetlana P. Golova,^b Vladimir S. Kuz'min,^c Lenor I. Khmel'nitskii^{*b} and Boris L. Korsunskii^a

^aInstitute of Chemical Physics in Chernogolovka, Russian Academy of Sciences, 142432 Chernogolovka, Moscow Region, Russian Federation. Fax: +7 096 51 53588

^bN. D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 117913 Moscow, Russian Federation. Fax: +7 095 135 5328


^cN. S. Kurnakov Institute of General and Inorganic Chemistry, Russian Academy of Sciences, 117907 Moscow, Russian Federation. Fax: +7 095 954 1279

Destructive nitration of the carboxyl group to the trinitromethyl entity has been accomplished for the first time and 2,4,6-tris(trinitromethyl)-1,3,5-triazine, hitherto unknown, has been synthesized.

Destructive nitration of carboxyl and carbonyl derivatives is used as a method of polynitromethyl group formation in organic compounds. Thus, interaction between methyl malonate and nitric acid leads to methyl dinitroacetate,¹ and nitration of 1,1,1-trinitroalkylketones with a mixture of concentrated nitric and sulfuric acids affords hexanitroethane.² No compound containing the trinitromethyl group has, however, been synthesized from carboxyl derivatives.

To prepare 2,4,6-tris(trinitromethyl)-1,3,5-triazine **1**, the destructive nitration of 2,4,6-tris(dimethoxycarbonylmethylene)- and 2,4,6-tris(diethoxycarbonylmethylene)-1,3,5-hexahydrotriazines, **2** and **3** respectively, readily obtained from cyanuric chloride **4** and the pertinent malonic ester sodium salt,³ was initially investigated.

The hydrolysis and nitration of **2** and **3** were intended to be carried out simultaneously by the use of a mixture of sulfuric and nitric acids. It was found, however, that the dicarboalkoxymethylene groups in compounds **2** and **3** did not transform into trinitromethyl groups under reaction conditions, but instead underwent exclusively mononitration to give 2,4,6-tris[(dimethoxycarbonyl)nitromethyl]- and 2,4,6-tris[(diethoxycarbonyl)nitromethyl]-1,3,5-triazines, **5** and **6**, respectively, Scheme 1.




Scheme 1 Reagents and conditions: i, 1 mmol **2** or **3**, 1 ml HNO₃ (d 1.5), 2 ml H₂SO₄ (d 1.84), 3 ml CHCl₃, 0 °C, 40 min.

These results led us to the conclusion that an appropriate hexacarbonic acid **7**, not its esters, should be used to synthesize **1**.

The *tert*-butyl group is known to be removed by hydrolysis more readily than methyl and ethyl groups in esters and ethers.^{4,5} 2,4,6-Tris(*di-tert*-butoxycarbonylmethylene)-1,3,5-hexahydrotriazine **8** was prepared in 50% yield from **4** and *tert*-butyl malonate sodium salt according to the procedure described in ref. 3. The hydrolysis of **8** was performed by trifluoroacetic acid. It is impossible to establish in which particular form, **A** or **B**, the acid **7** exists because of its insolubility in solvents used for taking NMR spectra.

The desired product **1** was obtained by nitration of **7** with a mixture of sulfuric and nitric acids, Scheme 2.

The structure of **1** was substantiated by elemental analysis, IR, ¹³C and ¹⁴N NMR spectroscopy as well as some chemical transformations. Thus 2,4,6-trimethoxy-1,3,5-triazine **9** and nitroform potassium salt **10** were formed from **1** and KOH or MeOK nearly quantitatively, Scheme 3.



Scheme 2 Reagents and conditions: i, 1 mmol **8**, 5 ml CF₃COOH, 0–5 °C, 30 min, 7 filtered; ii, 1 mmol **7**, 4 ml H₂SO₄ (d 1.84), 3 ml HNO₃ (d 1.5), 0–5 °C, kept 12 h at 18 °C.

Compound **1** is unstable in the air, and so is preferably stored as a solution in absolute CCl₄ at 5–10 °C. When an undried dioxane is added to the solution of **1** in absolute CCl₄, one trinitromethyl group is subjected to nucleophilic substitution by hydroxyl on account of the water present in the dioxane. The 2-hydroxy-4,6-bis(trinitromethyl)-1,3,5-triazine **11** formed was isolated as a stable complex **12** with dioxane in the ratio 2:1 mol, Scheme 4.

The structure of the complex **12** was determined by X-ray analysis of a single crystal[†] (Fig. 1). The triazine ring is planar,

[†] Crystal data for **12**: C₅HN₉O₁₃·0.5C₄H₈O₂, triclinic, space group $\bar{P}\bar{1}$, $a = 8.119(2)$, $b = 10.570(2)$, $c = 10.713(2)$ Å, $\alpha = 75.64(2)$, $\beta = 73.73(2)$, $\gamma = 72.04(2)$, $U = 826.7(3)$ Å³, $D_c = 1.764$ g cm⁻³, $Z = 2$. The intensities of 2916 reflections (2060 observed): I_{hkl} were measured on a Syntex-P2₁ diffractometer [graphite monochromated Mo-K α radiation ($\lambda = 0.71069$ Å)] using the $\theta/2\theta$ scan technique ($2\theta \leq 50^\circ$). The structure was solved by direct methods and refined by the full-matrix least-squares method, anisotropically for the non-hydrogen atoms. The final R_F was 0.067 for $1631 F_{hkl} \geq 4\sigma$. Atomic coordinates, bond lengths and angles and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre, see Notice to Authors, *Mendeleev Commun.*, 1995, issue 1.

1 M.p. 90–91 °C, ¹³C NMR (CDCl₃): 164.525 (C=N) and 120.579 [C(NO₂)₃]; ¹⁴N NMR (CDCl₃): -43.334 (NO₂); IR, v/cm⁻¹ (KBr pellets): 1600, 1565, 1520, 1420, 1300, 1250, 1145, 1100, 920, 835, 805, 750. CAUTION! Explosion hazardous.


5 M.p. 90–91 °C; ¹³C NMR (CDCl₃): 55.156 (Me), 99.358 (C *tert*), 158.997 (C=O), 169.012 (C=N); ¹⁴N NMR (CDCl₃): -21.771 (NO₂).

6 M.p. 42–44 °C; ¹H NMR (CDCl₃): 1.23 t (18H, Me), 4.40 q (12H, CH₂); ¹³C NMR (CDCl₃): 13.60 (Me), 65.22 (CH₂), 99.85 (C *tert*), 158.67 (C=O), 169.26 (C=N); ¹⁴N NMR (CDCl₃): -20.984 (NO₂).

7 M.p. 120 °C (decomp.), unstable in storage; IR, v/cm⁻¹ (KBr pellets): 3400 (OH), 1600 (C=O), 1540 (C=C).

8 M.p. 203–205 °C (decomp.); ¹H NMR (CDCl₃): 1.500 s (54H, Bu³), 12.935 s (3H, NH); ¹³C NMR (CDCl₃): 166.951 (=CN), 145.670 (C=O), 85.738 (C=C), 80.985 (C=O), 26.163 (Me); ¹⁵N NMR (CDCl₃): -277.318 (C=NH), ¹J_{N-H} 95.76 Hz, ³J_{N-H} 2.40 Hz (NH form).

12 M.p. 150–152 °C (decomp.); ¹H NMR (CDCl₃): 3.62 s (8H, CH₂), 12.01 br.s (1H, OH); ¹³C NMR (CDCl₃): 170.985 s (=COH), 164.197 s (C=N), 121.415 s [C(NO₂)₃], 66.696 t (CH₂); ¹⁴N NMR (CDCl₃): -40.0613 (NO₂), $\gamma_{\frac{1}{2}} = 4.5$ Hz.



Scheme 3 Reagents and conditions: i, 1 mmol **1**, 3 mmol KOH or MeOK, 5 ml abs. MeOH, 18 °C, 2 h.

the trinitromethyl groups have a propeller-like conformation and are oriented identically relative to the triazine plane. The intramolecular distances N(1)...N(11) 2.770(7), N(2)...N(13) 2.645(7), N(2)...N(22) 2.738(7) and N(3)...N(23) 2.709(7) Å are shorter than the two-fold Van-der-Waals radius (3.00 Å). The dioxane molecule in the chair conformation occupies the centrosymmetric position in the crystal.





Fig. 1 Molecular structure of **12**. (The second molecule **11** has a position symmetric relative to dioxane). The main bond lengths/Å: C_{cycle}—N_{cycle} 1.302–1.346 [mean 1.325(8)], C_{cycle}—C_{sp³} 1.519, 1.527 [mean 1.523(9)], C_{sp³}—N_{nitro} 1.510–1.550 [mean 1.528(9)], N—O 1.110–1.210 [mean 1.17(1)], C(1)—O(1) 1.298(8). H-Bond parameters: O(1)...O(12) 2.502(8), O(1)...H 0.98, H...O(12) 1.55, angle O(1)–H...O(12) 169°.



Scheme 4 Reagents and conditions: i, 1 mmol **1** in 10 ml abs. CCl₄, 5 ml dioxane, 18 °C, 1 h.

All new compounds prepared have satisfactory elemental analysis data and have been characterised by ¹H, ¹³C and ^{14/15}N NMR spectroscopy.

References

- 1 V. Grakauskas and A. M. Guest, *J. Org. Chem.*, 1978, **43**, 3485.
- 2 E. L. Golod and L. I. Bagal, *Zh. Org. Khim.*, 1994, **30**, 29 (in Russian).
- 3 G. M. Vakhitova and L. N. Yakhontov, *Khim. Geterotsikl. Soedin.*, 1980, 554 [*Chem. Heterocycl. Compd. (Engl. Transl.)*, 1980, 433].
- 4 R. Schwyzer and H. Kappeler, *Helv. Chim. Acta*, 1961, **44**, 1991.
- 5 G. S. Fonken and W. S. Johnson, *J. Am. Chem. Soc.*, 1952, **74**, 831.

Received: Moscow, 29th June 1994

Cambridge, 31st October 1994; Com. 4/04002C