

An efficient synthesis of 2-trimethylsilyloxy-4-neopentyl quinoline via 3-lithio-1-*tert*-butyllallene and phenyl isocyanate


Nina A. Nedolya,^{*a} Lambert Brandsma^{*b} and Boris A. Trofimov^a

^a Irkutsk Institute of Organic Chemistry, Siberian Branch of the Russian Academy of Sciences, 664033 Irkutsk, Russian Federation. Fax: +7 3952 35 6046; e-mail: admin@irioch.irkutsk.su^b Department of Preparative Organic Chemistry, Debye Institute, University of Utrecht, 3584 CH Utrecht, The Netherlands. E-mail: l.brandsma@chem.ruu.nl

A new method for the preparation of 2-trimethylsilyloxy-4-neopentyl quinoline via lithiated *tert*-butyllallene and phenyl isocyanate has been developed.

The principal starting compounds in the most important methods for the synthesis of quinolines from non-heterocyclic precursors are benzene derivatives.¹ Direct formation of the quinoline ring via isocyanates and organometallic compounds remains unknown. Only a few examples of these reactions leading to amides have been described.^{2,3}

As a part of our investigation of the reactions of heterocumulenes with organometallic compounds, leading to important heterocycles, including pyrrole and dihydropyridine derivatives,^{4–6} we have developed an efficient synthesis of quinolines, using readily-available starting reagents. In particular, we have shown for the first time that 2,4-disubstituted quinolines can be easily obtained in high yield by the interaction of lithiated allenes and acetylenes with isocyanates and isothiocyanates in a one-pot procedure. Thus, hitherto almost inaccessible 2-trimethylsilyloxy-4-neopentyl quinoline **5** has been synthesised in 75% yield by the addition of lithiated *tert*-butyllallene **2** to phenyl isocyanate in a tetrahydrofuran (THF)–hexane solution followed by trimethylsilylation of intermediate **3** and intramolecular cyclization of azatriene **4**.



Intermediate 1,3,4-azatriene **4** was isolated in 95% yield. The structure of the compounds **4** and **5** has been confirmed by IR, ¹H and ¹³C NMR spectroscopy and elemental analysis data.[†]

More complete results of this study will be given in the next paper.

We are grateful to R.-J. de Lang for spectroscopic measurements.

References

- 1 Obshchaya organicheskaya khimiya. T. 8. Azotsoderzhashchie geterotsikly, Khimiya, Moscow, 1985, pp. 196–255 (in Russian) (Comprehensive Organic Chemistry. The Synthesis and Reactions of Organic Compounds, eds. S. D. Barton and W. D. Ollis, Vol. 4. Heterocyclic Compounds, ed. P. G. Sammes, Pergamon Press, Oxford–New York–Toronto–Sydney–Paris–Frankfurt, 1979).

- 2 L. Brandsma, *Studies in Organic Chemistry*. 34. *Preparative Acetylenic Chemistry*, (2nd edn.), Elsevier, Amsterdam, Oxford, New York, Tokyo, 1988, pp. 99–100, 108, 299.
- 3 L. Brandsma and H. D. Verkruisje, *Preparative Polar Organometallic Chemistry*, Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, 1987, vol. 1, pp. 152–154.
- 4 N. A. Nedolya, R.-J. de Lang, L. Brandsma and B. A. Trofimov, *Abstracts of the 11th International Conference on Organic Synthesis*, Amsterdam, 1996, p. 251.
- 5 L. Brandsma, B. Trofimov, N. Nedolya and A. Mal'kina, *Abstracts of the 17th International Symposium on the Organic Chemistry of Sulfur*, Tsukuba, Japan, 1996, p. 233.
- 6 L. Brandsma, B. Trofimov, N. Nedolya and A. Mal'kina, *Abstracts of the 17th International Symposium on the Organic Chemistry of Sulfur*, Tsukuba, Japan, 1996, p. 234.

Received: Moscow, 2nd December 1996

Cambridge, 5th January 1997; Com. 6/08392G

[†] The reaction was performed under anhydrous conditions and in a nitrogen atmosphere. In a typical experiment, *tert*-butyllallene (0.11 mol) was introduced into a stirred solution of Bu^tLi (0.11 mol) in 70 ml THF and 70 ml hexane cooled to -40°C . After efficient stirring at -35°C during 30 min, the reaction mixture was cooled to -115°C and a solution of phenyl isocyanate (0.11 mol) in 20 ml THF was added in small portions at -100 to -95°C for 5 min. The reaction mixture was then treated with freshly-distilled trimethylchlorosilane (0.13 mol) at -80°C . The cooling bath was removed and when the temperature reached 0°C , dry work-up of the reaction mixture was carried out: removal of solvents on a rotary evaporator, dilution of the rest (a mixture of azatriene **4** with LiCl) with small portions of pentane, filtration of LiCl , removal of pentane on a rotary evaporator. 30 g (95%) of crude **4** (light yellow liquid) was isolated. ¹H NMR (90 MHz, CCl_4 , standard TMS), δ 5.52 (d, 1H, CH=), 5.32 (d, 1H, CH=), 7.25–6.65 (m, 5H, NPh), 1.05 (s, 9H, Bu^t), 0.30 (s, 9H, OSiMe_3).

After refluxing of azatriene **4** during 7–10 min in the presence of ca. 1 ml toluene and subsequent distillation *in vacuo* 23.7 g (75%) of quinoline **5** was obtained, bp 135 – 140°C (~ 0.5 mmHg), n_D^{20} 1.5315. IR, ν/cm^{-1} (film): 500, 530, 640, 690, 740, 760, 790, 850–860, 910, 940, 960, 1000, 1030, 1040, 1050, 1070, 1135, 1160, 1190, 1240, 1250, 1260, 1310, 1330, 1350, 1380, 1410, 1440, 1460, 1490, 1510, 1560, 1600, 1630, 1660, 2860, 2900, 2950, 3030, 3050. ¹H NMR (300 MHz, CDCl_3 , standard TMS), δ 6.97 (s, 1H, H-3), 7.86 (d, 1H, H-5), 7.38 (dt, 1H, H-6), 7.60 (dt, 1H, H-7), 8.01 (d, 1H, H-8), 2.96 (s, 2H, CH_2), 1.04 (s, 2H, CH_2), 1.04 (s, 9H, Bu^t), 0.52 (s, 9H, OSiMe_3). ¹³C NMR (75 MHz, CDCl_3), δ 160.32 (C-2), 116.28 (C-3), 144.86 (C-4), 124.82 (C-4a), 148.67 (C-8a), 128.66, 128.06, 127.72, 123.19 (C-5–8), 44.28 (CH_2), 32.73 [$\text{C}(\text{Me})_3$], 30.15 [$\text{C}(\text{Me})_3$], 0.68 (SiMe_3). Found: C 70.78, H 8.75, N 5.26, Si 10.08%; calc. for $\text{C}_{17}\text{H}_{25}\text{NOSi}$: C 71.08, H 8.71, N 4.88, Si 9.76%.