

Dolichyl phosphate derivatives with a fluorescent label at an internal isoprene unit

Natalia Ya. Grigorieva,^{*a} Olga A. Pinsker,^a Sergei D. Maltsev,^a Leonid L. Danilov,^a Vladimir N. Shibaev^a and Mark J. Jedrzejak^b

^a N. D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 117913 Moscow, Russian Federation.

Fax: + 7 095 135 5328; e-mail: ves@cacr.ioc.ac.ru

^b Department of Microbiology, The University of Alabama at Birmingham, Birmingham, Alabama 35294, USA

10.1070/MC2000v010n03ABEH001257

A general approach based on directed aldol condensation followed by reductive amination with fluorescent amines and phosphorylation was developed and illustrated by the synthesis of two dolichyl phosphate derivatives with the 1-aminonaphthalene fluorophore at the γ -isoprene unit of the chain.


The carbohydrate chains of glycoproteins are involved in many cellular activities¹ and are often structurally changed in diseases.² The biosynthesis of Asn-linked oligosaccharides on eukaryotic glycoproteins, the O-glycosylation of fungal glycoproteins and the formation of glycosylphosphatidylinositol anchors present in numerous membrane proteins require the participation of dolichyl phosphate (Dol-P), a derivative of a linear long-chain polyisoprene alcohol, and its glycosylated forms as key intermediates.³ The fluorescence methodology seems to be promising for studies of molecular mechanisms of the interaction of Dol-P and its glycosylated derivatives with biological membrane components. The synthesis of Dol-P derivatives with fluorescent labels at the ω -end of the chain was reported recently.⁴ Here, we describe a procedure for the incorporation of fluorophores in an internal isoprene unit of Dol-P.

The methodology is based on the preferential formation of thermodynamically more stable (*E*)-isomers of α,β -disubstituted

acroleins in directed aldol condensation, which was found previously⁵ and successfully used in the syntheses of dolichols, poly-prenols and their analogues.^{6,7} The key step in the synthetic strategy was the construction of dolichyl-like (*E*)-acroleins through the cross-condensation of two appropriate aldehydes, one of which was used as an aldimine. The resulting (*E*)-acrolein was subjected to reductive amination with a fluorescent amine, and the amino alcohol formed was phosphorylated.

This approach is illustrated by Scheme 1 where the synthesis of two Dol-P analogues (**5a,b**) with a 1-aminonaphthalene label at the γ -isoprene unit is shown. The aldehydoacetate **1**⁴ was used as an aldehyde component to build the key intermediates, (*E*)-acroleins **3a,b**. Condensation of **1** with aldimine **2a**⁸ deprotonated by treatment with LDA at -70°C leads to acetoxyacrolein **3a** (40%, after flash chromatography on SiO_2) which contains ~2% of the (*Z*)-isomer (¹H NMR data). In a similar manner, (*E*)-acetoxyacrolein **3b** [~3% of the (*Z*)-isomer, according to ¹H NMR data] was obtained in ~21% yield through the condensation of **1** with aldimine **2b** prepared in five synthetic steps (cf. ref. 9) from a mixture of polyprenols $\text{WT}_2\text{C}_n\text{-OH}$ isolated from birch tree¹⁰ with C_{35} - and C_{40} -prenols ($n = 4, 5$) as main components.[†]

The reductive amination of **3a,b** with 1-aminonaphthalene and NaBH_3CN (cf. ref. 11) led to amino alcohols **4a,b** in 37 and ~20% yields (after flash chromatography on SiO_2), respectively.[‡] To prepare fluorescent Dol-P derivatives, **4a,b** were subjected to phosphorylation with $\text{Bu}_4\text{N-H}_2\text{PO}_4/\text{CCl}_3\text{CN}$ ¹² under conditions essentially similar to those described for the synthesis of ω -labeled analogues.⁴ The resulting phosphates **5a,b**[§] were isolated as ammonium salts after purification by ion-exchange chromatography (DE-52, AcO^-) in 18 and 28% yields, respectively. As expected, they exhibited intense fluorescence with an excita-



Scheme 1 Reagents and conditions: i, LDA, Et_2O -hexane (1:1), $-10^\circ\text{C} \rightarrow 0^\circ\text{C}$, 30 min; ii, **1**, Et_2O , -70°C (2 h) $\rightarrow -20^\circ\text{C}$ (2.5 h); iii, 3.5% aq. HCl, 20°C , 3 h, then $\text{Ac}_2\text{O}/\text{Py}$ -DMAP, 20°C , 4 h; iv, 1-aminonaphthalene, $\text{NaBH}_3\text{CN}/\text{AcOH}$ -MeOH (for **4a**) or MeOH- Et_2O (for **4b**), 20°C , 92 h, then HCl to pH 2, 20°C , 10 min, then NaOH/MeOH, 20°C , 15 min; v, $\text{Bu}_4\text{N-H}_2\text{PO}_4$, $\text{CCl}_3\text{CN}/\text{CH}_2\text{Cl}_2$, 20°C , 48 h, chromatography on DE-52 (AcO^-).

[†] **3a**: ¹H NMR (CDCl_3) δ : 0.92 (d, 3H, MeC-10, J 6.5 Hz), 1.05–1.50 (m, 5H, H_2C -9,11, HC-10), 1.62 (s, 12H, *cis*-Me), 1.71 (s, 6H, *trans*-MeC-6,16'), 2.0 (m, 16H, $\text{CH}_2\text{C}=\text{C}$), 2.01 (s, 3H, MeCO), 2.25 (m, 4H, H_2C -5,1'), 2.45 (td, 2H, H_2C -4, $J_1 = J_2 = 7$ Hz), 4.09 (td, 2H, H_2C -12, J_1 7 Hz, J_2 2.5 Hz), 5.12 (m, 5H, HC=C), 6.43 (t, 1H, HC-3, J 7 Hz), 9.34 (s, 1H, HC-1). ¹³C NMR, δ : 15.9 (*cis*-Me), 17.5 (*cis*-MeC-16'), 19.3 (*MeC*-10), 20.9 (*MeCO*), 23.1 (*MeC*-6), 24.2 (C-1'), 25.5 (*trans*-MeC-16'), 25.2, 25.6, 26.0, 26.7, 27.0, 27.3 ($\text{CH}_2\text{CH}=\text{C}$), 29.5 (C-10), 30.6 (C-5), 35.4 (C-9), 37.1 (C-11), 39.6 [$\text{H}_2\text{CC}(\text{Me})=\text{C}$ of (*E*)-units], 62.8 (C-12), 123.2, 124.2, 124.35, 126.6 (HC=C), 131.1, (C-16'), 133.2, 134.8, 134.9, 136.0 (*MeC*=C), 143.3 (C-2), 154.3 (C-3), 171.0 (COMe), 194.9 (C-1).

3b: ¹H NMR (CDCl_3) δ : 0.93 (d, 3H, MeC-10, J 6.5 Hz), 1.10–1.60 (m, 5H, H_2C -9,11, HC-10), 1.59 (s, 9H, *cis*-Me), 1.68 (s, 19.5H, *trans*-MeC), 2.0 (m, 33H, $\text{CH}_2\text{C}=\text{C}$, MeCO), 2.20 (m, 4H, H_2C -5,1'), 2.45 (td, 2H, H_2C -4, $J_1 = J_2 = 7$ Hz), 4.11 (td, 2H, H_2C -12, J_1 7 Hz, J_2 2.5 Hz), 5.15 (m, 8.5H, HC=C), 6.45 (t, 1H, HC-3, J 7 Hz), 9.35 [s, 0.97H, HC-1 of (*E*)-isomer], 10.10 [s, 0.03H, HC-1 of (*Z*)-isomer]. ¹³C NMR, δ : 15.9 (*cis*-Me), 17.6 (*cis*-Me of ω -terminal unit), 19.3 (*MeC*-10), 20.9 (*MeCO*), 23.1 (*MeC*-6), 23.4 (*trans*-Me), 24.4 (C-1'), 25.6 (*trans*-Me of ω -terminal unit), 25.2, 26.3, 26.6, 26.7, 26.8, 27.0, 27.3 ($\text{CH}_2\text{CH}=\text{C}$), 29.4 (C-10), 30.5, 31.9, 32.1 [$\text{H}_2\text{CC}(\text{Me})=\text{C}$ of (*Z*)-units], 35.3 (C-9), 37.1 (C-11), 39.7 [$\text{H}_2\text{CC}(\text{Me})=\text{C}$ of (*E*)-units], 62.8 (C-12), 124.1, 124.15, 124.3, 124.4, 124.95, 126.6 (HC=C), 131.15, (MeC=C of ω -terminal unit), 133.2, 134.8, 135.1, 135.3, 135.95 (MeC=C), 143.6 (C-2), 154.45 (C-3), 171.0 (COMe), 194.9 (C-1).

tion/emission maximum at 340/410 nm (in *n*-heptane–2-propanol, 4:1) and are quite similar in this respect to the derivatives of Dol-P with a 1-naphthylamine residue at the ω -end of the chain⁴ (see ref. 13 for the excitation and emission spectra).

In preliminary biochemical tests, phosphate **5a** was found to serve as a substrate for recombinant yeast Dol-P-mannose synthase (EC 2.4.1.83) isolated from *Escherichia coli*.¹⁴

It is likely that the described approach to the incorporation of a fluorescent label at an internal isoprene unit of Dol-P can be employed for the preparation of a wide range of related derivatives starting from the corresponding aldehydes and aldimines

‡ **4a**: ¹H NMR (CDCl₃) δ : 0.91 (d, 3H, MeC-3, *J* 6.6 Hz), 1.10–1.60 (m, 5H, H₂C-2,4, HC-3), 1.65 (s, 12H, *cis*-Me), 1.72 (s, 6H, *trans*-Me), 2.05 (m, 18H, CH₂C=C), 2.27 (m, 4H, H₂C-9,12), 3.20 (s, 1H, NH), 3.63 (dt, 2H, H₂C-1, *J*₁ 1.5 Hz, *J*₂ 6.6 Hz), 3.87 (s, 2H, CH₂N), 5.14 (m, 5H, HC=C), 5.45 (t, 1H, HC-10, *J* 6.7 Hz). ¹³C NMR, δ : 16.0, 16.1 (*cis*-Me), 17.6 (*cis*-MeC-27), 19.5 (MeC-3), 23.4 (MeC-7), 25.1, 25.3, 26.2, 26.6, 26.7, 27.2 (CH₂CH=C), 25.7 (*trans*-MeC-27), 29.2, 29.4 (C-3,12), 31.8 (C-8), 37.4 (C-4), 39.7 [CH₂C(Me)=C of (*E*)-units], 39.8 (C-2), 50.0 (CN), 61.1 (C-1), 123.8, 124.2, 124.4, 125.5, 125.55, 127.1 (HC=C), 131.2 (C-27), 134.2, 134.6, 135.0, 135.4, 136.2 (MeC=C). Signals for 1-aminonaphthalene fragment are not shown.

4b: ¹H NMR (CDCl₃) δ : 0.90 (d, 3H, MeC-3, *J* 6.5 Hz), 0.95–1.55 (m, 5H, H₂C-2,4, HC-3), 1.57 (s, 9H, *cis*-Me), 1.65 (s, 19.5H, *trans*-Me), 2.01 (m, 32H, CH₂C=C), 2.20 (m, 4H, H₂C-9,12), 3.20 (s, 1H, NH), 3.70 (dt, 2H, H₂C-1, *J*₁ 1.5 Hz, *J*₂ 6.6 Hz), 3.87 (s, 2H, H₂CN), 5.13 (m, 8.5H, HC=C), 5.53 (t, 1H, HC-10, *J* 6.7 Hz). ¹³C NMR, δ : 16.0 (*cis*-Me), 17.7 (*cis*-Me of ω -terminal unit), 19.5 (MeC-3), 23.4 (*trans*-Me), 25.7 (*trans*-Me of ω -terminal unit), 25.1, 26.2, 26.4, 26.6, 26.7, 27.0 (CH₂CH=C), 29.1 (C-3), 29.6 (C-12), 31.8, 31.9, 32.2 [CH₂C(Me)=C of (*Z*)-units], 37.4 (C-4), 39.7 [CH₂C(Me)=C of (*E*)-units], 39.8 (C-2), 50.0 (CN), 61.0 (C-1), 123.8, 124.1, 124.2, 124.3, 124.7, 124.9, 125.6, 125.7, 127.1, 128.8 (HC=C), 131.2 (MeC=C of ω -terminal unit), 134.2, 134.6, 135.15, 135.2, 135.3, 135.7, 136.0 (MeC=C). Signals for 1-aminonaphthalene fragment are not shown.

§ **5a**: UV, λ_{\max} /nm: 252, 335; ϵ_{\max} : 18000, 5500. ESI-MS, *m/z*: 716 [M (free acid) – H].

5b: UV, λ_{\max} /nm: 252, 335; ϵ_{\max} : 18500, 5000; ³¹P NMR (CD₃OD–CDCl₃, 1:2) δ : 2.25.

¹H and ¹³C NMR spectra of **5a,b** are similar to the spectra of **4a,b** except for signals of HC-1 (3.90 and 3.85 for **5a** and **5b**, respectively) and C-1 (64.0 and 63.8 for **5a** and **5b**, respectively).

available from different terpenols.⁷ The use of these Dol-P analogues is of paramount importance for studies of the interaction of Dol-P with biological membrane components.

This work was supported by the U.S. Civilian Research and Development Foundation (grant no. RN1-404).

References

- 1 A. Varki, *Glycobiology*, 1993, **3**, 97.
- 2 I. Brockhausen, J. S. Schutzbach and W. Kuhns, *Acta Anatom.*, 1998, **161**, 36.
- 3 F. W. Hemming, in *New Comprehensive Biochemistry*, eds. J. Montreuil, H. Schachter and J. F. G. Vliegthart, Elsevier, Amsterdam, 1995, vol. 29a, p. 127.
- 4 V. N. Shibaev, V. V. Veselovsky, A. V. Lozanova, S. D. Maltsev, L. L. Danilov, W. T. Forsee, J. Xing, H. C. Cheung and M. J. Jedrzejak, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 189.
- 5 N. Ya. Grigorieva, E. P. Prokof'ev and A. V. Semenovskiy, *Dokl. Akad. Nauk SSSR*, 1979, **245**, 366 [*Dokl. Chem. (Engl. Transl.)*, 1979, **245**, 112].
- 6 N. Ya. Grigorieva, V. V. Veselovsky and A. M. Moiseenkov, *Khim.-Farm. Zh.*, 1987, **21**, 845 (in Russian).
- 7 N. Ya. Grigorieva and O. A. Pinsker, *Usp. Khim.*, 1994, **63**, 177 (*Russ. Chem. Rev.*, 1994, **63**, 169).
- 8 N. Ya. Grigorieva, I. M. Avrutov and A. V. Semenovskiy, *Tetrahedron Lett.*, 1983, **24**, 5531.
- 9 N. Ya. Grigorieva, O. A. Pinsker, E. D. Daeva and A. M. Moiseenkov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1991, 2325 (*Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1991, **40**, 2037).
- 10 A. R. Wellburn and F. W. Hemming, *Nature*, 1966, **212**, 1364.
- 11 A. Schepartz and R. Breslow, *J. Am. Chem. Soc.*, 1987, **109**, 1814.
- 12 L. L. Danilov, T. N. Druzhinina, N. A. Kalinchuk, S. D. Maltsev and V. N. Shibaev, *Chem. Phys. Lipids*, 1989, **51**, 191.
- 13 J. Xing, W. T. Forsee, E. Lamani, S. D. Maltsev, L. L. Danilov, V. N. Shibaev, J. S. Schutzbach, H. C. Cheung and M. J. Jedrzejak, *Biochemistry*, submitted.
- 14 J. S. Schutzbach, J. W. Zimmerman and W. T. Forsee, *J. Biol. Chem.*, 1993, **268**, 24190.

Received: 28th December 1999; Com. 99/1583