

Intermolecular and intramolecular cycloaddition reactions of 1-ethyl-1,2,4-triazinium salts with alkynes


Nataliya N. Mochulskaya, Anatoly A. Andreiko, Valery N. Charushin,* Boris V. Shulgin, Dmitry V. Raikov and Vladimir I. Solomonov

Department of Organic Chemistry, Urals State Technical University, 620002 Ekaterinburg, Russian Federation.
Fax: +7 3432 74 0458; e-mail: charushin@prm.uran.ru

10.1070/MC2001v011n01ABEH001370

Azomethyne ylides generated from 3-alkylthio substituted 1-alkyl-5-aryl-1,2,4-triazinium salts **4a–c** on treatment with triethylamine undergo 1,3-dipolar cycloaddition with dimethyl acetylenedicarboxylate to give pyrrolo[2,1-*f*][1,2,4]triazines, while 1-ethyl-5-phenyl-1,2,4-triazinium salts **1a,b** bearing the C≡C bond in the side-chain 3-alkynylthio substituent react with acetylenes to undergo either intermolecular 1,3-dipolar cycloaddition or intramolecular inverse electron demand Diels–Alder reactions.

Inverse electron demand Diels–Alder reactions of π -deficient azaromatics is a powerful synthetic methodology.¹ *N*-Alkyl-1,2,4-triazinium salts are appropriate substrates to be used in these (4 + 2) cycloaddition reactions. Indeed, we found that 1-ethyl-5-phenyl-1,2,4-triazinium salts bearing acetylenic 3-butynylthio (compound **1a**) or 4-pentynylthio (compound **1b**) substituents at C-3 can undergo intramolecular (4 + 2) cycloaddition reaction under very mild conditions, thus being transformed into thieno[2,3-*b*]-**3a** or thiopyrano[2,3-*b*]pyridines **3b** (Scheme 1).²



Scheme 1

On the other hand, *N*-alkylpyridinium and other azinium salts, especially those bearing the CH₂W fragment at the quaternary nitrogen (W is an electron-withdrawing group) are precursors of *N*-heterocyclic ylides, which can undergo 1,3-dipolar (3 + 2) cycloaddition reactions with both alkenes and acetylenes. The feature of *N*-alkyl-1,2,4-triazinium salts is that electron-withdrawing groups W in the *N*-alkyl fragment are not necessary to generate ylides, as found by the dimerisation into 4a,4b,9,10-tetrahydro-1,3,6,8a,10a-hexaazaphenanthrenes.^{3,4} In continuation of our studies on the behaviour of 1-alkyl-1,2,4-triazinium salts,^{5–7} we report that deprotonation of 1-ethyl-3-alkylthio-5-aryl-1,2,4-triazinium salts **4a–c** with triethylamine (TEA) results in azomethine ylides **5a–c** (non-stabilised species), which react *in situ* with dimethyl acetylenedicarboxylate (DMAD) to form pyrrolo[2,1-*f*][1,2,4]triazines **6a–c** in good yields (Scheme 2).[†] Attempts to use other alkynes, such as phenylacetylene, di(4-nitrophenyl)acetylene and di(pyridin-2-yl)acetylene, as dipolarophiles in the reaction with **4c** were unsuccessful.

When irradiated with UV light or pulsed electron beams, pyrrolo[2,1-*f*][1,2,4]triazines **6a–c** were found to exhibit intense luminescence giving emission in the region 450–650 nm with maxima at 500 nm and the light output comparable with that of stilbene.[‡]

It was interesting to examine the chemical behaviour of 3-alkynylthio-1,2,4-triazinium salts **1a,b** on treatment with DMAD, *i.e.*, when both types of cycloaddition reactions [(4 + 2) and (3 + 2)] can compete with each other. We found that, in case of a 4-pentynylthio substituent (*n* = 2), the intermolecular 1,3-dipolar cycloaddition reaction of **1b** with DMAD leading to

pyrrolotriazine **7** (47% yield) proceeds much faster than the intramolecular Diels–Alder reaction, while 3-butynylthio substituted 1-ethyl-1,2,4-triazinium salt **1a** (*n* = 1) undergoes predominantly the intramolecular ring transformation into thieno[2,3-*b*]pyridine **3a** in 44% yield (Scheme 3). Both reactions were carried out under identical conditions with triethylamine in absolute ethanol at room temperature. Although in both reactions we cannot exclude the formation of minor quantities of alternative products, moderate yields of **3a** and **7** can be

[†] All new compounds gave expected ¹H NMR spectra, mass spectra and satisfactory elemental analyses.

3a: Dimethyl acetylenedicarboxylate (0.08 g, 0.56 mmol) was added to a suspension of 1-ethyl-3-(3-butynylthio)-5-phenyl-1,2,4-triazinium tetrafluoroborate **1a** (0.18 g, 0.50 mmol) in 4 ml of absolute ethanol, and then triethylamine (0.06 g, 0.55 mmol) was added dropwise with stirring. The reaction mixture was stirred at room temperature for 36 h. After filtration, the solvent was removed *in vacuo*. Flash dry-column chromatography on silica gel (hexane–acetone, 4:1) gave 6-phenyl-2,3-dihydrothieno[2,3-*b*]pyridine (0.047 g, 44% yield) as white crystals, mp 82–83 °C (lit.,² mp 83 °C).


6a: Dimethyl acetylenedicarboxylate (0.21 g, 1.46 mmol) was added to a suspension of 5-(4-bromophenyl)-1-ethyl-3-propylthio-1,2,4-triazinium tetrafluoroborate (0.52 g, 1.22 mmol) in 4 ml of absolute ethanol, and then triethylamine (0.14 g, 1.34 mmol) was added dropwise with stirring. The reaction mixture was stirred at room temperature for 72 h to precipitate a yellow crystalline product, which was collected by filtration and recrystallised from ethanol to yield 0.23 g (40%) of 4-(4-bromophenyl)-5,6-dimethoxycarbonyl-7-methyl-2-propylthiopyrrolo[2,1-*f*]-1,2,4-triazine as a yellow product, mp 126–128 °C. ¹H NMR (CDCl₃) δ : 1.09 (t, 3H, SCH₂CH₂Me), 1.85 (m, 2H, SCH₂CH₂Me), 2.79 (s, 3H, Me), 3.20 (t, 2H, SCH₂CH₂Me), 3.47 (s, 3H, COOMe), 3.88 (s, 3H, COOMe), 7.54 (d, 2H, BrC₆H₄), 7.62 (d, 2H, BrC₆H₄). MS, *m/z* (%): 479 (58, [M + 1]⁺), 478 (16, M⁺), 477 (55), 464 (27), 462 (26), 446 (48), 444 (35), 438 (20), 437 (100), 436 (23), 435 (93).

Compounds **6b** and **6c** were obtained analogously to **6a**.

6b: yield 65%, yellow crystals, mp 108–110 °C. ¹H NMR (CDCl₃) δ : 1.48 (t, 3H, SCH₂Me), 2.80 (s, 3H, Me), 3.21 (q, 2H, SCH₂Me), 3.39 (s, 3H, COOMe), 3.87 (s, 3H, COOMe), 7.4–7.6 (m, 3H, Ph), 7.6–7.7 (m, 2H, Ph). MS, *m/z* (%): 386 (23, [M + 1]⁺), 385 (100, M⁺), 357 (21), 352 (41), 294 (32), 262 (20).


6c: yield 40%, yellow crystals, mp 165–167 °C. ¹H NMR ([²H₆]DMSO) δ : 2.61 (s, 3H, SMe), 2.76 (s, 3H, Me), 3.35 (s, 3H, COOMe), 3.82 (s, 3H, COOMe), 7.4–7.6 (m, 5H, Ph). MS, *m/z* (%): 372 (34, [M + 1]⁺), 371 (100, M⁺), 340 (22), 339 (35), 280 (35), 252 (22), 239 (21).

7: Dimethyl acetylenedicarboxylate (0.18 g, 1.26 mmol) was added to a suspension of 1-ethyl-3-(4-pentynylthio)-5-phenyl-1,2,4-triazinium tetrafluoroborate **1b** (0.39 g, 1.05 mmol) in 5 ml of absolute ethanol, and then triethylamine (0.12 g, 1.16 mmol) was added dropwise with stirring. The reaction mixture was allowed to stand at room temperature for 24 h. The solvent was removed *in vacuo*. Flash dry-column chromatography on silica gel (hexane–ethyl acetate, 4:1) gave 5,6-dimethoxycarbonyl-7-methyl-2-(4-pentynylthio)-4-phenylpyrrolo[2,1-*f*][1,2,4]triazine (0.208 g, 47%) as a yellow solid, mp 80–82 °C. ¹H NMR (CDCl₃) δ : 1.99 (t, 1H, C≡CH, *J* 2.8 Hz), 2.07 (m, 2H, SCH₂CH₂, *J* 7.0 Hz), 2.42 (td, 2H, CH₂C≡CH, *J* 2.8 Hz, *J* 7.0 Hz), 2.80 (s, 3H, Me), 3.32 (t, 2H, SCH₂, *J* 7.0 Hz), 3.39 (s, 3H, COOMe), 3.87 (s, 3H, COOMe), 7.4–7.6 (m, 3H, Ph), 7.6–7.7 (m, 2H, Ph). MS, *m/z* (%): 423 (24, M⁺), 396 (24), 395 (100), 384 (78), 371 (25), 364 (38), 357 (41), 267 (23).



Scheme 2

explained by side reactions of starting 1,2,4-triazinium salts **1a,b**, such as dequaternisation⁵ and dimerisation,^{3,4} as experimentally found by TLC. Thus, the conversion of 1-ethyl-3-alkynylthio-1,2,4-triazinium salts **1a,b** into thieno[2,3-*b*]-pyridine **3a** or pyrrolo[2,1-*f*][1,2,4]triazine **7** depends on the length of the side chain bearing the acetylenic fragment. The found phenomenon is consistent with well-known features of the intramolecular Diels–Alder reactions of alkynyl-substituted pyrazines, pyrimidines and 1,2,4-triazines.^{2,8–11} In these ring transformations, the formation of bridged intermediates like **2** (Scheme 1) with an annelated five-membered ring is more favourable as compared to cycloadducts **2** condensed with a six-membered ring.^{2,8–11} As a consequence, intramolecular Diels–Alder reactions leading to azines condensed with five-membered rings proceed much faster and require milder conditions.^{2,8–11}



Scheme 3

‡ For measuring the photoluminescence spectra, compounds **6a–c** and **7** were irradiated with monochromatic UV light at a wavelength of 310 nm followed by measuring emission with the second monochromator equipped with a photoelectric multiplier (1050 V) and an automatic potentiometer. The radioluminescence and scintillation characteristics of compounds **6a–c** and **7** were obtained using special nanosecond techniques. The test materials were excited by irradiation with pulsed electron beams at a high current density (pulse duration of 2 ns, electron energy of 100–200 keV, current density of 100–700 A cm^{–2}) using a Radan electron accelerator. The spectra in the range 300–800 nm were measured by a charge-connecting detector (CCD) equipped with a computer.

The above results demonstrate a new interesting property of 1,2,4-triazinium salts to undergo two modes of cycloaddition reactions with alkynes. Evidence for the structures of **3a**, **6a–c** and **7** is provided by ¹H NMR spectroscopy and mass spectrometry.[‡]

This work was supported by the US Civilian Research and Development Foundation (grant no. REC-005) and the Russian Foundation for Basic Research (grant nos. 00-03-32785a and 00-15-97390).

References

- 1 D. L. Boger and S. M. Weinreb, *Hetero Diels–Alder Methodology in Organic Syntheses*, Academic Press, New York, 1987.
- 2 V. N. Charushin, A. van Veldhuizen, H. C. van der Plas and C. H. Stam, *Tetrahedron*, 1989, **45**, 6499.
- 3 O. N. Chupakhin, B. V. Rudakov, S. G. Alexeev, V. N. Charushin and V. A. Chertkov, *Tetrahedron Lett.*, 1990, **31**, 7665.
- 4 B. V. Rudakov, S. G. Alexeev, V. N. Charushin, O. N. Chupakhin, V. A. Chertkov, G. G. Alexandrov and E. S. Klimov, *Zh. Org. Khim.*, 1992, **28**, 589 (*Russ. J. Org. Chem.*, 1992, **28**, 479).
- 5 O. N. Chupakhin, B. V. Rudakov, S. G. Alexeev, S. V. Shorshnev and V. N. Charushin, *Mendeleev Commun.*, 1992, 85.
- 6 V. N. Charushin, S. G. Alexeev, O. N. Chupakhin and H. C. van der Plas, *Adv. Heterocycl. Chem.*, 1989, **46**, 73.
- 7 O. N. Chupakhin, S. G. Alexeev, B. V. Rudakov and V. N. Charushin, *Heterocycles*, 1992, **33**, 931.
- 8 A. E. Frissen, *Intramolecular Inverse Electron Demand Diels–Alder Reaction of Pyrimidines*, Thesis, Wageningen, 1990, p. 117.
- 9 A. E. Frissen, A. T. M. Marcelis and H. C. van der Plas, *Tetrahedron*, 1987, **43**, 1589.
- 10 A. E. Frissen, A. T. M. Marcelis and H. C. van der Plas, *Tetrahedron*, 1989, **45**, 803.
- 11 A. E. Frissen, G. Geurtsen, A. T. M. Marcelis and H. C. van der Plas, *Tetrahedron*, 1990, **46**, 595.

Received: 1st September 2000; Com. 00/1696