

An alternative reaction of *ortho*-(*N*-benzylidene)aminophenol with chlorophosphites: formation of 2-(2'-alkoxy)-2-oxo-3-phenyl-5,6-benzo-1,4,2-oxazaphosphorinanes


Mudaris N. Dimukhametov,* Eugenia V. Bajandina, Elena Yu. Davydova, Alexey B. Dobrynin, Aidar T. Gubaidullin, Igor A. Litvinov and Vladimir A. Alfonsov

A. E. Arbuzov Institute of Organic and Physical Chemistry, Kazan Scientific Centre of the Russian Academy of Sciences, 420088 Kazan, Russian Federation. Fax: +7 8432 75 5322; e-mail: mudaris@iopc.kcn.ru

10.1070/MC2001v011n05ABEH001478

The reactions of *o*-(*N*-benzylidene)aminophenol with ethylene chlorophosphite and diethyl chlorophosphite in CHCl_3 in the absence of an external acceptor of HCl result in the formation of diastereomeric 2-(2'-alkoxy)-2-oxo-3-phenyl-5,6-benzo-1,4,2-oxazaphosphorinanes different in the configurations of phosphorus atoms.

The reactions of P^{III} halides with compounds which contain both a hydroxy group and an imino $\text{C}=\text{N}$ unit have not been widely studied. Several papers describe the formation of poly-cyclic phosphoranes *via* the reaction of $\text{Cl}-\text{P}^{\text{III}}$ derivatives with phenols containing an imino group in the presence of a base.^{1–3} We have observed an alternative reaction pathway in similar systems. Here we report the reactions of *o*-(*N*-benzylidene)aminophenol **1** with ethylene chlorophosphite and diethyl chlorophosphite. These reactions[†] were carried out in the absence of an external HCl acceptor, and they resulted in the formation of two stereoisomers (diastereomers **A** and **B**) of 2-(2'-alkoxy)-2-oxo-3-phenyl-5,6-benzo-1,4,2-oxazaphosphorinanes **4** and **5** differing only in the configuration at the phosphorus atom. The diastereomeric ratio **A**:**B** in both cases was approximately equal to 2:1.



Scheme 1

Diastereomers **4A** and **4B** (from the reaction of imine **1** with ethylene chlorophosphite) and diastereoisomer **5A** (from the reaction of imine **1** with diethyl chlorophosphite) were isolated by column chromatography on silica gel as air-stable colourless crystals. The structures of 2-(2'-chloroethoxy)-2-oxo-3-phenyl-5,6-benzo-1,4,2-oxazaphosphorinane diastereomers **4A** and **4B** were determined by X-ray[‡] diffraction analysis (Figure 1).

The structures of products **5A** and **5B** formed in the reaction of imine **1** with diethyl chlorophosphite were determined by comparing their ^1H NMR and IR spectra with those of diastereomers **4A** and **4B**.[§] In the ^1H NMR spectrum of diastereomer **4A**, the proton of the PCH fragment appears at δ 4.81 ppm, whereas the analogous proton of diastereomer **4B** occurs at a lower field (δ 4.94 ppm). In addition, the magnitude of the

geminal coupling $^2J_{\text{HP}}$ in **4A** is approximately twice lower than that for diastereomer **4B**.

The ^1H NMR spectrum of a mixture of the two diastereomers of 2-(2'-ethoxy)-2-oxo-3-phenyl-5,6-benzo-1,4,2-oxazaphosphor-

[‡] X-ray diffraction analysis. Crystallographic data for **4A** and **4B** at 20 °C: crystals of $\text{C}_{15}\text{H}_{15}\text{ClNO}_3\text{P}$ **4A** are orthorhombic, space group $P2_12_12_1$, $a = 8.857(6)$ Å, $b = 10.532(2)$ Å, $c = 16.40(1)$ Å, $V = 1529.5(1)$ Å³, $Z = 4$, $M = 323.72$, $d_{\text{calc}} = 1.41$ g cm⁻³, $\mu(\text{CuK}\alpha) = 3.327$ cm⁻¹, $F(000) = 672$; crystals of $\text{C}_{15}\text{H}_{15}\text{ClNO}_3\text{P}$ **4B** are monoclinic, space group $P2_1/c$, $a = 8.70(2)$ Å, $b = 16.698(6)$ Å, $c = 10.942(6)$ Å, $\beta = 102.10(5)^\circ$, $V = 1554.8(4)$ Å³, $Z = 4$, $M = 323.72$, $d_{\text{calc}} = 1.38$ g cm⁻³, $\mu(\text{CuK}\alpha) = 3.273$ cm⁻¹, $F(000) = 672$. Intensities of 1816 reflections for M10 and 2377 reflections for M11 were measured on an Enraf Nonius CAD-4 diffractometer at 20 °C (λ CuK α radiation, $\omega/2\theta$ scan technique, $2\theta_{\text{max}} < 144^\circ$ for M10 and 114° for M11), of which 1710 and 2068 were with $I > 3\sigma$ for M10 and M11, respectively.

The structures were solved by direct methods and difference Fourier syntheses using the SIR program⁴ and the MolEN package.⁵ All non-hydrogen atoms were refined anisotropically; H atoms located in ΔF maps were refined isotropically. The absolute crystal structure and the absolute configuration of a molecule of **4A** were determined by the Hamilton test ratio⁶ with a probability of 95%. The final agreement factors are R 0.034, R_w 0.044 based on 1643 reflections with $F^2 \geq 3\sigma$ for **4A** and R 0.034, R_w 0.046 based on 1955 reflections with $F^2 \geq 3\sigma$ for **4B**. Atomic coordinates, bond lengths, bond angles and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre (CCDC). For details, see 'Notice to Authors', *Mendeleev Commun.*, Issue 1, 2001. Any request to the CCDC for data should quote the full literature citation and the reference number 1135/96.

[§] ^1H and ^{31}P NMR spectra (solvent: CD_3CN) were measured on Bruker WM-250 (^1H , 250 MHz, TMS) and Bruker MSL-400 (^{31}P , 162 MHz, 85% H_3PO_4) instruments. IR spectra were recorded on a UR-20 spectrometer.

Compound **4**. Overall yield of diastereomers **4A** + **4B** is 73%.

Diastereomer **4A**: mp 149–150 °C. ^1H NMR, δ : 3.51 (m, 2H, CH_2Cl), 3.91 (m, 1H, OCH), 4.08 (m, 1H, OCH'), 4.81 (d, 1H, CHP, 2J 10.6 Hz), 6.80–7.04 (m, 4H, C_6H_4), 7.44–7.60 (m, 5H, Ph). ^{31}P NMR, δ : 10.6. IR (vaseline oil, KBr, ν/cm^{-1}): 1030, 1043 (P–O–C), 1222, 1252 (P=O), 3275 (N–H). Found (%): C, 55.74; H, 4.68; Cl, 10.48; N, 4.38; P, 9.31. Calc. for $\text{C}_{15}\text{H}_{15}\text{ClNO}_3\text{P}$ (%): C, 55.64; H, 4.64; Cl, 10.97; N, 4.33; P, 9.58.

Diastereomer **4B**: mp 90–91 °C. ^1H NMR, δ : 3.76 (m, 2H, CH_2Cl), 4.37 (m, 2H, OCH₂), 4.94 (d, 1H, CHP, 2J 20.4 Hz), 6.76–7.05 (m, 4H, C_6H_4), 7.39 (m, 5H, Ph). ^{31}P NMR, δ : 11.2. IR (vaseline oil, KBr, ν/cm^{-1}): 1038, 1088 (P–O–C), 1230, 1250 (P=O), 3253 (N–H). Found (%): C, 55.85; H, 4.41; Cl, 11.21; N, 4.19; P, 9.12. Calc. for $\text{C}_{15}\text{H}_{15}\text{ClNO}_3\text{P}$ (%): C, 55.64; H, 4.64; Cl, 10.97; N, 4.33; P, 9.58.

Compound **5**. Overall yield of diastereomers **5A** + **5B** is 79%.

Diastereomer **5A**: mp 138–139 °C. ^1H NMR, δ : 1.08 (t, 3H, Me, 3J 7.5 Hz), 3.83 (m, 1H, OCH), 3.97 (m, 1H, OCH'), 4.78 (d, 1H, CHP, 2J 11.4 Hz), 6.81–7.03 (m, 4H, C_6H_4), 7.39–7.62 (m, 5H, Ph). ^{31}P NMR, δ : 10.3. IR (vaseline oil, KBr, ν/cm^{-1}): 1035 (P–O–C), 1205, 1260 (P=O), 3292 (N–H). Found (%): C, 62.44; H, 5.60; N, 4.78; P, 10.68. Calc. for $\text{C}_{15}\text{H}_{16}\text{NO}_3\text{P}$ (%): C, 62.28; H, 5.54; N, 4.84; P, 10.78.

Mixture of diastereomers **5A** + **5B**, 1:1.3. Light-brown oil, n_D^{20} 1.5670. IR (KBr, ν/cm^{-1}): 1025, 1035, 1070 (P–O–C), 1210, 1223, 1260 (P=O), 3280, 3292 (N–H). Found (%): C, 62.02; H 5.16; N 4.19; P 9.12.

Diastereomer **5B**: ^1H NMR, δ : 1.26 (t, 3H, Me, 3J 7.1 Hz), 4.23 (m, 2H, OCH₂), 4.87 (d, 1H, CHP, 2J 19.5 Hz), 6.75–7.07 (m, 4H, C_6H_4), 7.27–7.78 (m, 5H, Ph). ^{31}P NMR, δ : 12.0.

[†] A typical experimental procedure was as follows: A solution of **1** (7.1 mmol) in CHCl_3 (5 ml) was added dropwise to a solution of **2** (7.1 mmol) in CHCl_3 (10 ml) in a dry argon atmosphere, with cooling (10 °C) and stirring. After stirring for 10 min at this temperature, the reaction mixture was warmed slowly to room temperature and additionally stirred for 2 h; then, it was evaporated to dryness in a vacuum. The residue was purified by column chromatography (silica gel; toluene–acetonitrile, 4:1).




Figure 1 Molecular structure of diastereomers **A,B** of compound **4**. For **4A**, selected bond lengths (Å) : P(2)–O(1) 1.569(2), P(2)–O(2) 1.466(2), P(2)–O(3) 1.572(2), P(2)–C(3) 1.815(2), N(4)–C(3) 1.450(3), N(4)–C(5) 1.394(3), O(1)–C(6) 1.407(3), C(5)–C(6) 1.402(3), C(3)–C(13) 1.511(3); selected bond angles (°): O(1)–P(2)–O(2) 113.0(1), O(1)–P(2)–O(3) 101.8(1), O(1)–P(2)–C(3) 103.0(1), O(2)–P(2)–O(3) 114.9(1), O(2)–P(2)–C(3) 115.5(1), O(3)–P(2)–C(3) 107.0(1), P(2)–O(1)–C(6) 124.5(2), P(2)–O(3)–C(11) 122.9(2), C(3)–N(4)–C(5) 117.6(2), P(2)–C(3)–N(4) 107.1(2), P(2)–C(3)–C(13) 110.4(2), N(4)–C(3)–C(13) 113.6(2), N(4)–C(5)–C(6) 121.1(2), O(1)–C(6)–C(5) 121.2(2), O(1)–C(6)–C(7) 116.8(2). For **4B**, selected bond lengths (Å): P(2)–O(1) 1.584(1), P(2)–O(2) 1.459(2), P(2)–O(3) 1.559(1), P(2)–C(3) 1.810(2), N(4)–C(3) 1.457(2), N(4)–C(5) 1.386(2), O(1)–C(6) 1.413(2), C(5)–C(6) 1.390(2), C(3)–C(13) 1.509(2); selected bond angles (°): O(1)–P(2)–O(2) 113.39(8), O(1)–P(2)–O(3) 102.83(7), O(1)–P(2)–C(3) 102.76(8), O(2)–P(2)–O(3) 115.74(7), O(2)–P(2)–C(3) 116.68(8), O(3)–P(2)–C(3) 103.65(8), P(2)–O(1)–C(6) 120.5(1), P(2)–O(3)–C(11) 121.6(1), C(3)–N(4)–C(5) 122.2(1), P(2)–C(3)–N(4) 105.9(1), P(2)–C(3)–C(13) 111.8(1), N(4)–C(3)–C(13) 111.7(1), N(4)–C(5)–C(6) 122.7(1), O(1)–C(6)–C(5) 120.7(1), O(1)–C(6)–C(7) 116.9(2).

inane **5** formed in the reaction of imine **1** with diethyl chlorophosphite shows two doublets at δ 4.76 (J 11.4 Hz) and 4.87 ppm (J 19.5 Hz). The ^1H NMR spectrum of the isolated product **5A** exhibits a doublet at δ 4.78 ppm (J 11.4 Hz). Based on this observation and the earlier result for diastereomers **4A** and **4B**, we can conclude that the isolated diastereomer of **5** has the structure of **5A**, which is analogous to diastereomer **4A**, *i.e.*, the alkoxy group in both cases adopts an axial orientation. The IR spectra of diastereomers **4A**, **4B**, **5A** and **5B** are similar with the most characteristic absorption band for the secondary amino group NH at 3250–3280 cm^{-1} .

The formation of 1,4,2-oxazaphosphorinanes **4** and **5** in the course of the reaction of **1** with chlorophosphites can be illustrated by Scheme 1. Initially formed iminium salts **2** undergo intramolecular cyclization by the nucleophilic attack of the P^{III} atom at the electrophilic carbon atom of the imonium group, resulting in the formation of quasiphosphonium salts **3**. The latter give final products **4** and **5** according to the second stage of the Arbuzov reaction.

References

- 1 A. Schmidpeter and J. H. Weinmaier, *Angew. Chem.*, 1975, **87**, 517.
- 2 A. Schmidpeter and J. H. Weinmaier, *Chem. Ber.*, 1978, **111**, 2086.
- 3 S. D. Harper and A. J. Arduengo, *J. Am. Chem. Soc.*, 1982, **104**, 2497.
- 4 A. Altomare, G. Cascarano, C. Giacovazzo and D. Viterbo, *Acta Crystallogr.*, 1991, **A47**, 744.
- 5 L. Straver and A. J. Schierbeek, *MolEN, Structure Determination System*, Nonius B.V., 1994, vol. 1, p. 180.
- 6 W. C. Hamilton, *Acta Crystallogr.*, 1965, **18**, 502.

Received: 22nd May 2001; Com. 01/1804