

First direct calculation of the partial quadrupole splitting of ligands for the prediction of Mössbauer spectra parameters in low-spin iron(II) complexes

Victor N. Nemykin,^{a,b} Anna E. Polshina,^{a,c} Ernst V. Polshin^d and Nagao Kobayashi^{*a}

^a Department of Chemistry, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan.

Fax: +81 222 17 7719; e-mail: victor_nemykin@yahoo.com

^b V. I. Vernadsky Institute of General and Inorganic Chemistry, National Academy of Sciences of Ukraine, 252680 Kiev, Ukraine.

Fax: +380 44 444 3070

^c Institute of Sorption and Endoecological Problems, National Academy of Sciences of Ukraine, 252680 Kiev, Ukraine.

Fax: +380 44 452 9327

^d Institute of Geochemistry, Mineralogy and Ore Formation, National Academy of Sciences of Ukraine, 252680 Kiev, Ukraine.

Fax: +380 44 444 0342

DOI: 10.1070/MC2000v010n02ABEH001207

A semi-empirical quantum mechanics method and a cone angle conception were used to factorise partial quadrupole splitting parameters for different ligands in axially coordinated macrocyclic complexes.

The ⁵⁷Fe Mössbauer spectroscopy is one of the most powerful techniques for probing the metal centre in inorganic and organometallic compounds of iron.¹ In spite of large progress in the theoretical analysis of Mössbauer spectra parameters on the basis of modern quantum theory methods, such as DFT and *ab initio*,² the partial quadrupole splitting (p.q.s.) and partial central shift (p.c.s.) models for the prediction of Mössbauer spectra parameters are still popular and useful in the case of low-spin iron(II) compounds.³ The p.q.s. parameters have been introduced, developed and successfully applied over the last 20 years.^{3,4} The p.q.s. model is attractive because of the following: (i) tabulated p.q.s. values already contain the sum of all ligand properties on which the experimental quadrupole splitting (Δ) depends; (ii) the experimental geometries of compounds are not required; (iii) experimental Δ values can be easily estimated 'by hand' [see, for instance, equations (1) and (2) for the estimation of Δ in *trans*-[FeA₂B₄] and *trans*-[FeACB₄] complexes].³ The application of equations (1) and (2) to the pcFeL₂, pcFeL'L'', nx₂FeL₂ and nx₂FeL'L' complexes, where [B] is pc/4, dmg/2 or nx/2, [A] is L or L', and [C] is L'', leads to the estimation of Δ values for the complexes of interest.

$$\Delta = 4\text{p.q.s.}(A) - 4\text{p.q.s.}(B) \quad (1)$$

$$\Delta = 2\text{p.q.s.}(A) + 2\text{p.q.s.}(C) - 4\text{p.q.s.}(B) \quad (2)$$

The p.q.s. values can be found from experimental Δ values³ or from structural parameters of the corresponding complexes.⁴ However, the factorisation of the p.q.s. of new ligands without a wide range of experimental data is a difficult problem. Here, we consider the relationship between the electronic and geometric structures of ligands and the p.q.s. values in low-spin iron(II) complexes. The first direct calculations[†] of p.q.s. given below can be easily used for the calculations of p.q.s. values of unknown ligands.

In general, the p.q.s. values include the following three main contributions:

(i) A 'lattice' contribution, which depends on the ligand charge or, more correctly, the ligand coordination atom (LCA) and on the distance between the LCA and the iron atom. This interaction leads to a change in the lattice contribution to p.q.s. Usually, for low-spin iron(II) complexes, this lattice contribution is small and can be neglected.¹ However, recent DFT calculations in a

Table 1 Ligand parameters estimated from experimental data and calculated p.q.s. values.

L ^a	−V _{MEP} /eV	V _b /eV	C×10 ²	T	p.q.s. (calc.)	p.q.s. (exp.) ^b [Ref.]
nh ₃	3.09	0.035	−1.0	93	−0.54	−0.52 [3(b), 4]
n-prnh ₂	3.24	0.036	−0.65	110	−0.46	−0.48 [4]
s-bunh ₂	3.28	0.035	−0.75	113	−0.44	−0.45 [this work ^c]
t-bunh ₂	3.35	0.036	−0.77	129	−0.36	−0.36 [this work ^c]
phnh ₂	2.45	0.036	1.4	110	−0.44	−0.43 [this work ^c]
mepip	3.13	0.036	−0.27	123	−0.39	−0.39 [this work ^c]
mph	3.02	0.038	−0.25	123	−0.39	−0.38 [this work ^c]
14en	3.84	0.03	−0.82	110	−0.5	−0.48 [this work ^c]
dabco	2.99	0.041	0.05	145	−0.27	−0.24 [this work ^c]
py	2.86	0.093	−0.87	100	−0.51	−0.48 [this work ^c]
4ohpy	2.99	0.091	−1.46	100	−0.51	−0.51 [this work ^c]
4chopy	2.53	0.11	−0.76	100	−0.5	−0.5 [this work ^c]
tz	2.2	0.219	−0.47	100	−0.51	−0.52 [this work ^c]
meim	3.32	0.079	−1.6	98	−0.58	−0.53 [4]
pzl	2.67	0.087	−1.4	98	−0.51	−0.51 [this work ^c]
tzl	2.91	0.096	−2.05	98	−0.51	−0.53 [this work ^c]
t-bunc	2.07	0.319	1.9	76	−0.68	−0.72 [3(d)]
phnc	2.04	0.362	1.6	76	−0.68	−0.7 [3(b)]
cn [−]	8.59	0.165	−2.5	76	−0.86	−0.84 [4]
co	1.51	0.633	5.5	78	−0.75	−0.74 [3(d)]
pet ₃	1.9	0.12	26.29	118	−0.59	−0.59 [this work ^c]
pobu ₃	0.59	0.368	32.96	109	−0.69	−0.68 [this work ^c]
tht	1.28	0.1	8.5	122	−0.4	−0.41 [this work ^c]
pc ^{2−}	11.97	0.06	−1.57	85	−0.93	−0.96 [4]
Hdmg [−]	12.07	0.048	−0.32	85	−0.94	−0.92 [4]
Hnx [−]						

^aAbbreviations: 14en, 1,4-diaminobutane; 4ohpy, 4-hydroxypyridine; 4chopy, pyridine-4-carbaldehyde; bu, butyl; dabco, diazabicyclooctane; meim, N-methylimidazole; mepip, 4-methylpiperidine; mph, morpholine; nh₃, ammonia; ph, phenyl; pr, propyl; py, pyridine; pzl, pyrazole; tht, tetrahydrothiophene; tz, 1,2,4,5-tetrazine; tzl, 1,2,4-triazole; ^bp.q.s. in mm s^{−1}. The p.q.s.(Cl[−]) = −0.27 mm s^{−1} scale has been used for all values, see ref. 3. ^cAll p.q.s. values from this work are based on the pcFeL₂, (Hnx)₂FeL₂ and (Hdmg)₂FeL₂ complexes, where pc^{2−} is the phthalocyanine dianion, Hnx[−] is the nioximate anion and Hdmg[−] is the dimethylgioximate anion.¹¹

local density approximation for the oepFe(pme₃)₂ complex^{2(a)} have shown that at least in some cases the lattice contribution can play a dominant role.

(ii) A 'valence' contribution, which arises from the σ - and π -interaction between iron and LCA atom orbitals and leads to a change of the effective population of iron atom orbitals. For example, the σ -donation of electron density by a ligand changes the effective population of 3d and 4p orbitals of the iron atom. The relationship between the σ - and π -donor and π -acceptor properties of ligands and the p.q.s. values have been widely discussed.³

(iii) A 'steric' contribution of the ligand, on which the metal–ligand distance and, correspondingly, the metal–ligand interaction depends. The influence of cone angle on p.q.s. parameters for phosphorus-containing ligands was discussed recently.^{3(f)}

[†] All computations were performed using the HyperChem 5.1 Pro program (HyperCube Inc.) on a Pentium PC. All ligand structures were fully optimised by the gradient Polak–Ribiere method at the semi-empirical AM1 level.⁷ The molecular electrostatic potential (V_{MEP}) and the 'molecular back-bonding potential' (V_b) were evaluated as described previously.⁵ The ligand resonance integrals were calculated in the NDO formalism.⁸ Steric parameters of axial ligands were evaluated from a cone angle conception.⁶ The effective van der Waals radius of the π -system was chosen as 1.7 Å.⁹ Multiparameter regression analysis was carried out by the Powell quadratic convergence method.¹⁰

Thus, an increase in the cone angle of a ligand probably leads to the elongation of the Fe–LCA bond. This elongated Fe–LCA bond decreases the overlap between the lone-pair orbital of the ligand and the 3d and 4p orbitals of iron and hence leads to a change in Δ . Note that this statement is true only for conformationally rigid ligands. From the above, we can conclude that the p.q.s. values for each ligand depend on the σ - and π -donor and π -acceptor properties, the LCA charge and the cone angle.

Recently, Fielder *et al.*⁵ have developed a model for the calculation of the σ - and π -donor and π -acceptor capabilities of ligands. In this model, the σ - and π -donor properties were evaluated from the molecular electrostatic potential function, V_{MEP} , while the π -acceptor properties were derived using the second-order perturbation theory, and we adopt these two parameters (V_{MEP} and V_b) for factorisation of the electronic structure of ligands. The ‘lattice’ contribution of ligands to p.q.s. can be roughly proportional to

$$C = Z_{\text{LCA}}/r^3 \quad (3)$$

Where Z_{LCA} is the LCA charge, and r is the probe atom–LCA distance. The steric factors of ligands were evaluated using the cone angle conception.⁶ The final expression for evaluating the p.q.s. values is the following:

$$\text{p.q.s. (calc.)} = a_0 + a_1 V_{\text{MEP}} + a_2 V_b + a_3 C + a_4 T, \quad (4)$$

where T is the cone angle of the ligand,⁶ C is the ‘lattice’ contribution of LCA and V_{MEP} and V_b were derived as described previously.⁵ The results obtained for 25 ligands are presented in Table 1. The final parameters are $a_0 = -0.96172 \text{ mm s}^{-1}$, $a_1 = -0.03752 \text{ mm s}^{-1} \text{ eV}^{-1}$, $a_2 = -0.20849 \text{ mm s}^{-1} \text{ eV}^{-1}$, $a_3 = -0.75555 \text{ mm s}^{-1} \text{ au}^{-1}$, $a_4 = 0.0057 \text{ mm s}^{-1} \text{ deg}^{-1}$ with the correlation coefficient 0.995 for 25 compounds with a root-mean-square error of 0.018 mm s^{-1} . Our correlation line combines different classes of ligands (alkyl- and arylamines, six- and five-membered heterocycles, isonitriles, carbon monoxide, sulfides,

phosphines and phosphites). An increase in the σ - and π -donor or π -acceptor properties of ligands leads to a decrease in the p.q.s. values, this is consistent with qualitative assumptions.^{3,4} As can be concluded from the results of the above regression analysis, an increase in the steric effect of ligands leads to an increase in the p.q.s. value, in accordance with experimental data.^{3(f)} It is interesting that the cone angle is more important for most of the ligands, as compared to V_{MEP} and V_b , which are important only for ligands with strong σ -donor and/or π -acceptor properties, while C is important only for phosphorus-containing ligands with a good agreement with recent DFT calculations. The values of Δ , both calculated from equations (1) and (2) and experimental, for macFeL_2 and $\text{macFeL}'\text{L}''$ complexes, where mac is pc^{2-} , Hdmg^- or Hnx^- , are presented in Table 2. The correlation coefficient $r^2 = 0.975$ with a root-mean-square error of 0.10 mm s^{-1} for 27 complexes has been observed. The computed Δ values are lower than the experimental data. Probably, this fact suggests that the cone angle conception for pc^{2-} and Hnx^- is not a good estimation. Really, the recomputed Δ values, in which experimental p.q.s. values were used for macrocyclic ligands, are in better agreement with experimental Δ values (Table 2).

Thus, we have proposed a new technique for the estimation of generally usable ligand p.q.s. values. The correlation coefficient obtained with a small root-mean-square error for p.q.s. values indicates that the proposed technique is a simple and correct tool for predicting Δ values in inorganic and organometallic compounds.

VNN is grateful to JSPS for financial support (grant no. P98418). This work was partially supported by INTAS (grant no. 97-0791).

References

- 1 *Chemical Mössbauer Spectroscopy*, ed. R. H. Herber, Plenum Press, New York, 1984.
- 2 (a) M. Grodzicki, H. Flint, H. Winkler, A. F. Walker and A. X. Trautwein, *J. Phys. Chem.*, 1997, **101**, 4202; (b) R. H. Havlin, N. Godbout, R. Salzmann, M. Wojdelski, W. Arnold, C. E. Schulz and E. Oldfield, *J. Am. Chem. Soc.*, 1998, **120**, 3144.
- 3 (a) L. M. D. R. S. Martins, M. T. Duarte, A. M. Galvao, C. Resende, A. J. L. Pombeiro, R. A. Henderson and D. J. Evans, *J. Chem. Soc., Dalton Trans.*, 1998, 3311; (b) G. M. Bancroft, *Coord. Chem. Rev.*, 1973, **11**, 47; (c) J. M. Bellerby, M. J. Mays and P. L. Sears, *J. Chem. Soc., Dalton Trans.*, 1976, 1232; (d) D. J. Evans, M. Jimenez-Tenorio and G. J. Leigh, *J. Chem. Soc., Dalton Trans.*, 1991, 1785; (e) G. M. Bancroft and E. T. Libbey, *J. Chem. Soc., Dalton Trans.*, 1973, 2103; (f) J. Silver, *Inorg. Chim. Acta.*, 1991, **184**, 235.
- 4 A. Y. Nazarenko, E. V. Polshin and Ya. Z. Voloshin, *Mendeleev Commun.*, 1993, 45.
- 5 S. S. Fielder, M. C. Osborne, A. B. P. Lever and W. J. Pietro, *J. Am. Chem. Soc.*, 1995, **117**, 6990.
- 6 C. A. Tolman, *Chem. Rev.*, 1977, **77**, 313.
- 7 M. J. S. Dewar, E. G. Zebisch, E. F. Healy and J. J. P. Stewart, *J. Am. Chem. Soc.*, 1985, **107**, 3902.
- 8 M. J. S. Dewar and W. Thiel, *J. Am. Chem. Soc.*, 1977, **99**, 4839.
- 9 A. J. Gordon and R. A. Ford, *The Chemist's Companion*, Wiley, New York, 1972.
- 10 R. P. Brent, *Algorithms for Minimization without Derivatives*, Prentice–Hall, Englewood Cliffs, New Jersey, 1973.
- 11 (a) B. W. Dale, R. J. R. Williams, P. R. Edwards and C. E. Johnson, *Trans. Farad. Soc.*, 1968, **64**, 620; (b) R. Taube, *Pure Appl. Chem.*, 1974, **38**, 427; (c) V. N. Nemykin, V. Y. Chernii, E. V. Polshin and S. V. Volkov, *Ukr. Khim. Zh.*, 1997, **63**, 75 (in Russian); (d) M. Hanack, U. Keppeler, A. Lange, A. Hirsch and R. Dieing, in *Phthalocyanines: Properties and Applications*, eds. A. B. P. Lever and C. C. Leznoff, VCH Publishers, New York, 1989, vol. 1, part 2.

Table 2 Calculated and experimental Δ values for macFeL_2 and $\text{macFeL}'\text{L}''$ complexes.

Complex	Δ (calc.)	Δ (calc.) ^a	Δ (exp.) [ref.]
$\text{pcFe}(n\text{-prnh}_2)_2$	1.88	2.00	1.97 [1(b)]
$\text{pcFe}(s\text{-bunh}_2)_2$	1.96	2.08	2.04 [1(c)]
$\text{pcFe}(t\text{-bunh}_2)_2$	2.32	2.40	2.38 [1(a)]
$\text{pcFe}(\text{phnh}_2)_2$	1.96	2.08	2.11 [1(b)]
$\text{pcFe}(\text{mepip})_2$	2.20	2.28	2.28 [1(c)]
$\text{pcFe}(\text{mph})_2$	2.20	2.28	2.31 [1(c)]
$\text{pcFe}(\text{dabco})_2$	2.68	2.76	2.89 [1(d)]
$\text{pcFe}(14\text{en})_2$	1.80	1.84	1.84 [1(c)]
$\text{pcFe}(\text{py})_2$	1.68	1.80	2.02 [1(d)]
$\text{pcFe}(4\text{chopy})_2$	1.70	1.84	1.84 [1(d)]
$\text{pcFe}(4\text{ohpy})_2$	1.72	1.84	1.80 [1(d)]
$\text{pcFe}(\text{tz})_2$	1.64	1.80	1.79 [1(d)]
$\text{pcFe}(\text{meim})_2$	1.60	1.72	1.71 [this work]
$\text{pcFe}(\text{pzl})_2$	1.68	1.80	1.79 [this work]
$\text{pcFe}(\text{tzl})_2$	1.64	1.80	1.73 [this work]
$\text{pcFe}(t\text{-bunc})_2$	0.96	1.12	0.80 [1(d)]
$\text{pcFe}(\text{phnc})_2$	0.96	1.12	0.68 [1(d)]
$\text{pcFe}(\text{co})_2$	0.72	0.84	0.82 [1(d)]
$\text{pcFe}(\text{pet}_3)_2$	1.40	1.48	1.47 [1(d)]
$\text{pcFe}(\text{pobu}_3)_2$	0.96	1.08	1.09 [this work]
$\text{pcFe}(\text{tht})_2$	2.12	2.24	2.20 [1(d)]
$\text{pcFe}(\text{nh}_3)(\text{co})$	1.12	1.24	1.02 [1(d)]
$\text{pcFe}(\text{py})(\text{co})$	1.20	1.34	1.19 [1(d)]
$(\text{Hnx})_2\text{Fe}(n\text{-bunh}_2)_2$	1.92	1.84	1.83 [1(a)]
$(\text{Hnx})_2\text{Fe}(\text{py})_2$	1.72	1.64	1.79 [1(a)]
$(\text{Hnx})_2\text{Fe}(4\text{chopy})_2$	1.76	1.68	1.68 [1(a)]
$(\text{Hdmg})_2\text{Fe}(\text{py})_2$	1.72	1.64	1.67 [1]

^aCalculated using p.q.s. values of -0.96 and -0.92 for pc^{2-} and Hnx^- , respectively.

Received: 22nd September 1999; Com. 99/1535