

CONTENTS

Altered Response of Insulin-Like Growth Factor–Binding Protein 1 to Nutritional Deprivation in Type 2 Diabetes Mellitus <i>Moirá S. Lewitt, Kerstin Hall, Peter Bang, and Kerstin Brismar</i>	275
Correlation Between the Adiponectin-Leptin Ratio and Parameters of Insulin Resistance in Patients With Type 2 Diabetes <i>Minoru Inoue, Eisuke Maehata, Masao Yano, Matsuo Taniyama, and Seiji Suzuki</i>	281
Expression Levels and Functional Aspects of the Hyaluronan Receptor CD44. Effects of Insulin, Glucose, Insulin-like Growth Factor I, or Growth Hormone on Human Arterial Smooth Muscle Cells <i>Kirsten Schultz, Lars Melholt Rasmussen, and Thomas Ledet</i>	287
GPR40 Gene Arg211His Polymorphism May Contribute to the Variation of Insulin Secretory Capacity in Japanese Men <i>Takeo Ogawa, Hiroshi Hirose, Kiichi Miyashita, Ikuo Saito, and Takao Saruta</i>	296
Propylthiouracil and Thiamazole Do Not Alter In Vitro Neutrophil Oxidative Burst <i>Elisa Maria de Sousa Russo-Carbolante, Ana Elisa Caleiro Seixas Azzolini, Ana Christina Morseli Polizello, Ana Isabel de Assis-Pandochi, and Yara Maria Lucisano-Valim</i>	300
Red Wine Consumption Improves Insulin Resistance But Not Endothelial Function in Type 2 Diabetic Patients <i>Raffaele Napoli, Domenico Cozzolino, Vincenzo Guardasole, Valentina Angelini, Emanuela Zarra, Margherita Matarazzo, Antonio Cittadini, Luigi Saccà, and Roberto Torella</i>	306
The Effects of Rosiglitazone and Metformin on the Plasma Concentrations of Resistin in Patients With Type 2 Diabetes Mellitus <i>Hye Seung Jung, Byung-Soo Youn, Young Min Cho, Kang-Yeol Yu, Hong Je Park, Chan Soo Shin, Seong Yeon Kim, Hong Kyu Lee, and Kyong Soo Park</i>	314
Carotid Artery Intima-Media Thickness and Lipid Profile in Adults With Growth Hormone Deficiency After Long-Term Growth Hormone Replacement <i>Débora Vieira Soares, Luciana Diniz Carneiro Spina, Rosane Resende de Lima Oliveira Brasil, Eliane Maria Carvalho da Silva, Priscila Marise Lobo, Elizabeth Salles, Cláudia Medina Coeli, Flávia Lúcia Conceição, and Mario Vaisman</i>	321
Altered Relationship Between Body Fat and Plasma Adiponectin in End-Stage Renal Disease <i>Tetsuo Shoji, Kayo Shinohara, Sawako Hatsuda, Eiji Kimoto, Shinya Fukumoto, Masanori Emoto, Hideki Tahara, Hidenori Koyama, Eiji Ishimura, Takami Miki, Tsutomu Tabata, and Yoshiki Nishizawa</i>	330
Isotopomer Spectral Analysis of Intermediates of Cholesterol Synthesis in Patients With Cerebrotendinous Xanthomatosis <i>Jacob J. Clarenbach, Bernhard Lindenthal, Maria Theresa Dotti, Antonio Federico, Joanne K. Kelleher, and Klaus von Bergmann</i>	335
High Serum Pentosidine Concentrations Are Associated With Increased Arterial Stiffness and Thickness in Patients With Type 2 Diabetes <i>Noboru Yoshida, Ki-ichi Okumura, and Yoshimasa Aso</i>	345
Relationship Between S-Adenosylmethionine, S-Adenosylhomocysteine, Asymmetric Dimethylarginine, and Endothelial Function in Healthy Human Subjects During Experimental Hyper- and Hypohomocysteinemia <i>Sagar Doshi, Ian McDowell, Jonathan Goodfellow, Sally Stabler, Rainer Boger, Robert Allen, Robert Newcombe, Malcolm Lewis, and Stuart Moat</i>	351
The Metabolism of Apolipoproteins (a) and B-100 Within Plasma Lipoprotein (a) in Human Beings <i>Jennifer L. Jenner, Leo J. Seman, John S. Millar, Stefania Lamon-Fava, Francine K. Welty, Gregory G. Dolnikowski, Santica M. Marcovina, Alice H. Lichtenstein, P. Hugh R. Barrett, Carl deLuca, and Ernst J. Schaefer</i>	361

(Contents continued)

Cellular Insulin Resistance in Epstein-Barr Virus-Transformed Lymphoblasts From Young Insulin-Resistant Japanese Men <i>Toshiyuki Morii, Yoichi Ohno, Hiroshi Hirose, Hiroshi Kawabe, Tsutomu Ogata, Keiichi Hirao, Takashi Eguchi, Tatsuya Maruyama, Yoshihiko Kanno, Matsuhiko Hayashi, Ikuo Saito, and Takao Saruta</i>	370
Soluble E-Selectin, Leptin, Triglycerides, and Insulin Resistance in Nonobese Japanese Type 2 Diabetic Patients <i>Ataru Taniguchi, Mitsuo Fukushima, Yoshikatsu Nakai, Akira Kuroe, Gen Yamano, Takiko Yanagawa, Minako Ohgushi, Michihiro Ohya, Satoru Yoshii, Yoshiro Taki, and Yutaka Seino</i>	376
The K469E Polymorphism of the Intercellular Adhesion Molecule-1 Gene Is Associated With Plasma Fibrinogen Level in Type 2 Diabetes <i>Hisayo Yokoyama, Hideki Tahara, Masanori Emoto, Shigehiko Fujiwara, Takahiro Araki, Kayo Shinohara, Sawako Hatsuda, Takaaki Maeno, Takuhito Shoji, Hidenori Koyama, Tetsuo Shoji, and Yoshiki Nishizawa</i>	381
Effect of Two α -Glucosidase Inhibitors, Voglibose and Acarbose, on Postprandial Hyperglycemia Correlates With Subjective Abdominal Symptoms <i>Tomomi Fujisawa, Hiroshi Ikegami, Kaori Inoue, Yumiko Kawabata, and Toshio Ogihara</i>	387
Changes in Blood Pressure and Plasma Catecholamine Levels During Prolonged Hyperinsulinemia <i>Werner Kern, Achim Peters, Jan Born, Horst L. Fehm, and Bernd Schultes</i>	391
Insulin Sensitivity Indices of Glucose and Free Fatty Acid Metabolism in Obese Children and Adolescents in Relation to Serum Lipids <i>Thomas Reinehr, Wieland Kiess, and Werner Andler</i>	397
Increased Diacylglycerol Acyltransferase Activity Is Associated With Triglyceride Accumulation in Tissues of Diet-Induced Insulin-Resistant Hyperlipidemic Hamsters <i>Adele Casaschi, Geoffrey K. Maiyoh, Khosrow Adeli, and Andre G. Theriault</i>	403
Influence of Calcium-Induced Workload Transitions and Fatty Acid Supply on Myocardial Substrate Selection <i>Antti Ala-Rämi, Mika Ylihautala, Petri Ingman, and Ilmo E. Hassinen</i>	410