

- [3] Lagrand WK, Visser CA, Hermes WT, et al. C-reactive protein as a cardiovascular risk factor. More than an epiphenomenon? *Circulation* 1999;100:96–102.
- [4] Chenillot O, Henny J, Steinmetz J, et al. High sensitivity C-reactive protein: biological variations and reference limits. *Clin Chem Lab Med* 2000;38:1003–11.
- [5] von Eckardstein A, Nofer JR, Assmann G. High density lipoproteins and arteriosclerosis. Role of cholesterol efflux and reverse cholesterol transport. *Arterioscler Thromb Vasc Biol* 2001;21:13–27.
- [6] Cockerill GW, Rye KA, Gamble JR, et al. High-density lipoproteins inhibit cytokine-induced expression of endothelial cell adhesion molecules. *Arterioscler Thromb Vasc Biol* 1995;15:1987–94.
- [7] Pirro M, Siepi D, Lupattelli G, et al. Plasma C-reactive protein in subjects with hypo/hyperalphalipoproteinemias. *Metabolism* 2003;52:432–6.
- [8] Multinational monitoring of trends and determinants in cardiovascular diseases: MONICA Project, manual of operations WHO/MNC 82.2, Nov 1983.

Jaroslav A. Hubacek
Dagmar Bobkova
Zdena Skodova
Vera Lanska
Vera Adamkova

*Institute for Clinical and Experimental Medicine
Cardiovascular Research Centre
Prague, Czech Republic
E-mail address: jaroslav.hubacek@medicon.cz*

DOI of original article 10.1016/j.metabol.2004.06.010
doi:10.1016/j.metabol.2005.06.001

Reply

To the Editor,

We read with interest that the findings of Hubacek et al further confirm ours showing an inverse association between high-density lipoprotein cholesterol (HDL-C) and C-reactive protein (CRP). In our study including 760 men and women, we found a strong inverse association, with CRP levels being 88% higher in those with HDL-C level lower than 1.1 mmol/L (42.5 mg/dL) than in those with HDL-C level higher than 1.6 mmol/L (62 mg/dL). Hubacek et al report about a similar association between CRP levels in white males with HDL-C level higher and lower than 40 mg/dL.

Gunilla Nordin Fredrikson
Jan Nilsson
*Department of Clinical Sciences
Malmö University Hospital
Lund University
SE-205 02 Malmö, Sweden*

doi:10.1016/j.metabol.2005.06.002

Influence of endurance training on energy intake, norepinephrine kinetics, and metabolic rate in older individuals. Poehlman E, Gardner A, Goran M. *Metabolism* 1992;41(September):941–8

Dear Dr Field,

In the above-referenced paper, we presented data purportedly showing that endurance training stimulates an increase in energy intake and resting metabolic rate, which are associated with an increase in sympathetic nervous system activity, as determined by norepinephrine kinetics in older individuals. This study purports to have extended an earlier study (*Am J Physiol* 1991;261:E233–9) where the energy intake levels were obtained from the subjects' diaries. In the referenced study, energy intake was also covertly monitored.

I now wish to report that I intentionally omitted a material data point for the norepinephrine results for 1 of the 7 subjects reported in Figure 4 and the associated text of the referenced *Metabolism* paper to make the association between increased sympathetic nervous system activity and endurance training appear more significant than was actually the case.

I also report that I intentionally omitted norepinephrine data in a 1994 paper (*J Appl Physiol* 1994;76(6):2281–7) that I have also asked to be retracted. Because both the 1992 and 1994 papers relied on results where data were intentionally omitted, the 1991 *J Appl Physiol* paper should not be relied upon.

I take sole responsibility for the intentional omission. My coauthors were unaware of my actions and I now publicly exonerate them. I request that you publish this letter of retraction.

Eric T. Poehlman
*Montreal, Quebec
Canada*

doi:10.1016/j.metabol.2005.04.003