

Erratum

In the article “Low sex hormone-binding globulin is associated with the metabolic syndrome in postmenopausal women” by Weinberg et al in the November 2006 issue

(*Metabolism* 2006;1473-80), the second vertical rule in [Table 3](#) on page 1477 appeared in the wrong place. The corrected table is shown below.

Table 3
ORs and 95% CIs for the metabolic syndrome by tertiles of endogenous sex hormone levels

Hormone	Number with/without metabolic syndrome	Crude OR (95% CI)	Multivariate adjusted OR (95% CI) ^a	Multivariate adjusted OR + BMI (95% CI) ^b	Number with/without metabolic syndrome and BMI <26.7 kg/m ²	Multivariate adjusted OR + BMI ^b among women with BMI <26.7 kg/m ² (95% CI)
Estradiol ^c						
Tertile 1	16/34	Referent	Referent	Referent	8/30	Referent
Tertile 2	34/40	1.8 (0.9-3.8)	2.3 (1.0-5.3)	1.6 (0.7-3.8)	14/32	1.3 (0.4-4.1)
Tertile 3	56/29	4.1 (1.9-8.6)	4.5 (2.0-10.1)	2.0 (0.8-4.8)	5/19	0.4 (0.1-2.0)
Testosterone						
Tertile 1	19/39	Referent	Referent	Referent	8/32	Referent
Tertile 2	30/34	1.8 (0.9-3.8)	1.8 (0.8-4.0)	1.4 (0.6-3.3)	5/26	0.6 (0.1-2.3)
Tertile 3	59/31	3.9 (1.9-7.8)	4.3 (2.0-9.4)	3.2 (1.4-7.3)	14/24	1.6 (0.5-5.4)
SHBG						
Tertile 3	7/36	Referent	Referent	Referent	2/32	Referent
Tertile 2	18/34	2.7 (1.0-7.3)	2.7 (0.9-7.5)	2.5 (0.8-7.3)	7/30	13.7 (0.7-21.3)
Tertile 1	83/34	12.6 (5.1-31.0)	11.6 (4.5-29.8)	7.3 (2.7-19.8)	18/20	0.5 (2.0-56.3)
FAI						
Tertile 1	4/36	Referent	Referent	Referent	2/31	Referent
Tertile 2	18/33	4.9 (1.5-16.0)	5.9 (1.7-20.1)	4.2 (1.2-14.7)	6/28	2.4 (0.4-14.0)
Tertile 3	86/35	22.1 (7.3-66.8)	22.9 (7.3-72.1)	12.6 (3.8-41.6)	19/23	7.7 (1.4-41.4)

Tertiles were based on the distribution in women without the metabolic syndrome.

^a Adjusted for age, smoking, physical activity, alcohol use, and the presence or absence of CVD during follow-up.

^b Adjusted for age, smoking, physical activity, alcohol use, the presence or absence of CVD during follow-up, and BMI.

^c There were 3 subjects missing data for estradiol.