This article was downloaded by: [Tomsk State University of Control Systems and Radio]

On: 19 February 2013, At: 11:41

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered

office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Molecular Crystals and Liquid Crystals Incorporating Nonlinear Optics

Publication details, including instructions for authors and subscription information:

http://www.tandfonline.com/loi/gmcl17

## Liquid Crystal Polymers: Conformation of Deuterated Parts by SANS

L. Noirez  $^a$  , J. P. Cotton  $^a$  , F. Hardouin  $^a$  , P. Keller  $^a$  , M. Lambert  $^a$  , F. Moussa  $^a$  & G. Pepy  $^a$ 

<sup>a</sup> Laboratoire Leon Brillouin. CEA, CEN de Saclay, F-91191, Gif-sur-Yvette, Cedex, France

Version of record first published: 22 Sep 2006.

To cite this article: L. Noirez, J. P. Cotton, F. Hardouin, P. Keller, M. Lambert, F. Moussa & G. Pepy (1990): Liquid Crystal Polymers: Conformation of Deuterated Parts by SANS, Molecular Crystals and Liquid Crystals Incorporating Nonlinear Optics, 180:1, 165-165

To link to this article: <a href="http://dx.doi.org/10.1080/00268949008025797">http://dx.doi.org/10.1080/00268949008025797</a>

### PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Mol. Cryst. Liq. Cryst., 1990, Vol. 180A, pp. 165 Reprints available directly from the publisher Photocopying permitted by license only © 1990 Gordon and Breach Science Publishers S.A. Printed in the United States of America

### Liquid Crystal Polymers: Conformation of Deuterated Parts by SANS

L. NOIREZ, J. P. COTTON, F. HARDOUIN, P. KELLER, M. LAMBERT, F. MOUSSA and G. PÉPY

presented by B. Farnoux, Laboratoire Leon Brillouin, CEA, CEN de Saclay, F-91191 Gif-sur-Yvette Cedex, France

Comb like liquid crystal polymers are polymers on each monomer is grafted a hanging liquid crystal molecule. The bulk state has the classical liquid crystal phases (nematic, smectic); what is the conformation of a chain in these phases is the problem to solve with the help of small angle neutron scattering. The values of the global sizes  $R_{\parallel}$  and  $R_{\perp}$  of a chain (in the direction parallel and perpendicular to the director) are discussed as a function of the chemical species and the temperature. Strange results are discussed in terms of segregation of labelled parts.

#### References

- 1. P. Keller, B. Carvalho, J. P. Cotton, M. Lambert and G. Pépy, J. Physique Lett., 46, 1065 (1985).
- F. Moussa, J. P. Cotton, F. Hardouin, P. Keller, M. Lambert, G. Pépy, M. Mauzac and H. Richard, J. Physique, 48, 1079 (1987).