

Purinoreceptors on Neuroglia

Alexei Verkhrasky · Oleg A. Krishtal ·
Geoffrey Burnstock

Received: 8 February 2009 / Accepted: 24 February 2009 / Published online: 13 March 2009
© Humana Press Inc. 2009

Abstract Purinergic transmission is one of the most ancient and widespread extracellular signalling systems. In the brain, purinergic signalling plays a unique role in integrating neuronal and glial cellular circuits, as virtually every type of glial cell possesses receptors to purines and pyrimidines. These receptors, represented by metabotropic P1 adenosine receptors, metabotropic P2Y purinoreceptors and ionotropic P2X purinoreceptors, control numerous physiological functions of glial cells and are intimately involved in virtually every form of neuropathology. In this essay, we provide an in depth overview of purinoreceptor distribution in two types of CNS glia—in astrocytes and oligodendrocytes—and discuss their physiological and pathophysiological roles.

Keywords ATP · Adenosine · P1 receptors · P2Y receptors · P2X receptors · Astrocytes · Oligodendrocytes

A. Verkhrasky
School of Biological Sciences, The University of Manchester,
1.124 Stopford Building, Oxford Road,
Manchester M13 9PT, UK

A. Verkhrasky (✉)
Institute of Experimental Medicine, ASCR,
Videnska 1083,
Prague 4 142 20, Czech Republic
e-mail: alex.verkhratsky@manchester.ac.uk

O. A. Krishtal
Bogomoletz Institute of Physiology,
Bogomoletz Str. 4,
Kiev-24, Ukraine

G. Burnstock
Autonomic Neuroscience Centre,
Royal Free and University College Medical School,
Rowland Hill Street,
London NW3 2PF, UK

Neuronal–Glial Circuitry

The mammalian brain, evolved through millions of years, is moulded from two functionally distinct cell populations, the electrically excitable neurones and electrically non-excitable glial cells. The neuroglia, the concept of which was introduced by Rudolf Virchow 150 years ago ([1] for historic overview, see [2, 3]), provides a three-dimensional canvas into which the synaptically connected neuronal networks are embedded [4, 5]. The main type of grey matter glia, the astroglia, divides, through the process of “tiling”, the brain parenchyma into relatively independent structural units, determined by the territories occupied by individual astrocytes [6–8]. Within these territories, astrocytes provide structural and functional support to neurones [9, 10], create neuronal–glial–vascular units [11] and enwrap the synaptic contacts making the tripartite synapses [12, 13], characteristic for the central nervous system (CNS). Functional importance of these individual domains acquired experimental attention only very recently, and yet, these local neuronal–glial units may appear to be the basic structural elements of the grey matter, which shape integrative processes in the CNS. Furthermore, the astroglia, being the main cellular element of brain homeostasis, is intimately involved in neuropathology, determining to a very great extent the progress and outcome of various diseases of the CNS [14–17].

Purinergic Signalling System


The purinergic signalling system, which utilises purines and pyrimidines as extracellular messengers [18–20], appeared very early in evolution and became exceptionally widespread in both the animal and plant kingdoms [21, 22].

Indeed, purine- and pyrimidine-mediated information transfer can be found in virtually every type of cell and tissue [23] and throughout every developmental stage [24]. Adenosine 5'-triphosphate (ATP), which is the principal purinergic signaller, is released from cells by several mechanisms, which include exocytosis, diffusion through "maxi" plasmalemmal channels and probably transporters or even lysosomes [20, 25, 26]. In addition, ATP is released from damaged cells, being a universal "danger" signal. The released ATP is rapidly degraded by numerous endonucleotidases [20, 27] that produces a trail of derivatives [adenosine diphosphate (ADP), adenosine monophosphate (AMP) and adenosine], which in turn also act as signalling molecules. At the receiving end, purines and pyrimidines activate several families of purinoceptors' broadly classified as metabotropic P1 adenosine receptors, metabotropic P2Y purinoceptors and ionotropic P2X purinoceptors [18, 28–33]. These receptors alone or in combination are expressed throughout living cells and tissues and mediate a remarkable variety of physiological and pathophysiological reactions.

Purinergic Signalling in Neuronal–Glial Networks

The purinergic signalling system plays a unique role in neuronal–glial interactions, as virtually every type of glial cell, be it the cells of ectodermal/neural origin (astrocytes and oligodendrocytes) or of mesodermal origin (microglia), displays sensitivity to ATP and its analogues (Fig. 1). In the nervous system, ATP is released from neurones, their axons and terminals as well as from neuroglia. Probably the predominant mechanism of neuronal ATP release in the CNS is vesicular [25, 34]. ATP is involved in synaptic transmission in many brain regions [20, 35], as it can be stored and released either on its own or together with other neurotransmitters such as glutamate, γ -aminobutyric acid (GABA), noradrenaline or acetylcholine (ACh). In addition, ATP is released from astrocytes and oligodendrocytes by not yet fully characterised pathways [36–38], which may include exocytosis, or diffusion through maxi-pore forming channels (such as hemichannels, pannexins, volume-sensitive anion channels or P2X₇ receptors). Regulated release of ATP from glial cells may play important roles in

Fig. 1 Omnipresence of purinergic signalling pathways in neuronal–glial circuits in the grey matter. The microarchitecture of the grey matter (as shown in the centre) is defined by astroglial domains, composed of astrocyte, neighbouring blood vessel encompassed by astroglial endfeet and neurones residing within astroglial territory. The microglial cells (each also having its own territory) are constantly surveying these domains spying for damage. ATP and its derivatives act as an extracellular signalling molecule at all levels of communications within neuronal–glial networks. Within the tripartite synapse (I), ATP, released during synaptic transmission, activates astrocytes receptors, which in turn initiate Ca^{2+} signals and Ca^{2+} waves in astroglial syncytium. Astroglial Ca^{2+} signals induce release of ATP, which feeds back to neurones via activation of pre- and postsynaptic P1 and P2 receptors. ATP released from astrocytes (II) triggers and maintains astroglial Ca^{2+} waves. Finally, ATP released from all types of neural cells control activation (III) of microglia



both glial–glial signalling (for example by initiating propagating Ca^{2+} waves [37, 39]) and in glial–neuronal communications (for example regulating synaptic plasticity [40]). Finally, massive release of ATP inevitably accompanies neural tissue damage, being thus ultimately involved in many forms of neuropathology.

Purinergic signalling in neuronal–glial communications has been discussed in many reviews [35, 41–50]; the pathophysiological importance of purinergic signalling for microglial activation was overviewed even more frequently [51–63]. In the present essay, we shall specifically focus on the purinergic receptors expressed in two major types of glial cells of neural origin, in astrocytes and oligodendrocytes, and consider their physiological and pathophysiological relevance.

P1 Receptors

P1 adenosine receptors are classical 7-transmembrane-spanning metabotropic receptors coupled to several families of G_i and G_o proteins. Four types of adenosine receptors (A_1 , A_{2A} , A_{2B} and A_3) with distinct pharmacological and functional properties were cloned [33]. As a rule, the A_1 and A_3 receptors exert an inhibitory effect on adenylyl cyclase (mediated through $G_{i/o}$ proteins), whereas A_{2A} and A_{2B} receptors activate cyclic AMP (cAMP) production via G_s proteins. A_1 and A_3 receptors also regulate phospholipase C (PLC) and thus inositol 1,4,5-trisphosphate (InsP_3) synthesis; in some cells, A_1 receptors were reported to activate K^+ and/or Ca^{2+} channels [33, 64–66].

Astrocytes

All four P1 receptors have been identified in astrocytes (Table 1) using various functional assays applied to both in vitro and in situ preparations [66]. Initially, the functional role for adenosine receptors was demonstrated in electrophysiological experiments on cultured rat astrocytes [67]; this study showed that adenosine hyperpolarised a subpopulation of astroglial cells, and this hyperpolarisation was antagonised by a P1 receptor antagonist 8-phenyltheophylline. Subsequently, adenosine receptors with pharmacological profiles corresponding to A_1 and A_2 types were identified in human foetal astrocytes [68]. In these cells, the A_1 receptors inhibited, whereas A_2 receptors potentiated synthesis of cAMP. In primary cultures of rat astrocytes, A_{2B} receptors stimulated adenylyl cyclase and their activation caused dose-dependent accumulation of cAMP [69]. In the same cultures, stimulation of A_1 receptors in type 1 (but not in type 2) astrocytes lead to an inhibition of cAMP production [70].

Expression of A_1 receptor-specific messenger RNA (mRNA) was demonstrated in rat-cultured astroglia [71]. Activation of A_1 receptors in rat-cultured astrocytes activated PLC; incidentally, this activation was observed only in cultures with high levels of A_1 receptor expression [72]; up-regulation of A_1 receptors synthesis potentiated A_1 -dependent PLC stimulation [71, 73].

In cortical astrocytes, acutely isolated from 4–12-day-old rats, adenosine triggered $[\text{Ca}^{2+}]_i$ responses, which were mediated through InsP_3 -induced Ca^{2+} release and were blocked by the selective A_{2B} antagonist alloxazine [74]. The sensitivity of acutely isolated cells to adenosine was much higher as compared with the same cells maintained in culture, thus indicating modified adenosine receptors expression in the in vitro conditions [74]. In astroglial cultures obtained from neonatal rat forebrains, stimulation of A_1 receptors triggered both intracellular Ca^{2+} release and Ca^{2+} entry and potentiated histamine-induced Ca^{2+} mobilisation [75]. Similarly, adenosine, acting through P1 receptors, triggered $[\text{Ca}^{2+}]_i$ elevation in the majority of astrocytes in acute rat hippocampal slices [76]. In astrocytes from acutely isolated mouse olfactory bulb slices, adenosine, which occurred following enzymatic degradation of ATP released from olfactory nerve terminals, induced $[\text{Ca}^{2+}]_i$ elevation via activation of A_{2A} receptors [77].

In cultured mouse astrocytes adenosine triggered $[\text{Ca}^{2+}]_i$ transients in ~85% of cells via activation of A_3 receptors as judged by a specific sensitivity to A_3 receptor antagonists [78]; incidentally, treatment with guanosine induced $[\text{Ca}^{2+}]_i$ responses as well, and these responses are likely to be also mediated through A_3 receptors.

In primary cultured astroglia, adenosine was shown to modulate the amplitude and/or kinetics of $[\text{Ca}^{2+}]_i$ transients originated through InsP_3 -induced Ca^{2+} release initiated by activation of metabotropic glutamate receptors, muscarinic ACh receptors or P2Y receptors [79–84]. The signalling systems involved, however, were different; in some cases, intracellular Ca^{2+} mobilisation was promoted through activation of A_1 receptors [79, 80, 82]; in others, Ca^{2+} release was up-regulated through A_{2B} receptors [81, 83], whereas stimulation of A_1 receptors suppressed the plateau phase of the Ca^{2+} signal, possibly through inhibition of store-operated Ca^{2+} entry [81]. Similarly, activation of A_1 receptors suppressed sustained Ca^{2+} influx following opening of P2X₇ receptors in cultured cortical astrocytes [85].

Astroglial adenosine receptors are also coupled with astrocyte-dependent regulation of extracellular glutamate. Activation of A_{2A} receptors in hippocampal astrocytes reduced glutamate uptake via inhibition of GLT-1 transporter and resulted in glutamate release from astrocytes through $[\text{Ca}^{2+}]_i$ and a protein kinase A-dependent pathway [86, 87]. This in turn potentiated neuronal activity in

Table 1 P1 adenosine receptors in neuroglia

Receptor type	Experimental preparation/species/technique	Properties/function	References
Astrocytes			
P1 receptors ^a	Cell culture/rat/electrophysiology	Cell hyperpolarization	[67]
	Hippocampal slices/Ca ²⁺ imaging	Initiation of [Ca ²⁺] _i transients due to Ca ²⁺ release from the ER	[76]
A ₁ receptor	Cell culture/rat	Inhibition of cAMP production	[69]
	Cell culture/rat	Activation of PLC	[71–73]
	Cell culture/rat forebrain/Ca ²⁺ imaging	Initiation of [Ca ²⁺] _i transients due to Ca ²⁺ release from the ER, activation of Ca ²⁺ entry and potentiation of histamine-induced Ca ²⁺ release	[75]
	Cell culture/rat cortex/Ca ²⁺ imaging	Potential of ACh-induced Ca ²⁺ signalling	[82]
	Cell culture/rat hippocampus/Ca ²⁺ imaging	Potential of glutamate (mGluRs)-induced Ca ²⁺ signalling	[79]
A _{2A} receptor	Cell culture/rat cortex/Ca ²⁺ imaging	Inhibition of P2X ₇ -mediated Ca ²⁺ influx	[85]
	Acute slices/mouse olfactory bulb/Ca ²⁺ imaging	Initiation of [Ca ²⁺] _i transients due to Ca ²⁺ release from the ER	[77]
	Cell culture/Slices/rat hippocampus/electrophysiology	Inhibition of astroglial glutamate transporter GLT-1 and activation of astroglial glutamate release	[86]
	Cell culture/rat striatum	Inhibition of astrogliosis	[90]
	Cell culture/rat	Inhibition of iNOS and NO production	[97]
A _{2B} receptor	Cell culture/rat	Stimulation of cAMP production	[69]
	Acutely isolated cells/rat cortex/Ca ²⁺ imaging	Initiation of [Ca ²⁺] _i transients due to Ca ²⁺ release from the ER	[74]
	Cell culture/rat cerebellum/Ca ²⁺ imaging	Potential of glutamate (P2Y)-induced Ca ²⁺ signalling	[83]
A ₃ receptor	Cell culture/mouse	Initiation of [Ca ²⁺] _i transients due to Ca ²⁺ release from the ER	[78]
	Cell culture/rat	Activation of apoptosis at intense stimulation; protective effects at low/moderate stimulation	[93–95]
	Cell culture/mouse	Up-regulation of CCL2 cytokine synthesis	[100]
Oligodendrocytes			
P1 receptors	Neuronal–glial co-culture/mouse/Ca ²⁺ imaging	Initiation of [Ca ²⁺] _i transients following axonal firing, inhibition of oligodendroglial precursors proliferation and promotion of differentiation and myelination	[101, 103]

^a This refers to experiments where P1 receptors subtypes were not identified

hippocampus due to an increase in glutamate concentration in the synaptic zones [86].

P1 Receptors in Neuropathology Brain injury results in a massive release of ATP (a substantial part of which is rapidly converted into adenosine) and adenosine; the latter can significantly increase in the cytosol upon conditions of cell stress and hypoxia. As a consequence, adenosine concentrations can attain 10–50 μ M [88]. Direct injections of adenosine or the adenosine analogue 5'-(*N*-cyclopropyl)-carboxamidoadenosine into the rat cortex triggered prominent reactive astrogliosis, which was antagonised by the A₂ receptor blocker 1,3-dipropyl-7-methylxanthine [89]. In primary rat striatal astroglial cultures, pharmacological blockade of A_{2A} receptors inhibited astrogliosis induced by basic fibroblast growth factor [90]. Short-term (3 h) treatment of cultured human astrocytes and astrocytoma cells with the

astrogliosis-promoting factor tumour necrosis factor (TNF)- α caused phosphorylation of A_{2B} receptors, which reduced their positive coupling to adenylyl cyclase and in turn suppressed A_{2B} receptor-mediated cAMP production [91, 92].

Adenosine receptors are also involved in the regulation of astroglia survival and death. Selective activation of A₃ but not A₁/A₂ receptors triggered apoptosis in primary cultured rat astrocytes and in C6 glioma cell line through down-regulation of BCL2 expression and increase in caspase 3 activity [93]. In fact, activation of A₃ receptors modulated astroglial cell death in a concentration-dependent manner; low doses of A₃ agonists induced reorganisation of the cytoskeleton and increased cell survival, whereas over-stimulation of A₃ receptors induced astroglial death [94, 95].

Adenosine also exerted a general glio-protective action in conditions of glucose deprivation by maintaining mito-

chondrial potential and ATP synthesis [96]. Activation of A_{2A} receptors was reported to inhibit inducible nitric oxide (NO) synthase expression and NO production in cultured rat astrocytes [97], which also may contribute to glio-protection. Specific activation of A_1 receptors had an anti-apoptotic effect in primary astroglial cultures treated with staurosporine; this effect was mediated through the phosphatidylinositol 3-kinase (PI3K) pathway [98]. Both A_1 and A_3 receptors are involved in protection of astrocytes against hypoxic conditions. Hypoxia-induced cytotoxicity was much more pronounced in astroglial cultures obtained from A_1 or A_3 receptors knockout mice [99]. Stimulation of A_3 receptors also increased the synthesis of neuroprotective cytokine CCL2 in cultured mouse astroglia [100].

Oligodendrocytes

All four P_1 adenosine receptors were identified in cultured oligodendrocytes (Table 1) and their precursors at the mRNA level [101]. Adenosine receptors play an important role in differentiation of cells from the oligodendroglial lineage. In experiments in vitro, in co-cultures of sensory neurones and oligodendroglial precursor cells (OPCs), adenosine and ATP released during action potentials triggered Ca^{2+} signals in immature oligodendrocytes (represented by both NG2-positive cells and O4-positive OPCs). This action was mediated through both P_2Y and P_1 adenosine receptors [101] and allowed OPCs to detect electrical activity of non-myelinated nerve fibres. Activation of adenosine receptors, however, had another specific action; it inhibited OPCs proliferation, promoted their differentiation and initiated myelination, thus constituting a signalling loop between axon activity and oligodendrocyte function [101]. Incidentally, myelination was also promoted by the astroglia-derived cytokine, leukaemia inhibitory factor, which was released following stimulation of astroglial P_2Y receptors [102]. All in all, ATP and adenosine released from axons represent a specific signalling system coordinating axonal/oligodendroglial development and interaction [103].

Adenosine receptors are also involved in various forms of oligodendrocyte pathology. Genetic deletion of A_1 adenosine receptors triggered severe demyelination and provoked the progressive-relapsing form of experimental allergic encephalomyelitis (EAE), which is generally considered to be a model for multiple sclerosis (MS) [104]. A substantial part of the pathology was associated with activation of microglia, although direct effects on oligodendrocytes can also play a relevant role. Similarly, sustained activation of A_1 receptors in early postnatal brain reduces expression of myelin basic protein and triggers white matter damage and ventriculomegaly [105].

P2X Receptors

Ionotropic ATP receptors, classified as P2X in 1985 [29, 32] are represented by archetypical ligand-gated cationic ($Na^+/K^+/Ca^{2+}$) channels, assembled in trimeric form by different subunits [106–108]. These subunits, encoded by distinct genes, are classified $P2X_1$ to $P2X_7$ according to historical order of cloning [31, 107]. Receptors, formed through homo- or heteromeric assembly of $P2X_1$ to $P2X_6$ subunits (so far, homomeric composition was shown for $P2X_{1-5}$ subunits, whereas $P2X_6$ subunits apparently cannot oligomerise; heteromeric compositions are represented by $P2X_{1/2}$, $P2X_{1/4}$, $P2X_{1/5}$, $P2X_{2/3}$, $P2X_{2/6}$ and $P2X_{4/6}$ channels), are activated by low micromolar ATP concentrations [20, 35, 109, 110]. In contrast $P2X_7$ receptors are much less sensitive to ATP (their full activation is reached at mM ATP concentrations) and demonstrate several properties, which readily distinguish them from other P2X receptors [110–112]. All P2X receptor subunits are expressed in the nervous system, although their expression varies in different regions of the peripheral nervous system and CNS.

Astrocytes

Surprisingly, our knowledge on the functional expression of P2X receptors in astrocytes is very limited (Table 2). Expression of various P2X subunits on the transcriptional level was found in several astroglial preparations. Studies on primary cultured rat cortical astrocytes identified the expression of mRNA for $P2X_{1-5}$ and $P2X_7$ receptors [113, 114]. In tissue extracts from rat nucleus accumbens, reverse transcriptase polymerase chain reaction (RT-PCR) revealed expression of all seven P2X mRNAs [115]. In freshly isolated retinal Müller cells, $P2X_3$, $P2X_4$, $P2X_5$ but not $P2X_7$ mRNAs were identified [116]. In contrast, $P2X_7$ receptor-specific mRNA was identified in Müller cells isolated from human retina [117]. In acutely isolated mouse cortical astrocytes, only $P2X_1$ - and $P2X_5$ -specific mRNAs were found [118]. Astroglial localisation of some P2X subunits was also corroborated by immunohistochemistry. In nucleus accumbens (where all seven receptors were present at transcriptional level), immunofluorescence revealed that only $P2X_{2-4}$ receptors were co-localised with glial fibrillary acidic protein (GFAP)-labelled astroglial profiles [115]. Mechanical lesion triggered up-regulation of $P2X_{1-4}$ and $P2X_7$ immunofluorescence in nucleus accumbens astrocytes [115]. Immunoreactivity for $P2X_1$ and $P2X_2$ receptors was detected in astroglial cells in cerebellum [119, 120]; in a similar way, $P2X_2$ receptors were found in spinal cord astrocytes [121], whereas $P2X_4$ receptors were identified in astrocytes from the brainstem [122]. In hippocampal astrocytes, immunostaining revealed expression of $P2X_{1-4}$, $P2X_6$ and $P2X_7$ subunits [123].

Table 2 P2X receptors in neuroglial cells

Receptor type	Experimental preparation/species/technique	Properties/function	References
Astrocytes			
P2X ₁₋₅ , P2X ₇	Cell culture/rat cortex/RT-PCR	Specific mRNAs detected	[113, 114]
P2X ₁₋₇	Tissue extracts/rat nucleus accumbens/ RT-PCR	Specific mRNAs detected	[115]
P2X ₂₋₄	Rat nucleus accumbens/immunostaining	Immunoreactivity detected	[115]
P2X ₃₋₅	Acutely isolated Müller cells/rat retina/ RT-PCR	Specific mRNAs detected	[116]
P2X ₁ , P2X ₅	Acutely isolated cells/mouse cortex/RT-PCR	Specific mRNAs detected	[118]
P2X ₁ , P2X ₂	Rat, guinea pig cerebellum/immunostaining	Immunoreactivity detected	[119, 120]
P2X ₄	Rat brainstem/immunostaining	Immunoreactivity detected	[122]
P2X ₁₋₄ , P2X ₆ , P2X ₇	Rat hippocampus/immunostaining	Immunoreactivity detected	[123]
P2X _{1/5}	Acute slices/mouse/electrophysiology	Specific currents through P2X _{1/5} heteromeric receptors	[118]
P2X?, P2X ₇ ?	Acutely isolated optic nerve/mouse/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with receptor mediated Ca ²⁺ entry	[133, 145]
P2X ₇	Acutely isolated Müller cells/human retina/ electrophysiology	Specific mRNAs and immunoreactivity as well as currents through P2X ₇ receptors were detected	[144]
P2X ₇	Cell culture/rat/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with receptor mediated Ca ²⁺ entry	[85, 113, 143]
P2X ₇	Cell culture/mouse cortex/electrophysiology	Specific currents through P2X ₇ receptors; activation of P2X ₇ receptors resulted in release of excitatory amino-acids	[138]
P2X ₇	Cell culture/mouse	Activate synthesis of endocannabinoid 2-arachidonoylglycerol	[150]
P2X ₇	Cell culture/human	Stimulation of NO production	[139]
P2X ₇	Cell culture/human	Regulation of NF-κB signalling	[142]
P2X ₇	Cell culture/rat	Up-regulation of P2Y receptors expression	[157]
P2X ₇	Cell culture/rat	Down-regulation of aquaporin-4 expression	[158]
Oligodendrocytes			
P2X _{1,2,3,4,7}	Cell culture/rat/Western blot	Specific proteins detected	[161, 162]
P2X ₇	Cell culture/rat/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with receptor mediated Ca ²⁺ entry	[162]
P2X ₇	Cell culture/rat optic nerve/electrophysiology	Specific currents through P2X ₇ receptors	[165]

At a functional level, however, the activity and role of astroglial P2X receptors remains virtually uncharacterized. ATP-induced depolarisation and membrane currents with accompanying [Ca²⁺]_i rises were recorded from cultured astrocytes [124, 125], although the subunit composition of underlying receptors was not investigated. In hippocampal astrocytes, voltage- and concentration-clamped in slices or in isolation, an exhaustive series of experiments failed to identify P2X-mediated currents [126], despite immunohistochemical evidence suggestive of astroglial expression of P2X subunits [123]. Similarly, ATP-induced currents were not observed in Bergmann glial cells in acute cerebellar slices [127]. Nonetheless, absence of ATP-induced currents in the in situ experiments is not conclusive, as complex geometry of glial cells, diffusional barriers and rapid degradation of ATP in slice tissue may prevent detection of functional responses.

In acutely isolated cortical astrocytes, P2X_{1/5} heteromeric receptor-mediated currents were discovered and characterised [118]. The P2X_{1/5} heteromeric receptors were initially described in heterologous expression systems [128–131]; surprisingly, cortical astrocytes remain the only “real” cells where operational P2X_{1/5} receptors were hitherto found. The P2X_{1/5} combination exhibits several unique features, which include a very high sensitivity to ATP (currents are activated at nM ATP concentrations), biphasic kinetics with distinct peak and steady-state components and very little desensitisation in response to the repetitive agonist applications. Using these P2X_{1/5} receptors, cortical astrocytes therefore are able to detect extremely low levels of extracellular ATP.

In acutely isolated optic nerves, ATP triggered large [Ca²⁺]_i transients in astrocytes, which, at least in part, were mediated through Ca²⁺ influx; these [Ca²⁺]_i responses demonstrated

sensitivity to the P2X receptor antagonist NF023 and could be mimicked by the P2X agonist α,β -methylene ATP ($\alpha\beta$ -meATP); yet, the subunit composition of the underlying P2X receptors remains unknown [132, 133].

Functional roles of astroglial P2X₇ receptors deserve special attention. The P2X₇ receptors are unique in their low ATP sensitivity (in all probability, it is ATP⁴⁻ that acts as a true agonist), almost complete absence of desensitisation and ability to produce large transmembrane pores upon intense stimulation [31, 110, 112]. The mechanism of the pore formation remains obscure [111]; it may involve the dilatation of the channel or activation of other proteins (such as, for example, pannexins [134]) closely associated with P2X₇ receptors. Expression and distribution of P2X₇ receptors in healthy brain remains controversial; initial experiments using *in situ* hybridization found that specific mRNA was restricted to the ependymal layer of the third ventricle [135]. Subsequent RT-PCR studies, however, detected P2X₇ mRNA in many areas of the brain including hippocampus, cortex and brain stem (see [112] for detailed account). Similarly, immunoreactivity for P2X₇ receptors was demonstrated in hippocampus, medulla oblongata, cerebellum, thalamus and amygdala [112]. At the same time, the specificity of many antibodies used for P2X₇ receptors immunostaining remains far from ideal [136], thus making many previous observations questionable. A recent *in-depth* analysis of the cellular distribution of P2X₇ mRNA in the rat brain using isotopic *in situ* hybridization found its presence in microglia, oligodendrocytes and neurones in many brain areas; yet it failed to detect any presence of P2X₇ mRNA in astroglia [137].

There is ample evidence for P2X₇ receptor expression in cultured astrocytes at both the transcriptional and protein levels [113, 114, 138–143]. Immunoreactivity for P2X₇ receptors was also reported for freshly isolated astrocytes and astrocytes in brain slices [123, 143]. On a functional level, experiments on cultured astrocytes demonstrated both P2X₇-mediated Ca²⁺ signalling [85, 113, 144] and characteristic P2X₇ ion currents [138]. Similarly, P2X₇ currents were identified in Müller cells freshly isolated from human retina [145]. In isolated optic nerve, activation of P2X₇ receptors underlie Ca²⁺ influx and astroglial release of ATP, which subsequently acted upon neighbouring glial cells [133, 146]. This astroglia-originated signalling was absent in a P2X₇ knockout model [133].

Activation of P2X₇ receptors in cultured astrocytes has numerous functional consequences. First and foremost, opening of P2X₇ channels triggers release of gliotransmitters glutamate, GABA, ATP and associated purines through exocytosis, P2X₇ associated transmembrane pore or through Cl⁻/HCO₃⁻-dependent mechanism of an as yet unidentified nature [111, 138, 144, 147, 148]. In hippo-

campal astrocytes, prolonged activation of P2X₇ receptors lead to a sustained glutamate release, which obviously may have pathological relevance [149]. High concentrations of ATP, acting most likely through P2X₇ receptors, were also shown to remarkably (~60 times) increase production of endocannabinoid 2-arachidonoylglycerol in cultured astroglia [150]. Furthermore, stimulation of P2X₇ receptors in cultured astrocytes or astroglial cell lines was shown to affect other signalling pathways, for example, modulate release of TNF- α [151], stimulate NO production [139, 152], induce phosphorylation of AKT [153] and p38MAPK/ERK1/ERK2 [154], stimulate transmembrane transport of NADH [155] and regulate NF- κ B signalling [142]. Stimulation of P2X₇ receptors increased production of lipid mediators of inflammation cysteinyl leukotrienes, this action being mediated through P2X₇-mediated Ca²⁺ signalling [156]. Activation of P2X₇ receptors in the astroglial cell line RBA-2 rapidly decreased glutamate uptake via Na⁺-dependent transporter and reduced expression and activity of glutamine synthetase [157]. Furthermore, P2X₇ receptors are involved in the control of expression of other purinoceptors and channels; in particular, P2X₇ stimulation up-regulates expression of P2Y₂ receptors [158] and down-regulates expression of aquaporin-4 [159] in cultured rat astrocytes.

Nonetheless, the crucial data on the functional expression of P2X₇ receptors in astroglial cells in undisturbed grey matter are yet to be obtained. Indeed, astrocytes in tissue culture are certainly different from the *in vivo* state, as the procedure for isolation triggers astrogliosis, which launches various programmes of functional remodelling, including, most likely, changes in P2X₇ receptors expression. There are certain hints that brain injury does induce expression of P2X₇ channels. For example, P2X₇ immunoreactivity in nucleus accumbens was observed only after mechanical damage [115], similarly focal cerebral ischemia resulted in an appearance of astroglial P2X₇ receptors in the rat cortex [160]. Vitreoretinopathy resulted in a significant increase in P2X₇ current density in freshly isolated human Müller cells [161]. The immunoreactivity for P2X₇ receptors was also found in reactive astrocytes from brain autopsies obtained from MS patients [139].

In conclusion, P2X₇ receptors are associated with astroglial responses to brain lesions and most likely constitute a part of global functional remodelling, which accompanies reactive astrogliosis. In this process, P2X₇ receptors play an important role regulating both pathologically relevant signalling events (for example, underlying massive Ca²⁺ influx or regulation of various kinases) and production and release of numerous inflammatory factors. The full pathological profile of astroglial P2X₇ receptors is still to be uncovered, although they might be considered as potentially important therapeutic targets.

Oligodendrocytes

Oligodendroglial precursor cells in purified postnatal cultures expressed P2X_{1,2,3,4,7} proteins [162, 163]. There is little evidence, however, about functional expression of P2X_{1–6} receptors in both OPCs and mature oligodendrocytes (Table 2). In the isolated optic nerve, the broad agonist of P2X receptors $\alpha\beta$ -meATP triggered a small $[Ca^{2+}]_i$ elevation, thus suggesting possible involvement of P2X_{1–6} receptors [164]. In contrast, in oligodendrocytes from corpus callosum slices, ATP failed to activate measurable currents [165].

At the same time, P2X₇ receptors may be operational in oligodendrocytes from other areas of the CNS and in white matter tracts. For example, functional P2X₇ receptors were found in the cells of oligodendroglial lineage in vitro. In cultured OPCs, the specific P2X₇ agonist 2',3'-O-(benzoyl-4-benzoyl)-ATP (BzATP) triggered large Ca^{2+} transients, which were effectively inhibited by the P2X₇ receptor agonist oxidised ATP (oxATP) [163]. In cultured oligodendrocytes from optic nerve ATP in high concentrations (EC₅₀, ~8.8 mM) and BzATP (EC₅₀, ~0.5 mM) triggered sustained inward currents. These currents were potentiated in divalent cation-free extracellular solutions and were inhibited by oxATP. In addition high concentrations of ATP and BzATP induced a rapid increase in $[Ca^{2+}]_i$, which was almost exclusively dependent on transmembrane Ca^{2+} entry [166]. These data taken together are indicative of activation of P2X₇ receptors [166]. In addition P2X₇ receptor immunoreactivity was detected in oligodendrocytes from the optic nerve and the spinal cord [166, 167]. Stimulation of P2X₇ receptors for 15 min (with 1 mM ATP or BzATP) induced significant oligodendroglial death in culture and in situ in the optic nerve.

The P2X₇-dependent death of oligodendrocytes may have pathophysiological relevance for demyelinating diseases and for MS in particular. Indeed, in EAE, which is considered a model for MS, treatment with the P2X₇ antagonists oxATP or brilliant blue G inhibited demyelination and restored axon conduction velocity [166, 167]. Moreover, the levels of P2X₇ expression appeared to be increased in white matter of MS patients [166]. These data may indicate the relevance of P2X₇ receptors as a therapeutic target for treatment of demyelinating diseases.

P2Y Receptors

Metabotropic P2Y purinoceptors are (similar to P1 receptors) 7-transmembrane domain G protein-coupled receptors [20, 45]. They can be broadly divided into the P2Y_{1,2,4,6,11} and P2Y_{12,13,14} groups based on phylogenetic similarity and G protein preference [30]. The P2Y_{1,2,4,6,11} are coupled

to G_q/G₁₁ proteins and regulate activity of PLC, thus controlling InsP₃-mediated Ca^{2+} release from the endoplasmic reticulum (ER) [30]. The P2Y_{12,13,14} receptors modulate ion channels and inhibit adenylyl cyclase via G_{i/o} proteins [30]. This general scheme has exceptions, and in some cases, the same receptor can couple to different G proteins [168] or exert effector action without any G proteins involvement [169].

Astrocytes

The majority of astrocytes studied in situ or in isolation express metabotropic P2Y purinoceptors (Table 3 and [41]). Primary cultured rat cortical astrocytes express mRNA for P2Y_{1,2,4,6,12,13} and UDP-glucose P2Y₁₄ receptors [41, 113, 114]. Similarly, in cerebrocortical glial cultures, RT-PCR found the predominant expression of P2Y_{1,4,6} mRNAs [170]. Spinal cord astrocytes predominantly express P2Y_{1,2}-specific mRNA [171]. In astrocytes freshly isolated from the CA1 area of 8–12-day-old rat hippocampi, P2Y₁ receptors were identified in about 50% of cells at both transcriptional and protein levels [172], although some cells also expressed P2Y_{2,4} receptors. The proportion of astrocytes expressing P2Y₂ receptors in CA1 area increased from ~5% at P8–P12 to ~38% at P25 [173]. Isolated rat Müller retinal glial cells expressed mRNA for P2Y_{1,2,4,6} receptors [174], expression of these receptors was further corroborated by immunostaining as well as by functional and pharmacological analysis [175]. Interestingly, in tiger salamander Müller cells, the palette of P2Y receptors was somewhat richer: They express functional P2Y_{1,2,6,11} and possibly P2Y₄ and P2Y₁₃ receptors [176]. In nucleus accumbens, the immunoreactivity for P2Y_{1,4} receptors in astroglial cells was detected; whereas in cortex, astrocytic profiles were positively stained for P2Y_{1,2,4,6} receptors [177].

Treatment of cultured astrocytes from various brain regions with ATP induced $[Ca^{2+}]_i$ transients, which were, as a rule, a consequence of P2Y receptor activation, production of InsP₃ and subsequent Ca^{2+} release from the ER [178–181]. Stimulation of rat brain astroglial cultures with ATP and UTP induced rapid and dose-dependent increase in PLC-dependent inositol phosphate production [182–184] accompanied with $[Ca^{2+}]_i$ increase due to intracellular Ca^{2+} release [185–188]. In rat striatal astrocytes in vitro ATP induced fast $[Ca^{2+}]_i$ transients, which were independent from extracellular Ca^{2+} and were inhibited by the SERCA blocker thapsigargin or by the PLC blocker U-73122 [189]. Likewise, ATP triggered $[Ca^{2+}]_i$ rise in pituitary folliculo-stellate cells (pituitary glia) and in rat neurohypophysial astrocytes through generation of InsP₃-induced Ca^{2+} release [190, 191]. The very same InsP₃-induced intracellular Ca^{2+} release occurred in cultured spinal

Table 3 P2Y receptors in neuroglial cells

Receptor type	Experimental preparation/species/technique	Properties/function	References
Astrocytes			
P2Y _{1,2,4,6,12, 13} and UDP-glucose P2Y ₁₄ receptor	Cell culture/rat/RT-PCR	Specific mRNAs detected	[41, 113, 114]
P2Y _{1,4,6}	Cell culture/rat cortex/RT-PCR	Specific mRNAs detected	[169]
P2Y _{1,2}	Cell culture/rat spinal cord/RT-PCR	Specific mRNAs detected	[170]
P2Y _{1,2,4}	Cell culture/rat hippocampus/RT-PCR, Western blot	Specific mRNAs and proteins detected; ~50% of cells express P2Y ₁ , some cells also express P2Y _{2,4}	[171]
P2Y _{1,2,4,6}	Acutely isolated Müller cells/rat retina/RT-PCR, immunostaining, electrophysiology	Specific mRNAs and immunoreactivity detected; stimulation of P2Y receptors triggered Ca ²⁺ -dependent K ⁺ currents	[173, 174]
P2Y _{1,2,4,6,11,13}	Acutely isolated Müller cells/tiger salamander/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER; receptors subtypes were identified using specific pharmacology	[175]
P2Y ^a	Cell culture/rat/biochemical assays, Ca ²⁺ imaging	Increase in InsP ₃ production; [Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[181–183, 185–187]
P2Y ^a	Cell culture/rat, striatum/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[188]
P2Y ^a	Cell culture/rat, neurohypophysis/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[189, 190]
P2Y ^a	Cell culture/rat, spinal cord	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[191, 192]
P2Y ^a	Acute slices/mouse, cerebellum, Bergmann glial cells/Ca ²⁺ imaging, electrophysiology	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[126, 203]
P2Y	Cell culture/rat, hippocampus/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[204]
P2Y ₁	Acute slices/mouse, olfactory bulb/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[77]
P2Y ₁ , P2Y ₄	Acutely isolated optic nerve/mouse/Ca ²⁺ imaging	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[163]
P2Y ₁	Cell culture/mouse, cortex	Stimulation of ATP release through volume-sensitive anion channels	[36]
P2Y ^a	Cell culture/rat	Stimulation of ATP release via exocytosis	[37, 208, 209, 211]
P2Y ^a	Cell culture/cell lines	Stimulation of ATP release through hemichannels	[213, 214]
P2Y ^a	Cell culture/rat	Stimulation of glutamate release via exocytosis	[218]
P2Y ₁	Cell culture/rat hippocampus; Acute slices/mouse hippocampus	Stimulation of glutamate release via exocytosis	[219]
P2Y ^a	Cell culture/rat	Promotion of astroglial differentiation	[230, 231]
P2Y ^a	Cell culture/rat, human	Initiation of astrogliosis	[234–237]
P2Y ^a	Retina preparations/rat, rabbit	Initiation of astrogliosis	[240–242]
P2Y ₁ , P2Y ₁₂	Rat, Nucleus accumbens, cortex/in vivo	Initiation of astrogliosis	[176, 245]
Oligodendrocytes			
P2Y _{1,2,4}	Cell culture/rat/Western blot	Specific proteins detected	[161, 162]
P2Y ₁	Rat, rabbit/brain/immunostaining	Immunoreactivity detected in oligodendrocytes throughout the CNS	[257]
P2Y ^a	Cell culture/mouse, rabbit, retina/Acute slices/mouse, corpus callosum/Ca ²⁺ imaging, electrophysiology	[Ca ²⁺] _i transients associated with Ca ²⁺ release of the ER	[164]

^a This refers to experiments where P2Y receptor subtypes were not identified

cord astrocytes [192, 193]. Sometimes, P2Y-mediated Ca²⁺ release from the ER in cultured astrocytes also triggered secondary store-operated Ca²⁺ entry [194].

Functionally, P2Y₁ and/or P2Y₂ receptors assume the leading role in triggering ATP-induced Ca²⁺ signalling in astroglia. In embryonic glial cultures, the agonist profile for

initiating [Ca²⁺]_i transients was 2-methylthioADP (2-MeSADP) > 2-methylthioATP (2-MeSATP) > ADP > ATP > adenosine 5'-O-(3-thiotriphosphate), which is characteristic for P2Y₁ receptor [170]. Similarly, P2Y₁ receptors drive ATP-induced [Ca²⁺]_i responses in astrocytes from the supraoptic nucleus in acutely isolated slice preparations

[195]. The ER-release component of ATP-induced $[Ca^{2+}]_i$ elevation in rat cortical astrocytes in culture was significantly inhibited by the P2Y₁ antagonist MRS2179 [113], although in the same cells, sugar nucleotides triggered $[Ca^{2+}]_i$ rise by activation of UDP-glucose P2Y₁₄ receptor [113].

P2Y₁ receptors played the major role in generating and maintaining propagating Ca^{2+} waves in hippocampal cultured astrocytes [37]. In cultured spinal cord astrocytes, however, ATP-mediated propagating Ca^{2+} waves required both P2Y₁ and P2Y₂ receptors, as pharmacological inhibition of either of them eliminated the wave propagation [196, 197]. Further investigations demonstrated that P2Y₁ and P2Y₂ receptors expressed in cultured dorsal spinal cord astrocytes have different functional kinetics. Activation of P2Y₁ receptors specifically resulted in $[Ca^{2+}]_i$ oscillations, which were the consequence of cyclic protein kinase C (PKC)-mediated depression of the said receptors [171, 198]. Incidentally, Ca^{2+} waves in astroglial syncytium can be generated by both InsP₃ diffusion through gap junctions and by a regenerative wave of ATP release. The contribution of these mechanisms differs between various brain regions and can be readjusted under various physiological conditions. In spinal cord astroglia, expression levels of connexin 43 regulate expression of P2Y receptors, which, in turn, maintain ATP-dependent Ca^{2+} wave propagation. Acute inhibition of Cx43 synthesis results in down-regulation of P2Y₁ receptor production and in an increase of P2Y₄ receptor expression, which in turn modified the mode of Ca^{2+} wave propagation [199]. Similar remodelling of P2Y receptor profile and Ca^{2+} wave propagation was also observed after treatment of astroglial cultures with the cytokine IL-1 β [200]. Interestingly, in the inflammatory conditions (for example, in astroglial cultures chronically treated with IL-1 β), activation of P2Y₂ receptors decreases gap junction communications [201].

Expression of P2Y₂ receptors in cultured rat astrocytes was up-regulated by guanosine and UTP; they increased both the levels of P2Y₂-specific mRNA and augmented P2Y₂-mediated $[Ca^{2+}]_i$ transients [202]. This regulatory action of guanosine and UTP involved ERK1-2/MAPK signalling cascade [202].

Functional P2Y receptors linked to InsP₃-induced Ca^{2+} release were also found in freshly isolated human Müller cells; Ca^{2+} released from the ER store activated Ca^{2+} -dependent (BK) K⁺ channels and Ca^{2+} -gated cationic channels [203].

Activation of metabotropic purinoceptors was also instrumental in inducing Ca^{2+} signalling in astroglial cells in situ. In Bergmann glial cells in cerebellar slices, ATP triggered $[Ca^{2+}]_i$ transients, which did not require extracellular Ca^{2+} and were inhibited by incubation with thapsigargin or by intracellular perfusion with the InsP₃ receptor

antagonist heparin, thus indicating involvement of a P2Y/PLC/InsP₃-dependent signalling cascade [127, 204]. Similarly, astroglial $[Ca^{2+}]_i$ transients mediated by P2Y receptors were observed in astrocytes from stratum radiatum region of mouse hippocampus [205]. In astrocytes imaged in acute mouse olfactory bulb slices, a significant part of ATP-induced $[Ca^{2+}]_i$ transients were mediated through MRS2179-sensitive P2Y₁ receptors [77]. In isolated optic nerve preparations, P2Y₁ and/or P2Y₄ receptors were responsible for the major part of $[Ca^{2+}]_i$ elevation following exposure to ATP [164].

P2Y Receptors and Regulation of Gliotransmitter Release

Stimulation of metabotropic ATP receptors triggers release of gliotransmitters from astroglia. In particular ATP stimulation of cultured astrocytes triggers release of ATP itself; this release occurs through several pathways. In astrocytes cultured from 1-day-old mouse cortex, for example, ATP-triggered ATP release was not affected by chelating $[Ca^{2+}]_i$ with BAPTA/AM but was inhibited by non-selective anion channel blockers [36]. Interestingly, stimulation of P2Y₁ receptors was found to activate volume-sensitive Cl[−] channels in cultured astrocytes, thus indicating a direct link between activation of purinoceptors and ATP release via anion channel [206]. Similar Ca^{2+} independency of ATP release was found in rat cultured astrocytes; in this study, inhibition of the ER by thapsigargin also did not affect ATP secretion [207]. ATP can be also released from astrocytes through hemichannels [208].

At the same time, there is ample evidence in favour of exocytotic ATP release from astroglia. Exposure of rat cultured cortical astrocytes to 10 μ M UTP triggered ATP release [209]. This release was inhibited by the P2 receptor antagonist suramin, by the inhibitor of ER Ca^{2+} accumulation thapsigargin, by the complex Golgi fragmenter brefeldin A, by cytoskeleton disruption with cytochalasin D and by the exocytosis inhibitor botulinum toxin A [209]. These data taken together suggested a role for InsP₃-mediated ER Ca^{2+} release triggering Ca^{2+} -regulated exocytosis [209]. Similarly, exocytotic ATP release was found to be the primary cause for generation of propagating Ca^{2+} signals in cultured rat hippocampal astrocytes [37]. The ATP-rich vesicles, which can be released following $[Ca^{2+}]_i$ elevation, were identified in rat cultured astrocytes; in fact, these astrocytes contained two pools of vesicles, containing either glutamate or ATP, with distinct properties [210]. The ATP release with subsequent activation of P2Y receptors was also the primary mechanism for Ca^{2+} wave propagation in corpus callosum slices [211]. Exocytotic $[Ca^{2+}]_i$ - and vSNARE-dependent ATP release was also detected in astroglial progenitors [212]. In addition, release of ATP from astrocytes constitutes a powerful mechanism for astroglia–microglia signalling [213].

P2Y-induced $[Ca^{2+}]_i$ rise may trigger ATP release through connexins/hemichannels; in particular, overexpression of connexins in glial cell lines (normally devoid of gap junctions) significantly increased $[Ca^{2+}]_i$ -regulated ATP release and produced propagating Ca^{2+} waves [214, 215].

ATP released from astroglia also acts as a source of adenosine, which rapidly builds up due to ATP degradation. This adenosine acts as an important gliotransmitter because it affects synaptic transmission through numerous synaptic and extrasynaptic P1 receptors expressed in neurones [216, 217]. In certain conditions, for example upon hypoxic stress, astrocytes are able to release adenosine from separate unidentified pools, which, in turn, suppresses overall synaptic activity and exerts a general cytoprotective action [218].

ATP, acting through metabotropic receptors, triggered release of glutamate and aspartate from cultured cortical astrocytes [219]. Secretion of excitatory amino acids in this preparation was inhibited by intracellular Ca^{2+} chelation and by thapsigargin, indicating a role for Ca^{2+} release from the ER [219]. The P2Y₁-mediated intracellular Ca^{2+} release also controls secretion of glutamate from astrocytes in culture and in situ in acute hippocampal slices [220]. This glutamate release occurs through exocytosis, which was directly shown by visualising fluorescence-labelled glutamatergic vesicles using total internal fluorescence reflection imaging. The P2Y₁ receptor activation also triggered release of TNF α and prostaglandins, which regulated glutamate release synergistically with intracellular Ca^{2+} signals [220]. In mouse prefrontal cortex, stimulation of astroglial P2Y₄ receptors triggered vesicular release of glutamate, which positively modulated NMDA receptors in layer V pyramidal neurones through activation of neuronal metabotropic glutamate receptors [221]. Furthermore, evidence exists suggesting that ATP may trigger glutamate release through activation or positive modulation of volume-regulated anion channels [222, 223].

In fact, co-release of glutamate and ATP/adenosine from astrocytes may provide for a coordinated regulation of synaptic transmission, where glutamate exerts general excitatory and adenosine general inhibitory action on synaptic transmission [48, 224, 225].

ATP also stimulates mobilisation of arachidonic acid and eicosanoid production in cultured astrocytes, an effect that depends on synergism between P2Y-mediated $[Ca^{2+}]_i$ elevation and direct coupling of subset of P2Y receptors with phospholipase A2 [226, 227]. Similarly stimulation of metabotropic P2Y receptors increased synthesis of prostaglandins [228]. Activation of P2Y₁ receptors induced expression of brain-derived neurotrophic factor in astroglial cell line [229]. Stimulation of P2Y₄ receptors in cultured cortical astrocytes induced a significant increase in expression and release of glycoprotein thrombospondin (TSP)-1, which is a potent stimulator of synaptogenesis [230].

Regulation of Growth and Differentiation In postnatal rat-cultured astrocytes, ATP and its analogues (α , β -meATP, β , γ -meATP, ADP β S, 2-MeSATP and UTP) promote astroglial differentiation and process growth, these being accompanied with *C-fos* and *C-jun* mobilisation and involving activation of phospholipase A2 [231, 232]. Interestingly, ATP effects on morphological differentiation were developmentally regulated [233]: In cultures from embryonic (E18) animals, ATP suppressed cAMP-dependent stellation via P2Y receptors, whereas in postnatal cells, it promoted differentiation (although most likely through P1 receptors).

P2Y Receptors in Neuropathology ATP triggered morphological gliosis in cultured rat and human astrocytes [234], which involved extracellular signal regulated protein kinases (ERK)1/2 mediated induction of cyclo-oxygenase-2 (COX-2), but was independent of $[Ca^{2+}]_i$ changes, thus suggesting a subset of P2Y receptors linked to ERK/COX-2 pathway [235–237]. In rat primary astroglial cultures, ATP, acting through P2Y receptors, stimulated astrocyte proliferative activity [238]. Activation of P2Y receptors also potentiated proliferation of astroglia induced by broad astroglial agent fibroblast growth factor 2 [239]. Mitogenic effects of ATP are mediated through the P2Y–PKC–ERK signalling pathway in a $[Ca^{2+}]_i$ -independent fashion [240]. Reactive gliosis in retina, induced by intravitreal injection of the proteolytic enzyme dispase, triggered an up-regulation of P2Y receptors and increased Ca^{2+} signalling [241, 242]. Similar up-regulation of P2Y receptors and proportion of cells demonstrating P2Y-induced Ca^{2+} signals was observed in rabbit Müller cells following mechanical detachment of retina from pigmented epithelium [243]. Inhibition of P2 receptors with suramin attenuated reactive gliosis of Müller cells in the same model [244].

Likewise, injection of the P2Y agonist 2-MeSATP into the rat nucleus accumbens triggered astrogliosis manifested by increase in GFAP immunoreactivity and astroglial hypertrophy [245]; these effects were alleviated by the non-selective P2 receptor antagonists PPADS and reactive blue 2. Detailed pharmacological analysis led to a suggestion that purines induced astrogliosis in the nucleus accumbens mostly through activation of P2Y₁ and P2Y₁₂ receptors [246]; whereas in cortex, P2Y₁ receptors take the leading role [177]. The P2Y₁ receptors may also be involved in sensitization to D-amphetamine; the latter triggered up-regulation of P2Y₁ receptors in vivo, which enhanced ATP-induced astrogliosis [247]. In primary mouse astroglial cultures, stimulation of P2Y₁ and P2Y₆ receptors led to a $[Ca^{2+}]_i$ -dependent activation of nuclear factor of activated T cells, a transcription factor believed to activate astrogliosis [248]. Astroglial P2Y₂ receptors are also coupled with $\alpha\beta 3/\beta 5$ integrin signalling pathways,

which control cytoskeletal remodelling and motility and are involved in regulation of various stages of astrogliosis [249, 250]. Activation of P2Y₂ and P2Y₄ receptors can also regulate astrogliosis via extracellular signal-regulated protein kinases (ERK) and signal transducer and activator of transcription 3 (STAT3) signalling cascade [251].

Brain lesions up-regulated P2Y expression in several regions of the brain, including cortex (where stab wound increased the immunoreactivity for P2Y_{1,2,4,6} receptors) and in nucleus accumbens, where lesion increased expression of P2Y_{1,4} proteins and triggered additional expression of P2Y_{2,6} receptors [177].

Metabotropic P2Y receptors are linked to several glioprotective pathways. Activation of P2Y₂ receptors was reported to inhibit cell death in astroglial cell line [252]. In hippocampal astroglial cultures, stimulation of P2Y₁ receptors protected the cells against hydrogen peroxide-induced oxidative damage [253]. Brief stimulation of P2Y receptors significantly increased resistance of astrocytes isolated from old mice to oxidative stress, this effect being connected with Ca²⁺ release from the ER store [254]. In human glial cell lines, the glioprotective action of ATP was mediated through P2Y₆ receptors [255].

Stimulation of recombinant P2Y₂ receptors expressed in an astrocytoma cell line activated several cell survival mechanisms, including up-regulation of anti-apoptotic proteins BCL-2 and BCL-x and down-regulation of pro-apoptotic factor BAX; in addition, P2Y₂ receptors stimulated expression of various neurotrophins, neuropeptides and growth factors known for their neuroprotective abilities [256].

Oligodendrocytes

Cultured rat OPCs express P2Y_{1,2,4} proteins (Table 3); the ATP-induced [Ca²⁺]_i elevation, however, is mediated mainly by P2Y₁ receptors (as suggested based on sensitivity to the P2Y₁ antagonist MRS2179 [257]). Using immunohistochemistry, P2Y₁ receptors were also localised in NG-positive glial cells in rat cortical sections [257]. Similarly, P2Y₁ immunoreactivity was detected in oligodendrocytes throughout the CNS [258]. Activation of P2Y receptors triggers intracellular Ca²⁺ release in cultured mature (O10 positive) and immature (O4 positive) oligodendrocytes but not in O4-negative precursor cells [165]. In oligodendrocytes in corpus callosum slices, ATP triggered robust [Ca²⁺]_i transients [165, 259], which originated exclusively through P2Y-activated/InsP₃-induced Ca²⁺ release from the ER stores [165]. Similarly, P2Y receptor-mediated ER Ca²⁺ release significantly contributes to Ca²⁺ signalling in oligodendrocytes from the optic nerve [260]. Stimulation of P2Y receptors control OPC migration and maturation in vitro [161].

Conclusions

The purinergic signalling system is, arguably, the main extracellular signalling system that integrates neuronal–glial and glial–glial circuits in the nervous system. Indeed, in the CNS, purines and pyrimidines mediate reciprocative signalling between neurones and astrocytes in the grey matter and between axons and oligodendrocytes in the white matter. In addition, purines and pyrimidines provide for multiple signalling pathways within glial syncytium, being responsible for propagating Ca²⁺ waves and for astroglial–oligodendroglial communications. Furthermore, the purinergic signalling system is intimately involved in neuropathology by mediating reactive astrogliosis, providing for glioprotection in stress conditions and assuming the main responsibility for microglial activation. The extended family of purinoceptors, universally expressed in glial cells, is coupled to numerous signalling cascades governing glial physiological and pathological responses. In-depth understanding of the molecular physiology and pathophysiology of these receptors may further our understanding of the integrative mechanisms within neural circuits and provide new strategies for curing neurological diseases.

Acknowledgements GB thanks the Leverhulme Trust for their support. AV and OK acknowledge financial support from INTAS; AV research is also supported by Alzheimer Research Foundation (UK) and by the Grant Agency of the Czech Republic. The authors thank Dr. Gillian E. Knight for excellent editorial assistance.

References

1. Virchow R (1858) Die Cellularpathologie in ihrer Begründung auf physiologische und pathologische Gewebelehre. Zwanzig Vorlesungen gehalten während der Monate Februar, März und April 1858 im pathologischen Institut zu Berlin. August Hirschwald, Berlin
2. Verkhratsky A (2006) Patching the glia reveals the functional organisation of the brain. *Pflügers Arch* 453:411–420
3. Kettenmann H, Verkhratsky A (2008) Neuroglia: the 150 years after. *Trends Neurosci* 31:653–659
4. Verkhratsky A, Toescu EC (2006) Neuronal–glial networks as substrate for CNS integration. *J Cell Mol Med* 10:826–836
5. Verkhratsky A (2009) Neuronismo y reticulismo: neuronal–glial circuits unify the reticular and neuronal theories of brain organization. *Acta Physiol (Oxf)* 195:111–122
6. Oberheim NA, Tian GF, Han X, Peng W, Takano T, Ransom B, Nedergaard M (2008) Loss of astrocytic domain organization in the epileptic brain. *J Neurosci* 28:3264–3276
7. Bushong EA, Martone ME, Jones YZ, Ellisman MH (2002) Protoplasmic astrocytes in CA1 stratum radiatum occupy separate anatomical domains. *J Neurosci* 22:183–192
8. Nedergaard M, Ransom B, Goldman SA (2003) New roles for astrocytes: redefining the functional architecture of the brain. *Trends Neurosci* 26:523–530
9. Haydon PG, Carmignoto G (2006) Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol Rev* 86:1009–1031

10. Magistretti PJ (2006) Neuron–glia metabolic coupling and plasticity. *J Exp Biol* 209:2304–2311
11. Simard M, Arcuino G, Takano T, Liu QS, Nedergaard M (2003) Signaling at the gliovascular interface. *J Neurosci* 23:9254–9262
12. Araque A, Parpura V, Sanzgiri RP, Haydon PG (1999) Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci* 22:208–215
13. Halassa MM, Fellin T, Haydon PG (2007) The tripartite synapse: roles for gliotransmission in health and disease. *Trends Mol Med* 13:54–63
14. Tian GF, Azmi H, Takano T, Xu Q, Peng W, Lin J, Oberheim N, Lou N, Wang X, Zielke HR, Kang J, Nedergaard M (2005) An astrocytic basis of epilepsy. *Nat Med* 11:973–981
15. Nedergaard M, Dirnagl U (2005) Role of glial cells in cerebral ischemia. *Glia* 50:281–286
16. Giaume C, Kirchhoff F, Matute C, Reichenbach A, Verkhratsky A (2007) Glia: the fulcrum of brain diseases. *Cell Death Differ* 14:1324–1335
17. Rodriguez JJ, Olabarria M, Chvatal A, Verkhratsky A (2009) Astroglia in dementia and Alzheimer's disease. *Cell Death Differ* 16:378–385
18. Burnstock G (2007) Physiology and pathophysiology of purinergic neurotransmission. *Physiol Rev* 87:659–797
19. Burnstock G (2007) Purine and pyrimidine receptors. *Cell Mol Life Sci* 64:1471–1483
20. Abbracchio MP, Burnstock G, Verkhratsky A, Zimmerman H (2009) Purinergic signalling in the nervous system: an overview. *Trends Neurosci* 32:19–29
21. Burnstock G (1996) Purinoceptors: ontogeny and phylogeny. *Drug Dev Res* 39:204–242
22. Burnstock G, Verkhratsky A (2009) Evolutionary origins of the purinergic signalling system. *Acta Physiologica (Oxf)* 195:415–447
23. Burnstock G, Knight GE (2004) Cellular distribution and functions of P2 receptor subtypes in different systems. *Int Rev Cytol* 240:31–304
24. Dale N (2008) Dynamic ATP signalling and neural development. *J Physiol* 586:2429–2436
25. Pankratov Y, Lalo U, Verkhratsky A, North RA (2006) Vesicular release of ATP at central synapses. *Pflugers Arch* 452:589–597
26. Zhang Z, Chen G, Zhou W, Song A, Xu T, Luo Q, Wang W, Gu XS, Duan S (2007) Regulated ATP release from astrocytes through lysosome exocytosis. *Nat Cell Biol* 9:945–953
27. Zimmermann H (2006) Ectonucleotidases in the nervous system. *Novartis Found Symp* 276:113–128 discussion 128–130, 233–117, 275–181
28. Ralevic V, Burnstock G (1998) Receptors for purines and pyrimidines. *Pharmacol Rev* 50:413–492
29. Khakh BS, Burnstock G, Kennedy C, King BF, North RA, Seguela P, Voigt M, Humphrey PP (2001) International union of pharmacology. XXIV. Current status of the nomenclature and properties of P2X receptors and their subunits. *Pharmacol Rev* 53:107–118
30. Abbracchio MP, Burnstock G, Boeynaems JM, Barnard EA, Boyer JL, Kennedy C, Knight GE, Fumagalli M, Gachet C, Jacobson KA, Weisman GA (2006) International Union of Pharmacology LVIII: update on the P2Y G protein-coupled nucleotide receptors: from molecular mechanisms and pathophysiology to therapy. *Pharmacol Rev* 58:281–341
31. North RA (2002) Molecular physiology of P2X receptors. *Physiol Rev* 82:1013–1067
32. Burnstock G, Kennedy C (1985) Is there a basis for distinguishing two types of P2-purinoreceptor? *Gen Pharmacol* 16:433–440
33. Fredholm BB, IJ AP, Jacobson KA, Klotz KN, Linden J (2001) International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol Rev* 53:527–552
34. Pankratov Y, Lalo U, Verkhratsky A, North RA (2007) Quantal release of ATP in mouse cortex. *J Gen Physiol* 129:257–265
35. North RA, Verkhratsky A (2006) Purinergic transmission in the central nervous system. *Pflugers Arch* 452:479–485
36. Anderson CM, Bergher JP, Swanson RA (2004) ATP-induced ATP release from astrocytes. *J Neurochem* 88:246–256
37. Bowser DN, Khakh BS (2007) Vesicular ATP is the predominant cause of intercellular calcium waves in astrocytes. *J Gen Physiol* 129:485–491
38. Evanko DS, Zhang Q, Zorec R, Haydon PG (2004) Defining pathways of loss and secretion of chemical messengers from astrocytes. *Glia* 47:233–240
39. Guthrie PB, Knappenberger J, Segal M, Bennett MV, Charles AC, Kater SB (1999) ATP released from astrocytes mediates glial calcium waves. *J Neurosci* 19:520–528
40. Bains JS, Oliet SH (2007) Glia: they make your memories stick!. *Trends Neurosci* 30:417–424
41. Abbracchio MP, Ceruti S (2006) Roles of P2 receptors in glial cells: focus on astrocytes. *Purinergic Signal* 2:595–604
42. Abbracchio MP, Verderio C (2006) Pathophysiological roles of P2 receptors in glial cells. *Novartis Found Symp* 276:91–103 discussion 103–112, 275–181
43. Fields RD, Stevens B (2000) ATP: an extracellular signaling molecule between neurons and glia. *Trends Neurosci* 23:625–633
44. Fields RD, Burnstock G (2006) Purinergic signalling in neuron–glia interactions. *Nat Rev Neurosci* 7:423–436
45. Fischer W, Krugel U (2007) P2Y receptors: focus on structural, pharmacological and functional aspects in the brain. *Curr Med Chem* 14:2429–2455
46. Franke H, Krugel U, Illes P (2006) P2 receptors and neuronal injury. *Pflugers Arch* 452:622–644
47. Inoue K, Koizumi S, Tsuda M (2007) The role of nucleotides in the neuron–glia communication responsible for the brain functions. *J Neurochem* 102:1447–1458
48. Fellin T, Sul JY, D'Ascenzo M, Takano H, Pascual O, Haydon PG (2006) Bidirectional astrocyte–neuron communication: the many roles of glutamate and ATP. *Novartis Found Symp* 276:208–217
49. Newman EA (2006) A purinergic dialogue between glia and neurons in the retina. *Novartis Found Symp* 276:193–202
50. Verkhratsky A, Steinhauser C (2000) Ion channels in glial cells. *Brain Res Brain Res Rev* 32:380–412
51. Franke H, Schepper C, Illes P, Krugel U (2007) Involvement of P2X and P2Y receptors in microglial activation in vivo. *Purinergic Signal* 3:435–445
52. Haas S, Brockhaus J, Verkhratsky A, Kettenmann H (1996) ATP-induced membrane currents in amoeboid microglia acutely isolated from mouse brain slices. *Neuroscience* 75:257–261
53. Illes P, Norenberg W, Gebicke-Haerter PJ (1996) Molecular mechanisms of microglial activation. B. Voltage- and purinoceptor-operated channels in microglia. *Neurochem Int* 29:13–24
54. Inoue K (2002) Microglial activation by purines and pyrimidines. *Glia* 40:156–163
55. Inoue K, Koizumi S, Tsuda M, Shigemoto-Mogami Y (2003) Signaling of ATP receptors in glia–neuron interaction and pain. *Life Sci* 74:189–197
56. Inoue K, Tsuda M, Koizumi S (2004) Chronic pain and microglia: the role of ATP. *Novartis Found Symp* 261:55–64
57. Inoue K (2006) ATP receptors of microglia involved in pain. *Novartis Found Symp* 276:263–272
58. Inoue K (2006) The function of microglia through purinergic receptors: neuropathic pain and cytokine release. *Pharmacol Ther* 109:210–226
59. Inoue K (2008) Purinergic systems in microglia. *Cell Mol Life Sci*. 65:3074–3080

60. Farber K, Kettenmann H (2006) Purinergic signaling and microglia. *Pflugers Arch* 452:615–621
61. Sperlgh B, Illes P (2007) Purinergic modulation of microglial cell activation. *Purinergic Signal* 3:117–127
62. Trang T, Beggs S, Salter MW (2006) Purinoceptors in microglia and neuropathic pain. *Pflugers Arch* 452:645–652
63. Tsuda M, Inoue K, Salter MW (2005) Neuropathic pain and spinal microglia: a big problem from molecules in “small” glia. *Trends Neurosci* 28:101–107
64. Boison D (2008) Adenosine as a neuromodulator in neurological diseases. *Curr Opin Pharmacol* 8:2–7
65. Abbracchio MP, Cattabeni F (1999) Brain adenosine receptors as targets for therapeutic intervention in neurodegenerative diseases. *Ann N Y Acad Sci* 890:79–92
66. Dare E, Schulte G, Karovic O, Hammarberg C, Fredholm BB (2007) Modulation of glial cell functions by adenosine receptors. *Physiol Behav* 92:15–20
67. Hosli L, Hosli E, Uhr M, Della Briotta G (1987) Electrophysiological evidence for adenosine receptors on astrocytes of cultured rat central nervous system. *Neurosci Lett* 79:108–112
68. Woods MD, Freshney RI, Ball SG, Vaughan PF (1989) Regulation of cyclic AMP formation in cultures of human foetal astrocytes by beta 2-adrenergic and adenosine receptors. *J Neurochem* 53:864–869
69. Peakman MC, Hill SJ (1994) Adenosine A_{2B}-receptor-mediated cyclic AMP accumulation in primary rat astrocytes. *Br J Pharmacol* 111:191–198
70. Peakman MC, Hill SJ (1996) Adenosine A₁ receptor-mediated inhibition of cyclic AMP accumulation in type-2 but not type-1 rat astrocytes. *Eur J Pharmacol* 306:281–289
71. Biber K, Fiebich BL, Gebicke-Harter P, van Calcar D (1999) Carbamazepine-induced upregulation of adenosine A₁-receptors in astrocyte cultures affects coupling to the phosphoinositol signaling pathway. *Neuropsychopharmacology* 20:271–278
72. Biber K, Klotz KN, Berger M, Gebicke-Harter PJ, van Calcar D (1997) Adenosine A₁ receptor-mediated activation of phospholipase C in cultured astrocytes depends on the level of receptor expression. *J Neurosci* 17:4956–4964
73. Biber K, Lubrich B, Fiebich BL, Boddeke HW, van Calcar D (2001) Interleukin-6 enhances expression of adenosine A₁ receptor mRNA and signaling in cultured rat cortical astrocytes and brain slices. *Neuropsychopharmacology* 24:86–96
74. Pilitsis JG, Kimelberg HK (1998) Adenosine receptor mediated stimulation of intracellular calcium in acutely isolated astrocytes. *Brain Res* 798:294–303
75. Peakman MC, Hill SJ (1995) Adenosine A₁ receptor-mediated changes in basal and histamine-stimulated levels of intracellular calcium in primary rat astrocytes. *Br J Pharmacol* 115:801–810
76. Porter JT, McCarthy KD (1995) Adenosine receptors modulate $[Ca^{2+}]_i$ in hippocampal astrocytes in situ. *J Neurochem* 65:1515–1523
77. Doengi M, Deitmer JW, Lohr C (2008) New evidence for purinergic signaling in the olfactory bulb: A_{2A} and P2Y₁ receptors mediate intracellular calcium release in astrocytes. *FASEB J* 22:2368–2378
78. Chen Y, Rathbone MP, Hertz L (2001) Guanosine-induced increase in free cytosolic calcium concentration in mouse astrocytes in primary cultures: does it act on an A₃ adenosine receptor? *J Neurosci Res* 65:184–189
79. Ogata T, Nakamura Y, Tsuji K, Shibata T, Kataoka K, Schubert P (1994) Adenosine enhances intracellular Ca^{2+} mobilization in conjunction with metabotropic glutamate receptor activation by t-ACPD in cultured hippocampal astrocytes. *Neurosci Lett* 170:5–8
80. Ogata T, Nakamura Y, Schubert P (1996) Potentiated cAMP rise in metabotropically stimulated rat cultured astrocytes by a Ca^{2+} -related A₁/A₂ adenosine receptor cooperation. *Eur J Neurosci* 8:1124–1131
81. Alloisio S, Cugnoli C, Ferroni S, Nobile M (2004) Differential modulation of ATP-induced calcium signalling by A₁ and A₂ adenosine receptors in cultured cortical astrocytes. *Br J Pharmacol* 141:935–942
82. Ferroni S, Marchini C, Ogata T, Schubert P (2002) Recovery of deficient cholinergic calcium signaling by adenosine in cultured rat cortical astrocytes. *J Neurosci Res* 68:615–621
83. Jimenez AI, Castro E, Mirabet M, Franco R, Delicado EG, Miras-Portugal MT (1999) Potentiation of ATP calcium responses by A_{2B} receptor stimulation and other signals coupled to G_s proteins in type-1 cerebellar astrocytes. *Glia* 26:119–128
84. Toms NJ, Roberts PJ (1999) Group 1 mGlu receptors elevate $[Ca^{2+}]_i$ in rat cultured cortical type 2 astrocytes: $[Ca^{2+}]_i$ synergy with adenosine A₁ receptors. *Neuropharmacology* 38:1511–1517
85. Nobile M, Monaldi I, Alloisio S, Cugnoli C, Ferroni S (2003) ATP-induced, sustained calcium signalling in cultured rat cortical astrocytes: evidence for a non-capacitative, P2X₇-like-mediated calcium entry. *FEBS Lett* 538:71–76
86. Nishizaki T, Nagai K, Nomura T, Tada H, Kanno T, Tozaki H, Li XX, Kondoh T, Kodama N, Takahashi E, Sakai N, Tanaka K, Saito N (2002) A new neuromodulatory pathway with a glial contribution mediated via A_{2A} adenosine receptors. *Glia* 39:133–147
87. Li XX, Nomura T, Aihara H, Nishizaki T (2001) Adenosine enhances glial glutamate efflux via A_{2A} adenosine receptors. *Life Sci* 68:1343–1350
88. Hagberg H, Andersson P, Lacarewicz J, Jacobson I, Butcher S, Sandberg M (1987) Extracellular adenosine, inosine, hypoxanthine, and xanthine in relation to tissue nucleotides and purines in rat striatum during transient ischemia. *J Neurochem* 49:227–231
89. Hindley S, Herman MA, Rathbone MP (1994) Stimulation of reactive astrogliosis in vivo by extracellular adenosine diphosphate or an adenosine A₂ receptor agonist. *J Neurosci Res* 38:399–406
90. Brambilla R, Cottini L, Fumagalli M, Ceruti S, Abbracchio MP (2003) Blockade of A_{2A} adenosine receptors prevents basic fibroblast growth factor-induced reactive astrogliosis in rat striatal primary astrocytes. *Glia* 43:190–194
91. Trincavelli ML, Marroni M, Tuscano D, Ceruti S, Mazzola A, Mitro N, Abbracchio MP, Martini C (2004) Regulation of A_{2B} adenosine receptor functioning by tumour necrosis factor α in human astroglial cells. *J Neurochem* 91:1180–1190
92. Trincavelli ML, Tonazzini I, Montali M, Abbracchio MP, Martini C (2008) Short-term TNF- α treatment induced A_{2B} adenosine receptor desensitization in human astroglial cells. *J Cell Biochem* 104:150–161
93. Appel E, Kazimirsky G, Ashkenazi E, Kim SG, Jacobson KA, Brodie C (2001) Roles of BCL-2 and caspase 3 in the adenosine A₃ receptor-induced apoptosis. *J Mol Neurosci* 17:285–292
94. Abbracchio MP, Ceruti S, Brambilla R, Franceschi C, Malorni W, Jacobson KA, von Lubitz DK, Cattabeni F (1997) Modulation of apoptosis by adenosine in the central nervous system: a possible role for the A₃ receptor. Pathophysiological significance and therapeutic implications for neurodegenerative disorders. *Ann N Y Acad Sci* 825:11–22
95. Abbracchio MP, Ceruti S, Brambilla R, Barbieri D, Camurri A, Franceschi C, Giammarioli AM, Jacobson KA, Cattabeni F, Malorni W (1998) Adenosine A₃ receptors and viability of astrocytes. *Drug Dev Res* 45:379–386
96. Choi JW, Yoo BK, Ryu MK, Choi MS, Park GH, Ko KH (2005) Adenosine and purine nucleosides prevent the disruption of mitochondrial transmembrane potential by peroxynitrite in rat primary astrocytes. *Arch Pharm Res* 28:810–815

97. Brodie C, Blumberg PM, Jacobson KA (1998) Activation of the A_{2A} adenosine receptor inhibits nitric oxide production in glial cells. *FEBS Lett* 429:139–142
98. D'Alimonte I, Ballerini P, Nargi E, Buccella S, Giuliani P, Di Iorio P, Caciagli F, Ciccarelli R (2007) Staurosporine-induced apoptosis in astrocytes is prevented by A₁ adenosine receptor activation. *Neurosci Lett* 418:66–71
99. Bjorklund O, Shang M, Tonazzini I, Dare E, Fredholm BB (2008) Adenosine A₁ and A₃ receptors protect astrocytes from hypoxic damage. *Eur J Pharmacol* 596:6–13
100. Wittendorp MC, Boddeke HW, Biber K (2004) Adenosine A₃ receptor-induced CCL2 synthesis in cultured mouse astrocytes. *Glia* 46:410–418
101. Stevens B, Porta S, Haak LL, Gallo V, Fields RD (2002) Adenosine: a neuron-glial transmitter promoting myelination in the CNS in response to action potentials. *Neuron* 36:855–868
102. Ishibashi T, Dakin KA, Stevens B, Lee PR, Kozlov SV, Stewart CL, Fields RD (2006) Astrocytes promote myelination in response to electrical impulses. *Neuron* 49:823–832
103. Fields RD (2006) Nerve impulses regulate myelination through purinergic signalling. *Novartis Found Symp* 276:148–158
104. Tsutsui S, Schnermann J, Noorbakhsh F, Henry S, Yong VW, Winston BW, Warren K, Power C (2004) A₁ adenosine receptor upregulation and activation attenuates neuroinflammation and demyelination in a model of multiple sclerosis. *J Neurosci* 24:1521–1529
105. Turner CP, Yan H, Schwartz M, Othman T, Rivkees SA (2002) A₁ adenosine receptor activation induces ventriculomegaly and white matter loss. *Neuroreport* 13:1199–1204
106. Nicke A, Baumert HG, Rettinger J, Eichele A, Lambrecht G, Mutschler E, Schmalzing G (1998) P2X₁ and P2X₃ receptors form stable trimers: a novel structural motif of ligand-gated ion channels. *EMBO J* 17:3016–3028
107. Khakh BS (2001) Molecular physiology of P2X receptors and ATP signalling at synapses. *Nat Rev Neurosci* 2:165–174
108. Roberts JA, Vial C, Digby HR, Agboh KC, Wen H, Atterbury-Thomas A, Evans RJ (2006) Molecular properties of P2X receptors. *Pflugers Arch* 452:486–500
109. Illes P, Alexandre Ribeiro J (2004) Molecular physiology of P2 receptors in the central nervous system. *Eur J Pharmacol* 483:5–17
110. Surprenant A, North RA (2009) Signaling at purinergic P2X receptors. *Annu Rev Physiol* 71:333–359
111. Duan S, Neary JT (2006) P2X₇ receptors: properties and relevance to CNS function. *Glia* 54:738–746
112. Sperlagh B, Vizi ES, Wirkner K, Illes P (2006) P2X₇ receptors in the nervous system. *Prog Neurobiol* 78:327–346
113. Fumagalli M, Brambilla R, D'Ambrosi N, Volonte C, Matteoli M, Verderio C, Abbracchio MP (2003) Nucleotide-mediated calcium signaling in rat cortical astrocytes: Role of P2X and P2Y receptors. *Glia* 43:218–203
114. Dixon SJ, Yu R, Panupinthu N, Wilson JX (2004) Activation of P2 nucleotide receptors stimulates acid efflux from astrocytes. *Glia* 47:367–376
115. Franke H, Grosche J, Schadlich H, Krugel U, Allgaier C, Illes P (2001) P2X receptor expression on astrocytes in the nucleus accumbens of rats. *Neuroscience* 108:421–429
116. Jabs R, Guenther E, Marquardt K, Wheeler-Schilling TH (2000) Evidence for P2X₃, P2X₄, P2X₅ but not for P2X₇ containing purinergic receptors in Müller cells of the rat retina. *Brain Res Mol Brain Res* 76:205–210
117. Pannicke T, Fischer W, Biedermann B, Schadlich H, Grosche J, Faude F, Wiedemann P, Allgaier C, Illes P, Burnstock G, Reichenbach A (2000) P2X₇ receptors in Müller glial cells from the human retina. *J Neurosci* 20:5965–5972
118. Lalo U, Pankratov Y, Wichert SP, Rossner MJ, North RA, Kirchhoff F, Verkhratsky A (2008) P2X₁ and P2X₅ subunits form the functional P2X receptor in mouse cortical astrocytes. *J Neurosci* 28:5473–5480
119. Loesch A, Burnstock G (1998) Electron-immunocytochemical localization of P2X₁ receptors in the rat cerebellum. *Cell Tissue Res* 294:253–260
120. Kanjhan R, Housley GD, Thorne PR, Christie DL, Palmer DJ, Luo L, Ryan AF (1996) Localization of ATP-gated ion channels in cerebellum using P2X₂R subunit-specific antisera. *Neuroreport* 7:2665–2669
121. Kanjhan R, Housley GD, Burton LD, Christie DL, Kippenberger A, Thorne PR, Luo L, Ryan AF (1999) Distribution of the P2X₂ receptor subunit of the ATP-gated ion channels in the rat central nervous system. *J Comp Neurol* 407:11–32
122. Ashour F, Deuchars J (2004) Electron microscopic localisation of P2X₄ receptor subunit immunoreactivity to pre- and post-synaptic neuronal elements and glial processes in the dorsal vagal complex of the rat. *Brain Res* 1026:44–55
123. Kukley M, Barden JA, Steinhauser C, Jabs R (2001) Distribution of P2X receptors on astrocytes in juvenile rat hippocampus. *Glia* 36:11–21
124. Magoski NS, Walz W (1992) Ionic dependence of a P₂-purinoceptor mediated depolarization of cultured astrocytes. *J Neurosci Res* 32:530–538
125. Walz W, Gimpl G, Ohlemeyer C, Kettenmann H (1994) Extracellular ATP-induced currents in astrocytes: involvement of a cation channel. *J Neurosci Res* 38:12–18
126. Jabs R, Matthias K, Grote A, Grauer M, Seifert G, Steinhauser C (2007) Lack of P2X receptor mediated currents in astrocytes and GluR type glial cells of the hippocampal CA1 region. *Glia* 55:1648–1655
127. Kirischuk S, Moller T, Voitenko N, Kettenmann H, Verkhratsky A (1995) ATP-induced cytoplasmic calcium mobilization in Bergmann glial cells. *J Neurosci* 15:7861–7871
128. Torres GE, Haines WR, Egan TM, Voigt MM (1998) Co-expression of P2X₁ and P2X₅ receptor subunits reveals a novel ATP-gated ion channel. *Mol Pharmacol* 54:989–993
129. Surprenant A, Schneider DA, Wilson HL, Galligan JJ, North RA (2000) Functional properties of heteromeric P2X_{1/5} receptors expressed in HEK cells and excitatory junction potentials in guinea-pig submucosal arterioles. *J Auton Nerv Syst* 81:249–263
130. Haines WR, Torres GE, Voigt MM, Egan TM (1999) Properties of the novel ATP-gated ionotropic receptor composed of the P2X₁ and P2X₅ isoforms. *Mol Pharmacol* 56:720–727
131. Le KT, Boue-Grabot E, Archambault V, Seguela P (1999) Functional and biochemical evidence for heteromeric ATP-gated channels composed of P2X₁ and P2X₅ subunits. *J Biol Chem* 274:15415–15419
132. James G, Butt AM (1999) Adenosine 5' triphosphate evoked mobilization of intracellular calcium in central nervous system white matter of adult mouse optic nerve. *Neurosci Lett* 268:53–56
133. Hamilton N, Vayro S, Kirchhoff F, Verkhratsky A, Robbins J, Gorecki DC, Butt AM (2008) Mechanisms of ATP- and glutamate-mediated calcium signaling in white matter astrocytes. *Glia* 56:734–749
134. Pelegrin P, Surprenant A (2006) Pannexin-1 mediates large pore formation and interleukin-1 β release by the ATP-gated P2X₇ receptor. *Embo J* 25:5071–5082
135. Collo G, Neidhart S, Kawashima E, Kosco-Vilbois M, North RA, Buell G (1997) Tissue distribution of the P2X₇ receptor. *Neuropharmacology* 36:1277–1283
136. Sim JA, Young MT, Sung HY, North RA, Surprenant A (2004) Reanalysis of P2X₇ receptor expression in rodent brain. *J Neurosci* 24:6307–6314
137. Yu Y, Ugawa S, Ueda T, Ishida Y, Inoue K, Kyaw Nyunt A, Umemura A, Mase M, Yamada K, Shimada S (2008) Cellular

- localization of P2X₇ receptor mRNA in the rat brain. *Brain Res* 1194:45–55
138. Duan S, Anderson CM, Keung EC, Chen Y, Swanson RA (2003) P2X₇ receptor-mediated release of excitatory amino acids from astrocytes. *J Neurosci* 23:1320–1328
 139. Narcisse L, Scemes E, Zhao Y, Lee SC, Brosnan CF (2005) The cytokine IL-1 β transiently enhances P2X₇ receptor expression and function in human astrocytes. *Glia* 49:245–258
 140. Hung AC, Sun SH (2002) The P2X₇ receptor-mediated phospholipase D activation is regulated by both PKC-dependent and PKC-independent pathways in a rat brain-derived Type-2 astrocyte cell line, RBA-2. *Cell Signal* 14:83–92
 141. Wang CM, Chang YY, Sun SH (2003) Activation of P2X₇ purinoceptor-stimulated TGF- β 1 mRNA expression involves PKC/MAPK signalling pathway in a rat brain-derived type-2 astrocyte cell line, RBA-2. *Cell Signal* 15:1129–1137
 142. John GR, Simpson JE, Woodroffe MN, Lee SC, Brosnan CF (2001) Extracellular nucleotides differentially regulate interleukin-1 β signaling in primary human astrocytes: implications for inflammatory gene expression. *J Neurosci* 21:4134–4142
 143. Panenka W, Jijon H, Herx LM, Armstrong JN, Feighan D, Wei T, Yong VW, Ransohoff RM, MacVicar BA (2001) P2X₇-like receptor activation in astrocytes increases chemokine monocyte chemoattractant protein-1 expression via mitogen-activated protein kinase. *J Neurosci* 21:7135–7142
 144. Ballerini P, Rathbone MP, Di Iorio P, Renzetti A, Giuliani P, D'Alimonte I, Trubiani O, Caciagli F, Ciccarelli R (1996) Rat astroglial P2_Z (P2X₇) receptors regulate intracellular calcium and purine release. *Neuroreport* 7:2533–2537
 145. James G, Butt AM (2002) P2Y and P2X purinoceptor mediated Ca²⁺ signalling in glial cell pathology in the central nervous system. *Eur J Pharmacol* 447:247–260
 146. Wang CM, Chang YY, Kuo JS, Sun SH (2002) Activation of P2X₇ receptors induced [(3) H]GABA release from the RBA-2 type-2 astrocyte cell line through a Cl(-)/HCO₃(-)-dependent mechanism. *Glia* 37:8–18
 147. Suadicani SO, Brosnan CF, Scemes E (2006) P2X₇ receptors mediate ATP release and amplification of astrocytic intercellular Ca²⁺ signaling. *J Neurosci* 26:1378–1385
 148. Fellin T, Pozzan T, Carmignoto G (2006) Purinergic receptors mediate two distinct glutamate release pathways in hippocampal astrocytes. *J Biol Chem* 281:4274–4284
 149. Walter L, Dinh T, Stella N (2004) ATP induces a rapid and pronounced increase in 2-arachidonoylglycerol production by astrocytes, a response limited by monoacylglycerol lipase. *J Neurosci* 24:8068–8074
 150. Kucher BM, Neary JT (2005) Bi-functional effects of ATP/P2 receptor activation on tumor necrosis factor- α release in lipopolysaccharide-stimulated astrocytes. *J Neurochem* 92:525–535
 151. Murakami K, Nakamura Y, Yoneda Y (2003) Potentiation by ATP of lipopolysaccharide-stimulated nitric oxide production in cultured astrocytes. *Neuroscience* 117:37–42
 152. Jacques-Silva MC, Rodnight R, Lenz G, Liao Z, Kong Q, Tran M, Kang Y, Gonzalez FA, Weisman GA, Neary JT (2004) P2X₇ receptors stimulate AKT phosphorylation in astrocytes. *Br J Pharmacol* 141:1106–1117
 153. Gendron FP, Neary JT, Theiss PM, Sun GY, Gonzalez FA, Weisman GA (2003) Mechanisms of P2X₇ receptor-mediated ERK1/2 phosphorylation in human astrocytoma cells. *Am J Physiol Cell Physiol* 284:C571–581
 154. Lu H, Burns D, Garnier P, Wei G, Zhu K, Ying W (2007) P2X₇ receptors mediate NADH transport across the plasma membranes of astrocytes. *Biochem Biophys Res Commun* 362:946–950
 155. Ballerini P, Ciccarelli R, Caciagli F, Rathbone MP, Werstiuk ES, Traversa U, Buccella S, Giuliani P, Jang S, Nargi E, Visini D, Santavenere C, Di Iorio P (2005) P2X₇ receptor activation in rat brain cultured astrocytes increases the biosynthetic release of cysteinyl leukotrienes. *Int J Immunopathol Pharmacol* 18:417–430
 156. Lo JC, Huang WC, Chou YC, Tseng CH, Lee WL, Sun SH (2008) Activation of P2X₇ receptors decreases glutamate uptake and glutamine synthetase activity in RBA-2 astrocytes via distinct mechanisms. *J Neurochem* 105:151–164
 157. D'Alimonte I, Ciccarelli R, Di Iorio P, Nargi E, Buccella S, Giuliani P, Rathbone MP, Jang S, Caciagli F, Ballerini P (2007) Activation of P2X₇ receptors stimulates the expression of P2Y₂ receptor mRNA in astrocytes cultured from rat brain. *Int J Immunopathol Pharmacol* 20:301–316
 158. Lee M, Lee SJ, Choi HJ, Jung YW, Frokiaer J, Nielsen S, Kwon TH (2008) Regulation of AQP4 protein expression in rat brain astrocytes: role of P2X₇ receptor activation. *Brain Res* 1195:1–11
 159. Franke H, Gunther A, Grosche J, Schmidt R, Rossner S, Reinhardt R, Faber-Zuschratter H, Schneider D, Illes P (2004) P2X₇ receptor expression after ischemia in the cerebral cortex of rats. *J Neuropathol Exp Neurol* 63:686–699
 160. Bringmann A, Pannicke T, Moll V, Milenkovic I, Faude F, Enzmann V, Wolf S, Reichenbach A (2001) Upregulation of P2X₇ receptor currents in Müller glial cells during proliferative vitreoretinopathy. *Invest Ophthalmol Vis Sci* 42:860–867
 161. Meomartini ME, Amadio S, Visentin S, Franchini L, Aloisi F, Volontà C, Agresti C (2003) Expression and functional analysis of P2 receptors in oligodendrocytes. *Glia* 43(Suppl. 2):59
 162. Agresti C, Meomartini ME, Amadio S, Ambrosini E, Volonte C, Aloisi F, Visentin S (2005) ATP regulates oligodendrocyte progenitor migration, proliferation, and differentiation: involvement of metabotropic P2 receptors. *Brain Res Brain Res Rev* 48:157–165
 163. James G, Butt AM (2001) P2X and P2Y purinoceptors mediate ATP-evoked calcium signalling in optic nerve glia in situ. *Cell Calcium* 30:251–259
 164. Kirischuk S, Scherer J, Kettenmann H, Verkhratsky A (1995) Activation of P2-purinoreceptors triggered Ca²⁺ release from InsP₃-sensitive internal stores in mammalian oligodendrocytes. *J Physiol* 483:41–57
 165. Matute C, Torre I, Perez-Cerda F, Perez-Samartin A, Alberdi E, Etxebarria E, Arranz AM, Ravid R, Rodriguez-Antiguedad A, Sanchez-Gomez M, Domercq M (2007) P2X₇ receptor blockade prevents ATP excitotoxicity in oligodendrocytes and ameliorates experimental autoimmune encephalomyelitis. *J Neurosci* 27:9525–9533
 166. Matute C (2008) P2X₇ receptors in oligodendrocytes: a novel target for neuroprotection. *Mol Neurobiol* 38:123–128
 167. White PJ, Webb TE, Boarder MR (2003) Characterization of a Ca²⁺ response to both UTP and ATP at human P2Y₁₁ receptors: evidence for agonist-specific signaling. *Mol Pharmacol* 63:1356–1363
 168. Erb L, Liu J, Ockerhausen J, Kong Q, Garrad RC, Griffin K, Neal C, Krugh B, Santiago-Perez LI, Gonzalez FA, Gresham HD, Turner JT, Weisman GA (2001) An RGD sequence in the P2Y₂ receptor interacts with α V β 3 integrins and is required for G(o)-mediated signal transduction. *J Cell Biol* 153:491–501
 169. Bennett GC, Ford AP, Smith JA, Emmett CJ, Webb TE, Boarder MR (2003) P2Y receptor regulation of cultured rat cerebral cortical cells: calcium responses and mRNA expression in neurons and glia. *Br J Pharmacol* 139:279–288
 170. Fam SR, Gallagher CJ, Salter MW (2000) P2Y₁ purinoceptor-mediated Ca²⁺ signaling and Ca²⁺ wave propagation in dorsal spinal cord astrocytes. *J Neurosci* 20:2800–2808
 171. Zhu Y, Kimelberg HK (2004) Cellular expression of P2Y and β -AR receptor mRNAs and proteins in freshly isolated astrocytes

- and tissue sections from the CA1 region of P8–12 rat hippocampus. *Brain Res Dev Brain Res* 148:77–87
172. Zhu Y, Kimelberg HK (2001) Developmental expression of metabotropic P2Y₁ and P2Y₂ receptors in freshly isolated astrocytes from rat hippocampus. *J Neurochem* 77:530–541
 173. Fries JE, Wheeler-Schilling TH, Kohler K, Guenther E (2004) Distribution of metabotropic P2Y receptors in the rat retina: a single-cell RT-PCR study. *Brain Res Mol Brain Res* 130:1–6
 174. Fries JE, Goczalik IM, Wheeler-Schilling TH, Kohler K, Guenther E, Wolf S, Wiedemann P, Bringmann A, Reichenbach A, Francke M, Pannicke T (2005) Identification of P2Y receptor subtypes in human Müller glial cells by physiology, single cell RT-PCR, and immunohistochemistry. *Invest Ophthalmol Vis Sci* 46:3000–3007
 175. Reifel Saltzberg JM, Garvey KA, Keirstead SA (2003) Pharmacological characterization of P2Y receptor subtypes on isolated tiger salamander Müller cells. *Glia* 42:149–159
 176. Franke H, Krugel U, Grosche J, Heine C, Hartig W, Allgaier C, Illes P (2004) P2Y receptor expression on astrocytes in the nucleus accumbens of rats. *Neuroscience* 127:431–441
 177. Kostyuk P, Verkhratsky A (1994) Calcium stores in neurons and glia. *Neuroscience* 63:381–404
 178. Verkhratsky A, Kettenmann H (1996) Calcium signalling in glial cells. *Trends Neurosci* 19:346–352
 179. Verkhratsky A, Orkand RK, Kettenmann H (1998) Glial calcium: homeostasis and signaling function. *Physiol Rev* 78:99–141
 180. Bernstein M, Behnisch T, Balschun D, Reymann KG, Reiser G (1998) Pharmacological characterisation of metabotropic glutamatergic and purinergic receptors linked to Ca²⁺ signalling in hippocampal astrocytes. *Neuropharmacology* 37:169–178
 181. Pearce B, Murphy S, Jeremy J, Morrow C, Dandona P (1989) ATP-evoked Ca²⁺ mobilisation and prostanoid release from astrocytes: P2-purinergic receptors linked to phosphoinositide hydrolysis. *J Neurochem* 52:971–977
 182. Chen CC, Chen WC (1996) ATP-evoked inositol phosphates formation through activation of P2U purinergic receptors in cultured astrocytes: regulation by PKC subtypes alpha, delta, and theta. *Glia* 17:63–71
 183. Ishimoto H, Nakahata N, Matsuoka I, Nakanishi H (1997) Effects of ATP on phosphoinositide hydrolysis and prostaglandin E2 generation in rabbit astrocytes. *J Pharm Pharmacol* 49:520–524
 184. Pearce B, Langley D (1994) Purine- and pyrimidine-stimulated phosphoinositide breakdown and intracellular calcium mobilisation in astrocytes. *Brain Res* 660:329–332
 185. Bruner G, Murphy S (1993) UTP activates multiple second messenger systems in cultured rat astrocytes. *Neurosci Lett* 162:105–108
 186. Jimenez AI, Castro E, Communi D, Boeynaems JM, Delgado EG, Miras-Portugal MT (2000) Coexpression of several types of metabotropic nucleotide receptors in single cerebellar astrocytes. *J Neurochem* 75:2071–2079
 187. Kastriasis CH, Salm AK, McCarthy K (1992) Stimulation of the P2Y purinergic receptor on type 1 astroglia results in inositol phosphate formation and calcium mobilization. *J Neurochem* 58:1277–1284
 188. Centemeri C, Bolego C, Abbracchio MP, Cattabeni F, Puglisi L, Burnstock G, Nicosia S (1997) Characterization of the Ca²⁺ responses evoked by ATP and other nucleotides in mammalian brain astrocytes. *Br J Pharmacol* 121:1700–1706
 189. Uchiyama M, Nakajima Y, Sakuma Y, Kato M (2001) Purinergic regulation of intracellular Ca²⁺ concentration of rat pituitary folliculo-stellate cells in primary culture. *J Neuroendocrinol* 13:378–385
 190. Troade JD, Thirion S, Petturiti D, Bohn MT, Poujeol P (1999) ATP acting on P2Y receptors triggers calcium mobilization in primary cultures of rat neurohypophysial astrocytes (pituicytes). *Pflugers Arch* 437:745–753
 191. Salter MW, Hicks JL (1995) ATP causes release of intracellular Ca²⁺ via the phospholipase C β /IP₃ pathway in astrocytes from the dorsal spinal cord. *J Neurosci* 15:2961–2971
 192. Salter MW, Hicks JL (1994) ATP-evoked increases in intracellular calcium in neurons and glia from the dorsal spinal cord. *J Neurosci* 14:1563–1575
 193. Koizumi S, Saito Y, Nakazawa K, Nakajima K, Sawada JI, Kohsaka S, Illes P, Inoue K (2002) Spatial and temporal aspects of Ca²⁺ signaling mediated by P2Y receptors in cultured rat hippocampal astrocytes. *Life Sci* 72:431–442
 194. Espallergues J, Solovieva O, Techer V, Bauer K, Alonso G, Vincent A, Hussy N (2007) Synergistic activation of astrocytes by ATP and norepinephrine in the rat supraoptic nucleus. *Neuroscience* 148:712–723
 195. Bennett MR, Buljan V, Farnell L, Gibson WG (2006) Purinergic junctional transmission and propagation of calcium waves in spinal cord astrocyte networks. *Biophys J* 91:3560–3571
 196. Gallagher CJ, Salter MW (2003) Differential properties of astrocyte calcium waves mediated by P2Y₁ and P2Y₂ receptors. *J Neurosci* 23:6728–6739
 197. Fam SR, Gallagher CJ, Kalia LV, Salter MW (2003) Differential frequency dependence of P2Y₁- and P2Y₂- mediated Ca²⁺ signaling in astrocytes. *J Neurosci* 23:4437–4444
 198. Suadicani SO, De Pina-Benabou MH, Urban-Maldonado M, Spray DC, Scemes E (2003) Acute downregulation of Cx43 alters P2Y receptor expression levels in mouse spinal cord astrocytes. *Glia* 42:160–171
 199. John GR, Scemes E, Suadicani SO, Liu JS, Charles PC, Lee SC, Spray DC, Brosnan CF (1999) IL-1 β differentially regulates calcium wave propagation between primary human fetal astrocytes via pathways involving P2 receptors and gap junction channels. *Proc Natl Acad Sci U S A* 96:11613–11618
 200. Meme W, Ezan P, Venance L, Glowinski J, Giaume C (2004) ATP-induced inhibition of gap junctional communication is enhanced by interleukin-1 β treatment in cultured astrocytes. *Neuroscience* 126:95–104
 201. Ballerini P, Di Iorio P, Caciagli F, Rathbone MP, Jiang S, Nargi E, Buccella S, Giuliani P, D'Alimonte I, Fischione G, Masciulli A, Romano S, Ciccarelli R (2006) P2Y₂ receptor up-regulation induced by guanosine or UTP in rat brain cultured astrocytes. *Int J Immunopathol Pharmacol* 19:293–308
 202. Bringmann A, Pannicke T, Weick M, Biedermann B, Uhlmann S, Kohen L, Wiedemann P, Reichenbach A (2002) Activation of P2Y receptors stimulates potassium and cation currents in acutely isolated human Müller (glial) cells. *Glia* 37:139–152
 203. Piet R, Jahr CE (2007) Glutamatergic and purinergic receptor-mediated calcium transients in Bergmann glial cells. *J Neurosci* 27:4027–4035
 204. Bowser DN, Khakh BS (2004) ATP excites interneurons and astrocytes to increase synaptic inhibition in neuronal networks. *J Neurosci* 24:8606–8620
 205. Darby M, Kuzmiski JB, Panenka W, Feighan D, MacVicar BA (2003) ATP released from astrocytes during swelling activates chloride channels. *J Neurophysiol* 89:1870–1877
 206. Wang Z, Haydon PG, Yeung ES (2000) Direct observation of calcium-independent intercellular ATP signaling in astrocytes. *Anal Chem* 72:2001–2007
 207. Stout CE, Costantin JL, Naus CC, Charles AC (2002) Intercellular calcium signaling in astrocytes via ATP release through connexin hemichannels. *J Biol Chem* 277:10482–10488
 208. Abdipranoto A, Liu GJ, Werry EL, Bennett MR (2003) Mechanisms of secretion of ATP from cortical astrocytes triggered by uridine triphosphate. *Neuroreport* 14:2177–2181

209. Coco S, Calegari F, Pravettoni E, Pozzi D, Taverna E, Rosa P, Matteoli M, Verderio C (2003) Storage and release of ATP from astrocytes in culture. *J Biol Chem* 278:1354–1362
210. Haas B, Schipke CG, Peters O, Sohl G, Willecke K, Kettenmann H (2006) Activity-dependent ATP-waves in the mouse neocortex are independent from astrocytic calcium waves. *Cereb Cortex* 16:237–246
211. Striedinger K, Meda P, Scemes E (2007) Exocytosis of ATP from astrocyte progenitors modulates spontaneous Ca^{2+} oscillations and cell migration. *Glia* 55:652–662
212. Verderio C, Matteoli M (2001) ATP mediates calcium signaling between astrocytes and microglial cells: modulation by IFN- γ . *J Immunol* 166:6383–6391
213. Cotrina ML, Lin JH, Alves-Rodrigues A, Liu S, Li J, Azmi-Ghadimi H, Kang J, Naus CC, Nedergaard M (1998) Connexins regulate calcium signaling by controlling ATP release. *Proc Natl Acad Sci U S A* 95:15735–15740
214. Cotrina ML, Lin JH, Lopez-Garcia JC, Naus CC, Nedergaard M (2000) ATP-mediated glia signaling. *J Neurosci* 20:2835–2844
215. Cunha RA (2008) Different cellular sources and different roles of adenosine: A_1 receptor-mediated inhibition through astrocytic-driven volume transmission and synapse-restricted A_{2A} receptor-mediated facilitation of plasticity. *Neurochem Int* 52:65–72
216. Pascual O, Casper KB, Kubera C, Zhang J, Revilla-Sanchez R, Sul JY, Takano H, Moss SJ, McCarthy K, Haydon PG (2005) Astrocytic purinergic signaling coordinates synaptic networks. *Science* 310:113–116
217. Martin ED, Fernandez M, Perea G, Pascual O, Haydon PG, Araque A, Cena V (2007) Adenosine released by astrocytes contributes to hypoxia-induced modulation of synaptic transmission. *Glia* 55:36–45
218. Jeremic A, Jeftinija K, Stevanovic J, Glavaski A, Jeftinija S (2001) ATP stimulates calcium-dependent glutamate release from cultured astrocytes. *J Neurochem* 77:664–675
219. Domercq M, Brambilla L, Pilati E, Marchaland J, Volterra A, Bezzi P (2006) P2Y_1 receptor-evoked glutamate exocytosis from astrocytes: control by tumor necrosis factor- α and prostaglandins. *J Biol Chem* 281:30684–30696
220. Wirkner K, Gunther A, Weber M, Guzman SJ, Krause T, Fuchs J, Koles L, Norenberg W, Illes P (2007) Modulation of NMDA receptor current in layer V pyramidal neurons of the rat prefrontal cortex by P2Y receptor activation. *Cereb Cortex* 17:621–631
221. Mongin AA, Kimelberg HK (2002) ATP potently modulates anion channel-mediated excitatory amino acid release from cultured astrocytes. *Am J Physiol Cell Physiol* 283:C569–578
222. Kimelberg HK (2004) Increased release of excitatory amino acids by the actions of ATP and peroxynitrite on volume-regulated anion channels (VRACs) in astrocytes. *Neurochem Int* 45:511–519
223. Fellin T, Pascual O, Haydon PG (2006) Astrocytes coordinate synaptic networks: balanced excitation and inhibition. *Physiology (Bethesda)* 21:208–215
224. Zhang JM, Wang HK, Ye CQ, Ge W, Chen Y, Jiang ZL, Wu CP, Poo MM, Duan S (2003) ATP released by astrocytes mediates glutamatergic activity-dependent heterosynaptic suppression. *Neuron* 40:971–982
225. Bruner G, Murphy S (1990) ATP-evoked arachidonic acid mobilization in astrocytes is via a P2Y -purinergic receptor. *J Neurochem* 55:1569–1575
226. Bruner G, Murphy S (1993) Purinergic P2Y receptors on astrocytes are directly coupled to phospholipase A_2 . *Glia* 7:219–224
227. Gebicke-Haerter PJ, Wurster S, Schobert A, Hertting G (1988) P2 -purinoceptor induced prostaglandin synthesis in primary rat astrocyte cultures. *Naunyn Schmiedeberg Arch Pharmacol* 338:704–707
228. Takasaki I, Takarada S, Tatsumi S, Azegami A, Yasuda M, Fukuchi M, Tabuchi A, Kondo T, Tabuchi Y, Tsuda M (2008) Extracellular adenosine 5'-triphosphate elicits the expression of brain-derived neurotrophic factor exon IV mRNA in rat astrocytes. *Glia* 56:1369–1379
229. Tran MD, Neary JT (2006) Purinergic signaling induces thrombospondin-1 expression in astrocytes. *Proc Natl Acad Sci U S A* 103:9321–9326
230. Abbracchio MP, Ceruti S, Langfelder R, Cattabeni F, Saffrey MJ, Burnstock G (1995) Effects of ATP analogues and basic fibroblast growth factor on astroglial cell differentiation in primary cultures of rat striatum. *Int J Dev Neurosci* 13:685–693
231. Bolego C, Ceruti S, Brambilla R, Puglisi L, Cattabeni F, Burnstock G, Abbracchio MP (1997) Characterization of the signalling pathways involved in ATP and basic fibroblast growth factor-induced astrogliosis. *Br J Pharmacol* 121:1692–1699
232. Abe K, Saito H (1999) Effect of ATP on astrocyte stellation is switched from suppressive to stimulatory during development. *Brain Res* 850:150–157
233. Abbracchio MP, Brambilla R, Ceruti S, Cattabeni F (1999) Signalling mechanisms involved in P2Y receptor-mediated reactive astrogliosis. *Prog Brain Res* 120:333–342
234. Brambilla R, Burnstock G, Bonazzi A, Ceruti S, Cattabeni F, Abbracchio MP (1999) Cyclo-oxygenase-2 mediates P2Y receptor-induced reactive astrogliosis. *Br J Pharmacol* 126:563–567
235. Brambilla R, Ceruti S, Malorni W, Cattabeni F, Abbracchio MP (2000) A novel gliotic P2 receptor mediating cyclooxygenase-2 induction in rat and human astrocytes. *J Auton Nerv Syst* 81:3–9
236. Brambilla R, Neary JT, Cattabeni F, Cottini L, D'Ippolito G, Schiller PC, Abbracchio MP (2002) Induction of COX-2 and reactive gliosis by P2Y receptors in rat cortical astrocytes is dependent on ERK1/2 but independent of calcium signalling. *J Neurochem* 83:1285–1296
237. Ciccarelli R, Di Iorio P, Ballerini P, Ambrosini G, Giuliani P, Tiboni GM, Caciagli F (1994) Effects of exogenous ATP and related analogues on the proliferation rate of dissociated primary cultures of rat astrocytes. *J Neurosci Res* 39:556–566
238. Neary JT, Kang Y, Shi YF, Tran MD, Wanner IB (2006) P2 receptor signalling, proliferation of astrocytes, and expression of molecules involved in cell–cell interactions. *Novartis Found Symp* 276:131–143
239. Neary JT, Kang Y, Bu Y, Yu E, Akong K, Peters CM (1999) Mitogenic signaling by ATP/ P2Y purinergic receptors in astrocytes: involvement of a calcium-independent protein kinase C, extracellular signal-regulated protein kinase pathway distinct from the phosphatidylinositol-specific phospholipase C/calcium pathway. *J Neurosci* 19:4211–4220
240. Francke M, Weick M, Pannicke T, Uckermann O, Grosche J, Goczalik I, Milenkovic I, Uhlmann S, Faude F, Wiedemann P, Reichenbach A, Bringmann A (2002) Upregulation of extracellular ATP-induced Müller cell responses in a dispass model of proliferative vitreoretinopathy. *Invest Ophthalmol Vis Sci* 43:870–881
241. Francke M, Uhlmann S, Pannicke T, Goczalik I, Uckermann O, Weick M, Hartig W, Wiedemann P, Reichenbach A, Bringmann A (2003) Experimental dispass-induced retinopathy causes up-regulation of P2Y receptor-mediated calcium responses in Müller glial cells. *Ophthalmic Res* 35:30–41
242. Uckermann O, Uhlmann S, Weick M, Pannicke T, Francke M, Reichenbach A, Wiedemann P, Bringmann A (2003) Upregulation of purinergic P2Y receptor-mediated calcium responses in glial cells during experimental detachment of the rabbit retina. *Neurosci Lett* 338:131–134
243. Uhlmann S, Bringmann A, Uckermann O, Pannicke T, Weick M, Ulbricht E, Goczalik I, Reichenbach A, Wiedemann P, Francke

- M (2003) Early glial cell reactivity in experimental retinal detachment: effect of suramin. *Invest Ophthalmol Vis Sci* 44:4114–4122
244. Franke H, Krugel U, Illes P (1999) P2 receptor-mediated proliferative effects on astrocytes in vivo. *Glia* 28:190–200
 245. Franke H, Krugel U, Schmidt R, Grosche J, Reichenbach A, Illes P (2001) P2 receptor-types involved in astrogliosis in vivo. *Br J Pharmacol* 134:1180–1189
 246. Franke H, Kittner H, Grosche J, Illes P (2003) Enhanced P2Y₁ receptor expression in the brain after sensitisation with d-amphetamine. *Psychopharmacology (Berl)* 167:187–194
 247. Perez-Ortiz JM, Serrano-Perez MC, Pastor MD, Martin ED, Calvo S, Rincon M, Tranque P (2008) Mechanical lesion activates newly identified NFATc1 in primary astrocytes: implication of ATP and purinergic receptors. *Eur J Neurosci* 27:2453–2465
 248. Wang M, Kong Q, Gonzalez FA, Sun G, Erb L, Seye C, Weisman GA (2005) P2Y nucleotide receptor interaction with α integrin mediates astrocyte migration. *J Neurochem* 95:630–640
 249. Weisman GA, Wang M, Kong Q, Chorna NE, Neary JT, Sun GY, Gonzalez FA, Seye CI, Erb L (2005) Molecular determinants of P2Y₂ nucleotide receptor function: implications for proliferative and inflammatory pathways in astrocytes. *Mol Neurobiol* 31:169–183
 250. Washburn KB, Neary JT (2006) P2 purinergic receptors signal to STAT3 in astrocytes: Difference in STAT3 responses to P2Y and P2X receptor activation. *Neuroscience* 142:411–423
 251. Burgos M, Neary JT, Gonzalez FA (2007) P2Y₂ nucleotide receptors inhibit trauma-induced death of astrocytic cells. *J Neurochem* 103:1785–1800
 252. Fujita T, Tozaki-Saitoh H, Inoue K (2008) P2Y₁ receptor signaling enhances neuroprotection by astrocytes against oxidative stress via IL-6 release in hippocampal cultures. *Glia* 57:244–257
 253. Wu J, Holstein JD, Upadhyay G, Lin DT, Conway S, Muller E, Lechleiter JD (2007) Purinergic receptor-stimulated IP₃-mediated Ca²⁺ release enhances neuroprotection by increasing astrocyte mitochondrial metabolism during aging. *J Neurosci* 27:6510–6520
 254. Kim SG, Soltysiak KA, Gao ZG, Chang TS, Chung E, Jacobson KA (2003) Tumor necrosis factor α -induced apoptosis in astrocytes is prevented by the activation of P2Y₆, but not P2Y₄ nucleotide receptors. *Biochem Pharmacol* 65:923–931
 255. Chorna NE, Santiago-Perez LI, Erb L, Seye CI, Neary JT, Sun GY, Weisman GA, Gonzalez FA (2004) P2Y receptors activate neuroprotective mechanisms in astrocytic cells. *J Neurochem* 91:119–132
 256. Agresti C, Meomartini ME, Amadio S, Ambrosini E, Serafini B, Franchini L, Volonte C, Aloisi F, Visentin S (2005) Metabotropic P2 receptor activation regulates oligodendrocyte progenitor migration and development. *Glia* 50:132–144
 257. Moran-Jimenez MJ, Matute C (2000) Immunohistochemical localization of the P2Y₁ purinergic receptor in neurons and glial cells of the central nervous system. *Brain Res Mol Brain Res* 78:50–58
 258. Bernstein M, Lyons SA, Moller T, Kettenmann H (1996) Receptor-mediated calcium signalling in glial cells from mouse corpus callosum slices. *J Neurosci Res* 46:152–163
 259. Butt AM (2006) Neurotransmitter-mediated calcium signalling in oligodendrocyte physiology and pathology. *Glia* 54:666–675