

Functions of Noncoding RNAs in Neural Development and Neurological Diseases

Shan Bian · Tao Sun

Received: 18 July 2011 / Accepted: 21 September 2011 / Published online: 4 October 2011
© Springer Science+Business Media, LLC 2011

Abstract The development of the central nervous system (CNS) relies on precisely orchestrated gene expression regulation. Dysregulation of both genetic and environmental factors can affect proper CNS development and results in neurological diseases. Recent studies have shown that similar to protein coding genes, noncoding RNA molecules have a significant impact on normal CNS development and on causes and progression of human neurological disorders. In this review, we have highlighted discoveries of functions of noncoding RNAs, in particular microRNAs and long noncoding RNAs, in neural development and neurological diseases. Emerging evidence has shown that microRNAs play an essential role in many aspects of neural development, such as proliferation of neural stem cells and progenitors, neuronal differentiation, maturation, and synaptogenesis. Misregulation of microRNAs is associated with some mental disorders and neurodegeneration diseases. In addition, long noncoding RNAs are found to play a role in neural development by regulating the expression of protein coding genes. Therefore, examining noncoding RNA-mediated gene regulations has revealed novel mechanisms of neural development and provided new insights into the etiology of human neurological diseases.

Keywords Neural development · Mental disorders · Neurodegeneration diseases · Noncoding RNAs · MicroRNAs (miRNAs) · Long noncoding RNAs (lncRNAs) · Neural stem cells

Introduction

Noncoding RNAs (ncRNAs) are functional RNA molecules that do not show protein translation capability. ncRNAs consist of ribosomal RNAs (rRNAs), transfer RNAs (tRNAs), piwi-interacting RNAs (piRNAs), microRNAs (miRNAs), and long noncoding RNAs (lncRNAs) and so on. These ncRNAs have been shown to play distinct but also conserved roles in normal development and pathological processes in invertebrates and vertebrates. Studies have reported that many miRNAs as well as lncRNAs are specifically expressed in the central nervous system (CNS) [1–5]. In this review, we will focus on revealing the functions of miRNAs and lncRNAs in CNS development and neurological disorders.

miRNAs are ~22 nucleotides (nt) highly conserved small noncoding RNAs found in almost all eukaryotic cells [6]. Like coding genes, miRNAs are mainly transcribed by the RNA Polymerase II (Pol II) into long primary miRNA transcripts (pri-miRNAs), while a subset of miRNAs has been shown to be transcribed by RNA Pol III [7]. Pri-miRNAs are cleaved by Drosha, a class 2 RNase III enzyme, to produce short hairpin stem-loop structures, known as precursor miRNAs (pre-miRNAs) [8]. Pre-miRNAs are then processed into about 20–25 nucleotides double-stranded mature miRNAs by RNase III enzyme Dicer [9]. The duplex undergoes unwinding and the strand with the weakest base pairing at its 5' terminus, together with Dicer and many associated proteins including Argonaute proteins, forms an active RNA-induced silencing complex (RISC) [10, 11]. A miRNA recognizes and binds to the non-coding portion of target messenger RNAs (mRNAs), typically in the 3'-untranslated region (3'-UTR), by imperfect complementary sequence recognition,

S. Bian · T. Sun (✉)
Department of Cell and Developmental Biology,
Weill Medical College of Cornell University,
1300 York Avenue, Box 60, New York, NY 10065, USA
e-mail: tas2009@med.cornell.edu

particularly the seed sequence of the mature miRNA [11, 12]. Once bound, the RISC can negatively regulate miRNA targets by degrading the mRNA or repressing its translation [6, 13]. miRNAs are able to target numerous mRNAs and so their regulatory potential is large. Moreover, several evidences indicate regulatory feedback loop between miRNA and their targets [14].

miRNAs are classified into intergenic and intragenic miRNAs, depending on their genomic location [15–17]. Intergenic miRNAs are located at intergenic regions (between genes). These miRNAs are usually transcribed using their own promoters. Intragenic miRNAs normally lie in intronic regions of coding genes with the same orientation. They are often transcribed together with the host genes. Additionally, some miRNAs can be co-transcribed as clusters [18].

Emerging evidence has demonstrated the potential functions of lncRNAs in many biological events. The majority of the human genome has previously been considered as “junk” DNA since only about 1.5% of the human genome, which occupies over 3 billion DNA base pairs, consists of protein-coding genes [19]. A recent study of a large-scale complementary DNA (cDNA) sequencing project has shown that only one fifth of transcripts of the human genome are protein-coding genes, while the rest at four fifths are RNA transcripts that do not encode proteins [20]. These RNA transcripts are normally longer than 200 nucleotides; thus, they are called lncRNAs. Except that no open reading frame was found within lncRNAs, they share many features with coding mRNAs such as 5' capping and usually contain exons and introns [21]. However, some studies point out that lncRNAs may or may not be polyadenylated and are often transcribed from either strand within a protein-coding gene [22]. Because of the huge number and different features of the genomic location and orientation of lncRNAs, the potential functions of lncRNAs have not yet been well characterized. However, it is unlikely that lncRNAs are “junk” transcripts. lncRNAs have been shown to modulate gene transcription via different mechanisms. For example, some *cis*-antisense lncRNAs complementary to protein-coding transcripts regulate the expression of the coding genes [23–27]; some lncRNAs modulate the function of transcription factors by acting as co-regulators [26, 28–32]. Moreover, some studies have shown that lncRNAs are involved in epigenetic regulation such as chromatin and histone modification and X-chromosome inactivation [33–39].

Therefore, noncoding RNAs such as miRNAs and lncRNAs participate in regulating gene expression in many ways. The underlying mechanisms of noncoding RNA functions in normal development and under disease conditions are beginning to be uncovered.

miRNA Functions in CNS Development, a Dicer Ablation Approach

Since Dicer is the key enzyme in miRNA processing, several studies have reported the global effects of miRNAs in CNS development by ablating Dicer and in turn blocking the biogenesis of all miRNAs in specific regions and at different developing stages in the CNS using tissue-specific Cre lines.

Conditional deletion of *Dicer* from the mouse cerebral cortex using the *Emx1-Cre* line results in a significant reduction in cortical size, which is caused by reduced neural stem cell and neural progenitor pool, increased apoptosis, and impaired neuronal differentiation [40–43]. *Dicer* ablation from the mouse CNS using the *Nestin-Cre* line has confirmed the cortical defects and revealed abnormal expansion and differentiation of oligodendrocytes in the spinal cord [43]. *Dicer* deletion from post-mitotic neurons in the cortex and the hippocampus using *calmodulin kinase II* (*CaMKII*) promoter-driven *Cre* transgenic mice results in smaller cortex, enhanced cortical cell death, impaired axonal path finding, and dendritic branching in the hippocampal CA1 region [44]. Ablation of *Dicer* in the hippocampus at different time points using *Emx1-Cre*, *Nestin-Cre*, and *Nex-Cre* lines has revealed miRNA functions in hippocampal progenitor proliferation and differentiation [45]. Conditional deletion of *Dicer* in the dopaminergic neurons in the basal ganglia using a *dopamine receptor-1* (*DR-1*) *Cre* line causes reduced brain size and mass and smaller striatal medium spiny neurons [46].

Since *Wnt1-Cre* is active in neural crest cells, apoptosis of neural crest-derived cells and subsequently loss of the enteric, sensory, and sympathetic neurons are reported in *Wnt1-Cre Dicer* conditional knockout mice [47]. Another investigation of *Wnt1-Cre*-mediated *Dicer* ablation has found malformation of the midbrain and cerebellum, reduced dopaminergic neuron differentiation, and malformation of the dorsal root ganglia [48]. Ablation of *Dicer* using the *Olig1-Cre* line does not affect motor neuron development, but reduces the number of astrocytes and oligodendrocyte progenitor cells (OPCs) in the spinal cord, and results in a great reduction of mature oligodendrocytes in the brain and an impaired myelination [49, 50]. Similarly, *Dicer* ablation using the 2', 3'-cyclic nucleotide 3' phosphodiesterase (*Cnp*)-*Cre* line, which is active in the oligodendrocyte lineage, causes a great reduction of OPCs and mature oligodendrocytes [51].

Conditional ablation of *Dicer* in specific regions in the CNS and in distinct cell lineages has demonstrated the importance of the miRNA-mediated pathway in CNS development and in the differentiation of different cell types. However, because Dicer is also involved in maintaining heterochromatin assembly, likely by the short interfering

RNA (siRNA) pathway, the CNS defects in *Dicer* knockout mice need to be carefully interpreted [52, 53]. Examining functions of individual miRNAs will help reveal the precise roles of miRNAs in CNS development.

miRNA Functions in the Development of Neural Stem Cells and Progenitors

Neural stem cells (NSCs) and neural progenitors (NPs) are identified in both embryonic and adult CNS in mammals [54]. NSCs can self-renew and differentiate into multiple lineages such as neurons, astrocytes, and oligodendrocytes [54, 55]. Both intrinsic factors, for instance transcription factors and epigenetic and post-transcriptional regulators, and extrinsic signals from the stem cell microenvironment play important roles in regulating self-renewal and the differentiation of NSCs [55–57]. Several reports have shown that miRNAs are also essential for the development of NSCs and NPs in the CNS in vivo and in cultures (Fig. 1).

let-7, the first identified miRNA [58], has been shown to regulate NSC proliferation and differentiation by targeting the nuclear receptor TLX and the cell cycle regulator cyclin D1 [59]. Over-expression of *let-7b* inhibits NSC proliferation and enhances differentiation, while knockdown of *let-7b* promotes NSC proliferation [59]. It appears that the expression levels of let-7 in NSCs are controlled by a feedback regulation of Lin-28, a pluripotency factor that controls miRNA processing in NSCs [60]. Lin-28 binds to the *let-7* precursor and inhibits its processing by Dicer. On the other hand, the expression of Lin-28 is repressed by let-7 and miR-125, allowing the maturation of let-7. This feedback loop reveals an autoregulation between miRNA let-7 and miR-125, and the transcription factor Lin-28 during NSC development [60] (Fig. 2a).


miR-124 is identified as a brain-enriched miRNA and its expression is upregulated during neuronal differentiation [61]. Over-expressing *miR-124* in cultured NSCs and in embryonic cortical NPs using lenti-virus and in utero electroporation, respectively, promotes neurogenesis and stimulates

cortical progenitor migration [62]. In the adult brain, NSCs are identified in the subventricular zone (SVZ). In cultured adult NSCs derived from the SVZ and in the SVZ in vivo, knocking down *miR-124* causes an increase of dividing cells and a decrease of post-mitotic neurons, while over-expressing *miR-124* enhances neuronal differentiation [63]. Moreover, miR-124 regulates neuronal differentiation by suppressing Sox9 expression in adult NSCs [63]. A recent study has shown that miR-124 regulates neuronal differentiation through a mutual inhibition mechanism of Ephrin-B1 [64]. In addition, miR-124 promotes differentiation of NPs by modulating a network of nervous system-specific alternative splicing through suppressing the expression of PTBP1, which encodes a global repressor of alternative pre-mRNA splicing [65]. Together miR-124 plays a general role in promoting the differentiation of embryonic and adult NSCs and NPs. It appears that miR-124 executes its function through repressing various targets (Fig. 3a).

miR-9 is another CNS-enriched miRNA. miR-9 is shown to inhibit NSC proliferation but promote differentiation through feedback regulation of a nuclear receptor TLX [66] (Fig. 2b). In human embryonic stem cell (ESC)-derived NPs, miR-9 is shown to have a positive effect on proliferation and to promote migration by directly targeting *Stmn1*, which increases microtubule instability [67]. The opposite effect of miR-9 on proliferation is perhaps caused by differential physical contacts of miR-9 with target genes and the different culture systems.

In the CNS of *Xenopus*, miR-9 function is required for the development of the forebrain and hindbrain through different mechanisms: progenitors lacking *miR-9* in the forebrain undergo apoptosis, but *miR-9*-deficient progenitors in the hindbrain fail to exit the cell cycle [68]. In zebrafish, miR-9 promotes differentiation of NPs that give rise to neurons at the midbrain–hindbrain domain and controls the organization of the midbrain–hindbrain boundary by targeting several genes in the Fibroblast growth factor (Fgf) signaling, such as *fgf8-1* and *fgfr1* [69]. In the chick spinal cord, miR-9 defines a subtype of motor neurons that project axons to the axial muscles by specifically targeting *FoxP1* [70]. In the mouse

Fig. 1 A list of representative miRNAs that function on neural stem cell self-renewal, neuronal differentiation, and neuronal maturation






Fig. 2 A feedback regulation of miRNAs and their target genes. Protein coding genes inhibit (a, b, d) or promote (c) the biogenesis of miRNAs, while miRNAs post-transcriptionally silence the output of these proteins

brain, miR-9 function is demonstrated by the generation of *miR-9-2* and *miR-9-3* double-knockout mice. *miR-9* double-mutants show reduced cortical layers, disordered migration of

interneurons, and misrouted thalamocortical axons and corticofugal axon projections, suggesting an important role of miR-9 in NP proliferation, differentiation, and migration




Fig. 3 miRNAs, such as miR-124 (a), miR-9 (b), miR-134 (c), and miR-137 (d), control the proliferation and apoptosis of neural stem cells and progenitors, neuronal differentiation, migration, and maturation and synaptogenesis by targeting various genes

during brain development [71]. Moreover, it appears that miR-9 regulates multiple target genes, including *Foxg1*, *Pax6*, and *Gsh2*, which have been shown to be essential in cortical development [71]. Therefore, miR-9 plays an important role in many aspects of CNS development, such as neuronal differentiation and migration, and cell type specification (Fig. 3b).

The above studies have demonstrated that the major role of let-7, miR-124, and miR-9 is to induce the differentiation of NSCs and NPs into specific cell types. miRNAs that promote proliferation of NSCs and NPs have also been identified (Fig. 1). miR-134 plays a role in enhancing the proliferation of cortical NPs by targeting double-cortin (*Dcx*) and/or Chordin-like 1 (*Chrdl-1*) [72] (Fig. 3c). miR-25 is shown to be a major player in the miR-106-25 cluster in neural development. Overexpression of *miR-25* but not *miR-106b* and *miR-93* promotes adult NP proliferation [73]. Interestingly, the expression of the miR-106-25 cluster is regulated by FoxO3, a transcription factor maintaining the NSC population [74].

miRNAs regulate target gene expression at a post-transcriptional level. Recent work has shown that miRNAs themselves are also controlled by epigenetic regulators in NSC development. miR-137, whose expression is co-regulated by DNA methyl-CpG-binding protein (MeCP2) and transcriptional factor Sox2, is shown to modulate adult NSC proliferation and cell fate determination by targeting *Ezh2*, a histone methyltransferase and Polycomb group protein [75] (Fig. 3d). Ectopic expression of *miR-137* in adult NSCs enhances proliferation, while knockdown of *miR-137* promotes the differentiation of adult NSCs [75]. miR-184 expression is suppressed by methyl-CpG binding protein 1, and miR-184 promotes adult NSC proliferation by repressing the expression of Numb-like (*Numb*) [76].

Taken together, self-renewal, proliferation, and differentiation of NSCs and NPs are controlled by complex gene regulation networks that consist of both protein-coding genes and noncoding miRNAs. During proliferation and differentiation of NSCs and NPs, one miRNA can have multiple target genes (Figs. 1 and 3). The availability of physical contacts and the binding affinity of a miRNA and its targets perhaps determine the interactions of the miRNA with the specific targets. The interactions of miRNAs and their target genes eventually produce proper protein output of key factors that control the precise specification of distinct cell types in different regions of the CNS and ensure accurate CNS functions.

miRNA Regulation of Neuronal Maturation and Synapse Formation

Neuronal maturation, neurite outgrowth, and synaptogenesis allow the assembly of a functional CNS to control complex

behaviors. miRNAs have been shown to play a critical role in these progresses.

miR-124 functions not only in neuronal differentiation but also in neurite outgrowth and synaptogenesis (Fig. 3a). In primary cortical neurons, ectopic miR-124 expression enhances neurite outgrowth [77]. *Cdc42* and *Rac1* belong to the Rho family and have been shown to regulate actin filaments and microtubules [78]. miR-124 regulates neurite outgrowth by inhibiting *Cdc42* expression and altering the localization of *Rac1* in mouse P19 embryonic carcinoma cells [77]. In the nervous system of *Aplysia californica*, miR-124 is highly expressed in the sensory motor synapse [79]. Over-expression of miR-124 results in a remarkable reduction of long-term facilitation of the synapses, while inhibition of miR-124 shows the opposite phenotype. The regulation of miR-124 on long-term plasticity is through targeting cAMP response element-binding (CREB), a cellular transcriptional factor [79]. Furthermore, miR-124a is shown to be required for hippocampal axogenesis through suppression of LIM homeobox 2 [80].

miR-137 has been reported to promote NSC proliferation and inhibit neuronal differentiation [75]. A recent study has shown that miR-137 negatively regulates neuronal maturation (Fig. 3d). Ectopic expression of *miR-137* inhibits neuronal maturation, dendritic morphogenesis, and spine development both in cultured hippocampal neurons and in mouse brains. Conversely, lack of *miR-137* promotes neuronal maturation [81]. miR-137 regulates neuronal maturation by targeting Mind bomb 1 (*Mib1*), an ubiquitin ligase, since the ectopic expression of *Mib1* enhances dendritic elongation and rescues the defects in neuronal maturation caused by miR-137 [81].

miR-133b is specifically expressed in dopaminergic neurons (DNs) in midbrains and is deficient in the midbrain of Parkinson's disease patients [82]. miR-133b is shown to inhibit the maturation and function of DNs by targeting paired-like homeodomain transcription factor *Pitx3*. In *Pitx3*-deficient mouse model, miR-133b expression is significantly reduced, suggesting that miR-133b expression is under the regulation of *Pitx3* [82, 83]. These reports indicate a negative feedback regulation between *Pitx3* and miR133b in balancing the development of DNs [82] (Fig. 2c).

Several studies have shown the importance of miRNAs in synaptogenesis. The *miR-132/miR-212* cluster is reported to mediate dendritic growth and spine formation [84]. Conditional deletion of the *miR-132/miR-212* cluster results in a great reduction in the dendrite length and spine density. Additionally, *miR-132* over-expression transgenic mice show decreased dendritic spine density and impaired novel object recognition memory [85]. As a predicted target of miR-132, MeCP2 is down-regulated in *miR-132* transgenic mice [85]. Furthermore, miR-132 is identified as a fragile X

mental retardation protein (FMRP)-associated miRNA and affects neuronal morphology and modifies synaptic strength in a FMRP-dependent manner [86].

miR-134 is a brain-specific miRNA localized to the synapto-dendritic compartment of rat hippocampal neurons. miR-134 is found to negatively regulate the size of dendritic spines, which are the post-synaptic sites of excitatory synaptic transmission, by targeting and repressing LIM domain kinase 1 [87] (Fig. 3c). Based on a large-scale screening in the synapto-dendritic compartment in primary hippocampal neurons, miR-138 is found to be highly expressed in synaptosomes, a biochemical fraction that is important for synaptic protein synthesis and storage [88]. In a single-cell fluorescent sensor assay, inhibition of miR-138 leads to a significant increase of spinal volume without changing the synaptic puncta number [89]. These functional regulations of miR-138 on synaptic development and synaptic transmission are related to its target acyl-protein thioesterase 1, whose activity is required for spine enlargement [89].

Functions of miRNAs in neurite outgrowth and synaptogenesis suggest that miRNAs may directly regulate the local translation of target proteins that are essential for these biological events [90]. The transportation of miRNAs to a subcellular location in neurons and the turnover of miRNAs on the binding sites of target genes during neuronal maturation and synaptogenesis still remain to be interesting research topics.

miRNAs in Neurological Disorders

Misregulation of miRNAs is associated with the pathogenesis of different human neurological diseases such as neurodevelopmental disorders, neurodegeneration diseases, and affective mental disorders (Table 1).

miRNAs have been proven to play a critical role in the etiology of human neuropsychiatric disorders. Variations of the *SLITRK1* gene in the 3'-UTR binding site for human miR-189 are detected in DNA samples of patients with Tourette's syndrome, a developmental neuropsychiatric disorder [91]. The Fragile X syndrome, a common human mental retardation, is caused by a loss of RNA-binding protein FMRP. FMRP is found to be associated with the miRNA processing pathway [92]. miR-19b, miR-302b*, and miR-323-3p have been reported to repress the expression of FMRP, suggesting a role of miRNAs in the pathogenesis of Fragile X syndrome [93]. A double-stranded RNA-binding protein DGCR8 is involved in miRNA processing and functions as an essential Drosha co-factor [94]. *DGCR8* is commonly deleted in patients with DiGeorge syndrome, characterized by a variety of defects, including schizophrenia and microcephaly [95]. Abnormal miRNA biogenesis may contribute to the etiology of DiGeorge syndrome [96].

Schizophrenia is a syndrome of reality distortion along with alterations in thought, emotional processing, volition, and cognition. Several miRNAs are found to be more highly expressed in the prefrontal cortex of schizophrenic patients than normal subjects [97]. A report has shown that miR-181b is upregulated, while its target genes calcium sensor gene visinin-like 1 (VSNL1) and ionotropic AMPA glutamate receptor subunit (GRIA2), which are associated with schizophrenia, are downregulated in the superior temporal gyrus of schizophrenic patient brain samples [98]. *N*-Methyl-D-aspartate (NMDA) glutamate receptor signaling regulates neurotransmission and synaptic plasticity and executes many brain functions. Dysfunction of the NMDA receptor signaling pathway results in cognitive disorders, which show symptoms of schizophrenia and autism. miR-219 is shown to specifically target calcium/calmodulin-dependent protein kinase II γ subunit (CaMKII γ), a component of the NMDA receptor signaling [99]. Knockdown of *miR-219* in the mouse brain alters the behavioral responses associated with disrupted NMDA receptor signaling [99]. Furthermore, autism spectrum disorder (ASD) is a neural developmental disorder featured in an impaired communication ability and social interaction, as well as repetitive or restricted behavior. miRNA expression profiles of ASD patients have been performed to explore a potential role of miRNAs in the pathogenesis of autism [100–102].

Alzheimer's disease (AD) is a neurodegenerative disease that is the most common form of dementia. It includes changes in memory and functions of the hippocampus. Comparisons in the hippocampus of fetal, adult, and AD brains have identified miRNAs that are highly expressed in the fetal hippocampus but differentially regulated in AD patients, for instance, an upregulation of miR-9 and miR-128 in AD patients [103]. Another expression profiling analysis of miRNAs in short post-mortem interval human brains has revealed an upregulation of miR-9, miR-125b, and miR-146a in the temporal lobe in the AD patient brains [104].

Several miRNAs are found to be downregulated in AD brain tissues, such as miR-15a and the miR-107 and miR-29a/b-1 cluster [105–107]. The expression of miR-107 is greatly decreased even in the early stages of AD patients [105]. β -Site amyloid precursor protein-cleaving enzyme (BACE1), which contributes to the accumulation of amyloid- β peptides (A β), the key molecule in the pathogenesis of AD, has been identified as the target of miR-107 [105]. During AD progression, the expression of BACE1 is upregulated but miR-107 expression is decreased [105]. Moreover, the miR-29a/b-1 cluster is significantly decreased in AD patients accompanied by an abnormally high expression of BACE1. In a cell culture system, the miR-29a/b-1 cluster is shown to negatively regulate the

Table 1 miRNAs involved in neurological disorders

Neurological disorders	miRNAs	Proven targets	Potential mechanisms	References
Tourette's syndrome	miR-189	SLITRK1	Mutation at the binding site for miR-189 in the target 3'UTR	[91]
Fragile X syndrome	miR-19b, miR-302b*, miR-323-3p	FMRP	Repression of FMR1 expression level	[93]
Schizophrenia	miR-181b	VSNL1	Upregulation of miR-181b in schizophrenia	[98]
	miR-219	GRIA2	NMDA signaling and miR-219 forming a feedback loop in schizophrenia	[99]
		CaMKII γ		
Alzheimer's disease	miR-107	BACE1	Downregulation of miR-107 and increase of its target BACE1 in AD	[105]
	miR-29a/b	BACE1	Downregulation of miR-29a/b and increase of its target BACE1 in AD	[106]
	miR-298, miR-328	BACE1	Repression of BACE1 expression and accumulation of A β	[108]
	miR-101	APP	Repression of APP expression and accumulation of A β	[109]
	miR-146a	CFH	Downregulation of miR-146a in AD	[110]
Huntington's disease	miR-9	REST	Upregulation of miR-9/9* in HD	[114]
	miR-9*	CoREST		
	miR-34b	?	A potential biomarker for HD	[115]
Parkinson's disease	miR-7, miR-153	α -Synuclein	Repression of α -synuclein accumulation	[116]
Prion disease	miR-433	FGF20	A SNP in the miR-433 binding site of FGF20 3'UTR in AD	[118]
	miR-191	EGR1	De-regulation after infection with mouse-adapted scrapie	[119]
	miR-342-3p, miR-320, let-7b, miR-328, miR-128, miR-139-5p, miR-146a, miR-337-3p, miR-338-3p	?		
Down's syndrome	miR-155, miR-802	MeCP2	Upregulation of miR-155 and miR-802 in DS and suppression of MeCP2	[124]
	miR-125b	CDKN2A	Upregulation of miR-125b in DS and suppression of CDKN2A	[125]
Stroke	miR-21	FASLG	Upregulation of miR-125b in stroke and suppression of FASLG	[127]
	miR-124a	JAG1	Downregulation of miR-124a after stroke	[128]
Epilepsy	miR-146a	?	Upregulation after TLE	[129]
Depression and bipolar disorder	miR-182	ADCY6, CLOCK, DSIP	Polymorphism in the rs76481776 of the pre-miR-182	[130]
	miR-30e	?	Polymorphism in the ss178077483 of the pre-miR-30e	[131]
	miR-134	?	A potential biomarker for bipolar disorders	[133]
	miR-16	SERT	Upregulation of miR-16 after SSRI treatment	[134]
	miR-34a	GRM7	Downregulation of miR-34a after treatment using mood stabilizers	[135]

expression of BACE1 and the accumulation of A β peptide [106]. In addition, miR-298 and miR-328 are reported to repress the expression of BACE1 [108]. Furthermore, the expression of amyloid precursor protein (APP) and accumulation of A β is negatively regulated by miR-101 in cultured hippocampal neurons, suggesting a potential role of miR-101 in AD progression [109]. miR-146a, which is regulated by NF- κ B signaling, is upregulated in AD brains, and miR-146a targets complement factor H (CFH), a repressor of the inflammatory response of the brain [110].

Huntington's disease (HD) is a genetic disorder with programmed degeneration of brain neurons. Patients have behavioral and motor symptoms and cognitive decline leading to dementia. The expression profile of miRNAs in two transgenic mouse models of HD has shown that miRNAs such as miR-22, miR-29c, and miR-128 are downregulated [111]. In human samples, the expression of miR-9/9*, miR-29b, and miR-124a is downregulated, while miR-132 is upregulated with increasing HD grade compared to controls [112]. Previous studies indicate that aberrant cellular distribution of RE1-silencing transcription factor (REST) is one of the molecular mechanisms underlying HD etiology. The REST repressor complex, including mSin3, REST co-repressor 1 (CoREST), and MeCP2, regulate the expression levels of miR-9/9*, miR-124a, and miR-132 [113]. Interestingly, miR-9 and miR-9* have targeting effects on REST and CoREST, respectively, suggesting that REST silencing complex and miR-9/9* form a negative feedback loop during HD development [114] (Fig. 2d). Furthermore, miR-34b is elevated in plasma from HD patients even before the symptom onsets, suggesting miR-34b to be a potential biomarker for HD diagnosis [115].

Parkinson's disease (PD) results from a loss of dopaminergic neurons in the substantia nigra. It involves abnormalities in movement variably accompanied by sensory, mood, and cognitive changes. Alpha-synuclein accumulation is related to the pathogenesis of PD. A study has shown that miR-7 and miR-153 negatively regulate α -synuclein in primary neurons [116]. In the 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine-induced neurotoxin PD mouse model, miR-7 is downregulated, suggesting that miR-7 may protect cells from oxidative stress [117]. Moreover, a recent study has demonstrated that a single nucleotide polymorphism (SNP) in the 3'-UTR of FGF20 destroys a target site for miR-433. Thus, miR-433 is linked to an increased risk of AD by indirectly causing overexpression of α -synuclein [118].

miRNAs are also associated with prion-induced neurodegeneration [119, 120]. In the mouse brains infected with mouse-adapted scrapie, miRNAs such as miR-342-3p, miR-320, and let-7b are greatly upregulated, while miR-337-3p and miR-338-3p are downregulated during the disease process [119]. Among potential targets of these miRNAs

predicted by computational algorithms, 119 genes are also de-regulated during the prion-induced neurodegeneration [119].

Down's syndrome (DS), also called trisomy 21, is a genetic disease caused by an extra copy of the entire or part of the human chromosome 21. DS patients usually have physical growth abnormality, heart defects, leukemia, low-cognitive ability, and Alzheimer's disease [121]. Since DS is caused by an extra copy of chromosome 21, which leads to overexpression of many genes located on it, it is possible that miRNAs located on chromosome 21 play a role during the DS pathological process by negatively regulating the expression of target genes and subsequently contribute to the symptoms of DS. Indeed miRNAs that are located on chromosome 21, including let-7c, miR-99a, miR-125b-2, miR-155, and miR-802, are found upregulated in the fetal hippocampus and heart samples from DS patients, while their target genes are found to be decreased [122, 123]. Gain and loss of function of *miR-155* and *miR-802* have shown that they target MeCP2, which is downregulated in the DS brain [124]. In addition, an upregulation of miR-125b and downregulation of its target cyclin-dependent kinase inhibitor 2A (CDKN2A) are found in advanced AD and DS brains [125].

Cerebral infarction (stroke) is related to a disturbance in the blood supply to the brain. miRNA expression profiling in young stroke patients has shown 138 upregulated miRNAs and 19 downregulated miRNAs [126]. miR-21 is found to be upregulated in neurons in the ischemic boundary zone after stroke in rat brains. miR-21 protects ischemic neurons, after oxygen and glucose deprivation, from cell death by targeting tumor necrosis factor α family member FAS ligand gene (FASLG), a cell death inducer [127]. Recent study has shown that miR-124a is downregulated in neural progenitor cells of the SVZ after stroke [128]. By targeting Notch signaling ligand Jagged-1 (JAG1), miR-124a impairs precursor proliferation and promotes neuronal differentiation after stroke [128].

In the temporal lobe epilepsy (TLE) rat model as well as in human TLE, the expression of miR-146a is found to be upregulated and is expressed in reactive astrocytes in the hippocampus, mainly in the regions where neuronal cell loss and reactive gliosis occur [129]. This observation implies the possible involvement of miR-146a in the modulation of astroglial inflammatory response during the TLE processes and provides a potential target for the clinical treatment of TLE.

Depression is an affective mental disorder with a low mood, a loss of interests in activities, and sometimes a cognitive impairment. A study has shown an association of the T allele of the rs76481776 polymorphism of pre-miR-182 with late insomnia in major depression (MD) patients [130]. Overexpression of *miR-182* causes reduction of miR-182 targeting genes adenylate cyclase type 6 (ADCY6), circadian

Locomotor Output Cycles Kaput (CLOCK) and Delta sleep-inducing peptide (DSIP), which are involved in molecular clock regulation, suggesting that miR-182 might contribute to the dysregulation of circadian rhythms in MD patients with insomnia [130]. The polymorphism in the ss178077483 of pre-miR-30e is also found to be associated with a higher risk of MD disorder [131].

A miRNA expression profiling analysis in the prefrontal cortex of patients with bipolar disorder (BD), also known as manic–depressive disorder, has identified differentially expressed miRNAs [132]. miR-134 is suggested as a biomarker of mania episodes in BD, based on an analysis of expression levels of miRNAs in the circulating blood before and after the treatment of bipolar mania patients [133]. In addition, chronic treatment with selective serotonin reuptake inhibitor (SSRI) fluoxetine in mice results in an upregulation of miR-16 and downregulation of SERT, which is a target of miR-16, suggesting that miRNAs participate in the therapeutic action of SSRI [134]. Several miRNAs are found to be dysregulated after chronic treatment with mood stabilizers lithium and valproate [135]. Among these miRNAs, miR-34a expression is reduced after the treatment, while its target metabotropic glutamate receptor 7 (GRM7) is upregulated [135]. These studies suggest a role of miRNAs in the clinical therapy of antidepressant medications.

In summary, these studies indicate that similar to coding genes, misregulation of miRNAs also contributes to the pathogenesis of neurodegeneration diseases and mental disorders (Table 1). miRNA expression profiles using patient samples have shown that the expression levels of many miRNAs are altered. Thus, it is a daunting task to figure out major miRNAs that are directly associated with the causes and progression of neurological diseases. Moreover, it is likely that neurological disorders are the result of altered networks of gene expression regulations of coding genes and noncoding RNAs.

lncRNAs in CNS Development

Based on in situ hybridization data from the Allen Brain Atlas (<http://www.brain-map.org/>), 849 lncRNAs from diverse locations in the genome have shown distinct expression patterns in different regions, cell types, and subcellular localization in adult mouse brains [4]. Similar to miRNAs, lncRNAs are also classified into intergenic and intragenic populations. Even though the functions of lncRNAs in CNS development are largely unclear, studies of intragenic lncRNAs that are associated with protein-coding genes in the genome are emerging. Bioinformatic investigation of genomic locations and comparisons of expression patterns of lncRNAs and their adjacent protein coding transcripts have

implied a potential *cis*-regulation mechanism of lncRNAs in modulating the transcription of coding genes in the developing and adult mouse brains [136].

Nkx2.2 antisense (Nkx2.2as) is an antisense lncRNA to *Nkx2.2* gene, which is expressed in the developing mammalian forebrain and is required for oligodendrocyte development [137] (Fig. 4a). Ectopic expression of *Nkx2.2as* in NSCs in cultures induces oligodendrocyte differentiation through an upregulation of the *Nkx2.2* mRNA level, suggesting that *Nkx2.2as* regulates NSC differentiation, at least partially, by promoting *Nkx2.2* expression [25].

Embryonic ventral forebrain-1 (Evf1) is a 2.7-kb lncRNA transcribed upstream of the mouse *Dlx-6* gene [138]. As an alternatively spliced form of *Evf1*, *Evf2* is




Fig. 4 Regulatory functions of representative long noncoding RNAs on the expression of protein coding genes. **a** *Nkx2.2 antisense (Nkx2.2as)* is an antisense lncRNA to *Nkx2.2* gene and promotes *Nkx2.2* expression. **b** *Evf2* is transcribed from the intergenic region between the *Dlx-5* and *Dlx-6* loci and is overlapped with *Dlx-5/6* enhancer i (*ei*) and enhancer ii (*eii*) sequences. *Evf2* acts as a transcriptional co-activator of *Dlx-2* and activates the *Dlx5/6* enhancer (modified the figure from [26]). **c** *Sox2 overlapping transcript Sox2OT* is a lncRNA containing *Sox2* gene and shares the same transcriptional orientation with *Sox2*. *Sox2OT* is expressed in the neurogenic region of the adult mouse brain

transcribed from the intergenic region between the *Dlx-5* and *Dlx-6* loci and is overlapped with the conserved *Dlx-5/6* intergenic enhancer [26, 139] (Fig. 4b). *Dlx-6* is a homeobox containing transcription factor and plays an important role in forebrain neurogenesis [140]. Induced by the Sonic hedgehog (Shh) signaling pathway, *Evf2* has been proven to function as a transcriptional co-activator of *Dlx-2* and activates the *Dlx5/6* enhancer during forebrain development [26]. Deletion of *Evf2* results in a reduction of GABAergic interneurons and impaired synaptic inhibition in the developing hippocampus [141].

Sox2 is a transcription factor and plays a key role in the maintenance of the undifferentiating state of embryonic and adult NSCs [142]. *Sox2 overlapping transcript (Sox2OT)* is a lncRNA containing *Sox2* gene and shares the same transcriptional orientation with *Sox2* (Fig. 4c). Similar to *Sox2*, *Sox2OT* is stably expressed in mouse embryonic stem cells and down-regulated during differentiation. *Sox2OT* is expressed in the neurogenic region of the adult mouse brain and is dynamically regulated during vertebrate CNS development, implying its role in regulating NSC self-renewal and neurogenesis [4, 143].

Retinal noncoding RNA 2 (RNCR2), an intergenic lncRNA also known as *Gomafu* and *Miat*, is an abundant polyadenylated RNA in the developing retina [144]. *RNCR2* is highly expressed in both mitotic and post-mitotic retinal progenitors. Knockdown of *RNCR2* leads to an increase of amacrine cells and Müller glial cells in post-natal retina. Mis-localization of *RNCR2* from nuclear to cytoplasm photocopies the effects caused by *RNCR2* knockdown, suggesting that *RNCR2* is required for retinal precursor cell specification [145].

lncRNAs in Neurological Disorders

Scattered studies have shown that lncRNAs are also involved in neurobiological disorders. *Sox2OT* has been identified as one of the potential biomarkers of AD for both early and late stages [146]. *BC200 RNA* is a 200-nt noncoding RNA that is expressed in the nervous system [147]. In the human nervous system, *BC200 RNA* is expressed by a subpopulation of neurons analogous to *BC1 RNA*, a 152-nt noncoding RNA localized in the dendrite-rich neuropil areas [147]. *BC200 RNA* and *BC1* are likely regulators of local protein expression in the post-synaptic dendritic microdomains [148]. A study has shown that *BC200 RNA* is upregulated in the frontal cortex of the AD brains compared to the age-matched controls. The expression level of *BC200 RNA* is increased in parallel with the AD progression [149]. However, in another independent work in a rigorously controlled study of short post-

mortem AD brains, *BC200 RNA* is found to be significantly decreased [150].

Human accelerated region 1 (HAR1) belongs to HAR family, which contains a set of 49 segments of the human genome that are conserved through vertebrate evolution but strikingly different in humans [151]. *HAR1* is located on the long arm of the chromosome 20 and overlaps with lncRNA *HAR1F*, which is expressed in Cajal–Retzius cells in the human fetal cortex [151]. The expression of *HAR1* is inhibited by transcription repressor REST in brain samples of HD patients, suggesting a potential role of HAR1 in neurodegeneration disorders [152].

Perspectives

It has been well accepted that noncoding RNAs, in particular miRNAs, play a critical role in neural development and functions. Identification of region and cell type-specific miRNAs in the CNS will help determine the actions of individual miRNAs in CNS development. Moreover, an investigation of how specific expression of miRNAs is regulated during development by identifying miRNA promoters and binding transcription factors will reveal the transcriptional logic of miRNA expression and functions. Because one miRNA has multiple targets, it remains a challenge to dissect the networks of miRNA targets in CNS development.

The association of miRNAs and human neurological diseases still requires comprehensive investigation. Because of the fast development of miRNA synthesis and delivery techniques, it is promising to use miRNAs as diagnostic biomarkers and as new therapeutic tools for neurological disorders. Moreover, to better understand the etiology of miRNA-mediated human neurological diseases, one needs to develop proper genetic tools to generate animal models with altered miRNA expressions.

There is no doubt that examining lncRNAs functions in neural development and under disease conditions will remain an exciting research focus. Genomewide RNA sequencing and bioinformatic analysis will help identify region and cell type-specific lncRNAs in the CNS and their association with protein-coding genes in the genome. Examining how lncRNAs manipulate gene expression will continue to reveal novel gene regulation mechanisms in neural development.

Acknowledgments We thank Dr. Dolores Malaspina, Dr. Rajiv Ratan, and Dr. Quanhong Ma for critical reading of the manuscript and providing thoughtful comments. Owing to space limitations, we apologize for being unable to cite many excellent papers in this field. This work was supported by Ellison Medical Foundation (T. S.), an award from Hirsch/Weill-Caulier Trust (T. S.), and an R01-MH083680 grant from NIH/NIMH (T. S.).

References

- Kapsimali M, Kloosterman WP, de Bruijn E, Rosa F, Plasterk RH, Wilson SW (2007) MicroRNAs show a wide diversity of expression profiles in the developing and mature central nervous system. *Genome Biol* 8(8):R173
- Bak M, Silahtaroglu A, Moller M, Christensen M, Rath MF, Skryabin B, Tommerup N, Kauppinen S (2008) MicroRNA expression in the adult mouse central nervous system. *RNA* 14(3):432–444
- Landgraf P, Rusu M, Sheridan R, Sewer A, Iovino N, Aravin A, Pfeffer S, Rice A, Kamphorst AO, Landthaler M et al (2007) A mammalian microRNA expression atlas based on small RNA library sequencing. *Cell* 129(7):1401–1414
- Mercer TR, Dingler ME, Sunkin SM, Mehler MF, Mattick JS (2008) Specific expression of long noncoding RNAs in the mouse brain. *Proc Natl Acad Sci U S A* 105(2):716–721
- Belgard TG, Marques AC, Oliver PL, Abaan HO, Sirey TM, Hoerder-Suabedissen A, Garcia-Moreno F, Molnar Z, Margulies EH, Ponting CP (2011) A transcriptomic atlas of mouse neocortical layers. *Neuron* 71(4):605–616
- Khraiwesh B, Arif MA, Seumel GI, Ossowski S, Weigel D, Reski R, Frank W (2010) Transcriptional control of gene expression by microRNAs. *Cell* 140(1):111–122
- Borchert GM, Lanier W, Davidson BL (2006) RNA polymerase III transcribes human microRNAs. *Nat Struct Mol Biol* 13(12):1097–1101
- Gregory RI, Chendrimada TP, Shiekhattar R (2006) MicroRNA biogenesis: isolation and characterization of the microprocessor complex. *Methods Mol Biol* 342:33–47
- Lund E, Dahlberg JE (2006) Substrate selectivity of exportin 5 and Dicer in the biogenesis of microRNAs. *Cold Spring Harb Symp Quant Biol* 71:59–66
- Rana TM (2007) Illuminating the silence: understanding the structure and function of small RNAs. *Nat Rev Mol Cell Biol* 8(1):23–36
- Neilson JR, Sharp PA (2008) Small RNA regulators of gene expression. *Cell* 134(6):899–902
- Wang XJ, Reyes JL, Chua NH, Gaasterland T (2004) Prediction and identification of *Arabidopsis thaliana* microRNAs and their mRNA targets. *Genome Biol* 5(9):R65
- Pratt AJ, MacRae IJ (2009) The RNA-induced silencing complex: a versatile gene-silencing machine. *J Biol Chem* 284(27):17897–17901
- Gao FB (2008) Posttranscriptional control of neuronal development by microRNA networks. *Trends Neurosci* 31(1):20–26
- Lau NC, Lim LP, Weinstein EG, Bartel DP (2001) An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science* 294(5543):858–862
- Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T (2001) Identification of novel genes coding for small expressed RNAs. *Science* 294(5543):853–858
- Lee RC, Ambros V (2001) An extensive class of small RNAs in *Caenorhabditis elegans*. *Science* 294(5543):862–864
- Altuvia Y, Landgraf P, Lithwick G, Elefant N, Pfeffer S, Aravin A, Brownstein MJ, Tuschl T, Margalit H (2005) Clustering and conservation patterns of human microRNAs. *Nucleic Acids Res* 33(8):2697–2706
- Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K, Doyle M, FitzHugh W et al (2001) Initial sequencing and analysis of the human genome. *Nature* 409(6822):860–921
- Kapranov P, Cheng J, Dike S, Nix DA, Duttagupta R, Willingham AT, Stadler PF, Hertel J, Hackermuller J, Hofacker IL et al (2007) RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science* 316(5830):1484–1488
- Carninci P, Kasukawa T, Katayama S, Gough J, Frith MC, Maeda N, Oyama R, Ravasi T, Lenhard B, Wells C et al (2005) The transcriptional landscape of the mammalian genome. *Science* 309(5740):1559–1563
- Birney E, Stamatoyannopoulos JA, Dutta A, Guigo R, Gingeras TR, Margulies EH, Weng Z, Snyder N, Dermitzakis ET, Thurman RE et al (2007) Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 447(7146):799–816
- Pasman E, Laurendeau I, Heron D, Vidaud M, Vidaud D, Bieche I (2007) Characterization of a germ-line deletion, including the entire INK4/ARF locus, in a melanoma–neural system tumor family: identification of ANRIL, an antisense noncoding RNA whose expression coclusters with ARF. *Cancer Res* 67(8):3963–3969
- Kotake Y, Nakagawa T, Kitagawa K, Suzuki S, Liu N, Kitagawa M, Xiong Y (2010) Long non-coding RNA ANRIL is required for the PRC2 recruitment to and silencing of p15(INK4B) tumor suppressor gene. *Oncogene* 30(16):1956–1962
- Tochitani S, Hayashizaki Y (2008) Nkx2.2 antisense RNA over-expression enhanced oligodendrocytic differentiation. *Biochem Biophys Res Commun* 372(4):691–696
- Feng J, Bi C, Clark BS, Mady R, Shah P, Kohtz JD (2006) The Evf-2 noncoding RNA is transcribed from the Dlx-5/6 ultra-conserved region and functions as a Dlx-2 transcriptional coactivator. *Genes Dev* 20(11):1470–1484
- Yu W, Gius D, Onyango P, Muldoon-Jacobs K, Karp J, Feinberg AP, Cui H (2008) Epigenetic silencing of tumour suppressor gene p15 by its antisense RNA. *Nature* 451(7175):202–206
- Wang X, Arai S, Song X, Reichart D, Du K, Pascual G, Tempst P, Rosenfeld MG, Glass CK, Kurokawa R (2008) Induced ncRNAs allosterically modify RNA-binding proteins in *cis* to inhibit transcription. *Nature* 454(7200):126–130
- Shamovsky I, Ivannikov M, Kandel ES, Gershon D, Nudler E (2006) RNA-mediated response to heat shock in mammalian cells. *Nature* 440(7083):556–560
- Willingham AT, Orth AP, Batalov S, Peters EC, Wen BG, Aza-Blanc P, Hogenesch JB, Schultz PG (2005) A strategy for probing the function of noncoding RNAs finds a repressor of NFAT. *Science* 309(5740):1570–1573
- Yang Z, Zhu Q, Luo K, Zhou Q (2001) The 7SK small nuclear RNA inhibits the CDK9/cyclin T1 kinase to control transcription. *Nature* 414(6861):317–322
- Nguyen VT, Kiss T, Michels AA, Bensaude O (2001) 7SK small nuclear RNA binds to and inhibits the activity of CDK9/cyclin T complexes. *Nature* 414(6861):322–325
- Sanchez-Elsner T, Gou D, Kremmer E, Sauer F (2006) Noncoding RNAs of trithorax response elements recruit *Drosophila* Ash1 to Ultrabithorax. *Science* 311(5764):1118–1123
- Mazo A, Hodgson JW, Petruk S, Sedkov Y, Brock HW (2007) Transcriptional interference: an unexpected layer of complexity in gene regulation. *J Cell Sci* 120(Pt 16):2755–2761
- Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Bruggmann SA, Goodnough LH, Helms JA, Farnham PJ, Segal E et al (2007) Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* 129(7):1311–1323
- Denisenko O, Shnyreva M, Suzuki H, Bomsztyk K (1998) Point mutations in the WD40 domain of Eed block its interaction with Ezh2. *Mol Cell Biol* 18(10):5634–5642
- Wutz A, Rasmussen TP, Jaenisch R (2002) Chromosomal silencing and localization are mediated by different domains of Xist RNA. *Nat Genet* 30(2):167–174

38. Mancini-Dinardo D, Steele SJ, Levorso JM, Ingram RS, Tilghman SM (2006) Elongation of the *Kcnqlot1* transcript is required for genomic imprinting of neighboring genes. *Genes Dev* 20 (10):1268–1282
39. Tsai MC, Manor O, Wan Y, Mosammaparast N, Wang JK, Lan F, Shi Y, Segal E, Chang HY (2010) Long noncoding RNA as modular scaffold of histone modification complexes. *Science* 329(5992):689–693
40. De Pietri Tonelli D, Pulvers JN, Haffner C, Murchison EP, Hannon GJ, Huttner WB: miRNAs are essential for survival and differentiation of newborn neurons but not for expansion of neural progenitors during early neurogenesis in the mouse embryonic neocortex. *Development* 2008, 135(23):3911–3921
41. Kawase-Koga Y, Low R, Otaegi G, Pollock A, Deng H, Eisenhaber F, Maurer-Stroh S, Sun T (2010) RNAase-III enzyme Dicer maintains signaling pathways for differentiation and survival in mouse cortical neural stem cells. *J Cell Sci* 123(Pt 4):586–594
42. Andersson T, Rahman S, Sansom SN, Alsio JM, Kaneda M, Smith J, O'Carroll D, Tarakhovsky A, Livesey FJ (2010) Reversible block of mouse neural stem cell differentiation in the absence of dicer and microRNAs. *PLoS One* 5(10):e13453
43. Kawase-Koga Y, Otaegi G, Sun T (2009) Different timings of Dicer deletion affect neurogenesis and gliogenesis in the developing mouse central nervous system. *Dev Dyn* 238 (11):2800–2812
44. Davis TH, Cuellar TL, Koch SM, Barker AJ, Harfe BD, McManus MT, Ullian EM (2008) Conditional loss of Dicer disrupts cellular and tissue morphogenesis in the cortex and hippocampus. *J Neurosci* 28(17):4322–4330
45. Li Q, Bian S, Hong J, Kawase-Koga Y, Zhu E, Zheng Y, Yang L, Sun T: Timing specific requirement of microRNA function is essential for embryonic and postnatal hippocampal development. *PLoS One* 2011, in press
46. Cuellar TL, Davis TH, Nelson PT, Loeb GB, Harfe BD, Ullian E, McManus MT (2008) Dicer loss in striatal neurons produces behavioral and neuroanatomical phenotypes in the absence of neurodegeneration. *Proc Natl Acad Sci U S A* 105(14):5614–5619
47. Zehir A, Hua LL, Maska EL, Morikawa Y, Cserjesi P (2010) Dicer is required for survival of differentiating neural crest cells. *Dev Biol* 340(2):459–467
48. Huang T, Liu Y, Huang M, Zhao X, Cheng L (2010) Wnt1-cre-mediated conditional loss of Dicer results in malformation of the midbrain and cerebellum and failure of neural crest and dopaminergic differentiation in mice. *J Mol Cell Biol* 2(3):152–163
49. Zheng K, Li H, Zhu Y, Zhu Q, Qiu M (2010) MicroRNAs are essential for the developmental switch from neurogenesis to gliogenesis in the developing spinal cord. *J Neurosci* 30(24):8245–8250
50. Zhao X, He X, Han X, Yu Y, Ye F, Chen Y, Hoang T, Xu X, Mi QS, Xin M et al (2010) MicroRNA-mediated control of oligodendrocyte differentiation. *Neuron* 65(5):612–626
51. Budde H, Schmitt S, Fitzner D, Opitz L, Salinas-Riester G, Simons M (2010) Control of oligodendroglial cell number by the miR-17-92 cluster. *Development* 137(13):2127–2132
52. Fukagawa T, Nogami M, Yoshikawa M, Ikeno M, Okazaki T, Takami Y, Nakayama T, Oshimura M (2004) Dicer is essential for formation of the heterochromatin structure in vertebrate cells. *Nat Cell Biol* 6(8):784–791
53. Kanellopoulou C, Muljo SA, Kung AL, Ganesan S, Drapkin R, Jenuwein T, Livingston DM, Rajewsky K (2005) Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes Dev* 19(4):489–501
54. Temple S (2001) The development of neural stem cells. *Nature* 414(6859):112–117
55. Gage FH (2000) Mammalian neural stem cells. *Science* 287 (5457):1433–1438
56. Doe CQ (2008) Neural stem cells: balancing self-renewal with differentiation. *Development* 135(9):1575–1587
57. Arsenijevic Y (2003) Mammalian neural stem-cell renewal: nature versus nurture. *Mol Neurobiol* 27(1):73–98
58. Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR, Ruvkun G (2000) The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 403(6772):901–906
59. Zhao C, Sun G, Li S, Lang MF, Yang S, Li W, Shi Y (2010) MicroRNA let-7b regulates neural stem cell proliferation and differentiation by targeting nuclear receptor TLX signaling. *Proc Natl Acad Sci U S A* 107(5):1876–1881
60. Rybak A, Fuchs H, Smirnova L, Brandt C, Pohl EE, Nitsch R, Wulczyn FG (2008) A feedback loop comprising lin-28 and let-7 controls pre-let-7 maturation during neural stem-cell commitment. *Nat Cell Biol* 10(8):987–993
61. Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, Tuschl T (2002) Identification of tissue-specific microRNAs from mouse. *Curr Biol* 12(9):735–739
62. Maiorano NA, Mallamaci A (2009) Promotion of embryonic cortico-cerebral neuronogenesis by miR-124. *Neural Dev* 4:40
63. Cheng LC, Pastrana E, Tavazoie M, Doetsch F (2009) miR-124 regulates adult neurogenesis in the subventricular zone stem cell niche. *Nat Neurosci* 12(4):399–408
64. Arvanitis DN, Jungas T, Behar A, Davy A (2010) Ephrin-B1 reverse signaling controls a posttranscriptional feedback mechanism via miR-124. *Mol Cell Biol* 30(10):2508–2517
65. Makeyev EV, Zhang J, Carrasco MA, Maniatis T (2007) The MicroRNA miR-124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Mol Cell* 27(3):435–448
66. Zhao C, Sun G, Li S, Shi Y (2009) A feedback regulatory loop involving microRNA-9 and nuclear receptor TLX in neural stem cell fate determination. *Nat Struct Mol Biol* 16(4):365–371
67. Delaloy C, Liu L, Lee JA, Su H, Shen F, Yang GY, Young WL, Ivey KN, Gao FB (2010) MicroRNA-9 coordinates proliferation and migration of human embryonic stem cell-derived neural progenitors. *Cell Stem Cell* 6(4):323–335
68. Bonev B, Pisco A, Papalopulu N (2011) MicroRNA-9 reveals regional diversity of neural progenitors along the anterior-posterior axis. *Dev Cell* 20(1):19–32
69. Leucht C, Stigloher C, Wizenmann A, Klafke R, Folchert A, Bally-Cuif L (2008) MicroRNA-9 directs late organizer activity of the midbrain–hindbrain boundary. *Nat Neurosci* 11(6):641–648
70. Otaegi G, Pollock A, Hong J, Sun T (2011) MicroRNA miR-9 modifies motor neuron columns by a tuning regulation of FoxP1 levels in developing spinal cords. *J Neurosci* 31(3):809–818
71. Shibata M, Nakao H, Kiyonari H, Abe T, Aizawa S (2011) MicroRNA-9 regulates neurogenesis in mouse telencephalon by targeting multiple transcription factors. *J Neurosci* 31(9):3407–3422
72. Gaughwin P, Ciesla M, Yang H, Lim B, Brundin P: Stage-specific modulation of cortical neuronal development by Mmu-miR-134. *Cereb Cortex* 2011
73. Brett JO, Renault VM, Rafalski VA, Webb AE, Brunet A (2011) The microRNA cluster miR-106b/25 regulates adult neural stem/progenitor cell proliferation and neuronal differentiation. *Aging (Albany NY)* 3(2):108–124
74. Renault VM, Rafalski VA, Morgan AA, Salih DA, Brett JO, Webb AE, Villeda SA, Thekkat PU, Guillerey C, Denko NC et al

- (2009) FoxO3 regulates neural stem cell homeostasis. *Cell Stem Cell* 5(5):527–539
75. Szulwach KE, Li X, Smrt RD, Li Y, Luo Y, Lin L, Santistevan NJ, Li W, Zhao X, Jin P (2010) Cross talk between microRNA and epigenetic regulation in adult neurogenesis. *J Cell Biol* 189(1):127–141
 76. Liu C, Teng ZQ, Santistevan NJ, Szulwach KE, Guo W, Jin P, Zhao X (2010) Epigenetic regulation of miR-184 by MBD1 governs neural stem cell proliferation and differentiation. *Cell Stem Cell* 6(5):433–444
 77. Yu JY, Chung KH, Deo M, Thompson RC, Turner DL (2008) MicroRNA miR-124 regulates neurite outgrowth during neuronal differentiation. *Exp Cell Res* 314(14):2618–2633
 78. Hall A (1998) Rho GTPases and the actin cytoskeleton. *Science* 279(5350):509–514
 79. Rajasethupathy P, Fiumara F, Sheridan R, Betel D, Puthanveettil SV, Russo JJ, Sander C, Tuschl T, Kandel E (2009) Characterization of small RNAs in *Aplysia* reveals a role for miR-124 in constraining synaptic plasticity through CREB. *Neuron* 63(6):803–817
 80. Sanuki R, Onishi A, Koike C, Muramatsu R, Watanabe S, Muranishi Y, Irie S, Uneo S, Koyasu T, Matsui R et al: miR-124a is required for hippocampal axogenesis and retinal cone survival through Lhx2 suppression. *Nat Neurosci*, 14(9):1125–1134
 81. Smrt RD, Szulwach KE, Pfeiffer RL, Li X, Guo W, Pathania M, Teng ZQ, Luo Y, Peng J, Bordey A et al (2010) MicroRNA miR-137 regulates neuronal maturation by targeting ubiquitin ligase mind bomb-1. *Stem Cells* 28(6):1060–1070
 82. Kim J, Inoue K, Ishii J, Vanti WB, Voronov SV, Murchison E, Hannon G, Abeliovich A (2007) A MicroRNA feedback circuit in midbrain dopamine neurons. *Science* 317(5842):1220–1224
 83. Nunes I, Tomvmasian LT, Silva RM, Burke RE, Goff SP (2003) Pitx3 is required for development of substantia nigra dopaminergic neurons. *Proc Natl Acad Sci U S A* 100(7):4245–4250
 84. Magill ST, Cambronne XA, Luikart BW, Lioy DT, Leighton BH, Westbrook GL, Mandel G, Goodman RH (2010) microRNA-132 regulates dendritic growth and arborization of newborn neurons in the adult hippocampus. *Proc Natl Acad Sci U S A* 107(47):20382–20387
 85. Hansen KF, Sakamoto K, Wayman GA, Impey S, Obrietan K (2010) Transgenic miR132 alters neuronal spine density and impairs novel object recognition memory. *PLoS One* 5(11):e15497
 86. Edbauer D, Neilson JR, Foster KA, Wang CF, Seeburg DP, Batterton MN, Tada T, Dolan BM, Sharp PA, Sheng M (2010) Regulation of synaptic structure and function by FMRP-associated microRNAs miR-125b and miR-132. *Neuron* 65(3):373–384
 87. Schrott GM, Tuebing F, Nigh EA, Kane CG, Sabatini ME, Kiebler M, Greenberg ME (2006) A brain-specific microRNA regulates dendritic spine development. *Nature* 439(7074):283–289
 88. Bai F, Witzmann FA (2007) Synaptosome proteomics. *Subcell Biochem* 43:77–98
 89. Siegel G, Obernosterer G, Fiore R, Oehmen M, Bicker S, Christensen M, Khudayberdiyev S, Leuschner PF, Busch CJ, Kane C et al (2009) A functional screen implicates microRNA-138-dependent regulation of the depalmitoylation enzyme APT1 in dendritic spine morphogenesis. *Nat Cell Biol* 11(6):705–716
 90. Banerjee S, Neveu P, Kosik KS (2009) A coordinated local translational control point at the synapse involving relief from silencing and MOV10 degradation. *Neuron* 64(6):871–884
 91. Abelson JF, Kwan KY, O'Roak BJ, Back DY, Stillman AA, Morgan TM, Mathews CA, Pauls DL, Rasin MR, Gunel M et al (2005) Sequence variants in SLITRK1 are associated with Tourette's syndrome. *Science* 310(5746):317–320
 92. Jin P, Zarnescu DC, Ceman S, Nakamoto M, Mowrey J, Jongens TA, Nelson DL, Moses K, Warren ST (2004) Biochemical and genetic interaction between the fragile X mental retardation protein and the microRNA pathway. *Nat Neurosci* 7(2):113–117
 93. Yi YH, Sun XS, Qin JM, Zhao QH, Liao WP, Long YS (2010) Experimental identification of microRNA targets on the 3' untranslated region of human FMR1 gene. *J Neurosci Methods* 190(1):34–38
 94. Han J, Lee Y, Yeom KH, Nam JW, Heo I, Rhee JK, Sohn SY, Cho Y, Zhang BT, Kim VN (2006) Molecular basis for the recognition of primary microRNAs by the Drosha-DGCR8 complex. *Cell* 125(5):887–901
 95. Digilio MC, Marino B, Cappa M, Cambiaso P, Giannotti A, Dallapiccola B (2001) Auxological evaluation in patients with DiGeorge/velocardiofacial syndrome (deletion 22q11.2 syndrome). *Genet Med* 3(1):30–33
 96. Stark KL, Xu B, Bagchi A, Lai WS, Liu H, Hsu R, Wan X, Pavlidis P, Mills AA, Karayiorgou M et al: Altered brain microRNA biogenesis contributes to phenotypic deficits in a 22q11-deletion mouse model. *Nat Genet* 2008
 97. Perkins DO, Jeffries CD, Jarskog LF, Thomson JM, Woods K, Newman MA, Parker JS, Jin J, Hammond SM (2007) microRNA expression in the prefrontal cortex of individuals with schizophrenia and schizoaffective disorder. *Genome Biol* 8(2):R27
 98. Beveridge NJ, Tooney PA, Carroll AP, Gardiner E, Bowden N, Scott RJ, Tran N, Dedova I, Cairns MJ (2008) Dysregulation of miRNA 181b in the temporal cortex in schizophrenia. *Hum Mol Genet* 17(8):1156–1168
 99. Kocerha J, Faghihi MA, Lopez-Toledano MA, Huang J, Ramsey AJ, Caron MG, Sales N, Willoughby D, Elmen J, Hansen HF et al (2009) MicroRNA-219 modulates NMDA receptor-mediated neurobehavioral dysfunction. *Proc Natl Acad Sci U S A* 106(9):3507–3512
 100. Abu-Elneel K, Liu T, Gazzaniga FS, Nishimura Y, Wall DP, Geschwind DH, Lao K, Kosik KS (2008) Heterogeneous dysregulation of microRNAs across the autism spectrum. *Neurogenetics* 9(3):153–161
 101. Sarachana T, Zhou R, Chen G, Manji HK, Hu VW (2010) Investigation of post-transcriptional gene regulatory networks associated with autism spectrum disorders by microRNA expression profiling of lymphoblastoid cell lines. *Genome Med* 2(4):23
 102. Seno MM, Hu P, Gwadry FG, Pinto D, Marshall CR, Casallo G, Scherer SW (2011) Gene and miRNA expression profiles in autism spectrum disorders. *Brain Res* 1380:85–97
 103. Lukiw WJ (2007) Micro-RNA speciation in fetal, adult and Alzheimer's disease hippocampus. *Neuroreport* 18(3):297–300
 104. Sethi P, Lukiw WJ (2009) Micro-RNA abundance and stability in human brain: specific alterations in Alzheimer's disease temporal lobe neocortex. *Neurosci Lett* 459(2):100–104
 105. Wang WX, Rajeev BW, Stromberg AJ, Ren N, Tang G, Huang Q, Rigoutsos I, Nelson PT (2008) The expression of microRNA miR-107 decreases early in Alzheimer's disease and may accelerate disease progression through regulation of beta-site amyloid precursor protein-cleaving enzyme 1. *J Neurosci* 28(5):1213–1223
 106. Hebert SS, Horre K, Nicolai L, Papadopoulou AS, Mandemakers W, Silahtaroglu AN, Kauppinen S, Delacourte A, De Strooper B (2008) Loss of microRNA cluster miR-29a/b-1 in sporadic Alzheimer's disease correlates with increased BACE1/beta-secretase expression. *Proc Natl Acad Sci U S A* 105(17):6415–6420
 107. Hebert SS, Papadopoulou AS, Smith P, Galas MC, Planel E, Silahtaroglu AN, Sergeant N, Buee L, De Strooper B (2010) Genetic ablation of Dicer in adult forebrain neurons results in

- abnormal tau hyperphosphorylation and neurodegeneration. *Hum Mol Genet* 19(20):3959–3969
108. Boissonneault V, Plante I, Rivest S, Provost P (2009) MicroRNA-298 and microRNA-328 regulate expression of mouse beta-amyloid precursor protein-converting enzyme 1. *J Biol Chem* 284(4):1971–1981
 109. Vilardo E, Barbato C, Ciotti M, Cogoni C, Ruberti F (2010) MicroRNA-101 regulates amyloid precursor protein expression in hippocampal neurons. *J Biol Chem* 285(24):18344–18351
 110. Lukiw WJ, Zhao Y, Cui JG (2008) An NF-kappaB-sensitive micro RNA-146a-mediated inflammatory circuit in Alzheimer disease and in stressed human brain cells. *J Biol Chem* 283(46):31315–31322
 111. Lee ST, Chu K, Im WS, Yoon HJ, Im JY, Park JE, Park KH, Jung KH, Lee SK, Kim M et al (2011) Altered microRNA regulation in Huntington's disease models. *Exp Neurol* 227(1):172–179
 112. Johnson R, Buckley NJ (2009) Gene dysregulation in Huntington's disease: REST, microRNAs and beyond. *Neuromolecular Med* 11(3):183–199
 113. Conaco C, Otto S, Han JJ, Mandel G (2006) Reciprocal actions of REST and a microRNA promote neuronal identity. *Proc Natl Acad Sci USA* 103(7):2422–2427
 114. Packer AN, Xing Y, Harper SQ, Jones L, Davidson BL (2008) The bifunctional microRNA miR-9/miR-9* regulates REST and CoREST and is downregulated in Huntington's disease. *J Neurosci* 28(53):14341–14346
 115. Gaughwin PM, Ciesla M, Lahiri N, Tabrizi SJ, Brundin P, Bjorkqvist M (2011) Hsa-miR-34b is a plasma-stable microRNA that is elevated in pre-manifest Huntington's disease. *Hum Mol Genet* 20(11):2225–2237
 116. Doxakis E (2010) Post-transcriptional regulation of alpha-synuclein expression by mir-7 and mir-153. *J Biol Chem* 285(17):12726–12734
 117. Junn E, Lee KW, Jeong BS, Chan TW, Im JY, Mouradian MM (2009) Repression of alpha-synuclein expression and toxicity by microRNA-7. *Proc Natl Acad Sci U S A* 106(31):13052–13057
 118. Wang G, van der Walt JM, Mayhew G, Li YJ, Zuchner S, Scott WK, Martin ER, Vance JM (2008) Variation in the miRNA-433 binding site of FGF20 confers risk for Parkinson disease by overexpression of alpha-synuclein. *Am J Hum Genet* 82(2):283–289
 119. Saba R, Goodman CD, Huzarewich RL, Robertson C, Booth SA (2008) A miRNA signature of prion induced neurodegeneration. *PLoS One* 3(11):e3652
 120. Pogue AI, Dua P, Eicken C, Hill JM, Lukiw WJ: Up-regulation of microRNA-146a (miRNA-146a), a marker for inflammatory neurodegeneration, in sporadic Creutzfeldt–Jakob disease (sCJD) and Gerstmann–Straussler Scheinker (GSS) syndrome. *Journal of Toxicology and Environmental Health* 2011, in press
 121. Roizen NJ, Patterson D (2003) Down's syndrome. *Lancet* 361(9365):1281–1289
 122. Kuhn DE, Nuovo GJ, Martin MM, Malana GE, Pleister AP, Jiang J, Schmittgen TD, Terry AV Jr, Gardiner K, Head E et al (2008) Human chromosome 21-derived miRNAs are overexpressed in down syndrome brains and hearts. *Biochem Biophys Res Commun* 370(3):473–477
 123. Elton TS, Sansom SE, Martin MM (2010) Trisomy-21 gene dosage over-expression of miRNAs results in the haploinsufficiency of specific target proteins. *RNA Biol* 7(5):540–547
 124. Kuhn DE, Nuovo GJ, Terry AV Jr, Martin MM, Malana GE, Sansom SE, Pleister AP, Beck WD, Head E, Feldman DS et al (2010) Chromosome 21-derived microRNAs provide an etiological basis for aberrant protein expression in human Down syndrome brains. *J Biol Chem* 285(2):1529–1543
 125. Pogue AI, Cui JG, Li YY, Zhao Y, Culicchia F, Lukiw WJ (2010) Micro RNA-125b (miRNA-125b) function in astrogliosis and glial cell proliferation. *Neurosci Lett* 476(1):18–22
 126. Tan KS, Arumugam A, Sepramaniam S, Lim KY, Setyowati KD, Wang CW, Jeyaseelan K (2009) Expression profile of MicroRNAs in young stroke patients. *PLoS One* 4(11):e7689
 127. Buller B, Liu X, Wang X, Zhang RL, Zhang L, Hozeska-Solgot A, Chopp M, Zhang ZG (2010) MicroRNA-21 protects neurons from ischemic death. *FEBS J* 277(20):4299–4307
 128. Liu XS, Chopp M, Zhang RL, Tao T, Wang XL, Kassis H, Hozeska-Solgot A, Zhang L, Chen C, Zhang ZG (2011) MicroRNA profiling in subventricular zone after stroke: MiR-124a regulates proliferation of neural progenitor cells through Notch signaling pathway. *PLoS One* 6(8):e23461
 129. Aronica E, Fluiter K, Iyer A, Zurolo E, Vreijling J, van Vliet EA, Baayen JC, Gorter JA (2010) Expression pattern of miR-146a, an inflammation-associated microRNA, in experimental and human temporal lobe epilepsy. *Eur J Neurosci* 31(6):1100–1107
 130. Saus E, Soria V, Escaramis G, Vivarelli F, Crespo JM, Kagerbauer B, Menchon JM, Urretavizcaya M, Gratacos M, Estivill X (2010) Genetic variants and abnormal processing of pre-miR-182, a circadian clock modulator, in major depression patients with late insomnia. *Hum Mol Genet* 19(20):4017–4025
 131. Xu Y, Liu H, Li F, Sun N, Ren Y, Liu Z, Cao X, Wang Y, Liu P, Zhang K (2010) A polymorphism in the microRNA-30e precursor associated with major depressive disorder risk and P300 waveform. *J Affect Disord* 127(1–3):332–336
 132. Kim AH, Reimers M, Maher B, Williamson V, McMichael O, McClay JL, van den Oord EJ, Riley BP, Kendler KS, Vladimirov VI (2010) MicroRNA expression profiling in the prefrontal cortex of individuals affected with schizophrenia and bipolar disorders. *Schizophr Res* 124(1–3):183–191
 133. Rong H, Liu TB, Yang KJ, Yang HC, Wu DH, Liao CP, Hong F, Yang HZ, Wan F, Ye XY et al (2011) MicroRNA-134 plasma levels before and after treatment for bipolar mania. *J Psychiatr Res* 45(1):92–95
 134. Baudry A, Mouillet-Richard S, Schneider B, Launay JM, Kellermann O (2010) miR-16 targets the serotonin transporter: a new facet for adaptive responses to antidepressants. *Science* 329(5998):1537–1541
 135. Zhou R, Yuan P, Wang Y, Hunsberger JG, Elkahouloun A, Wei Y, Damschroder-Williams P, Du J, Chen G, Manji HK (2009) Evidence for selective microRNAs and their effectors as common long-term targets for the actions of mood stabilizers. *Neuropsychopharmacology* 34(6):1395–1405
 136. Ponjavic J, Oliver PL, Lunter G, Ponting CP (2009) Genomic and transcriptional co-localization of protein-coding and long non-coding RNA pairs in the developing brain. *PLoS Genet* 5(8):e1000617
 137. Price M, Lazzaro D, Pohl T, Mattei MG, Ruther U, Olivo JC, Duboule D, Di Lauro R (1992) Regional expression of the homeobox gene Nkx-2.2 in the developing mammalian forebrain. *Neuron* 8(2):241–255
 138. Kohtz JD, Fishell G (2004) Developmental regulation of EVF-1, a novel non-coding RNA transcribed upstream of the mouse Dlx6 gene. *Gene Expr Patterns* 4(4):407–412
 139. Zerucha T, Stuhmer T, Hatch G, Park BK, Long Q, Yu G, Gambarotta A, Schultz JR, Rubenstein JL, Ekker M (2000) A highly conserved enhancer in the Dlx5/Dlx6 intergenic region is the site of cross-regulatory interactions between Dlx genes in the embryonic forebrain. *J Neurosci* 20(2):709–721
 140. Wang Y, Dye CA, Sohal V, Long JE, Estrada RC, Roztocil T, Lufkin T, Deisseroth K, Baraban SC, Rubenstein JL (2010) Dlx5 and Dlx6 regulate the development of parvalbumin-expressing cortical interneurons. *J Neurosci* 30(15):5334–5345

141. Bond AM, Vangompel MJ, Sametsky EA, Clark MF, Savage JC, Disterhoft JF, Kohtz JD (2009) Balanced gene regulation by an embryonic brain ncRNA is critical for adult hippocampal GABA circuitry. *Nat Neurosci* 12(8):1020–1027
142. Pevny L, Placzek M (2005) SOX genes and neural progenitor identity. *Curr Opin Neurobiol* 15(1):7–13
143. Amaral PP, Neyt C, Wilkins SJ, Askarian-Amiri ME, Sunkin SM, Perkins AC, Mattick JS (2009) Complex architecture and regulated expression of the *Sox2ot* locus during vertebrate development. *RNA* 15(11):2013–2027
144. Blackshaw S, Harpavat S, Trimarchi J, Cai L, Huang H, Kuo WP, Weber G, Lee K, Fraioli RE, Cho SH et al (2004) Genomic analysis of mouse retinal development. *PLoS Biol* 2(9):E247
145. Raponavoli NA, Poth EM, Blackshaw S (2010) The long noncoding RNA *RNCR2* directs mouse retinal cell specification. *BMC Dev Biol* 10:49
146. Arisi I, D'Onofrio M, Brandi R, Felsani A, Capsoni S, Drovandi G, Felici G, Weitschek E, Bertolazzi P, Cattaneo A (2011) Gene expression biomarkers in the brain of a mouse model for Alzheimer's disease: mining of microarray data by logic classification and feature selection. *J Alzheimers Dis* 24(4):721–738
147. Tiedge H, Chen W, Brosius J (1993) Primary structure, neural-specific expression, and dendritic location of human BC200 RNA. *J Neurosci* 13(6):2382–2390
148. Wang H, Iacoangeli A, Popp S, Muslimov IA, Imataka H, Sonenberg N, Lomakin IB, Tiedge H (2002) Dendritic BC1 RNA: functional role in regulation of translation initiation. *J Neurosci* 22(23):10232–10241
149. Mus E, Hof PR, Tiedge H (2007) Dendritic BC200 RNA in aging and in Alzheimer's disease. *Proc Natl Acad Sci U S A* 104(25):10679–10684
150. Lukiw WJ, Handley P, Wong L, Crapper McLachlan DR (1992) BC200 RNA in normal human neocortex, non-Alzheimer dementia (NAD), and senile dementia of the Alzheimer type (AD). *Neurochem Res* 17(6):591–597
151. Pollard KS, Salama SR, Lambert N, Lambot MA, Coppens S, Pedersen JS, Katzman S, King B, Onodera C, Siepel A et al (2006) An RNA gene expressed during cortical development evolved rapidly in humans. *Nature* 443(7108):167–172
152. Johnson R, Richter N, Jauch R, Gaughwin PM, Zuccato C, Cattaneo E, Stanton LW (2010) The Human Accelerated Region 1 noncoding RNA is repressed by REST in Huntington's disease. *Physiol Genomics* 41:269–274