

# Stimulation by Glutamate Receptors of Arachidonic Acid Release Depends on the $\text{Na}^+/\text{Ca}^{2+}$ Exchanger in Neuronal Cells

ALINE DUMUIS, MICHÈLE SEBBEN, LAURENT FAGNI, LAURENT PRÉZEAU, OLIVIER MANZONI, EDWARD J. CRAGOE, JR., and JOËL BOCKAERT

Centre CNRS-INSERM de Pharmacologie-Endocrinologie, 34094 Montpellier Cedex 5, France (A.D., M.S., L.F., L.P., O.M., J.B.), and Nacogdoches, Texas 75963-1548 (E.J.C.)

Received January 21, 1993; Accepted March 29, 1993

## SUMMARY

In primary cultures of striatal neurons, stimulation of *N*-methyl-D-aspartic acid (NMDA) receptors or associative activation (but not separate activation) of (RS)- $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors and metabotropic glutamate receptors (mGluR) strongly increased arachidonic acid (AA) release via activation of phospholipase  $\text{A}_2$  ( $\text{PLA}_2$ ). Depolarizing agents, such as veratridine, were as potent as NMDA in stimulating AA release. However, increasing the intracellular  $\text{Ca}^{2+}$  concentration via voltage-sensitive  $\text{Ca}^{2+}$  channels did not result

in a significant stimulation of  $\text{PLA}_2$ . Substitution of sodium by lithium, a monovalent cation that does not participate in the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger activity but permeates ionotropic glutamate receptor channels, blocked AA release induced by veratridine or AMPA plus mGluR agonists. It also reduced the NMDA-induced AA release, to a lesser extent. The contribution of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger to the activation of  $\text{PLA}_2$  after veratridine, NMDA receptor, or AMPA receptor plus mGluR stimulation was confirmed by using a selective inhibitor of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger.

Glu, like other neurotransmitters, activates channel receptors (ionotropic receptors) and receptors coupled to guanine nucleotide-binding proteins (metabotropic receptors) (1–4). Three families of ionotropic Glu receptors have already been described, the NMDA, AMPA, and KAI receptors. KAI and AMPA receptor channels are permeable to monovalent cations ( $\text{Na}^+$  and  $\text{K}^+$ ) and increase  $[\text{Ca}^{2+}]_i$  mainly via depolarization-induced opening of VSCC (1, 5). A minority of AMPA/KAI receptors are permeable to  $\text{Ca}^{2+}$  (6). In contrast to most KAI and AMPA receptors, NMDA receptor channels appear to be largely permeable to  $\text{Ca}^{2+}$  ions, in addition to  $\text{Na}^+$  and  $\text{K}^+$  ions (7). One class of metabotropic receptors activates phospholipase C and thus inositol trisphosphate formation, as well as intracellular  $\text{Ca}^{2+}$  release (3, 6, 8–12).

In addition to these primary ionic and enzymatic events, at least two other messengers have been associated with Glu receptor activation, namely AA and NO (13–15). A third, cGMP, is also produced as a consequence of guanylate cyclase activation by NO (15–20). AA and NO are interesting messen-

gers because they can easily diffuse in and out of the cell and therefore behave as intra- and intercellular messengers. Therefore, in addition to acting within the cells where they are produced, there is evidence that they can also act as retrograde messengers by transducing information from the postsynaptic to the presynaptic element (21–27). Such a function appears to be of crucial importance in long term potentiation, which is likely to represent an elementary event underlying learning and memory (28, 29).

We have demonstrated that AA is produced in striatal neurons by combined stimulation of AMPA receptors and either mGluR or NMDA receptors (13, 14). The requirement of a combined stimulation of AMPA and NMDA receptors for releasing AA can be easily explained. Indeed, NMDA receptor channels are blocked by  $\text{Mg}^{2+}$  ions at resting membrane potential and AMPA-induced depolarization is required to suppress the  $\text{Mg}^{2+}$  blockage (30, 31). It is believed that, once activated, NMDA receptor channels lead to sufficient  $\text{Ca}^{2+}$  entry to stimulate phospholipase  $\text{A}_2$  (7). However, we have no indication as to the molecular mechanisms underlying the associative stimulation of AMPA receptors and mGluR leading to AA release (14). In this report, we provide evidence for a crucial role of  $\text{Ca}^{2+}$  influx via the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger in this associative process.

This work was supported by grants from CNRS, INSERM, and Bayer France/Troponwerke (Germany), by Direction des Recherches, Etudes, et Techniques Contract 91/161, and by the International Human Frontier Science Program Organization.

**ABBREVIATIONS:** Glu, glutamate; NMDA, *N*-methyl-D-aspartic acid; AMPA,  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; KAI, kainic acid; QA, quisqualic acid; *t*-ACPD, (*trans*)-1-amino-cyclopentyl-1,3-dicarboxylic acid; AA, arachidonic acid; CBDMB, 5-(*N*-4-chlorobenzyl)-2',4'-dimethylbenzamil; MGCMA, 5-[*N*-methyl-*N*-(guanidinocarbonylmethyl)]amiloride; TTX, tetrodotoxin; VSCC, voltage-sensitive  $\text{Ca}^{2+}$  channels; PTX, *Bordetella pertussis* toxin; NO, nitric oxide; HEPES, *N*-[ $\alpha$ -hydroxyethyl]piperazine-*N'*-[2-ethanesulfonic acid]; EGTA, ethylene glycol bis( $\beta$ -aminoethyl ether)-*N,N,N',N'*-tetraacetic acid; mGluR, metabotropic glutamate receptor(s); PKC, protein kinase C;  $[\text{Ca}^{2+}]_i$ , intracellular calcium concentration;  $[\text{Na}^+]_i$ , intracellular sodium concentration.

## Materials and Methods

**Primary neuronal cultures.** Primary cultures of striatal neurons were prepared as described previously (32). Striata were removed from 14–15-day-old mouse embryos (Iffa Credo, Lyon, France) and cells were gently dissociated in culture medium and plated ( $1.5 \times 10^6$ /dish) in 35-mm-diameter Costar (Cambridge, MA) culture dishes (six-well clusters, 2 ml/well). The dishes had been previously coated with poly-L-ornithine (15  $\mu$ g/ml, *M*, 40,000) and culture medium containing 10% fetal calf serum. Culture was then performed in a defined medium (32). Under these conditions, the cultures have been shown to be highly enriched in neurons that form many mature and functional synapses (32). Cultures were maintained at 37° in a humidified atmosphere in 6% CO<sub>2</sub>/94% air.


**Measurement of [<sup>3</sup>H]AA release.** Striatal neurons, after 13 days *in vitro*, were incubated overnight with 0.25  $\mu$ Ci/ml [<sup>3</sup>H]AA (0.5  $\mu$ Ci/well; specific activity, 166 Ci/mmol; CEA, Saclay, France). As reported previously (13), >90% of the radioactivity was incorporated into the cells. Eighteen to 24 hr later, cells were washed five times (at 5-min intervals) with 2 ml/well of Krebs bicarbonate buffer (3.5 mM KCl, 0.75 mM CaCl<sub>2</sub>, 124 mM NaCl, 1.25 mM KH<sub>2</sub>PO<sub>4</sub>, 26.3 mM NaHCO<sub>3</sub>, 10 mM glucose, pH 7.4) containing 0.05% bovine serum albumin (fatty acid free); the buffer was saturated with 95% O<sub>2</sub>/5% CO<sub>2</sub>. Neurons were then exposed to the stimulating medium for 15 min. The radioactivity released in the medium (<sup>3</sup>H-med) and that remaining in the cells (<sup>3</sup>H-cell) were determined. The percentage of radioactivity released from the cells was calculated as  $100 \times [\text{<sup>3</sup>H-med}/(\text{<sup>3</sup>H-med} + \text{<sup>3</sup>H-cell})]$ . For the sodium substitution experiments, the Krebs bicarbonate buffer was replaced with modified Krebs buffer, in which sodium chloride could be totally substituted by equimolar concentrations of lithium chloride (3.5 mM KCl, 0.75 mM CaCl<sub>2</sub>, 124 mM NaCl or 124 mM LiCl, 1.25 mM KHPO<sub>4</sub>, 10 mM KHCO<sub>3</sub>, 10 mM glucose, pH 7.4); 0.05% bovine serum albumin (fatty acid free) was added and the buffer was saturated with 95% O<sub>2</sub>/5% CO<sub>2</sub>.

**Electrophysiological techniques.** Recordings were performed at room temperature from 2-week-old cultured neurons, using the whole-cell configuration of the patch-clamp technique. External solution contained 140 mM NaCl, 5 mM KCl, 2 mM CaCl<sub>2</sub>, 2 mM MgCl<sub>2</sub>, 0.3  $\mu$ M TTX, and 10 mM HEPES, pH 7.4. In experiments using NMDA, MgCl<sub>2</sub> was omitted and 1  $\mu$ M glycine was added to the solution. In all cases, internal medium contained 140 mM CsCl, 0.5 mM CaCl<sub>2</sub>, 5 mM EGTA, and 10 mM HEPES, pH 7.2. Recording pipettes filled with this internal medium had impedance ranging from 3 to 5 M $\Omega$ . Drug solutions were adjusted to pH 7.4 and were applied to individual cells using a rapid perfusion system.

**Materials.** The chemicals used in the present study were obtained from the following sources. NMDA, AMPA, and QA were purchased from Euromedex (Strasbourg, France). Veratridine and TTX were from Sigma (L'Isle d'Abeau, France). *t*-ACPD was from Tocris Neuramin (Essex, England). Staurosporine was from Calbiochem (Meudon, France). [<sup>3</sup>H]AA (specific radioactivity, 166 Ci/mmol) was from CEA. CBDMB and MGCMA were from Dr. E. J. Cragoe (Nacogdoches, TX).

## Results

**Evidence that mGluR stimulation must be associated with Na<sup>+</sup> entry to trigger AA release.** Separate stimulations of AMPA receptors and mGluR with their respective specific agonists, AMPA and *t*-ACPD, resulted in a weak stimulation of AA release (Fig. 1A). In contrast, combined stimulation of these receptors was very active in triggering AA release (Fig. 1A) (14). In a previous paper (33), we fully characterized the radioactivity released from neuronal cells in culture, after prelabeling with [<sup>3</sup>H]AA and after drug treatment leading to a large increase in [Ca<sup>2+</sup>]<sub>i</sub> (Glu and ionomycin). Only [<sup>3</sup>H]AA (and not cyclooxygenase or lipoxygenase derivatives) was detected (33). Because AMPA receptor stimulation is known to




**Fig. 1.** Sodium dependence of AMPA- plus *t*-ACPD- and veratridine-induced stimulation of AA release. **A**, Effect of 30  $\mu$ M AMPA and 300  $\mu$ M *t*-ACPD, alone or together, on [<sup>3</sup>H]AA release in striatal neurons after 13 days *in vitro*. **B**, Effect of neuronal depolarization induced by 56 mM K<sup>+</sup> alone or associated with 300  $\mu$ M *t*-ACPD on [<sup>3</sup>H]AA release. **C**, Effect of neuronal depolarization induced by 10  $\mu$ M veratridine alone or associated with 300  $\mu$ M *t*-ACPD in the presence or absence of 0.5  $\mu$ M TTX. Neurons were exposed to 0.5  $\mu$ M TTX 20 min before and during the 15-min stimulation period with either 10  $\mu$ M veratridine, 300  $\mu$ M *t*-ACPD, or 10  $\mu$ M veratridine plus 300  $\mu$ M *t*-ACPD. In **A**, **B**, and **C**, neurons were stimulated with the indicated concentrations of agonists and the percentage of radioactivity released from the cells was calculated as indicated in Materials and Methods. Results are expressed as percentage of stimulation over basal release (basal release,  $0.60 \pm 0.12\%$  of the total radioactivity in the cells). In histograms, each bar is the mean  $\pm$  standard error of four separate experiments, each performed in triplicate.

increase Na<sup>+</sup> conductance, depolarize neurons, and open VSCC leading to an increase of [Ca<sup>2+</sup>]<sub>i</sub>, we tried to determine which of these events triggered AA release when associated with mGluR stimulation. Obviously, depolarization was not a crucial event because K<sup>+</sup> (56 mM) alone or in association with *t*-ACPD application was ineffective in triggering AA release (Fig. 1B). In contrast, veratridine alone (a Na<sup>+</sup> channel opener) was a good stimulator of AA release and was even stronger when associated with mGluR stimulation (Fig. 1C). The effects of veratridine were suppressed by TTX, suggesting that they were due to Na<sup>+</sup> entry via voltage-sensitive Na<sup>+</sup> channels (Fig. 1C). Ouabain (100  $\mu$ M), an agent that increased [Na<sup>+</sup>]<sub>i</sub> and depolarized neurons, also induced AA release in combination with mGluR stimulation (120% of basal release) (data not shown).

**Requirement for extracellular Ca<sup>2+</sup> to trigger AA release.** As shown in Fig. 2, AA release induced by combined stimulation of AMPA receptors and mGluR was strictly dependent on external Ca<sup>2+</sup>. This was shown by applying either the specific agonists AMPA and *t*-ACPD together or QA, a mixed AMPA and mGluR agonist. The AA released by veratridine was also dependent on external Ca<sup>2+</sup> (Fig. 2).

**Inhibition of the veratridine-induced, AMPA- plus *t*-ACPD-induced, and part of the NMDA-induced AA release by blockade of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger.** Li<sup>+</sup> does not participate in the Na<sup>+</sup>/Ca<sup>2+</sup> exchange (34). However, Li<sup>+</sup> permeates sodium channels and ionotropic Glu receptor channels



**Fig. 2.**  $\text{Ca}^{2+}$  dependence of  $[^3\text{H}]\text{AA}$  release induced by either veratridine, AMPA plus  $t\text{-ACPD}$ , or QA.  $[^3\text{H}]\text{AA}$  release was induced by the indicated agents in Krebs bicarbonate buffer;  $\text{Ca}^{2+}$  (0.75 mM) was either present or omitted during the washing and 15-min stimulation periods. Results are expressed as percentage of radioactivity released from the cells. In the histogram, each vertical bar represents the mean  $\pm$  standard error of four independent experiments performed in triplicate.

(35). As shown in Fig. 3A, the veratridine-induced AA release was completely blocked when  $\text{Na}^+$  was replaced by  $\text{Li}^+$ . Note that under these conditions the basal production of AA was increased (Fig. 3A). CBDMB, an amiloride analogue that inhibits the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger with high potency and specificity ( $K_i$  about 1  $\mu\text{M}$ ) (36, 37), completely suppressed the veratridine-induced AA release, in a dose-dependent manner ( $\text{EC}_{50} = 562 \pm 53 \text{ nM}$ ) (Fig. 3D). Replacing  $\text{Na}^+$  with  $\text{Li}^+$ , as well as a concentration of CBDMB (30  $\mu\text{M}$ ) high enough to completely block the veratridine effect, also suppressed the AMPA plus  $t\text{-ACPD}$  stimulation of AA release (Fig. 3, B and D). The production of inositol monophosphate stimulated by  $t\text{-ACPD}$  was


similar in the presence of  $\text{Na}^+$  or  $\text{Li}^+$  (data not shown). These treatments also reduced the NMDA-induced AA release, but to a lesser extent (Fig. 3, B and D).

The specificity of the CBDMB effect was strengthened by the fact that another amiloride analogue, MGCMA, which is a poor  $\text{Na}^+/\text{Ca}^{2+}$  exchanger blocker but a good  $\text{Na}^+/\text{H}^+$  blocker (38, 39), was ineffective in blocking veratridine-, NMDA-, and AMPA- plus  $t\text{-ACPD}$ -induced AA release (Fig. 4A). Moreover, CBDMB (20  $\mu\text{M}$ ) did not significantly alter the current induced by either NMDA (50  $\mu\text{M}$ ) or AMPA (50  $\mu\text{M}$ ) (Fig. 4B). Indeed, variations induced by CBDMB in NMDA and AMPA currents were  $+5 \pm 3\%$  (eight experiments) and  $-7 \pm 5\%$  (four experiments), respectively (mean  $\pm$  standard deviation). These variation values are in the range of fluctuation observed under cortical conditions in a given cell.


**Absence of VSCC antagonist effect on veratridine-, NMDA-, and AMPA- plus  $t\text{-ACPD}$ -induced AA release.** Nifedipine (10  $\mu\text{M}$ ), an L-type VSCC channel blocker,  $\omega$ -conotoxin (10 nM), an N-type VSCC channel blocker, and cadmium (0.1 mM), a nonspecific VSCC channel blocker, had no significant effect on AA release induced by veratridine, NMDA, or AMPA plus  $t\text{-ACPD}$  (Table 1).

**Role of PKC.** Staurosporine (0.2  $\mu\text{M}$ ), a known inhibitor of PKC, did not inhibit AMPA- plus  $t\text{-ACPD}$ -, QA-, or NMDA-induced AA release. In contrast, staurosporine potentiated AMPA plus  $t\text{-ACPD}$  and QA responses by 72% and 130%, respectively, without affecting NMDA- or veratridine-induced AA release (Fig. 5).

**Absence of effect of PTX on the release of AA induced**



**Fig. 3.** Implication of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger in triggering  $[^3\text{H}]\text{AA}$  release. A, The percentage of radioactivity released from the cells was measured in the presence of NaCl and after substitution of NaCl with LiCl. B, NMDA- or AMPA- plus  $t\text{-ACPD}$ -induced AA release, expressed as percentage of radioactivity released from the cells, in the presence of NaCl and after substitution of NaCl with LiCl. C, Inhibition of veratridine-induced  $[^3\text{H}]\text{AA}$  release by increasing concentrations of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger inhibitor CBDMB. Neurons were preincubated for 20 min with increasing concentrations of CBDMB (0.01–60  $\mu\text{M}$ ) before and during the 15-min stimulation with 10  $\mu\text{M}$  veratridine. Results are expressed as percentage of the maximal veratridine-induced AA release obtained in the absence of CBDMB minus basal release. Each point is the mean  $\pm$  standard error of three separate experiments performed in duplicate. The maximum veratridine-induced  $[^3\text{H}]\text{AA}$  release was  $2.05 \pm 0.15\%$  and the basal release was  $0.55 \pm 0.06\%$  of the total radioactivity in the cells. D, Effect of CBDMB on 100  $\mu\text{M}$  NMDA- or 30  $\mu\text{M}$  AMPA- plus 300  $\mu\text{M}$   $t\text{-ACPD}$ -induced  $[^3\text{H}]\text{AA}$  release. Neurons were preincubated in the absence or presence of 30  $\mu\text{M}$  CBDMB for 20 min before stimulation and then during the 15-min stimulation period with either NMDA or AMPA plus  $t\text{-ACPD}$ . Results are expressed as percentage of radioactivity released from the cells. In histograms A, B, and D, each vertical bar represents the mean  $\pm$  standard error of four independent experiments, each performed in triplicate. \*\*, The  $t$  test analysis revealed a significant difference between basal AA release on one hand and AA release induced by replacing  $\text{Na}^+$  with  $\text{Li}^+$  or by adding CBDMB on the other hand, with  $p < 0.001$ .




**Fig. 4.** Specificity of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger inhibitor CBDMB. **A**, Absence of effect of the Na<sup>+</sup>/H<sup>+</sup> exchanger inhibitor MGCMA on [<sup>3</sup>H]AA release induced by either veratridine, NMDA, or AMPA plus *t*-ACPD. Neurons were preincubated with or without 30 μM MGCMA for 20 min before stimulation and then during the 15-min stimulation period. Results are expressed as percentage of radioactivity released from the cells. In the histogram, each vertical bar represents the mean + standard error of four independent experiments performed in triplicate. **B**, Absence of effect of CBDMB on NMDA or AMPA receptors. Membrane potential was held at -60 mV and brief applications (horizontal bars) of either NMDA or AMPA (50 μM for both agonists) were performed in the absence (a) or presence (b) of CBDMB (30 μM). The NMDA and AMPA currents were recorded from two different cells. Similar results were obtained in 10 other cells (seven with NMDA and three with AMPA).

**TABLE 1**  
**Absence of effect of VSCC antagonists on AA release induced by different agents**

Neurons were exposed to VSCC antagonists (nifedipine, ω-conotoxin, or cadmium) for 20 min before and during the 15-min stimulation period with either AMPA plus *t*-ACPD, NMDA, or veratridine. Results are expressed as percentage of stimulation over basal release. The basal release was measured in the absence of antagonists and was 0.58 ± 0.07% of the total radioactivity in the cells.

	Stimulation of AA release			
	Control	Nifedipine (10 μM)	ω-Conotoxin (10 nM)	Cd <sup>2+</sup> (100 μM)
	%			
AMPA (30 μM) + <i>t</i> -ACPD (300 μM)	212 ± 21	182 ± 16	170 ± 19	230 ± 28
NMDA (100 μM)	173 ± 18	184 ± 13	169 ± 10	152 ± 17
Veratridine (10 μM)	147 ± 15	121 ± 10	130 ± 11	166 ± 13



**Fig. 5.** Absence of PKC activation induced by the different agents evoking AA release. The effect of a PKC inhibitor, staurosporine, was tested on the [<sup>3</sup>H] AA release evoked by either 30 μM AMPA plus 300 μM *t*-ACPD, 10 μM QA, 100 μM NMDA, or 10 μM veratridine. Staurosporine (0.2 μM) was added to neurons during a 20-min preincubation period and then with the stimulating agents for 15 min. Results are expressed as percentage of radioactivity released from the cells. Each vertical bar of the histogram is the mean + standard error of four independent experiments performed in triplicate.

by NMDA, AMPA plus *t*-ACPD, or veratridine. Pretreatment of neuronal cultures for 24 hr with 0.1 μg/ml PTX affected neither 100 μM NMDA-, 30 μM AMPA- plus 300 μM *t*-ACPD-, nor 10 μM veratridine-induced AA release (data not shown).

## Discussion

One important conclusion of this study is that Ca<sup>2+</sup> entry through VSCC is unable to stimulate phospholipase A<sub>2</sub> in striatal neurons in primary cultures. This is clearly indicated by the following results: (a) high concentrations (up to 100 mM) of K<sup>+</sup>, which increased Ca<sup>2+</sup> levels in striatal neuronal cell bodies and main dendrites up to 600 nM (data not shown), were unable to increase AA release, (b) AA release induced by veratridine was completely blocked when Na<sup>+</sup> was replaced by Li<sup>+</sup>, a situation in which VSCC should be fully activated, and (c) Cd<sup>2+</sup>, nifedipine, and ω-conotoxin did not inhibit veratridine-induced AA release. These results indicate that VSCC and phospholipase A<sub>2</sub> are likely to be located in different neuronal compartments and that the compartment in which phospholipase A<sub>2</sub> is expressed should also contain NMDA receptors, as well as AMPA receptors and mGluR.

In contrast to the absence of a role for VSCC in mediating the stimulation of AA release, Ca<sup>2+</sup> entry via the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger was an important source of Ca<sup>2+</sup> to trigger this effect after veratridine or AMPA plus *t*-ACPD stimulation. It also represented an important component of the Ca<sup>2+</sup> required for NMDA receptor-induced stimulation of AA release. The Ca<sup>2+</sup> entering through the NMDA receptor channels certainly constituted another source of Ca<sup>2+</sup> for NMDA receptor-induced AA release. The role of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger was demonstrated by using Na<sup>+</sup>-Li<sup>+</sup> substitution, a condition under which the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger is not operative. We also used CBDMB, a blocker of the exchanger. The specificity of this blocker was demonstrated by the lack of effect of another amiloride analogue (MGCMA) that is a specific blocker of the Na<sup>+</sup>/H<sup>+</sup> exchanger and by the absence of a direct effect of CBDMB on NMDA and AMPA receptor channel activities. Blockade of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger by Li<sup>+</sup> or CBDMB, which is known to increase intracellular Ca<sup>2+</sup> under resting conditions (probably by suppressing the Ca<sup>2+</sup> extrusion) (37), also increases basal AA release. This further indicates the importance

of  $\text{Ca}^{2+}$  accumulation in the vicinity of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger for triggering AA release.

The mechanism by which veratridine, AMPA plus *t*-ACPD, and NMDA stimulated  $\text{Ca}^{2+}$  entry via the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger is likely to be the following. The  $\text{Na}^+/\text{Ca}^{2+}$  exchanger uses an electrochemical gradient for  $\text{Na}^+$  to "pump"  $\text{Ca}^{2+}$  out and  $\text{Na}^+$  into the cell, with a stoichiometry of 3  $\text{Na}^+$  to 1  $\text{Ca}^{2+}$  (the exchanger stoichiometry is  $n = 3$ ) (40). The depolarization and the  $\text{Na}^+$  entry induced by veratridine, AMPA plus *t*-ACPD, or NMDA applications reduced the  $\text{Na}^+$  electrochemical potential, and the reversal potential of the exchanger, given as  $E_{\text{NaCa}} = (nE_{\text{Na}} - 2E_{\text{Ca}})/(n - 2)$ , became lower than the membrane potential ( $V_m$ ). Under these conditions, the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger "pumped"  $\text{Ca}^{2+}$  into the cell (40, 41). The  $[\text{Ca}^{2+}]_i$  reached under these conditions is given by

$$[\text{Ca}^{2+}]_i = [\text{Ca}^{2+}]_o \exp \frac{FV_m(n-2)}{RT} \frac{([\text{Na}^+]_i)^n}{[\text{Na}^+]_o}$$

where  $R$  is the gas constant,  $T$  is the absolute temperature, and  $F$  is Faraday's constant (42).

Why doesn't the stimulation of AMPA receptors alone induce the reversal of the exchanger and why is the associative stimulation of mGluR required? One of the reasons could be that the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, as demonstrated in squid axons and heart cells, requires a minimal  $[\text{Ca}^{2+}]_i$  for activation, even when it is pumping  $\text{Ca}^{2+}$  out of the cell (43). This is certainly the consequence of the presence of a calmodulin-like sequence within the transporter (44). Therefore, by releasing  $\text{Ca}^{2+}$  from intracellular stores, mGluR probably raise  $[\text{Ca}^{2+}]_i$  to "help" the activation of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger and its reversal by  $\text{Na}^+$  entry triggered by AMPA receptor stimulation. Veratridine stimulation likely produced a situation similar to that of AMPA receptor stimulation, because mGluR potentiate veratridine-induced AA release. However, the reduction in the  $\text{Na}^+$  electrochemical gradient was likely to be higher after veratridine, rather than AMPA, application and therefore veratridine alone was sufficient to reverse the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. Following the same hypothesis, it is possible that NMDA receptors do not require mGluR stimulation to activate the  $\text{Ca}^{2+}$  influx via the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger because NMDA channels carry enough  $\text{Ca}^{2+}$  into the cell to raise  $\text{Ca}^{2+}$  to a level sufficient to occupy the exchanger intracellular  $\text{Ca}^{2+}$  sites.

Another hypothesis to explain the role of the mGluR could be the existence of a coupling between these receptors and the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. Such a coupling might exist in Purkinje cells (45), but its nature is still unknown. Our experiments exclude a role for PKC or for a guanine nucleotide-binding protein sensitive to PTX in this coupling. In view of the diversity of metabotropic receptors (6, 10–12, 46–48), several other coupling mechanism remain possible. The fact that staurosporine, a widely used PKC inhibitor, potentiated AMPA-plus mGluR-triggered AA release might reflect PKC-mediated inhibition of metabotropic receptors under physiological conditions. Indeed, we previously reported that mGluR stimulation induces PKC translocation and that PKC activation desensitizes mGluR receptors (49).

The compartment in which the reversal of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger takes place certainly has a limited volume and is possibly restricted to ionic diffusion, which can explain a rapid rise in  $[\text{Na}^+]_i$ . The existence of such restricted areas in small

axons has been recently invoked to explain the reversal of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger that is responsible for anoxic injury in mammalian white matter (42).

This report sheds new light on the importance of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger in the mechanism of action of Glu receptors. The role of this exchanger has certainly been previously underestimated, although one report indicated that in cerebellar slices the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger mediated a major part of the NMDA-induced cGMP accumulation and therefore the NMDA-induced NO synthesis.

AA has been shown to block Glu uptake. This may lead to an increase in Glu concentration in the synaptic cleft (50). AA also regulates PKC activity in neurons (51, 52) and increases NMDA receptor channel activity (53). Therefore the functional association described here between the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, phospholipase  $\text{A}_2$ , and several Glu receptors to trigger AA release is possibly of physiological relevance. The recent observation that AA, in association with mGluR activation, stimulates Glu release in synaptosomes via PKC activation certainly reinforces this possibility (54).

#### Acknowledgments

The authors would like to thank A. Turner-Madeuf for excellent secretarial assistance and M. Passama and L. Charvet for the artwork.

#### References

- Collingridge, G. L., and R. A. J. Lester. Excitatory amino acid receptors in the vertebrate central nervous system. *Pharmacol. Rev.* **40**:143–210 (1989).
- Monaghan, D. T., R. J. Bridges, and C. W. Cotman. The excitatory amino acid receptors: their classes, pharmacology and distinct properties in the function of the central nervous system. *Annu. Rev. Pharmacol. Toxicol.* **29**:365–402 (1989).
- Schoepp, D. D., J. Bockaert, and F. Sladeczek. Pharmacological and functional characteristics of metabotropic excitatory amino acid receptors. *Trends Pharmacol. Sci.* **11**:508–515 (1990).
- Watkins, J. C., P. Krogsgaard-Larsen, and T. Honoré. Structure-activity relationship in the development of excitatory amino acid receptor agonists and competitive antagonists. *Trends Pharmacol. Sci.* **11**:25–33 (1990).
- Murphy, S. W., and R. J. Miller. Two distinct glutamate receptors regulate  $\text{Ca}^{2+}$  homeostasis in hippocampal neurons *in vitro*. *Mol. Pharmacol.* **35**:671–680 (1989).
- Nakanishi, S. Molecular diversity of glutamate receptors and implications for brain function. *Science (Washington D. C.)* **258**:597–603 (1992).
- Mayer, M. L., and R. J. Miller. Excitatory amino acid receptors, second messengers and regulation of intracellular calcium in mammalian neurones. *Trends Pharmacol. Sci.* **11**:254–259 (1990).
- Sladeczek, F., J.-P. Pin, M. Récasens, J. Bockaert, and S. Weiss. Glutamate stimulates inositol phosphate formation in striatal neurones. *Nature (Lond.)* **317**:717–719 (1985).
- Nicoletti, F., M. J. Iadarola, J. T. Wroblewski, and E. Costa. Excitatory amino acid recognition sites coupled with inositol phospholipid metabolism: developmental changes and interaction with  $\alpha 1$ -adrenoreceptors. *Proc. Natl. Acad. Sci. USA* **83**:1931–1935 (1986).
- Masu, M., Y. Tanabe, K. Tsuchida, R. Shigemoto, and S. Nakanishi. Sequence and expression of a metabotropic glutamate receptor. *Nature (Lond.)* **349**:760–765 (1991).
- Houamed, K. M., J. L. Kuijper, T. L. Gilbert, B. A. Haldeman, P. J. O'Hara, E. R. Mulvihill, W. Almers, and F. S. Hagen. Cloning, expression, and gene structure of a G-protein-coupled glutamate receptor from rat brain. *Science (Washington D. C.)* **252**:1318–1321 (1991).
- Pin, J.-P., C. Waeber, L. Prézeau, J. Bockaert, and S. F. Heinemann. Alternative splicing generates metabotropic glutamate receptors inducing different patterns of calcium release in *Xenopus* oocytes. *Proc. Natl. Acad. Sci. USA* **89**:10331–10335 (1992).
- Dumuis, A., M. Sebben, L. Haynes, J. P. Pin, and J. Bockaert. NMDA receptors activate the arachidonic acid cascade system in striatal neurones. *Nature (Lond.)* **336**:68–70 (1988).
- Dumuis, A., J. P. Pin, K. Oomagari, M. Sebben, and J. Bockaert. Arachidonic acid released from striatal neurones by joint stimulation of ionotropic and metabotropic quisqualate receptors. *Nature (Lond.)* **347**:182–184 (1990).
- Garthwaite, J., S. L. Charles, and R. Chess-Williams. Endothelium-derived relaxing factor on activation of NMDA receptors suggest role as intercellular messenger in the brain. *Nature (Lond.)* **336**:385–388 (1988).
- Garthwaite, J., E. Southam, and M. Anderton. A kainate receptor linked to nitric oxide synthesis from arginine. *J. Neurochem.* **53**:1952–1954 (1989).
- Garthwaite, J. Glutamate, nitric oxide and cell-cell signalling in the nervous system. *Trends Neurosci.* **14**:60–67 (1991).

18. Bredt, D. S., and S. Snyder. Nitric oxide mediates glutamate-linked enhancement of cGMP levels in the cerebellum. *Proc. Natl. Acad. Sci. USA* **86**:9030-9033 (1989).
19. Marin, P., M. Lafon-Cazal, and J. Bockaert. A nitric oxide-synthase activity selectively stimulated by NMDA receptors via protein kinase C activation in mouse striatal neurons. *Eur. J. Neurosci.* **4**:425-432 (1992).
20. Marin, P., J.-F. Quignard, M. Lafon-Cazal, and J. Bockaert. Non-classical glutamate receptors, blocked by both NMDA and non-NMDA antagonists, stimulate nitric oxide production in neurons. *Neuropharmacology* **32**:29-36 (1993).
21. Williams, J., M. Errington, M. Lynch, and T. Bliss. Arachidonic acid induces a long-term activity-dependent enhancement of synaptic transmission. *Nature (Lond.)* **341**:739-742 (1989).
22. O'Dell, T. J., R. Hawkins, E. Kandel, and O. Arancio. Tests of the roles of diffusible substances in long-term potentiation: evidence for nitric oxide as a possible early retrograde messenger. *Proc. Natl. Acad. Sci. USA* **88**:11285-11289 (1992).
23. Böhme, G. A., C. Bon, J. M. Stutzman, A. Doble, and J.-C. Blanchard. Possible involvement of nitric oxide in long-term potentiation. *Eur. J. Pharmacol.* **199**:379-381 (1991).
24. Schuman, E. M., and D. V. Madison. A requirement for the intracellular messenger nitric oxide in long-term potentiation. *Science (Washington D. C.)* **254**:1503-1512 (1991).
25. Bon, C., G. A. Böhme, A. Doble, J. M. Stutzman, and J.-C. Blanchard. A role for nitric oxide in long-term potentiation. *Eur. J. Neurosci.* **4**:420-424 (1992).
26. Izumi, Y., D. B. Clifford, and C. F. Zorumski. Inhibition of long-term potentiation by NMDA-mediated nitric oxide release. *Science (Washington D. C.)* **257**:1273-1276 (1992).
27. Haley, J. E., G. L. Wilcox, and P. F. Chapman. The role of nitric oxide in hippocampal long-term potentiation. *Neuron* **8**:211-216 (1992).
28. Kennedy, M. Regulation of synaptic transmission in the central nervous system: long-term potentiation. *Cell* **59**:777-787 (1989).
29. Collingridge, G. L., and W. Singer. Excitatory amino acid receptors and synaptic plasticity. *Trends Pharmacol. Sci.* **11**:290-296 (1990).
30. Nowak, L., P. Bregestovski, P. Ascher, A. Herbert, and A. Prochiantz. Magnesium gates glutamate-activated channels in mouse central neurons. *Nature (Lond.)* **307**:462-465 (1984).
31. Mayer, M. L., G. L. Westbrook, and P. B. Guthrie. Voltage-dependent block by Mg<sup>2+</sup> of NMDA responses in spinal cord neurones. *Nature (Lond.)* **309**:261-263 (1984).
32. Weiss, S., J.-P. Pin, M. Sebben, D. Kemp, F. Sladeczek, J. Gabrion, and J. Bockaert. Synaptogenesis of cultured striatal neurones in serum-free medium: a morphological and biochemical study. *Proc. Natl. Acad. Sci. USA* **83**:2238-2242 (1986).
33. Oomagari, K., B. Buisson, A. Dumuis, J. Bockaert, and J.-P. Pin. Effect of glutamate and ionomycin on the release of arachidonic acid, prostaglandins and HETEs from cultured neurons and astrocytes. *Eur. J. Neurosci.* **3**:928-939 (1991).
34. Ledvora, R. F., and C. Hegyvary. Dependence of Na<sup>+</sup>/Ca<sup>2+</sup> exchange on monovalent cations. *Biochim. Biophys. Acta* **729**:123-136 (1983).
35. Mayer, M. L., and G. L. Westbrook. The physiology of excitatory amino acids in the vertebrate central nervous system. *Prog. Neurobiol.* **28**:197-276 (1987).
36. Kleyman, T. R., and E. J. Cragoe, Jr. Amiloride and its analogs as tools in the study of ion transport. *J. Membr. Biol.* **105**:1-21 (1988).
37. Brown, L., E. J. C. Cragoe, Jr., K. C. Abel, W. M. Simon, and J. R. Bourke. Amiloride analogues induce response in isolated rat cardiovascular tissues by inhibition of Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **344**:220-224 (1991).
38. Vigne, P., C. Frelin, E. J. Cragoe, Jr., and M. Lazdunski. Structure-activity relationships of amiloride and certain of its analogues in relation to blockade of the Na<sup>+</sup>/H<sup>+</sup> exchange system. *Mol. Pharmacol.* **25**:131-136 (1984).
39. Tagliatela, M., L. M. T. Canzoniero, E. J. Cragoe, Jr., G. F. Di Renzo, and L. Annunziato. Na<sup>+</sup>/Ca<sup>2+</sup> exchange activity in central nerve endings. II. Relationship between pharmacological blockade by amiloride analogues and dopamine release from tuberoinfundibular hypothalamic neurons. *Mol. Pharmacol.* **38**:393-400 (1990).
40. Blaustein, M. P. Calcium transport and buffering in neurons. *Trends Neurosci.* **11**:438-443 (1988).
41. Lederer, W. J., E. Niggli, and W. Hadley. Sodium-calcium exchange in excitable cells. *Science (Washington D. C.)* **248**:283 (1990).
42. Stys, P. K., S. G. Waxman, and B. R. Ransom. Ionic mechanisms of anoxic injury in mammalian CNS white matter: role of Na<sup>+</sup> channels and Na<sup>+</sup>-Ca<sup>2+</sup> exchanger. *J. Neurosci.* **12**:430-439 (1992).
43. Kimura, J., A. Noma, and H. Inasawa. Na<sup>+</sup>/Ca<sup>2+</sup> exchange current in mammalian heart cells. *Nature (Lond.)* **319**:596-597 (1986).
44. Nicoll, D. A., S. Longoni, and K. D. Philipson. Molecular cloning and functional expression of cardiac sarcolemmal Na<sup>+</sup>-Ca<sup>2+</sup> exchanger. *Science (Washington D. C.)* **250**:562-564 (1990).
45. Knöpfel, T., and C. Staub. Activation of metabotropic excitatory amino acid receptors induces an inward current in cerebellar Purkinje cells. *Pflüger's Arch. (Suppl. 4)* R12 (1991).
46. Tanabe, Y., M. Masu, T. Ishii, R. Shigemoto, and S. Nakanishi. A family of metabotropic glutamate receptors. *Neuron* **8**:169-179 (1992).
47. Manzoni, O., L. Prezeau, F. Sladeczek, and J. Bockaert. *trans*-ACPD inhibits cAMP formation via a pertussis toxin-sensitive G-protein. *Eur. J. Pharmacol.* **225**:357-358 (1992).
48. Prezeau, L., O. Manzoni, V. Homburger, F. Sladeczek, K. Curry, and J. Bockaert. Characterization of a metabotropic glutamate receptor: direct negative coupling to adenylate cyclase and involvement of a pertussis toxin-sensitive G protein. *Proc. Natl. Acad. Sci. USA* **89**:8040-8044 (1992).
49. Manzoni, O. J. J., F. Finiels-Marlier, I. Sassetti, J. Bockaert, C. Le Peuch, and F. A. J. Sladeczek. The glutamate receptor of the Qp-type activates protein kinase-C and is regulated by protein kinase-C. *Neurosci. Lett.* **109**:146-151 (1990).
50. Barbour, B., M. Szatkowski, N. Ingledew, and P. Atwell. Arachidonic acid induces a prolonged inhibition of glutamate uptake into glial cells. *Nature (Lond.)* **342**:918-920 (1989).
51. Sekiguchi, K., M. Tsukuda, K. Ogita, U. Kikkawa, and Y. Nishizuka. Three distinct forms of rat brain protein kinase C: differential response to unsaturated fatty acids. *Biochem. Biophys. Res. Commun.* **145**:797-802 (1987).
52. Keyser, D. O., and B. E. Alger. Arachidonic acid modulates hippocampal calcium current via protein kinase C and oxygen radicals. *Neuron* **5**:545-553 (1990).
53. Miller, B., M. Sarantis, S. F. Traynelis, and D. Atwell. Potentiation of NMDA receptor currents by arachidonic acid release. *Nature (Lond.)* **355**:722-725 (1992).
54. Herrero, I., T. Miras-Portugal, and J. Sanchez-Prieto. Positive feedback of glutamate exocytosis by metabotropic presynaptic receptor stimulation. *Nature (Lond.)* **360**:163-166 (1992).

Send reprint requests to: Dr. Aline Dumuis, Centre CNRS-INSERM de Pharmacologie-Endocrinologie, Rue de la Cardonille, 34094 Montpellier Cedex 5, France.