
Previous Editors

- **1965** Avram Goldstein, *Editor*. Stanford University.
- **1968** Paul Talalay, *Editor*; Donald S. Coffey, *Associate Editor*. Johns Hopkins University.
- 1971 Steven E. Mayer, *Editor*; Palmer W. Taylor, *Associate Editor*. University of California at San Diego.
- **1975** George I. Drummond, *Editor*; H. Joseph Goren, *Associate Editor*. University of Calgary.
- **1978** Norman Kirschner, *Editor*; Theodore A. Slotkin, *Associate Editor*. Duke University.
- Joel G. Hardman, *Editor*; Lee Limbird, F. Peter Guengerich, *Associate Editors*. Vanderbilt University.
- William A. Catterall, *Editor*; Joseph A. Beavo, Mont R. Juchau, Neil M. Nathanson, Daniel R. Storm, Frank F. Vincenzi, *Associate Editors*. University of Washington, Seattle.
- T. Kendall Harden, Editor, Raymond J. Dingledine, R. L. Juliano, James W. Putney, Jr., Kenneth D. Tew, Ronald G. Thurman, Associate Editors. University of North Carolina at Chapel Hill.



Williams & Wilkins Quick Copies. Tell us today. Read it tomorrow.

You can get copies of any article published in any Williams & Wilkins journal. It's fast – because we're the primary source, we can fax an article to you within 24 hours of when we receive your request. It costs less – there are no copyright fees and the cost of faxing is included in the price. It's affordable – look at the table to the right.

RATES	S, PER AR	TICLE
1-5	pages	\$10
6 - 10	pages	\$12
11-15	pages	\$15
16 - 20	pages	\$18
21 - 25	pages	\$20
26 +	pages	\$25


(\$20, Federal Express)

Fax, phone, or mail your citation to:

Williams & Wilkins Quick Copies

428 East Preston Street Baltimore, MD 21202-3993 Call 1-800-354-9685 (toll free) Fax 1-410-528-4312





The Only Comprehensive Assay System for Receptor Signaling.

THE CYTOSENSOR SYSTEM:
FUNCTIONAL RESPONSES IN MINUTES



The **Cytosensor Microphysiometer System** is the latest tool used for drug discovery and signal transduction research! Using a patented biosensor, the system can monitor the receptor-mediated responses from living cells in minutes, producing functional dose responses in just a few hours. Because of the unique biosensor, receptor activation can be studied even without previous knowledge of the signal transduction pathway. Thus, a single assay system can be used to evaluate a variety of receptor families. Previously time-consuming experiments can be performed quickly and easily, making it possible to obtain EC₅₀ and IC₅₀ values in just 4-7 hours; without radioactivity. And novel receptor responses, not seen with other methods, have been observed.

1. Science 257, 1906-1912. (1992)
2. J. NIH Bespearch 5, 69, (1993)

Call us for an evaluation of your cells and effector agents or for a detailed list of scientific references.



G PROTEIN-LINKED RECEPTORS



TYROSINE KINASE RECEPTORS



LIGAND-GATED ION CHANNELS



HEMATOPOIETIC RECEPTORS



U.S.A. 1-800-400-9060 (FAX: 408-747-3001) Germany: 49-898-54-5050 (FAX: 49-898-54-2238) United Kingdom: 44-293-619579 (FAX: 44-293-619586)