

PERSPECTIVE

New Determinants of Receptor-Effector Coupling: Trafficking and Compartmentation in Membrane Microdomains

RENNOLDS S OSTROM

Department of Pharmacology, University of California, San Diego, La Jolla, California

Received January 2, 2002; accepted January 3, 2002

This article is available online at <http://molpharm.aspetjournals.org>

The multicomponent modular nature of G protein-coupled receptor (GPCR) systems provides cells with numerous potential combinations by which to transduce signals. A typical cell appears to express a dozen or so different GPCR genes (of which nearly a thousand exist in the human genome), several different combinations of G protein subunits and multiple isoforms of effector molecules that can be activated by each type of G protein. The differential expression of these various proteins allows modulation of signals at many levels, resulting in messages that are customized for a specific cell type. Much of the effort to understand signaling via these complicated cellular networks has focused on defining the linear progression of molecular interactions involved in a given pathway. This approach has yielded important information regarding the types of pathways (i.e., the class of G protein and cognate downstream effectors) that are characteristically activated by a given receptor and, thereby, the types of alterations in cell function that are elicited. The current dogma is that high-affinity protein-protein interactions determine the identity of the G protein with which a particular GPCR interacts and, in turn, dictates the biochemical pathways that are activated by that receptor. However, numerous observations in various cells and tissues have indicated that different receptors coupling to the same G protein in a single cell can elicit different biochemical or cellular responses (Hayes and Brunton, 1982; Buxton and Brunton, 1983; Harper et al., 1985; Graeser and Neubig, 1993; Xu et al., 1996; Steinberg and Brunton, 2001). The classical view that GPCR signal transduction is one-dimensional cannot readily account for these observations. Two additional dimensions must be incorporated into our conceptual models: 1) the compartmentation of receptors and effector molecules in subcellular compartments and microdomains of the plasma membrane, and 2) the movement, or translocation, of receptors between cellular compartments (trafficking).

Compartmentation and Caveolae

The concept of receptor translocation is certainly not a new one. The decades old observations that binding sites are lost and responses desensitize following agonist exposure were explained, at least in part, by internalization and sequestration of receptors (Clark et al., 1985; Kassis et al., 1986). A more contemporary idea is that cellular plasma membranes do not uniformly express GPCR effector molecules but do so in specific membrane microdomains (i.e., compartmentation) (Neubig, 1994; Anderson, 1998; Okamoto et al., 1998; Ostrom et al., 2000). Incorporation of these two concepts with classical views of GPCR signaling is opening the door to a new understanding of the biology of signal transduction. We are just now gaining the knowledge and expertise to study the morphology and structural mechanisms by which cells compartment signaling components to begin to assess how the localization and translocation of a receptor and its effectors can influence its signaling.

Recent investigations have identified caveolae as key microdomains of the plasma membrane that appear to concentrate (perhaps preassemble) certain components of signal transduction pathways and, in some cases, serve as sites of internalization. Caveolae differ biochemically from another specialized region of the plasma membrane implicated in receptor-trafficking, clathrin-coated pits. Caveolae are enriched in sphingolipid and cholesterol, making these lipid domains more buoyant than other portions of the cell and facilitating their isolation using sucrose density centrifugation. Clathrin-coated pits and caveolae serve as sites that transport different types of molecules and thus appear to represent similar but distinct pathways for internalization of GPCR. A growing list of signaling molecules has been found to reside in caveolae or to closely associate (i.e., immunoprecipitate) with caveolins (Shaul and Anderson, 1998; Ostrom

ABBREVIATIONS: GPCR, G protein-coupled receptors, B₁R, bradykinin B₁ receptor; B₂R, bradykinin B₂ receptor; cPLA₂, cytosolic phospholipase A₂; MAP kinase, mitogen-activated protein kinase; AC, adenylyl cyclase; PLC, phospholipase C; PIP₂, phosphatidylinositol; GFP, green fluorescent protein.

et al., 2000). Various GPCR and receptor tyrosine kinases have been localized in caveolae or caveolin-rich cellular fractions along with many of the molecules critical for transducing the signals initiated by these types of receptors: e.g., G proteins, adenylyl cyclase, protein kinase C, nitric oxide synthase and the components of the mitogen-activated protein (MAP) kinase cascade (extracellular signal-regulated protein kinase, MAP kinase kinase, Raf, and Ras) (Okamoto et al., 1998; Shaul and Anderson, 1998; Ostrom et al., 2000). Some GPCR are enriched or excluded from caveolae whereas certain GPCR reportedly translocate out of or into caveolae upon activation by an agonist. Therefore, caveolae appear to act as centers that concentrate certain signaling molecules while excluding others, making them key domains that mediate compartmentation within the plasma membrane.

Translocation without Sequestration

In this issue, a report by Sabourin et al. (2002) describes a novel variation on the theme of receptor trafficking via caveolae. These workers report studies in which the rabbit bradykinin B₁ receptor (B₁R) was fused with a fluorescent protein, and trafficking of the chimeric protein was then studied following agonist exposure. The authors observe that upon activation by an agonist, B₁R translocated to caveolin-rich membrane microdomains but did not subsequently appear in intracellular compartments. These results are consistent with a recent report by Lamb et al., who studied trafficking of the human B₁R and B₂R (Lamb et al., 2001). However, these latter investigators noted a 10-fold decrease in agonist affinity of their B₁R-GFP construct relative to the wild-type receptor, whereas Sabourin et al. (2002) detected no functional change when GFP was fused to the rabbit B₁R. Lamb et al. (2001) were also unable to assess subcellular localization of their B₁R-GFP construct due to limited expression levels. The trafficking of the B₁R contrasts with that of the B₂R: in studies by Sabourin et al. (2002) and in previous reports from other groups, B₂R-GFP chimeras translocated to caveolin-rich membranes upon activation but then rapidly internalized (de Weerd and Leeb-Lundberg, 1997; Haasemann et al., 1998; Lamb et al., 2001). Trafficking of the B₁R also contrasts with other examples of caveolae-related GPCR translocation, including cardiac β_2 -adrenergic and A₁ adenosine receptors that move out of caveolin-rich membranes upon agonist exposure and, in the case of β_2 -adrenergic receptors, internalize via clathrin-coated pits (Cao et al., 1998; Lasley et al., 2000; Rybin et al., 2000; Ostrom et al., 2001). Taken together, these findings make evident that many permutations of receptor localization (caveolar or noncaveolar), translocation (into/out of caveolae or no translocation), and internalization (positive or negative, caveolae or coated pits) are employed in GPCR biology. Why and how such permutations occur is not yet clear.

What could be the purpose of a GPCR translocating to a vesicular domain if it does not undergo endocytosis? Perhaps the answer can be found in considering what receptor translocation accomplishes. Classically, receptor movement to a specific domain has been considered a means to associate with specialized regions of the plasma membrane capable of mediating endocytosis of the receptor as part of the desensitization/resensitization mechanism. However, recent evidence demonstrates a second key role for such movement: localization of the receptor promotes coupling to a particular

signaling pathway. For example, cardiac β_2 -adrenergic receptors couple sequentially to the stimulation of cAMP production and then to the activation of MAP kinases—a switching that appears to require endocytosis of the receptor (Daaka et al., 1998). Therefore, some cells utilize GPCR translocation as a means for gaining signaling diversity. This is likely a conserved theme in biology.

Localization as a Determinant of Signaling

An interesting implication of the report by Sabourin et al. (2002) is that the signaling of B₁R mirrors its trafficking and contrasts with that of both the trafficking and signaling of the B₂R. Both kinin receptors couple to the G_{q/11} family of G proteins and the activation of phospholipase C (PLC, via G_{q/11}) and cytosolic phospholipase A₂ (cPLA₂) (Farmer and Burch, 1992). The increases in intracellular Ca²⁺ concentrations stimulated by both receptors desensitize following exposure to agonist, but B₁R signaling through cPLA₂ continues unabated (Mathis et al., 1996; Zhou et al., 2000). As illustrated in a hypothetical schematic (Fig. 1), both receptors translocate from noncaveolar regions of the plasma membrane to caveolin-rich membranes upon agonist activation. This change in location likely facilitates linkage of activated receptor-G protein complex with effector enzyme (PLC) and substrate (phosphatidylinositol, PIP₂), both of which are enriched in caveolae (Pike and Casey, 1996; Pike and Miller, 1998). PLC-mediated production of the second messenger inositol trisphosphate leads to Ca²⁺ mobilization, whereas MAP kinase activation (via G_{βγ} and diacylglycerol) contributes to the activation of cPLA₂ (Qiu et al., 1998). At this point the signaling of the two receptor subtypes appears to diverge. B₂Rs are phosphorylated on the cytoplasmic tail and subsequently desensitize, terminating all signaling (Faussner et al., 1998; Pizard et al., 1999). These receptors then rapidly undergo sequestration via internalization of the caveolar vesicle to the endosomal compartment where the resensitization process presumably begins (Krueger et al., 1997). Meanwhile, B₁Rs are not similarly phosphorylated and remain in caveolar membranes on the cell surface where they can continue to interact with extracellular agonist and activate MAP kinase signaling (the upstream components of this pathway are enriched in caveolae) and cPLA₂ activity (Xing et al., 1997; Faussner et al., 1998). Therefore, the different trafficking of these two GPCR leads to distinct signaling characteristics, especially in terms of the kinetics of the responses.

The concept of location and translocation as determinants of GPCR coupling to signal transduction pathways is a relatively new one. Recent data obtained with cardiac myocytes demonstrate that the coupling of three different endogenous receptors to a particular effector enzyme is determined primarily by the colocalization of the receptor and effector (Ostrom et al., 2001). In this case, the effector enzyme (adenylyl cyclase type 6, AC6) is almost exclusively expressed in caveolin-rich membranes. The β_1 - and β_2 -adrenergic receptors are significantly enriched in these same microdomains and couple to the activation of AC6, whereas a prostanoïd receptor is excluded from caveolin-rich membrane fractions and cannot activate AC6 (despite its ability to activate G_s). Interestingly, the ability of β_2 -adrenergic receptors to activate AC6 in these cells is diminished in comparison to β_1 -adrenergic receptors. This less efficacious coupling by the β_2 -adrenergic receptor is

attributed to its translocation out of caveolae upon agonist exposure (presumably to sequester in clathrin-coated pits) while the β_1 -adrenergic receptors remain colocalized with AC6 in caveolae. Therefore, GPCR coupling to a given effector pathway is dependent upon both receptor-G protein coupling as well as the physical proximity of the G protein to a suitable effector.

Key Questions

Several questions are raised by the present report by Sa-bourin et al. (2002) that will require further study. Do kinin receptors translocate to caveolin-rich membranes in all cells, particularly cells endogenously expressing these receptors? Is caveolar localization of kinin receptors dependent only upon protein sequence of the receptors or does it require other molecules that may be expressed in a cell-specific manner (such as caveolins or other unknown proteins)? B_2 Rs can translocate to caveolae in multiple cell types, including smooth muscle (de Weerd and Leeb-Lundberg, 1997), but similar studies on endogenous B_1 Rs have not been reported. What exactly are the determinants of protein localization and sequestration in caveolae? The differential state of phosphorylation of the B_2 R versus the B_1 R appear to be a critical determinant of kinin receptor desensitization (Faussner et al., 1998; Pizard et al., 1999), but little is known about the mechanisms regulating trafficking in caveolae. Do the B_1 R and B_2 R translocate to the same caveolar domains? The fact that both kinin receptors translocate to caveolae but only the B_2 R undergoes internalization argues for different subpopulations of caveolae-like regions. One such region may be detergent-insoluble lipid rafts, which are present in the plasma membrane of all cells and represent domains that are biochemically similar to, but morphologically distinct from, caveolae. Some G proteins partition to rafts whereas others prefer caveolae, but the differences between these structures (other than the expression of caveolins) are not clear (Oh and Schnitzer, 2001). It is also possible that B_2 Rs are only transient occupants of caveolae and internalize in a different type of vesicle, such as clathrin-coated pits (Liu et al., 1996). Do B_2 Rs initiate signals that induce internalization of caveolar vesicles while B_1 Rs do not? Finally, do the kinin receptors generate any signal in noncaveolar domains because PLC and PIP_2 are reportedly enriched in caveolae (Pike and Casey, 1996; Pike and Miller, 1998)? Given that PLC and PIP_2 are not completely excluded from noncaveolar membranes and the kinetics of receptor translocation are probably too slow to account for the rapidity of signal onset, it is likely that some signaling occurs before these receptors translocate to caveolae (Fig. 1). These and other questions will likely serve as the basis for investigations of kinin receptors and other GPCR in the near future.

Localization and compartmentation of the components of GPCR signal transduction represent new complexities in receptor signaling, forcing us to move beyond one-dimensional characterization of signaling pathways and consider the three-dimensional organization of these networks within the context of cell structure. A challenge for the future is to merge the evolving biochemical data with more direct morphological documentation of the fate of receptors and their various signaling partners, a challenge made more formidable by the number of molecules involved and the cell-specific, sometimes dynamic, nature of their localization.

Acknowledgments

I am indebted to Drs. Paul Insel and Laurence Brunton for their helpful comments and discussion.

References

- Anderson R (1998) The caveolae membrane system. *Annu Rev Biochem* 67:199–225.
- Buxton IL and Brunton LL (1983) Compartments of cyclic AMP and protein kinase in mammalian cardiomyocytes. *J Biol Chem* 258:10233–10239.




Fig. 1. Schematic diagram illustrating the localization and translocation of kinin receptors and the potential effect on their signaling. A, both kinin receptor subtypes (B_1 R and B_2 R) are predominantly localized in non-caveolar domains of the plasma membrane of unstimulated cells. B, upon exposure to agonist (triangles), both kinin receptor subtypes translocate to caveolin-rich domains, probably after activating the G protein and generating a signal via PLC (a membrane-associated protein) in non-caveolar domains. Since PLC and its substrate, PIP_2 , are enriched in caveolae, the bulk of the signaling by both receptors likely occurs in this domain (Pike and Casey, 1996; Pike and Miller, 1998). C, the B_1 R does not internalize, but remains in caveolin-rich domains where it can couple to effectors of different signaling pathways (e.g., the MAP kinase cascade) that are expressed in this location. In contrast, the B_2 R internalizes rapidly, inducing only transient signals in the caveolin-rich domain. Thus, signals generated by the B_1 R are more prolonged and likely result in more pronounced signaling via the effectors colocalized in caveolin-rich domains.

- Cao TT, Mays RW, and von Zastrow M (1998) Regulated endocytosis of G-protein-coupled receptors by a biochemically and functionally distinct subpopulation of clathrin-coated pits. *J Biol Chem* **273**:24592–24602.
- Clark RB, Friedman J, Prashad N, and Ruoho AE (1985) Epinephrine-induced sequestration of the beta-adrenergic receptor in cultured S49 WT and cyc- lymphoma cells. *J Cyclic Nucleotide Protein Phosphor Res* **10**:97–119.
- Daaka Y, Luttrell LM, Ahn S, Della Rocca GJ, Ferguson SS, Caron MG, and Lefkowitz RJ (1998) Essential role for G protein-coupled receptor endocytosis in the activation of mitogen-activated protein kinase. *J Biol Chem* **273**:685–688.
- de Weerd WF and Leeb-Lundberg LM (1997) Bradykinin sequesters B2 bradykinin receptors and the receptor-coupled $G\alpha$ subunits $G\alpha_q$ and $G\alpha_i$ in caveolae in DDT1 MF-2 smooth muscle cells. *J Biol Chem* **272**:17858–17866.
- Farmer SG and Burch RM (1992) Biochemical and molecular pharmacology of kinin receptors. *Annu Rev Pharmacol Toxicol* **32**:511–536.
- Faussner A, Proud D, Towns M, and Bathon JM (1998) Influence of the cytosolic carboxyl termini of human B1 and B2 kinin receptors on receptor sequestration, ligand internalization, and signal transduction. *J Biol Chem* **273**:2617–2623.
- Graeser D and Neubig RR (1993) Compartmentation of receptors and guanine nucleotide-binding proteins in NG108–15 cells: lack of cross-talk in agonist binding among the alpha 2-adrenergic, muscarinic, and opiate receptors. *Mol Pharmacol* **43**:434–443.
- Haasemann M, Cartaud J, Muller-Esterl W, and Dunia I (1998) Agonist-induced redistribution of bradykinin B2 receptor in caveolae. *J Cell Sci* **111**:917–928.
- Harper JF, Haddox MK, Johanson RA, Hanley RM, and Steiner AL (1985) Compartmentation of second messenger action: immunocytochemical and biochemical evidence. *Vitam Horm* **42**:197–252.
- Hayes JS and Brunton LL (1982) Functional compartments in cyclic nucleotide action. *J Cyclic Nucleotide Res* **8**:1–16.
- Kassir S, Olasmaa M, Sullivan M, and Fishman PH (1986) Desensitization of the beta-adrenergic receptor-coupled adenylate cyclase in cultured mammalian cells. Receptor sequestration versus receptor function. *J Biol Chem* **261**:12233–12237.
- Krueger KM, Daaka Y, Pitcher JA, and Lefkowitz RJ (1997) The role of sequestration in G protein-coupled receptor resensitization. Regulation of beta2-adrenergic receptor dephosphorylation by vesicular acidification. *J Biol Chem* **272**:5–8.
- Lamb ME, De Weerd WF, and Leeb-Lundberg LM (2001) Agonist-promoted trafficking of human bradykinin receptors: arrestin- and dynamin-independent sequestration of the B2 receptor and bradykinin in HEK293 cells. *Biochem J* **355**:741–750.
- Lasley R, Prakash N, Uittenbogaard A, and Smart E (2000) Activated cardiac adenosine A1 receptors translocate out of caveolae. *J Biol Chem* **275**:4417–4421.
- Liu P, Ying Y, Ko YG, and Anderson RG (1996) Localization of platelet-derived growth factor-stimulated phosphorylation cascade to caveolae. *J Biol Chem* **271**:10299–10303.
- Mathis SA, Criscimagna NL, and Leeb-Lundberg LM (1996) B1 and B2 kinin receptors mediate distinct patterns of intracellular Ca^{2+} signaling in single cultured vascular smooth muscle cells. *Mol Pharmacol* **50**:128–139.
- Neubig RR (1994) Membrane organization in G-protein mechanisms. *FASEB J* **8**:939–946.
- Oh P and Schnitzer JE (2001) Segregation of heterotrimeric G proteins in cell surface microdomains G(q) binds caveolin to concentrate in caveolae, whereas g(i) and g(s) target lipid rafts by default. *Mol Biol Cell* **12**:685–698.
- Okamoto T, Schlegel A, Scherer PE, and Lisanti MP (1998) Caveolins, a family of scaffolding proteins for organizing “preassembled signaling complexes” at the plasma membrane. *J Biol Chem* **273**:5419–5422.
- Ostrom RS, Gregorian C, Drenan RM, Xiang Y, Regan JW, and Insel PA (2001) Receptor number and caveolar co-localization determine receptor coupling efficiency to adenylyl cyclase. *J Biol Chem* **276**:42063–42069.
- Ostrom RS, Post SR, and Insel PA (2000) Stoichiometry and compartmentation in G protein-coupled receptor signaling implications for therapeutic interventions involving Gs. *J Pharmacol Exp Ther* **294**:407–412.
- Pike LJ and Casey L (1996) Localization and turnover of phosphatidylinositol 4,5-bisphosphate in caveolin-enriched membrane domains. *J Biol Chem* **271**:26453–26456.
- Pike LJ and Miller JM (1998) Cholesterol depletion delocalizes phosphatidylinositol bisphosphate and inhibits hormone-stimulated phosphatidylinositol turnover. *J Biol Chem* **273**:22298–22304.
- Pizard A, Blaukat A, Muller-Esterl W, Alhenc-Gelas F, and Rajerison RM (1999) Bradykinin-induced internalization of the human B2 receptor requires phosphorylation of three serine and two threonine residues at its carboxyl tail. *J Biol Chem* **274**:12738–12747.
- Qiu ZH, Gijon MA, de Carvalho MS, Spencer DM, and Leslie CC (1998) The role of calcium and phosphorylation of cytosolic phospholipase A2 in regulating arachidonic acid release in macrophages. *J Biol Chem* **273**:8203–8211.
- Rybin VO, Xu X, Lisanti MP, and Steinberg SF (2000) Differential targeting of beta-adrenergic receptor subtypes and adenylyl cyclase to cardiomyocyte caveolae. A mechanism to functionally regulate the cAMP signaling pathway. *J Biol Chem* **275**:41447–41457.
- Sabourin T, Bastien L, Bachvarov DR, and Marceau F (2002) Agonist-induced translocation of the kinin B1 receptor to caveolae-related rafts. *Mol Pharmacol* **61**:546–553.
- Shaul PW and Anderson RG (1998) Role of plasmalemmal caveolae in signal transduction. *Am J Physiol* **275**:L843–L851.
- Steinberg SF and Brunton LL (2001) Compartmentation of g protein-coupled signaling pathways in cardiac myocytes. *Annu Rev Pharmacol Toxicol* **41**:751–773.
- Xing M, Tao L, and Insel PA (1997) Role of extracellular signal-regulated kinase and PKC alpha in cytosolic PLA2 activation by bradykinin in MDCK-D1 cells [published erratum appears in *Am J Physiol* 1997 Aug;273(2 Pt 1):section C following table of contents]. *Am J Physiol* **272**:C1380–C1387.
- Xu X, Zeng W, Diaz J, and Muallem S (1996) Spatial compartmentalization of Ca^{2+} signaling complexes in pancreatic acini. *J Biol Chem* **271**:24684–24690.
- Zhou X, Prado GN, Taylor L, Yang X, and Polgar P (2000) Regulation of inducible bradykinin B1 receptor gene expression through absence of internalization and resensitization. *J Cell Biochem* **78**:351–362.

Address correspondence to: Rennolds S Ostrom, Department of Pharmacology, University of California, San Diego, 9500 Gilman Drive, 0636, La Jolla, CA 92093-0636. E-mail: rostrom@ucsd.edu