

The Virally Encoded Fungal Toxin KP4 Specifically Blocks L-Type Voltage-Gated Calcium Channels

MATTHEW J. GAGE, STANLEY G. RANE, GREGORY H. HOCKERMAN, and THOMAS J. SMITH

Donald Danforth Plant Science Center, St. Louis, Missouri (M.J.G., T.J.S.); Fujisawa Research Institute of America, Northwestern University/Evanston Research Park, Evanston, Illinois (S.G.R.); Medicinal Chemistry/Molecular Pharmacology, Purdue University, West Lafayette, Indiana (G.H.H.)

Received September 20, 2001; accepted January 16, 2002

This article is available online at <http://molpharm.aspetjournals.org>

ABSTRACT

KP4 is a virally encoded fungal toxin secreted by the P4 killer strain of *Ustilago maydis*. Previous studies demonstrated that this toxin inhibits growth of the target fungal cells by blocking calcium uptake rather than forming channels, as had been suggested previously. Unexpectedly, this toxin was also shown to inhibit voltage-gated calcium channel activity in mammalian cells. We used whole-cell patch-clamp techniques to further characterize this activity against mammalian cells. KP4 is shown to specifically block L-type calcium channels with weak voltage dependence to the block. Because KP4 activity is abrogated by calcium, KP4 probably binds competitively with

calcium to the channel exterior. Finally, it is shown that chemical reagents that modify lysine residues reduce KP4 activity in both patch-clamp experiments on mammalian cells and in fungal killing assays. Because the only lysine residue is K42, this residue seems to be crucial for both mammalian and fungal channel activity. Our results defining the type of mammalian channel affected by this fungal toxin further support our contention that KP4 inhibits fungal growth by blocking transmembrane calcium flux through fungal calcium channels, and imply a high degree of structural homology between these fungal and mammalian calcium channels.

Interstrain inhibition in *Ustilago maydis* was discovered during heterokaryon experiments (Puhalla, 1968). The inhibitory factors (killer toxins) were shown to be secreted proteins encoded by double-stranded RNA mycoviruses (Hankin and Puhalla, 1971). Killer toxins have been identified in eight genera of yeast (Young, 1987b), but the killer toxins of *U. maydis* are the only ones known in a filamentous fungus.

KP4 is a single polypeptide of 105 amino acids produced by the UMV4 virus that infects the P4 strain of *U. maydis* (Park et al., 1994). It is the only *U. maydis* toxin not processed by Kex2p, and there is no sequence similarity to other toxins (Ganesa et al., 1991; Park et al., 1994). Although most of the yeast toxins are acidic (Bussey, 1972) and the KP6 and KP1 toxins have neutral pI values (Levine et al., 1979), KP4 is extremely basic, with a pI >9.0 (Ganesa et al., 1989). KP4 is an α/β sandwich protein with a relatively compact structure (Gu et al., 1995). From a tenuous structural similarity to the scorpion toxin AaHII from *Androctonus australis*, it was suggested (Gu et al., 1995) and then subsequently shown (Gage et al., 2001) that KP4 inhibits calcium uptake in fungal cells and that KP4 effects on fungal cells are reversible. To further support this structural similarity, Lys58 in AaHII

and the analogous lysine in KP4, Lys42, were both shown to be crucial for activity (Gage et al., 2001).

Calcium is a ubiquitous signaling molecule that plays an important role in the life cycle of both mammalian and fungal cells. In mammalian cells, calcium is involved in processes such as gene expression (Bean, 1989), neuronal migration, and neurotransmitter release (Catterall, 1998). In fungi, calcium is involved in, but not limited to, bud formation (Davis, 1995), hyphal elongation (Jackson and Heath, 1993), and cAMP regulation (Iida et al., 1990a). In both mammalian and fungal systems, cytosolic calcium levels are normally maintained between 100 and 200 nM, whereas extracellular calcium concentrations are normally between 0.1 and 10 mM (Halachmi and Eilam, 1989; Iida et al., 1990b; Clapham, 1995). This calcium gradient is maintained through a series of Ca^{2+} channels, antiporters, and pumps (Tsien and Tsien, 1990; Cunningham and Fink, 1994).

In mammals, there are two primary classes of calcium channels: low-voltage-activated and high-voltage-activated (HVA) channels. The primary low-voltage-activated channel is the T-type channel, which is found in a wide range of cell types (Catterall, 1998). HVA channels include L-, N-, P-, Q-, and R-type channels (Catterall, 1998). L-type channels are the primary HVA channel type in muscle and cardiac cells, whereas N-, P-, Q-, and R-type calcium channels are located primarily in neuronal cells (Catterall, 1995). L-type channels

This work was supported by National Institutes of Health grant GM10704 (to T.J.S.).

ABBREVIATIONS: HVA, high-voltage-activated; DMEM, Dulbecco's modified Eagle's medium; FBS, fetal bovine serum.

also play a role in hormone release in endocrine cells (Milani et al., 1990) and in gene expression in neurons (Bean, 1989). N-, P-, and Q- type channels play roles in neurotransmitter release (Catterall, 1998).

Relative to the mammalian systems, very little information is known about calcium channels in fungi. Two genes that have been identified as possible calcium channels in *Saccharomyces cerevisiae* are MIDI and CCHI. Both the CCHI and MIDI gene products have been shown to be involved in calcium import in *S. cerevisiae* (Iida et al., 1994; Fisher et al., 1997; Locke et al., 2000). The sequence of CCHI is similar to the α_1 subunit of animal voltage-gated calcium channels. MIDI does not have any sequence similarity to known ion channels, but has been reported to be a stretch-activated cation channel (Kanzaki et al., 1999). It has been suggested that the CCHI and MIDI gene products interact to regulate calcium import (Fisher et al., 1997; Locke et al., 2000).

Animal toxins have played a key role in the characterization of mammalian voltage-gated calcium channels. L-type calcium channels were first identified using dihydropyridine compounds and a large number of organic compounds are known to modulate the function of L-type channels. In contrast, N- and P- type calcium channels were first identified using, respectively, ω -conotoxin GVIA from the cone snail *Conus geographus* and ω -agatoxin IVA from the spider *Agelenopsis aperta* [reviewed by Olivera (1994)]. N-type calcium channels were the first dihydropyridine-insensitive channels identified (Fox et al., 1987). Since the discovery of ω -conotoxin GVIA, many other peptide toxins that target calcium channels have been identified in the venoms of mollusks, arthropods, and snakes. The differential specificity of these toxins has played an important role in increasing understanding of the function and pharmacology of calcium channels.

Here we continue these studies and demonstrate that KP4 specifically blocks L-type voltage-gated calcium channels. This result clearly eliminates the possibility that KP4 affects mammalian calcium channels in a nonspecific manner. We further show that KP4 acts in a weakly voltage-dependent fashion. Finally, chemical modification studies demonstrate a tight correlation between KP4 activity against fungal and mammalian cells. The most likely modification site of these reagents is K42 and therefore also supports our contention of the structural homology between KP4 and AaHII.

Materials and Methods

KP4 Purification

KP4 was purified as reported previously (Gu et al., 1994, 1995). In brief, the toxin was isolated from the supernatant of the KP4 toxin expressing strains of *U. maydis* (or *S. cerevisiae*) grown in complete *U. maydis* media (2.5% bacto-peptone, 1% dextrose, 0.15% ammonium nitrate, 0.1% yeast extract) for 7 to 10 days. Cells were removed by centrifugation at 10,000g for 30 min. The supernatant was stirred overnight with CM Sephadex-25 beads (Amersham Biosciences, Piscataway, NJ) that were equilibrated with 25 mM sodium acetate, pH 5.5. The toxin was eluted with 1 M NaCl using a Pharmacia GradiFrac system. The eluant was concentrated using a Minitan II Ultrafiltration System (Millipore, Bedford, MA) with 1-kDa cutoff membranes and dialyzed against a 10 mM malonic acid, pH 6.0 buffer. KP4 was then purified

using a high-resolution cation-exchange chromatography (Mono-S; Amersham Biosciences) matrix attached to a fast-performance liquid chromatography system in the same buffer and using NaCl for elution. The toxin was further purified with size exclusion chromatography using an Amersham Biosciences Superdex-75 gel filtration column. Toxin activity was tested throughout the purification using the killing-zone activity assay described below and purity was assessed using Homogenous 20 SDS gels on an Amersham Biosciences Phastgel system. Only a single band representing KP4 was observable when silver staining was used to observe the protein bands.

Killing-Zone Activity Assay

KP4-sensitive P2 cells were grown overnight in complete *U. maydis* media. P2 cells (~1 ml/100 ml) were added to warm complete *U. maydis* media containing 2% bacto-agar and poured into 100- × 20-mm culture dishes. Once the agar solidified, wells were cut into the agar and 10 μ l of the test solutions were added to each well. The plates were then incubated at 30°C for ~36 h. KP4 activity presents a clear zone around the point of application.

Cell Culture

PC12 cells. wtPDGF-R expressing PC12 cells (Vaillancourt et al., 1995) were cultured on rattail collagen-coated plastic tissue culture dishes. Cells were grown in Dulbecco's modified Eagle's medium (DMEM; Invitrogen, Carlsbad, CA), with 12.5% heat inactivated horse serum derived from platelet-poor plasma (to prevent differentiation; Sigma-Aldrich, St. Louis, MO) and 2.5% fetal bovine serum (FBS; Atlas Biological, Fort Collins, CO) at 37°C with 5% CO₂. Cell medium was exchanged every other day and cells were passaged weekly. Cells were plated in 35-mm rattail collagen-coated dishes for electrophysiology. Cells were differentiated by addition of fresh 100 ng/ml nerve growth factor to the cell medium every other day for 7 days.

GH₃ Cells. GH₃ cells were cultured on plastic tissue culture dishes. Cells were grown in DMEM/Ham's F-12 medium (Invitrogen) with 10% FBS (Atlas Biological, Fort Collins, CO) at 37°C with 5% CO₂. Cell medium was exchanged every other day and cells were passaged weekly. Cells were plated in 35-mm tissue culture dishes for electrophysiology.

tsA-201 Cells. tsA-201 cells, a subclone of the human embryonic kidney line 293 that expresses simian virus 40 T antigen were cultured on plastic tissue culture dishes. Cells were grown in DMEM/Ham's F-12 medium (Invitrogen) with 10% FBS (Atlas Biological, Fort Collins, CO) at 37°C with 5% CO₂. Cell medium was exchanged every other day and cells were passaged biweekly.

Expression

Wild-type Ca_v2.1 (de Weille et al., 1991), Ca_v1.2 (Snutch et al., 1991), and Ca_v2.3 (Soong et al., 1993) channel subunits were expressed with Ca_v β_{1b} (Pragnell et al., 1991) and Ca_v $\alpha_2\delta$ (Ellis et al., 1988) channel subunits and enhanced green fluorescent protein (CLONTECH, Palo Alto, CA) in tsA-201 cells. Cells were transfected using the reagent Geneporter2 (Gene Therapy Systems, San Diego, CA) and an equimolar ratio of the three channel subunit cDNAs along with 0.8 μ g of enhanced green fluorescent protein for a total of 4 μ g of DNA. Transfected cells were detected by fluorescence at 510 nm with excitation at 480 nm.

Electrophysiology

A standard whole-cell bath solution was used for the PC12 and GH₃ experiments: 10 mM BaCl₂, 135 mM tetraethylam-

Downloaded from molpharm.aspetjournals.org by guest on December 1, 2012

whole-cell current block. This indicates that KP4 might have different affinities for the different types of calcium channels producing more or less block depending on the proportion of channel subtypes in a given cell line. If KP4 were nonspecific, the level of modulation should be dependent on KP4 concentration and independent of channel type.

The magnitude of KP4 effects correlated with the known expression levels of L-type calcium channels in the various cell lines tested, indicating that KP4 might specifically bind L-type calcium channels. To test this hypothesis, the effects of 14 μM KP4 on whole-cell calcium currents in PC12 cells were measured in the presence of 5 μM nimodipine, a dihydropyridine calcium channel blocker. Nimodipine (5 μM) was also placed in the micropipet applicator with the 14 μM KP4 to prevent dilution of the local nimodipine concentration. If KP4 specifically blocks L-type calcium channels, then KP4 should have no measurable effect in the presence of nimodipine. Table 1 summarizes the results of this experiment (representative raw data traces for differentiated PC12 cell experiments are shown in Fig. 1). KP4 (14 μM) has no apparent effect on cells treated with 5 μM nimodipine. This suggests that KP4 may be specific for L-type calcium channels.

To further demonstrate that KP4 is specific for L-type calcium channels, we looked at KP4 specificity versus heterologously expressed individual calcium channel subtypes. Three different calcium channel subtypes were tested: $\text{Ca}_v1.2$, an L-type calcium channel; $\text{Ca}_v2.1$, a P/Q-type calcium channel; and $\text{Ca}_v2.3$, an R-type calcium channel. In the case of expressed $\text{Ca}_v1.2$ channels (Fig. 2A), 1 μM KP4 blocks 52% of the current. The IC_{50} of KP4 for expressed calcium channels was determined to be 1.24 μM (Fig. 2D). However, concentrations of up to 100 μM KP4 have no effect on either $\text{Ca}_v2.1$ or $\text{Ca}_v2.3$ (Fig. 2, B and C). KP4 clearly is specific for L-type channels of the Ca_v1 family.

Mechanism of KP4 Block of $\text{Ca}_v1.2$. Phenylalkylamine and benzothiazepine compounds often act in a voltage-dependent manner (Hering et al., 1997; Motoike et al., 1999). Voltage-dependent block has several characteristics: shifts in the steady-state inactivation curve, changes in the $\tau_{\text{inactivation}}$, and use-dependent block. Drugs that block in a voltage-dependent manner do so with different efficacy depending on the holding potential.

By ascertaining whether KP4 affected calcium currents in a voltage-dependent manner, the pharmacological effects of




Fig. 2. Effect of KP4 on heterologously expressed calcium channels. Effects of 1 μM KP4 on the whole cell calcium current from heterologously expressed $\text{Ca}_v1.2$ channels (Figure A). KP4 blocks 54% of the current ($n = 8$). Effects of 100 μM KP4 on the whole cell calcium current from heterologously expressed $\text{Ca}_v2.1$ (Figure B; $n = 7$) and $\text{Ca}_v2.3$ (Figure C; $n = 8$) channels, respectively. KP4 clearly does not affect the current in either $\text{Ca}_v2.1$ and $\text{Ca}_v2.3$ channels. D, the dose-response curve of KP4 on $\text{Ca}_v1.2$ channels. Each point represents the results from three to nine experiments. The IC_{50} for KP4 is 1.24 μM .

KP4 could be compared with these drugs. In these first studies, the effects of KP4 on steady state inactivation of α_{1C} calcium channels were measured. As shown in Fig. 3A, 1 μ M KP4 induces a negative shift in the steady state inactivation with a shift in the half-inactivation voltage ($V_{1/2}$) from -19.8 ± 2.1 mV to -28.5 ± 3.8 mV ($n = 5$). For comparison, verapamil and diltiazem induce negative shifts of 30 mV (Cai et al., 1997), which is much larger than the ~ 9 mV shift caused by KP4.

In the next set of experiments, it was ascertained whether KP4 blocked calcium channels in a use-dependent manner. Use-dependent block is an important feature of therapeutically useful calcium channel antagonists. Phenylalkylamines, such as verapamil and D600, show an accumulation

of block on the order of 50 to 80% during repetitive depolarization (Johnson et al., 1996). A 1-Hz train of 20 100-ms depolarizations to +10 mV was applied in the presence of 1 μ M KP4 (Fig. 3B). KP4 induces a 10% accumulation of block under these conditions. This is smaller than the accumulation of block observed for verapamil and D600.

Finally, if KP4 acted in a voltage-dependent manner, changes in the holding potential should affect the activity of KP4. As Fig. 3C shows, a 30 mV shift in holding potential from -90 mV to -60 mV causes an increase in block of from 53 ± 5 to $62 \pm 2\%$, or a change of $\sim 9\%$. In contrast, the block caused by verapamil increases $\sim 50\%$ when the holding potential is shifted from -100 to -60 mV (Cai et al., 1997). From these experiments, KP4 seems to act in a weakly volt-




Fig. 3. Effects of KP4 on the biophysical properties of $\text{Ca}_v1.2$ channels. For these experiments, tsA 201 cells were transfected with $\text{Ca}_v1.2$, $\text{Ca}_v\beta_{1b}$, and $\text{Ca}_v\alpha_2\delta$ channel subunit cDNA. Barium currents were evoked by 100-ms depolarizations to various voltages ($-50 \sim +50$ mV) from a holding potential of -60 mV in the presence or absence of 1 μ M KP4. A, steady-state inactivation curve from a representative cell before and after acute treatment with 1 μ M KP4. Absolute currents have been normalized to the maximum current. Treatment with 1 μ M KP4 causes a shift in the ($V_{1/2}$) of ~ 9 mV toward more negative potentials. The error on the control cells was ± 2 mV and ± 4 mV for the KP4 treated cells. B, use-dependent block with 1 μ M KP4. Accumulation of block by 1 μ M KP4 was measured with a 1 Hz train. An accumulation of block of $\sim 10\%$ is seen. C, effects of a shift in holding potential on block of calcium current by 1 μ M KP4. KP4 (1 μ M) blocks $53.87 \pm 4.71\%$ of the absolute current at a holding potential of -90 mV. This increases to $62.26 \pm 2.11\%$ when the holding potential is shifted to -60 mV. These results indicate that KP4 acts in a weakly voltage-dependent manner. D, KP4 does not modify the activation of $\text{Ca}_v1.2$ channels. For this figure, the channel conductance (g_{Ba}) was calculated using the equation: $g_{\text{Ba}} = I_{\text{Ba}} / (E - E_{\text{rev}})$, where I_{Ba} is the peak barium current evoked by the test potential E , and E_{rev} is the estimated $\text{Ca}_v1.2$ channel reversal potential. The conductance-voltage relationships are plotted as relative conductance ($g_{\text{Ba}} / \text{max } g_{\text{Ba}}$) versus test potentials. The data were fit to the equation: $g_{\text{Ba}} / \text{max } g_{\text{Ba}} = 1 / (1 + \exp [-(V - V_{g0.5})/k])$, where V is the test potential used to evoke barium current, $V_{g0.5}$ is the potential where channel conductance is half-maximal [0.77 ± 0.50 mV in the absence (\bullet) and 1.83 ± 1.67 mV in the presence (\circ) of 1 μ M KP4], and k is the slope factor (6.1 in the absence and 7.2 in the presence of 1 μ M KP4). Results shown are mean values \pm S.E. ($n = 6$). Because both curves are essentially identical, KP4 does not seem to modify the voltage dependence of activation.

age dependent manner. Figure 3D shows that KP4 does not alter the voltage dependence of channel activation. This is in contrast to hanatoxin modulation of K^+ channels (Swartz and MacKinnon, 1997) and α - (Calahan, 1975) and β - (Jonas et al., 1986) scorpion toxin modulation of sodium channels.

Calcium Effects on Current Modulation of KP4. In *U. maydis*, the activity of KP4 is abrogated by increasing extracellular calcium concentrations (Gage et al., 2001). As little as 10 mM $CaCl_2$ reduces the killing activity of KP4 and 100 mM $CaCl_2$ completely abrogates its effects. If the fungal and mammalian channels are functionally homologous, then the effects of KP4 on mammalian calcium channels should also be abrogated by extracellular calcium. To test for this, KP4 activity was measured in a bath solution, where the normal concentration of 10 mM barium chloride was replaced with 5 mM calcium chloride. Table 1 shows the modulation of the peak current in GH_3 cells treated with 14 μM KP4 in both a Ba^{2+} and Ca^{2+} bath. When Ba^{2+} is replaced by Ca^{2+} in the bath solution, KP4 activity is abolished (Table 1, raw data traces shown in Fig. 4). This is consistent with our finding in *U. maydis* that KP4 effects are abrogated by exogenous calcium and further demonstrates that the effects of KP4 on mammalian and fungal calcium channels are analogous.

Chemical Modification of Lysine 42. K42 has been shown to play an important role in KP4 activity (Gage et al., 2001). In the case of the scorpion toxin AaHII, chemical modification of lysine residues abrogated its channel blocking activity (Sampieri and Habersetzer-Rochat, 1978). To further test the structural/functional homology between AaHII and KP4 and to again demonstrate that the effects of KP4 on fungal calcium channels is analogous to its effects on mammalian channels, similar chemical modification studies were performed on KP4. Acetic anhydride modifies primary amines, the hydroxyl group of tyrosine, the thiol group of cysteine, and the amino group of histidine. Under these reaction conditions, histidine and cysteine spontaneously deacetylate and the tyrosine can be deacetylated with hy-

droxylamine. This leaves only acetylated primary amines. The degree of modification can be determined fluorometrically (Fig. 5A) from the ratio of the slopes of the modified and the unmodified KP4. Eighty-five percent of the primary amines (1.7 modification sites per KP4) were modified in Modified KP4 sample 1, which was used for fungal killing assays. 65% of the primary amines (1.3 modification sites per KP4) were modified in Modified KP4 sample 2, which was used for electrophysiology.

Acetylation clearly abrogates KP4 activity against both fungal and mammalian cells. The effects of acetylation on KP4 activity were examined using the fungal killing assay (Fig. 5B). Modified toxin is $\sim 85\%$ less active than the unmodified KP4, consistent with the degree of modification. At a concentration of 36 μM , acetylated KP4 has very weak activity, whereas there is no apparent activity at 3.6 μM . To determine the effects of chemical modification on KP4's inhibition of mammalian calcium channels, GH_3 calcium currents were measured. Modified KP4 (14 μM) blocked $17 \pm 4\%$ of the calcium current (Fig. 5C). This is 33% of the block found for wild-type KP4. Because 65% of the primary amines are modified, the decrease in block correlates well with the degree of modification. These results clearly demonstrate that KP4 activity against fungal cells correlates well with inhibition of mammalian calcium channels. Furthermore, because it is more than likely that the sole lysine, K42, is being modified by the acetic anhydride, K42 seems to be crucial for toxin activity. All of these results together strongly suggest a structural/functional homology between KP4 and the scorpion toxin AaHII and between fungal and mammalian L-type calcium channels.

Discussion

The results presented here demonstrate that the virally encoded fungal toxin KP4 inhibits mammalian L-type voltage-gated calcium channels. Although it might seem unusual




Fig. 4. Effects of Ca^{2+} on the activity of KP4. A, representative calcium current traces from GH_3 cells with 10 mM Ba^{2+} in the extracellular bath solution. B, representative traces from GH_3 cells with 5 mM Ca^{2+} in the extracellular bath solution. KP4 effects are abrogated by the presence of Ca^{2+} in the bath solution.

that the activity of a calcium channel blocker crosses animal and fungal kingdoms, it is not without precedence because venoms from mollusks, arthropods, and snakes contain peptides that inhibit mammalian voltage-gated calcium channels. What is unusual here, however, is that KP4 is the first toxin shown to act across such phylogenetically divergent organisms yet at a specific molecular target.

As shown here, KP4 specifically targets L-type calcium channels. Both undifferentiated and differentiated PC12 cells express multiple calcium channel types, L, N, and P/Q, and both have modest L type current components (Usovich et al., 1990; Lievano et al., 1994). KP4 blocks 20% of the total whole-cell calcium current in undifferentiated PC12 cells, and this is consistent with the contribution of the L-type current to the total in these cells. The percentage contribution of L current to total calcium current is reduced in differentiated cells because they have a relatively larger proportion of N type current. Consistent with this reduced L current percentage KP4 was observed to block less of the total current in differentiated cells. When PC12 cells are treated with the L-type channel blocker, nimodipine, the cells become insensitive to KP4. This is strong evidence that KP4

targets L-type calcium channels. When KP4 activity is tested against heterologously expressed calcium channels, KP4 is active against $\text{Ca}_v1.2$ but not against $\text{Ca}_v2.1$ and $\text{Ca}_v2.3$ channels. Therefore, we conclude that KP4 is specific for L-type calcium channels. What is particularly interesting is that, of all of the calcium channels found in mammalian cells, these results suggest that it is the L-type calcium channels that most closely resemble fungal channels being targeted by KP4.

From the studies on the pharmacological effects of KP4 on mammalian cells, it is also apparent that KP4 blocks $\text{Ca}_v1.2$ in a manner distinct from other calcium channel blocking agents. Unlike the small-molecule calcium channel inhibitors, verapamil, and D600, KP4 induces only a small shift in the voltage-dependence of inactivation ($V_{1/2}$) but not a significant change in $\tau_{\text{inactivation}}$. An increase in the frequency of stimulation from 0.05 to 1 Hz results in only a slight increase in block of $\text{Ca}_v1.2$ by KP4. Finally, KP4 demonstrates a very small but significant preference for block of $\text{Ca}_v1.2$ channels held at more depolarized membrane potentials. Therefore, KP4 seems to act in a weakly voltage-dependent manner. The characteristics of $\text{Ca}_v1.2$ channel block by KP4 are thus dif-




Fig. 5. Effects on the activity of KP4 from chemical modification. **A**, determination of the degree of chemical modification of KP4 samples used for fungal killing assays and electrophysiology. The degree of modification is determined by the ratio of the slopes. Modified KP4 sample 1, used for fungal killing assays, was 85% modified. Modified KP4 sample 2, used for electrophysiology, was 65% modified. **B**, fungal killing assay demonstrating the effects of modification on activity. These results show that KP4 is ~85% less active than native KP4, consistent with the degree of modification. **C**, current-voltage curve of a representative GH₃ cell treated with 14 μM modified KP4. **D**, raw data traces showing the effects of 14 μM KP4 on the whole cell calcium current of GH₃ cells. Modified KP4 blocks $17.46 \pm 4.11\%$ of the raw current ($n = 6$). Modified KP4 blocks whole-cell calcium current ~33% as well as native KP4, consistent with the degree of modification.

ferent from those of phenylalkylamine and benzothiazepine drugs but similar to those reported for block of $\text{Ca}_v1.2$ by the spider toxin ω -agatoxin IIIA (Cohen et al., 1992). These observations, along with the abrogation of KP4 block by high extracellular Ca^{2+} concentrations, suggest that KP4 may bind to the extracellular side of the $\text{Ca}_v1.2$ pore region, much like charybdotoxin block of K^+ channels (MacKinnon and Miller, 1989) or tetrodotoxin block of Na^+ channels (Terlau et al., 1991).

KP4 displays a similar potency to the polypeptide toxin calciseptine, from black mamba venom (de Weille et al., 1991), and is more potent than the ω -conotoxin TxVII (Fainzilber et al., 1996), both of which are specific for L-type calcium channels. Calciseptine also shows a similar sensitivity to calcium as reported here for KP4 (De Weille et al., 1991; Kini et al., 1998). As with calciseptine block of $\text{Ca}_v1.2$, KP4 does not affect the voltage-dependence of activation but slightly shifts the voltage-dependence of inactivation of the channel (Teramoto et al., 1996). Therefore, the mode of $\text{Ca}_v1.2$ inhibition by KP4 seems very similar to that of calciseptine.

The activity of KP4 against mammalian $\text{Ca}_v1.2$ channels closely parallels KP4 antifungal activity. This has been shown by the abrogation of KP4 activity against both mammalian and fungal cells by chemical modification and extracellular calcium. This implies that the fungal channels targeted by KP4 are most similar to these mammalian voltage-gated channels. A potential calcium channel called CCHI has been recently characterized in *S. cerevisiae* (Fisher et al., 1997; Locke et al., 2000). CCHI shows homology to mammalian voltage-gated calcium channels. It is possible that *U. maydis* has a homolog to CCHI and this channel may be the target for KP4.

The level of channel recognition by KP4 may be similar to that observed in the AaHII scorpion toxin. It has been hypothesized that a crucial lysine on AaHII inserts into the pore of the sodium channel via columbic interactions (Darbon et al., 1983). We propose that K42 in KP4 acts in a similar manner by perhaps interacting with the four glutamic acids believed to be responsible for coordinating calcium entry into the pore (Yang et al., 1993). Further studies on KP4 should elucidate the commonalities and differences between these fungal and mammalian calcium channels, which will not only address the evolutionary development of calcium channels but will also help define the role and regulation of fungal calcium channels.

References

- Bean BP (1989) Classes of calcium channels in vertebrate cells. *Annu Rev Physiol* **51**:367–384.
- Bussey H (1972) Effects of yeast killer factor on sensitive cells. *Nat New Biol* **235**:73–75.
- Cai D, Mülle JG, and Yue DT (1997) Inhibition of recombinant Ca^{2+} channels by benzothiazepines and phenylalkylamines: class-specific pharmacology and underlying molecular determinants. *Mol Pharmacol* **51**:872–881.
- Calahan ME (1975) Mapping the receptor site for hanatoxin, a gated modifier of voltage-dependent K^+ channels. *J Physiol (Lond)* **244**:511–534.
- Catterall WA (1995) Structure and Function of Voltage-Gated Ion Channels. *Annu Rev Biochem* **64**:493–531.
- Catterall WA (1998) Structure and function of neuronal Ca^{2+} channels and their role in neurotransmitter release. *Cell Calcium* **24**:307–323.
- Clapham DE (1995) Calcium signaling. *Cell* **80**:259–268.
- Cohen CJ, Ertel EA, Smith MM, Venema VJ, Adams ME, and Leibowitz MD (1992) High affinity block of myocardial L-type calcium channels by the spider toxin omega-AGA-toxin IIIA: advantages over 1,4-dihydropyridines. *Mol Pharmacol* **42**:947–951.
- Cunningham KW and Fink GR (1994) Ca^{2+} transport in *Sacchromyces cerevisiae*. *J Exp Biol* **196**:157–166.
- Darbon H, Jover E, Couraud F, and Rochat H (1983) α -Scorpion neurotoxin derivatives suitable as potential markers of sodium channels. *Int J Pept Protein Res* **22**:179–186.
- Davis TN (1995) Calcium in *Sacchromyces cerevisiae*. *Adv Second Messenger Phosphoprotein Res* **30**:339–358.
- de Weille JR, Schweitz H, Maes P, Tartar A, and Lazdunski M (1991) Calciseptine, a peptide isolated from black mamba venom, is a specific blocker of the L-type calcium channel. *Proc Natl Acad Sci USA* **88**:2437–2440.
- Ellis SB, Williams ME, Ways NR, Brenner R, Sharp AH, Leung AT, Campbell KP, McKenna E, Koch WJ, Hui A, Schwartz A, and Harpold MM (1988) Sequence and expression of mRNAs encoding the $\alpha 1$ and $\alpha 2$ subunits of a DHP-sensitive calcium channel. *Science (Wash DC)* **241**:1661–1664.
- Fainzilber M, Lodder JC, Van der Schors RC, Li KW, Yu Z, Burlingame AL, Geraerts WPM, and Kits KS (1996) A novel hydrophobic ω -conotoxin blocks molluscan dihydropyridine-sensitive calcium channels. *Biochemistry* **35**:8748–8752.
- Fisher M, Schnell N, Chattaway J, Davis P, Dixon G, and Sanders D (1997) The *Sacchromyces cerevisiae* CCH1 gene is involved in calcium influx and mating. *FEBS Lett* **419**:259–262.
- Fox AP, Nowycky MC, and Tsien RW (1987) Kinetic and pharmacological properties distinguishing three types of calcium currents in chick sensory neurons. *J Physiol* **394**:149–172.
- Gage MJ, Bruenn J, Fisher M, Sanders D, and Smith TJ (2001) KP4 fungal toxin inhibits growth in *Ustilago maydis* by blocking calcium uptake. *Mol Microbiol* **41**:775–786.
- Ganesa C, Chang Y-H, Flurkey WH, and Bozarth RF (1989) Purification and molecular properties of the toxin coded by *Ustilago maydis* virus P4. *Biophys Biochem Res Commun* **162**:651–657.
- Ganesa C, Flurkey WH, Randhawa ZI, and Bozarth RF (1991) *Ustilago maydis* virus P4 killer toxin: Characterization, partial amino terminus sequence, and evidence for glycosylation. *Arch Biochem Biophys* **286**:195–200.
- Gu F, Khimani A, Rane SG, Flurkey WH, Bozarth RF, and Smith TJ (1995) Structure and function of a virally encoded fungal toxin from *Ustilago maydis*: a fungal and mammalian Ca^{2+} channel inhibitor. *Structure* **3**:805–814.
- Gu F, Sullivan TS, Che Z, Ganesa C, Flurkey WH, Bozarth RF, and Smith TJ (1994) The characterization and crystallization of a virally encoded *Ustilago Maydis* KP4 toxin. *J Mol Biol* **243**:792–795.
- Halachmi D and Eilam Y (1989) Cytosolic and vacuolar Ca^{2+} concentrations in yeast measured with Ca^{2+} -sensitive dye Indo-1. *FEBS Lett* **256**:55–61.
- Hankin L and Puhalla JE (1971) Nature of a factor causing interstrain lethality in *Ustilago maydis*. *Phytopathology* **61**:50–53.
- Hering S, Aczel S, Kraus RL, Berjukow S, Striessnig J, and Timin EN (1997) Molecular mechanism of use-dependent calcium channel block by phenylalkylamines: role of inactivation. *Proc Natl Acad Sci USA* **94**:13323–13328.
- Iida H, Nakamura H, Ono T, Okumura MS, and Anraku Y (1994) *MIDI*, a novel *Sacchromyces cerevisiae* gene encoding a plasma membrane protein, is required for Ca^{2+} influx and mating. *Mol Cell Biol* **14**:8259–8271.
- Iida H, Sakaguchi S, Yagawa Y, and Anraku Y (1990a) Cell cycle control by Ca^{2+} in *Saccharomyces cerevisiae*. *J Biol Chem* **265**:21216–21222.
- Iida H, Yagawa Y, and Anraku Y (1990b) Essential role for induced Ca^{2+} influx followed by $[\text{Ca}^{2+}]_i$ rise in maintaining viability of yeast cells late in the mating pheromone response pathway. *J Biol Chem* **265**:13391–13399.
- Jackson SL and Heath IB (1993) Roles of calcium ions in hyphal growth. *Microbiol Rev* **57**:367–382.
- Johnson BD, Hockerman GH, Scheuer T, and Catterall WA (1996) Distinct effects of mutations in transmembrane segment V6 on block of L-type calcium channels by structurally similar phenylalkylamines. *Mol Pharmacol* **50**:1388–1400.
- Jonas P, Vogel W, Arantes EC, and Giglio JR (1986) Toxin γ of the scorpion *Tityus serrulatus* modifies both activation and inactivation of sodium permeability of nerve membrane. *Pflüg Arch Eur J Physiol* **407**:92–99.
- Kanzaki M, Nagasawa M, Kojima I, Sato C, Naruse K, Sokabe M, and Iida H (1999) Molecular Identification of a Eukaryotic Stretch-Activated Nonselective Cation Channel. *Science (Wash DC)* **285**:882–886.
- Kini RM, Caldwell RA, Wu QY, Baumgarten CM, Heher JJ, and Evans HJ (1998) Flanking proline residues identify the L-type Ca^{2+} channel binding site of calciseptine and FS2. *Biochem* **37**:9058–9063.
- Levine R, Koltin Y, and Kandel J (1979) Nuclease activity associated with the *Ustilago maydis* virus induced killer proteins. *Nucleic Acids Res* **6**:3717–3732.
- Lievano A, Bolden A, and Horn R (1994) Calcium channels in excitable cells: divergent genotypic and phenotypic expression of alpha 1-subunits. *Am J Physiol* **267**:C411–C424.
- Locke EG, Bonilla M, Liang L, Takita Y, and Cunningham KW (2000) A homolog of voltage-gated Ca^{2+} channels stimulated by depletion of secretory Ca^{2+} in yeast. *Mol Cell Biol* **20**:6686–6694.
- MacKinnon R and Miller C (1989) Mutant potassium channels with altered binding of charybdotoxin, a pore-blocking peptide inhibitor. *Science (Wash DC)* **245**:1382–1385.
- Milani D, Malgaroli A, Guidolin D, Fasolato C, Skaper SD, Meldolesi J, and Pozzan T (1990) Ca^{2+} channels and intracellular Ca^{2+} stores in neuronal and neuroendocrine cells. *Cell Calcium* **11**:191–199.
- Motoike HK, Bodi I, Nakayama H, Schwartz A, and Varadi G (1999) A region in IVS5 of the human cardiac L-type calcium channel is required for the use-dependent block by phenylalkylamines and benzothiazepines. *J Biol Chem* **274**:9409–9420.
- Olivera BM (1994) Calcium channel diversity and neurotransmitter release: The ω -conotoxins and ω -agatoxins. *Annu Rev Biochem* **63**:823–867.
- Park CM, Bruenn JA, Ganesa C, Flurkey WF, Bozarth RF, and Koltin Y (1994) Structure and heterologous expression of *Ustilago maydis* viral toxin KP4. *Mol Microbiol* **11**:155–164.
- Pragnell M, Sakamoto J, Jay SD, and Campbell KP (1991) Cloning and tissue-specific expression of the brain calcium channel beta-subunit. *FEBS Lett* **291**:253–258.

- Puhalla JE (1968) Compatibility reactions on solid medium and interstrain inhibition in *Ustilago maydis*. *Genetics* **60**:461–474.
- Sampieri F and Habersetzer-Rochat C (1978) Structure-function relationships in scorpion neurotoxins: identification of the superreactive lysine in toxin I of *Androctonus australis* Hector. *Biochim Biophys Acta* **535**:100–109.
- Snutch TP, Tomlinson WJ, Leonard JP, and Gilbert MM (1991) Distinct calcium channels are generated by alternative splicing and are differentially expressed in the mammalian CNS. *Neuron* **7**:45–57.
- Soong T, Stea A, Hodson CD, Dubel SJ, Vincent SR, and Snutch TP (1993) Structure and functional expression of a member of the low voltage-activated calcium channel family. *Science (Wash DC)* **260**:1133–1136.
- Stocks SJ, Jones AJM, Ramey CW, and Brooks DE (1986) A fluorometric assay of the degree of modification of protein primary amines with polyethylene glycol. *Analyt Biochem* **154**:232–234.
- Swartz KJ and MacKinnon R (1997) Mapping the receptor site for hanatoxin, a gated modifier of voltage-dependent K⁺ channels. *Neuron* **18**:675–682.
- Teramoto N, Ogata R, Okabe K, Kameyama A, Kameyama M, Watanabe TX, Kuriyama H, and Kitamura K (1996) Effects of calciseptine on unitary barium channel currents in guinea-pig portal vein. *Pflug Arch Eur J Physiol* **432**:462–470.
- Terlau H, Heinemann SH, Stuhmer W, Pusch M, Conti F, Imoto K, and Numa S (1991) Mapping the site of block by tetrodotoxin and saxitoxin of sodium channel II. *FEBS Lett* **293**:93–96.
- Tsien RW and Tsien RY (1990) Calcium channels, stores and oscillations. *Annu Rev Cell Biol* **6**:715–760.
- Usovich MM, Porzig H, Becker C, and Reuter H (1990) Differential expression by nerve growth factor of two types of Ca²⁺ channels in rat pheochromocytoma cell lines. *J Physiol* **426**:95–116.
- Vaillancourt RR, Heasley LE, Zamarripa J, Storey B, Valius M, Kazlauskas A, and Johnson GL (1995) Mitogen-activated protein kinase activation is insufficient for growth factor receptor-mediated PC12 cell differentiation. *Mol Cell Biol* **15**:3644–3653.
- Yang J, Ellinor PT, Sather WA, Zhang J-F, and Tsien RW (1993) Molecular determinants of Ca²⁺ selectivity and ion permeation in L-type Ca²⁺ channels. *Nature (Lond)* **366**:158–161.
- Young T (1987) Killer yeasts, in *The Yeasts* (Rose AH and Harrison JS eds) pp 131–164, Academic Press, Orlando, FL.

Address correspondence to: Dr. Thomas J. Smith, Donald Danforth Plant Science Center, 975 North Warson Road, St. Louis, MO 63132. E-mail: tsmith@danforthcenter.org
