

Evaluation of Muscarinic Agonist-Induced Analgesia in Muscarinic Acetylcholine Receptor Knockout Mice

ALOKESH DUTTARROY,¹ JESUS GOMEZA, JAI-WEI GAN, NASIR SIDDIQUI, ANTHONY S. BASILE, W. DEAN HARMAN, PHILIP L. SMITH, CHRISTIAN C. FELDER, ALLAN I. LEVEY, and JÜRGEN WESS

Laboratory of Bioorganic Chemistry, National Institute of Diabetes Digestive and Kidney Diseases, Bethesda, Maryland (A.D., J.G., J.-W.G., N.S., A.S.B., J.W.); Department of Chemistry, University of Virginia, Charlottesville, Virginia (W.D.H., P.L.S.); Lilly Research Laboratories, Eli Lilly & Company, Indianapolis, Indiana (C.C.F.); and Department of Neurology, Emory University School of Medicine, Atlanta, Georgia (A.I.L.)

Received February 21, 2002; accepted July 23, 2002

This article is available online at <http://molpharm.aspetjournals.org>

ABSTRACT

Centrally active muscarinic agonists display pronounced analgesic effects. Identification of the specific muscarinic acetylcholine receptor (mAChR) subtype(s) mediating this activity is of considerable therapeutic interest. To examine the roles of the M₂ and M₄ receptor subtypes, the two G_i/G_o-coupled mAChRs, in mediating agonist-dependent antinociception, we generated a mutant mouse line deficient in both M₂ and M₄ mAChRs [M₂/M₄ double-knockout (KO) mice]. In wild-type mice, systemic, intrathecal, or intracerebroventricular administration of centrally active muscarinic agonists resulted in robust analgesic effects, indicating that muscarinic analgesia can be mediated by both spinal and supraspinal mechanisms. Strikingly, muscarinic agonist-induced antinociception was totally abolished in M₂/M₄ double-KO mice, independent of the route of application. The nonselective muscarinic agonist oxotremorine showed reduced analgesic potency in M₂ receptor single-KO

mice, but retained full analgesic activity in M₄ receptor single-KO mice. In contrast, two novel muscarinic agonists chemically derived from epibatidine, CMI-936 and CMI-1145, displayed reduced analgesic activity in both M₂ and M₄ receptor single-KO mice, independent of the route of application. Radioligand binding studies indicated that the two CMI compounds, in contrast to oxotremorine, showed >6-fold higher affinity for M₄ than for M₂ receptors, providing a molecular basis for the observed differences in agonist activity profiles. These data provide unambiguous evidence that muscarinic analgesia is exclusively mediated by a combination of M₂ and M₄ mAChRs at both spinal and supraspinal sites. These findings should be of considerable relevance for the development of receptor subtype-selective muscarinic agonists as novel analgesic drugs.

A large body of evidence indicates that muscarinic acetylcholine receptor (mAChR) agonists as well as cholinesterase inhibitors can produce powerful analgesic effects in many different species including man (reviewed by Green and Kitchen, 1986; Pert, 1987; Hartvig et al., 1989; Eisenach, 1999). The antinociceptive effects of these agents are predicted to be mediated by activation of mAChRs present at both spinal and supraspinal sites (Green and Kitchen, 1986; Pert, 1987; Hartvig et al., 1989; Eisenach, 1999). Muscarinic agonists have been shown to be as efficacious as full opioid agonists in stringent antinociceptive tests, including the mouse tail-flick assay (Harris et al., 1969; Howes et al., 1969; Green and Kitchen, 1986; Pert, 1987). In addition, several studies (Widman et al., 1985; Swedberg et al., 1997; Petry et

al., 1998) suggest that the use of muscarinic analgesics is less likely to lead to addiction, physical dependence, and tolerance, severe side effects that limit the use of classic opioid analgesics. Identification of the mAChR subtype(s) mediating these potent analgesic effects is therefore of considerable therapeutic interest.

Molecular cloning studies have revealed the existence of five molecularly distinct mAChR subtypes (M₁–M₅; Caulfield, 1993; Wess, 1996). Multiple mAChRs have been detected in spinal cord, thalamus, and several other regions of the CNS known to be involved in the transmission of pain impulses (Levey et al., 1991; Levey, 1993; Vilaro et al., 1993; Wei et al., 1994; Wolfe and Yasuda, 1995; Höglund and Baghdoyan, 1997). Pharmacological studies have yielded contradictory results regarding the molecular identity of the mAChR subtypes mediating the analgesic effects of muscarinic agonists. For example, it has been proposed that muscarinic antinociception is mediated by M₁ (Bartolini et al.,

This research was supported by a Cooperative Research and Development Agreement between the National Institute of Diabetes and Digestive and Kidney Diseases and the Eli Lilly Research Laboratories.

¹ Current address: Human Genome Sciences, Inc., Rockville, MD 20850.

ABBREVIATIONS: mAChR, muscarinic acetylcholine receptor; KO, knockout; %MPE, percentage of maximum possible effect; CMI-936, 2-exo[5-(3-methyl-1,2,4-oxadiazolyl)]-[2.2.1]-7-azabicycloheptane; CMI-1145, 2-exo[5-(3-amino-1,2,4-oxadiazolyl)]-[2.2.1]-7-azabicycloheptane; CHO, Chinese hamster ovary; NMS, N-methylscopolamine; QNB, quinuclidinyl benzilate; CI, confidence interval.

1992; Ghelardini et al., 2000), M_1 and/or M_2 (Gillberg et al., 1989; Iwamoto and Marion, 1993), M_1 and/or M_3 (Naguib and Yaksh, 1997), M_2 (Ma et al., 2001), M_3 (Honda et al., 2000), or M_4 (Shannon et al., 1997; Ellis et al., 1999) mAChRs. It is likely that the limited receptor subtype selectivity of the currently available muscarinic agonists and antagonists that were used in these *in vivo* studies is the primary reason for these discrepant results (Caulfield, 1993; Wess, 1996). To overcome these difficulties, we and others generated mutant mouse strains in which specific mAChR genes were inactivated via gene targeting techniques (Hamilton et al., 1997; Gomeza et al., 1999a,b; Matsui et al., 2000; Yamada et al., 2001).

We recently demonstrated that the centrally active, non-subtype-selective muscarinic agonist oxotremorine (administered s.c.) showed clearly reduced analgesic potency in M_2 receptor-deficient mice (Gomeza et al., 1999a), suggesting that the M_2 receptor subtype plays a key role in mediating muscarinic antinociception. However, we also noted that oxotremorine-dependent antinociception was not abolished in the M_2 receptor KO mice (Gomeza et al., 1999a), indicating that non- M_2 mAChRs also participate in this activity. Because the M_4 receptor subtype couples to similar G proteins (G_i/G_o family) as the M_2 receptor (Caulfield, 1993; Wess, 1996), we speculated that the analgesic responses remaining in the M_2 receptor KO mice might be mediated by M_4 mAChRs. To test this hypothesis, we generated a mutant mouse strain lacking both M_2 and M_4 mAChRs (M_2/M_4 double-KO mice).

In the present study, we demonstrate that M_2/M_4 double-KO mice do not display any significant analgesic effects after systemic (s.c.), intrathecal (i.t.), or intracerebroventricular (i.c.v.) administration of muscarinic agonists. In contrast, considerable analgesic activity remained in M_2 and M_4 receptor single-KO mice, independent of the route of application. We also noted that the magnitude of analgesic responses remaining in the M_2 and M_4 single-KO mice was critically dependent on the M_2/M_4 receptor selectivity profile of the individual agonists used. These results provide unambiguous evidence that activation of both M_2 and M_4 receptors fully accounts for muscarinic-agonist dependent antinociception at both spinal and supraspinal sites.

Materials and Methods

Generation of M_2/M_4 mAChR Double-KO Mice. The generation of homozygous M_2 receptor KO [genetic background, 129/J1 (50%) \times CF1 (50%)] and M_4 receptor KO [129SvEv (50%) \times CF1 (50%)] mice has been described previously (Gomeza et al., 1999a,b). To generate mice deficient in both M_2 and M_4 mAChRs, homozygous M_2 receptor KO mice were mated with homozygous M_4 receptor KO mice. The resulting F1 compound heterozygotes were then intercrossed to generate F2 mice of nine possible M_2/M_4 genotypes. According to Mendelian inheritance, 1 of 16 of the F2 pups were predicted to be homozygous for both the M_2 and the M_4 receptor gene disruptions (M_2/M_4 double-KO mice). Similarly, 1 of 16 of the F2 pups were predicted to carry two copies of the wild-type M_2 and M_4 receptor genes. The M_2/M_4 double-KO mice were interbred to generate mice used for the experiments described in this study. In parallel, the wild-type F2 mice were interbred to obtain wild-type control mice. Thus, both M_2/M_4 double-KO mice and the corresponding wild-type control mice had an equivalent genetic background [129/J1 (25%) \times 129SvEv (25%) \times CF1 (50%)]. Mouse genotyping was car-

ried out by Southern blotting and polymerase chain reaction analysis of mouse tail DNA as described previously (Gomeza et al., 1999a,b).

M_2 and M_4 receptor single-KO mice were maintained by interbreeding homozygous mutant mice of the F2 generation. In parallel, wild-type littermates of the F2 generation were interbred to generate wild-type control mice of the matching genetic background [M_2 , wild-type and KO, 129/J1 (50%) \times CF1 (50%); M_4 , wild-type and KO, 129SvEv (50%) \times CF1 (50%)].

Unless stated otherwise, all experiments were performed during the light cycle using male mice that were at least 8 weeks old. In all experiments, KO mice and age-matched wild-type mice of the proper genetic background were run in parallel. All animal studies were conducted according to the National Institutes of Health guidelines for standard animal care and usage.

Analgesia Assays. Antinociceptive responses were quantitated by using the tail-flick and hot plate methods, after s.c., i.t., and i.c.v. administration of muscarinic agonists. For analgesia measurements after s.c. administration of drugs, a cumulative dose-response protocol was employed (Wenger, 1980; Adams et al., 1990; Duttaroy et al., 1997). Intrathecal or i.c.v. injections were carried out using a single dose of agonist that induced maximum analgesia in all mice tested (see below) (dose per mouse: oxotremorine, 10 μ g; CMI-936 and CMI-1145, 20 μ g). These doses were chosen based on pilot studies using several different agonist doses (doses per mouse: oxotremorine, 1, 10, and 20 μ g; CMI-936 and CMI-1145, 10, 20, and 50 μ g).

The tail-flick test was carried out using a modified version (radiant heat method) of the procedure described by D'Amour and Smith (1941). Briefly, the mouse was placed on the surface of the tail-flick analgesia meter (Pamotor, Burlingame, CA), and radiant heat was applied from a halogen lamp focused on the dorsal surface of its tail (2–3 cm from the base of the tail). Movement of the tail activated a photocell that turned off the stimulus light and stopped a reaction timer that recorded the time. The intensity of the radiant heat was adjusted so that baseline tail-flick occurred within 2 to 4 s. Withdrawal latency was measured immediately before (baseline) and 30 min after administration of drugs (Ellis et al., 1999; Gomeza et al., 1999a,b); a 10-s cut-off time was imposed to prevent tissue damage. The hot plate test (Woolfe and Macdonald, 1944) was performed on the same set of mice using an electronically controlled hot plate analgesia meter (Columbus Instrument, Columbus, OH). Responses were measured after placing mice on a 55°C hot plate before (baseline, 5–10 s) and 30-min after drug injection; the cut-off time was 30 s. Response latency to the first hind-paw response was recorded. The hind-paw response was either a paw lick or a foot shake, whichever occurred first. Mice that failed to respond within the respective cut-off times were defined as "analgesic".

In the case of the cumulative dose-response procedure, mice were injected s.c. with a low starting dose of drugs (oxotremorine, 0.01 mg/kg; CMI-936 and CMI-1145, 10 mg/kg) and tested 30 min after injection. Mice that were not analgesic (i.e., that responded within the specified cut-off time) were injected with another dose of drug (increment dose) and retested 30 min later, essentially as described by Duttaroy et al. (1997). This procedure was continued until all mice were analgesic (except for the M_2/M_4 double-KO mice, which showed no antinociception, and for the M_4 KO mice after CMI-936 administration due to the toxicity of cumulative CMI-936 doses greater than 680 mg/kg). Cumulative dose (s.c.) ranges were: oxotremorine, 0.01 to 2.62 mg/kg; CMI-936, 10 to 680 mg/kg; CMI-1145, 20 to 1,000 mg/kg. ED₅₀ values (\pm S.D.) for cumulative dose-response curves were calculated via nonlinear regression analysis (Prism v. 3.0; GraphPad Software, San Diego, CA).

For control purposes, we also carried out a series of analgesia studies using a more conventional single dose-response protocol. In this protocol, groups of mice ($n = 5$ –8/dose/genotype) were injected (s.c.) once with a single dose of oxotremorine (dose range, 0.01–1.0 mg/kg) or vehicle and tested for antinociception 30 min later in the tail-flick and hot plate tests. ED₅₀ values (\pm 95% confidence inter-

vals) obtained from a single experiment were calculated via nonlinear regression analysis (Prism).

After i.t. and i.c.v. injection of drugs, antinociceptive responses were expressed as percentage of maximum possible effect (%MPE): tail-flick assay, %MPE = $100 \times [(\text{postdrug latency} - \text{baseline latency}) / (10 - \text{baseline latency})]$; hot plate assay, %MPE = $100 \times [(\text{postdrug latency} - \text{baseline latency}) / (30 - \text{baseline latency})]$.

Drugs and Drug Administration. Oxotremorine sesquifumarate was purchased from Sigma (St Louis, MO). CMI-936 and CMI-1145 were synthesized as described previously (Ellis et al., 1999). Oxotremorine was dissolved in 0.9% NaCl, whereas CMI-936 and CMI-1145 were first dissolved in DMSO and then diluted in 0.9% NaCl to reduce the final concentration of DMSO to less than 0.1%. Subcutaneous injections (injection volume, body weight in grams \times 0.01 ml) were performed using a 1-cc syringe attached to a 26.5-gauge needle. Intrathecal injections (5 μ l/mouse) into the L5/L6 region of the mouse spinal cord were carried out as described previously (Hylden and Wilcox, 1980), using a 30.5-gauge needle attached to 25-gauge tubing that was connected to a Hamilton syringe (0.5 ml). Intracerebroventricular injections (2.5 μ l/mouse) were given freehand (Haley and McCormick, 1957) 0.3 mm posterior, 1.0 mm lateral, and 2.5 mm ventral to the bregma below the outer surface of the skull into the right lateral ventricle using a similarly sized needle and syringe as used for i.t. injections. Mice were anesthetized with pentobarbital (25 mg/kg s.c.) and injected i.t. or i.c.v. with muscarinic agonists 25 to 30 min after pentobarbital injection. Mice recovered from the pentobarbital anesthesia within 10 to 15 min after the i.t. or i.c.v. injections of muscarinic agonists. In control experiments, wild-type mice ($n = 5/\text{group}$) were injected i.t. or i.c.v. with vehicle. These control studies showed that the i.t. or i.c.v. injections themselves had no significant effect on response latencies in the two analgesia assays (data not shown).

For control purposes, we also carried out a series of i.c.v. and i.t. injection experiments using inhalational anesthesia (isoflurane/oxygen; 4:96) (Fortec model; Cyprane, Keighley, UK). Specifically, we assessed the analgesic effects of oxotremorine, administered i.t. or i.c.v. (10 μ g per mouse), in M_2 single-KO and M_2/M_4 double-KO mice and the two corresponding wild-type control groups ($n = 5/\text{group}$), using isoflurane/oxygen as the anesthetizing agent. For the gas anesthesia procedure, mice were placed in a plastic chamber provided with a nozzle and attached to a vacuum line that delivered isoflurane/oxygen (4:96) at a constant rate. Once anesthetized, mice were removed from the chamber and placed on a plane surface to perform the i.c.v. or i.t. injections. Mice recovered from the gas anesthesia within 3 to 5 min after the i.t. or i.c.v. injections of oxotremorine.

Radioligand Binding Studies. To determine the overall density of mAChRs in mouse spinal cord, whole spinal cord tissue was removed and frozen immediately on dry ice. Tissue samples were homogenized by hand with 20 strokes of a Dounce tissue grinder in 0.32 M sucrose, 5 mM Tris-HCl, pH 7.5, and 1 mM phenylmethylsulfonyl fluoride. Membranes were prepared, and ligand binding experiments were carried out using a saturating concentration (2 nM) of the nonselective muscarinic antagonist, N -[^3H]methylscopolamine ([^3H]NMS; 82 Ci/mmol; PerkinElmer Life Science, Boston, MA), essentially as described previously (Dörje et al., 1991; Gomeza et al., 1999a,b). In addition, competition-binding assays were carried out to determine the binding affinities of different muscarinic agonists (oxotremorine, CMI-936, and CMI-1145) at cloned M_2 and M_4 mAChRs. Membranes were prepared from CHO-K1 cell lines individually expressing the cloned human M_2 and M_4 mAChRs and incubated with eight different concentrations of muscarinic agonist in the presence of 50 pM [^3H]NMS, following the protocol described by Dörje et al. (1991). In saturation binding assays, six different concentrations of the radioligand, [^3H]NMS, ranging from 10 to 3000 pM, were tested. Binding buffer consisted of 25 mM sodium phosphate, pH 7.4, and 5 mM MgCl_2 . Incubations were carried out for 3 h at room temperature (22°C). Nonspecific binding was determined in

the presence of 10 μ M atropine. Competition binding data were analyzed using a nonlinear regression curve fitting procedure (Prism: sigmoidal dose-response, variable slope, bottom plateau = 0). IC_{50} values were converted to K_i values by using the Cheng-Prusoff equation (Cheng and Prusoff, 1973).

Immunoprecipitation Assays. For immunoprecipitation studies, mAChR subtype-specific rabbit polyclonal antisera were raised against nonconserved regions of the third cytoplasmic loops of the mouse mAChR proteins (Gomeza et al., 1999a,b; Yamada et al., 2001), following a procedure similar to that described by Levey et al. (1991). Membrane homogenates were prepared (see above) from different areas of the mouse brain and then incubated for 1 h with 2 nM of the nonselective muscarinic antagonist, [^3H]quinuclidinyl benzilate ([^3H]QNB; 81.5 Ci/mmol; PerkinElmer Life Science), washed thoroughly, and solubilized with 1% digitonin, followed by immunoprecipitation of solubilized [^3H]QNB-labeled receptors with receptor subtype-selective antisera, essentially as described previously (Gomeza et al., 1999a,b; Yamada et al., 2001).

Immunohistochemistry. Mice were fixed by perfusion with 4% paraformaldehyde. Frozen sections of cervical spinal cord were processed for immuno-peroxidase staining using an M_2 receptor-specific rat monoclonal antibody (1:500; Levey et al., 1995) or an M_4 receptor-specific rabbit polyclonal antibody (0.75 mg/ml; Levey et al., 1991). After incubation with biotinylated secondary antibody (rabbit anti-rat or goat anti-rabbit) and avidin-biotinylated enzyme complex reagent (Vector Laboratories, Burlingame, CA), sections were developed with diaminobenzidine as described in detail previously (Levey et al., 1991, 1995).

Statistics. Statistical significance between two or more groups was determined by Student's t tests or one-way analysis of variance using post hoc t tests (Bonferroni's method).

Results

Generation of M_2/M_4 mAChR Double-KO Mice. To study the roles of M_2 and M_4 mAChRs in muscarinic agonist-induced analgesia, we generated mice containing targeted disruptions of both the M_2 and M_4 mAChR genes. Initially, we crossed homozygous M_2 receptor KO mice (Gomeza et al., 1999a) with homozygous M_4 receptor KO mice (Gomeza et al., 1999b) to generate compound heterozygotes ($M_2 \pm, M_4 \pm$). The compound heterozygotes were then intermated to obtain homozygous M_2 and M_4 receptor mutant mice (M_2/M_4 double-KO mice). M_2/M_4 double-KO mice were obtained at a frequency of about 6% (the predicted Mendelian frequency is 1 of 16 or 6.25%), indicating that the absence of functional M_2 and M_4 receptors did not lead to an increase in embryonic or postnatal mortality. The inactivation of the M_2 and M_4 receptor genes and the absence of M_2 and M_4 receptor protein in the M_2/M_4 double-KO mice was confirmed by Southern and polymerase chain reaction analyses and immunoprecipitation studies (data not shown), respectively, in a fashion identical to that described for the M_2 and M_4 single-KO mice (Gomeza et al., 1999a,b). The M_2/M_4 double-KO mice showed no obvious morphological or behavioral abnormalities and did not differ from their wild-type littermates in overall health, fertility, and longevity.

We carried out a set of immunoprecipitation studies to verify that inactivation of the M_2 and M_4 mAChR genes did not lead to compensatory changes in the expression levels of the M_1 and M_3 receptor subtypes, the two major G_q -coupled mAChRs [M_5 receptor protein is barely detectable in rat (Yasuda et al., 1993) or mouse brain (M. Yamada and J. Wess, unpublished results)]. Specifically, membrane preparations derived from different areas of the mouse brain (wild-

type and M_2/M_4 double-KO) were incubated with a saturating concentration (2 nM) of the nonselective muscarinic antagonist [3 H]QNB to radioactively label all mAChRs. [3 H]QNB-labeled receptors were solubilized with 1% digitonin and then immunoprecipitated using M_1 and M_3 receptor subtype-selective antisera (Yamada et al., 2001). As shown in Fig. 1, both antisera immunoprecipitated similar amounts of M_1 and M_3 receptors in tissues derived from wild-type and M_2/M_4 double-KO mice, indicating that the lack of functional M_2 and M_4 receptors did not lead to compensatory changes in the expression levels of the M_1 and M_3 receptor proteins.

Analgesic Effects after Systemic Administration of Muscarinic Agonists. In the first set of experiments, wild-type and mAChR mutant mice received s.c. injections of different centrally active muscarinic agonists. Drug-induced antinociceptive responses were quantitated using the tail-flick and hot plate assays (D'Amour and Smith, 1941; Woolfe and Macdonald, 1944), employing a cumulative dose-response protocol (Wenger, 1980; Adams et al., 1990; Duttaroy et al., 1997). The antinociceptive properties of muscarinic agonists were assessed in M_2 single-KO, M_4 single-KO, and M_2/M_4 double-KO mice. In each case, wild-type animals of the corresponding genetic background served as controls (see *Materials and Methods*).

In wild-type mice, the nonselective muscarinic agonist oxotremorine produced profound analgesic effects in both the tail-flick and hot plate tests (ED_{50} , ~ 0.03–0.18 mg/kg s.c.; Fig. 2; Table 1). In M_2 KO mice, as reported previously (Gomez et al., 1999a), the analgesic potency of oxotremorine was significantly reduced in the tail-flick and hot plate assays (13- and 3-fold, respectively) (Fig. 2, A and D; Table 1). However, oxotremorine retained the ability to elicit maximum analgesic responses in the M_2 KO mice in both tests (Fig. 2, A and D). In contrast, oxotremorine showed similar analgesic potencies in M_4 KO mice and the corresponding wild-type control mice (Fig. 2, B and E; Table 1). Strikingly, oxotremorine was completely devoid of antinociceptive activity in M_2/M_4 double-KO mice in both analgesia tests used (Fig. 2, C and F; Table 1).




Fig. 1. Expression levels of M_1 and M_3 mAChRs in wild-type and M_2/M_4 double-KO mice as determined in immunoprecipitation studies. Membranes were prepared from the indicated brain regions derived from wild-type and M_2/M_4 double-KO mice, followed by labeling of mAChRs with 2 nM [3 H]QNB and solubilization of labeled receptors with 1% digitonin. [3 H]QNB-labeled M_1 or M_3 mAChRs were immunoprecipitated with receptor subtype-specific rabbit antisera as described under *Materials and Methods*. Cx, cerebral cortex; Hc, hippocampus; Str, striatum; OB, olfactory bulb; Cer, cerebellum; BS, brain stem. Data are presented as means \pm S.D. ($n = 3$).

To examine whether the results obtained with oxotremorine could be mimicked by other centrally active muscarinic agonists, we carried out analogous studies using two novel muscarinic agonists, CMI-936 and CMI-1145, which are chemically derived from epibatidine (Ellis et al., 1999). A recent study (Ellis et al., 1999) showed that the analgesic effects of these compounds could be partially prevented by injecting mice with the MT-3 snake toxin, which can block M_4 receptors with relatively high selectivity (Jolkkonen et al., 1994; Liang et al., 1996), suggesting that the CMI compounds might induce analgesia preferentially via activation of M_4 mAChRs. However, the in vitro receptor selectivity profile of these compounds was not reported and the possibility cannot be excluded that metabolic degradation of the snake toxin in vivo might have resulted in peptide fragments capable of blocking non- M_4 mAChRs (e.g., M_2). To address this question, we analyzed the analgesic properties of CMI-936 and CMI-1145 in M_2 KO, M_4 KO, and M_2/M_4 double-KO mice and the corresponding wild-type control animals. In addition, we also characterized the receptor subtype selectivity profiles of the two CMI compounds at cloned M_2 and M_4 mAChRs (see below).

In wild-type mice, s.c. administration of CMI-936 and CMI-1145 caused dose-dependent antinociceptive effects in both analgesia assays used (range of ED_{50} values, 48–180 mg/kg s.c.) (Figs. 3 and 4; Table 1). In M_2 KO mice, as observed with oxotremorine, both drugs showed significantly reduced (~2–6-fold) antinociceptive potencies (Figs. 3, A and D, and 4, A




Fig. 2. Antinociceptive responses of wild-type and mAChR KO mice after systemic (s.c.) administration of oxotremorine. A–C, tail-flick assay. D–F, hot plate assay. M_2 KO (A, D), M_4 KO (B, E), and M_2/M_4 double-KO mice (C, F) and the corresponding wild-type (WT) control mice ($n = 5$ –6/group) were injected s.c. with increasing doses of oxotremorine, using a cumulative dose-response protocol. Data are plotted as cumulative oxotremorine dose versus percent mice displaying full analgesia, as described under *Materials and Methods*. The dose-response curves shown are representative of two independent experiments which gave similar results.

and D; Table 1). However, in contrast to oxotremorine, which displayed essentially unchanged analgesic activity in M_4 KO mice (Fig. 2, B and E), CMI-936 and CMI-1145 also exhibited significantly reduced analgesic potencies (~2–5-fold) in M_4 KO mice (Figs. 3, B and E, and 4, B and E; Table 1). Strikingly, the two epibatidine analogs, like oxotremorine, were completely devoid of analgesic activity in M_2/M_4 double-KO mice in both the tail-flick and hot plate assays (Figs. 3, C and F, and 4, C and F; Table 1).

To verify experimentally that the key results obtained after systemic administration of muscarinic agonists were not affected by the use of the cumulative dose-response protocol, we carried out an additional set of studies using the more conventional single dose-response protocol. Specifically, we employed a standard single dose protocol to assess oxotremorine-mediated analgesic responses in M_2 single-KO and M_2/M_4 double-KO mice and the two corresponding wild-type groups. For these studies, groups of mice ($n = 5$ –7/dose/genotype) were injected s.c. with different doses of oxotremorine (0.01, 0.03, 0.1, 0.3, and 1.0 mg/kg) or vehicle, and analgesia responses were recorded 30 min later. This analysis showed that oxotremorine was significantly less potent in both the tail-flick (~13-fold reduction in potency) and hot plate tests (~4-fold reduction in potency) in the M_2 single-KO mice, compared with the corresponding wild-type group. The following ED_{50} values were obtained: tail-flick assay, M_2 WT, 0.025 mg/kg [95% confidence interval (CI), 0.011 to 0.086], M_2 KO, 0.32 mg/kg (95% CI, 0.21 to 0.63); hot plate assay, M_2 WT, 0.12 mg/kg (95% CI, 0.07 to 0.22), M_2 KO, 0.49 mg/kg (95% CI, 0.26 to 0.55). However, at the highest dose used (1 mg/kg s.c.), oxotremorine was still able to induce maximum analgesia (100% MPE) in the M_2 KO mice. On the other hand, oxotremorine-mediated analgesia responses were totally abolished in M_2/M_4 double-KO mice (data not shown). These findings obtained by using a standard single dose-

response protocol closely mimicked those obtained by application of the cumulative dose-response protocol (Fig. 2; Table 1), indicating that the outcome of the analgesia tests was not affected by the nature of the dosing protocol used.

Moreover, to exclude the possibility that the hybrid background of the mAChR wild-type and KO mice may have affected the outcome of the analgesia experiments, we compared oxotremorine-mediated analgesic responses in the 129SvEv (Taconic, Germantown, NY) and CF-1 (Charles River Laboratories, Wilmington, MA) parental mouse strains, using a standard single dose-response protocol ($n = 6$ –8/dose). These studies showed that oxotremorine ED_{50} values did not differ significantly between the two parental mouse strains in either the tail-flick test [CF-1, 0.037 mg/kg (95% CI, 0.021 to 0.086); 129SvEv, 0.029 mg/kg (95% CI, 0.019 to 0.075)] or hot plate assay [CF-1, 0.059 (95% CI, 0.037 to 0.094); 129SvEv, 0.057 mg/kg (95% CI, 0.041 to 0.091)]. Because the two parental strains did not differ in their sensitivity to muscarinic agonist-induced antinociception, it is very unlikely that the key conclusions drawn from the analysis of the mAChR mutant mice used in the present study were significantly affected by the mixed genetic background of the mice used.

Analgesic Effects after Intrathecal and Intracerebroventricular Administration of Muscarinic Agonists. We next wanted to assess the relative contribution of spinal and supraspinal mechanisms to muscarinic agonist-induced analgesic effects. Toward this goal, oxotremorine, CMI-936, and CMI-1145 were administered either i.t. or i.c.v. to wild-type and mAChR KO mice, and analgesic responses were measured using the tail-flick and hot plate tests. In wild-type mice, all three agonists (oxotremorine, 10 μ g/mouse; CMI-936 and CMI-1145, 20 μ g/mouse) produced maximum analgesia in both the i.t. and i.c.v. injection experiments (Figs. 5–7). After i.t. or i.c.v. administration, the

TABLE 1

Analgesic potencies of muscarinic agonists administered systemically (s.c.) to wild-type and mAChR mutant mice

M_2 KO, M_4 KO, and M_2/M_4 double-KO mice and the matching wild-type (WT) control animals were injected s.c. with increasing doses of oxotremorine, CMI-936, or CMI-1145. Analgesic responses were measured using the tail-flick and hot plate assays, employing a cumulative dose-response protocol as described under *Materials and Methods*. ED_{50} values are presented as means \pm S.D. from two independent experiments [($n = 5$ –6/group/experiment)].

	Tail-Flick Assay		Hot Plate Assay	
	ED_{50} mg/kg	ED_{50} -KO/ ED_{50} -WT	ED_{50} mg/kg	ED_{50} -KO/ ED_{50} -WT
Oxotremorine				
WT	0.03 \pm 0.01		0.18 \pm 0.05	
M_2 KO	0.41 \pm 0.29**	13	0.55 \pm 0.22*	3.1
WT	0.03 \pm 0.01		0.13 \pm 0.01	
M_4 KO	0.03 \pm 0.01	1.0	0.16 \pm 0.09	1.2
WT	0.03 \pm 0.01		0.04 \pm 0.02	
M_2/M_4 KO	No analgesia		No analgesia	
CMI-936				
WT	60 \pm 7		160 \pm 22	
M_2 KO	125 \pm 21*	2.1	225 \pm 42*	1.4
WT	75 \pm 36		180 \pm 20	
M_4 KO	370 \pm 173*	4.9	380 \pm 170*	2.1
WT	56 \pm 9		130 \pm 30	
M_2/M_4 KO	No analgesia		No analgesia	
CMI-1145				
WT	48 \pm 28		107 \pm 61	
M_2 KO	141 \pm 96*	2.9	650 \pm 257*	6.1
WT	120 \pm 41		180 \pm 10	
M_4 KO	201 \pm 35*	1.6	380 \pm 41*	2.1
WT	80 \pm 1		170 \pm 6	
M_2/M_4 KO	No analgesia		No analgesia	

* $p < 0.05$; ** $p < 0.01$ (KO versus WT; Student's t test).

analgesic activity of oxotremorine was significantly reduced (by about 50–90%) in M_2 KO mice in both assay systems used (Fig. 5A), as observed after s.c. administration (see above). On the other hand, oxotremorine-induced antinociceptive responses were not significantly affected in M_4 KO mice, independent of the route of application (i.t. or i.c.v.; Fig. 5B). In M_2/M_4 receptor double-KO mice, oxotremorine, administered i.t. or i.c.v., was completely devoid of analgesic activity (Fig. 5C). The very small residual responses seen in Fig. 5C were not significantly different from the corresponding preinjection values (data not shown).

As observed with oxotremorine, i.t. or i.c.v. administration of CMI-936 and CMI-1145 resulted in a significant reduction in the magnitude of analgesic responses (by ~40–60%) in M_2 KO mice in both the tail-flick and hot plate tests (Figs. 6A and 7A). However, in contrast to oxotremorine, the two epibatidine derivatives, when administered i.t. or i.c.v., also showed pronounced reductions in analgesic activity (by ~50–80%) in M_4 KO mice in both assays used (Figs. 6B and 7B). As found with oxotremorine, the analgesic responses to i.t. or i.c.v. CMI-936 and CMI-1145 were totally abolished in M_2/M_4 double-KO mice (Figs. 6C and 7C).

To exclude the possibility that the use of pentobarbital as the anesthetic agent had major effects on the outcome of the i.c.v. and i.t. injection experiments, we repeated several key experiments using inhalational anesthesia (isoflurane/oxygen; 4:96). Specifically, we assessed the analgesic effects of oxotremorine, administered i.t. or i.c.v. (10 μ g per mouse), in M_2 single-KO and M_2/M_4 double-KO mice and the two corre-

sponding wild-type control groups ($n = 5$ /group), using isoflurane/oxygen as the anesthetizing agent. These experiments showed that the analgesic activity of oxotremorine was significantly reduced (~65–75% reduction in MPE) in M_2 KO mice in both tail-flick and hot plate tests. The following responses were obtained (WT versus M_2 single KO): a) tail-flick, i.c.v., $96.5 \pm 5.2/36.6 \pm 12.2\%$ MPE; i.t., $91.6 \pm 10.2/24.3 \pm 11.1\%$ MPE; b) hot plate, i.c.v., $100 \pm 0/35.0 \pm 9.3\%$ MPE; i.t., $87.2 \pm 9.4/26.0 \pm 15.2\%$ MPE. In M_2/M_4 double-KO mice, oxotremorine showed only residual analgesic responses




Fig. 3. Analgesic responses of wild-type and mACHR KO mice to systemic (s.c.) administration of CMI-936. A–C, tail-flick assay. D–F, hot plate assay. M_2 KO (A, D), M_4 KO (B, E), and M_2/M_4 double-KO mice (C, F) and the corresponding wild-type (WT) control mice ($n = 5$ –6/group) were injected s.c. with increasing doses of CMI-936, using a cumulative dose-response protocol. Data are plotted as cumulative CMI-936 dose versus percent mice displaying full analgesia, as described under *Materials and Methods*. The dose-response curves shown are representative of two independent experiments that gave similar results.




Fig. 4. Antinociceptive responses of wild-type and mACHR KO mice after systemic (s.c.) administration of CMI-1145. A–C, tail-flick assay. D–F, hot plate assay. M_2 KO (A, D), M_4 KO (B, E), and M_2/M_4 double-KO mice (C, F) and the corresponding wild-type (WT) control mice ($n = 5$ –6/group) were injected s.c. with increasing doses of CMI-1145, using a cumulative dose-response protocol. Data are plotted as cumulative CMI-1145 dose versus percent mice displaying full analgesia, as described under *Materials and Methods*. The dose-response curves shown are representative of two independent experiments which gave similar results.




Fig. 5. Antinociceptive responses of wild-type and mACHR KO mice after i.t. or i.c.v. administration of oxotremorine. A, M_2 KO and wild-type control mice. B, M_4 KO and wild-type control mice. C, M_2/M_4 double-KO and wild-type control mice. Mice ($n = 5$ –6/group) were injected i.t. or i.c.v. with a single dose (10 μ g/mouse) of oxotremorine, and analgesic effects were determined using the tail-flick (TF) and hot plate (HP) assays. Antinociceptive effects were expressed as %MPE as described under *Materials and Methods*. Two independent experiments gave similar results. Data are plotted as means \pm S.D. (*, $p < 0.05$).

(reduction of MPE by $\geq 90\%$) which were not significantly different from the corresponding preinjection values (data not shown). The following responses were obtained (WT versus M_2/M_4 double-KO): a) tail-flick, i.c.v., $100 \pm 0/7.3 \pm 6.9\%$ MPE; i.t., $96.8 \pm 12.2/8.6 \pm 5.7\%$ MPE. b) hot plate, i.c.v., $100 \pm 0/10.0 \pm 4.6\%$ MPE; i.t., $95.7 \pm 11.4/6.9 \pm 8.0\%$ MPE. These results were similar to those obtained in studies in which pentobarbital was used as the anesthetic agent (Fig. 5), indicating that the key findings obtained in the i.c.v. and i.t. injection experiments were independent of the nature of the anesthetic agent used (pentobarbital versus isoflurane/oxygen).

Agonist Affinities for Cloned M_2 and M_4 mAChRs. Unlike oxotremorine, CMI-936 and CMI-1145 showed decreased analgesic effects not only in M_2 KO but also in M_4 KO mice, perhaps because of preferential binding of the CMI agents to M_4 receptors. To test this hypothesis, we determined the binding affinities of oxotremorine, CMI-936, and CMI-1145 for human M_2 and M_4 mAChRs individually expressed in stable CHO-K1 cell lines (Dörje et al., 1991). This analysis showed that oxotremorine displayed similar affinities for M_2 and M_4 receptors (Fig. 8A; Table 2). In contrast, CMI-936 and CMI-1145 displayed markedly higher affinities




Fig. 6. Analgesic responses of wild-type and mAChR KO mice after i.t. or i.c.v. administration of CMI-936. A, M_2 KO and wild-type control mice. B, M_4 KO and wild-type control mice. C, M_2/M_4 double-KO and wild-type control mice. Mice ($n = 5-6$ /group) were injected i.t. or i.c.v. with a single dose ($20 \mu\text{g}/\text{mouse}$) of CMI-936, and analgesic effects were determined using the tail-flick (TF) and hot plate (HP) assays. Antinociceptive effects were expressed as %MPE, as described under *Materials and Methods*. Two independent experiments gave similar results. Data are plotted as means \pm S.D. (*, $p < 0.05$).




Fig. 7. Analgesic responses of wild-type and mAChR KO mice to i.t. or i.c.v. administration of CMI-1145. A, M_2 KO and wild-type control mice. B, M_4 KO and wild-type control mice. C, M_2/M_4 double-KO and wild-type control mice. Mice ($n = 5-6$ /group) were injected i.t. or i.c.v. with a single dose ($20 \mu\text{g}/\text{mouse}$) of CMI-1145, and analgesic effects were determined using the tail-flick (TF) and hot plate (HP) assays. Antinociceptive effects were expressed as %MPE as described under *Materials and Methods*. Two independent experiments gave similar results. Data are plotted as means \pm S.D. (*, $p < 0.05$).

(15.5- and 6.5-fold, respectively) for M_4 than for M_2 receptors (Fig. 8, B and C; Table 2).

[^3H]NMS Binding Studies with Spinal Cord Tissue. The results of the tail-flick and i.t. injection experiments indicated that spinal cord M_2 and M_4 mAChRs play important roles in muscarinic agonist-induced analgesic responses. To quantitate spinal M_2 and M_4 receptor densities, membrane homogenates prepared from mouse whole spinal cord were incubated with a saturating concentration (2 nM) of the nonselective muscarinic antagonist [^3H]NMS. As shown in Fig. 9, spinal cord preparations from M_2 KO and M_2/M_4 double-KO mice showed a striking reduction (by about 90%) in the number of specific [^3H]NMS binding sites, compared with preparations from the corresponding wild-type strains. On the other hand, the number of [^3H]NMS binding sites detectable in spinal cord tissue from M_4 KO mice did not differ significantly from the corresponding wild-type value (Fig. 9). These observations clearly indicate that the M_2 subtype represents the predominant mAChR species in mouse spinal cord. In contrast, the levels of spinal cord M_4 receptors seem to be too low ($<10\%$ of the total mAChR population in mouse spinal cord) to be reliably detectable in [^3H]NMS binding assays.

Immunohistochemical Analysis of mAChR Expression in Mouse Spinal Cord. To complement the [^3H]NMS binding assays, we also carried out immunohistochemical studies to map the anatomical localization of M_2 and, if possible, M_4 receptors in the mouse spinal cord. As shown in




Fig. 8. Displacement of specific [^3H]NMS binding to cloned M_2 and M_4 mAChRs by the three muscarinic agonists used in this study. A, oxotremorine. B, CMI-936. C, CMI-1145. Competition binding assays were carried out with membranes prepared from CHO-K1 cell lines stably expressing the cloned human M_2 and M_4 mAChRs, using 50 pM of [^3H]NMS (for details, see *Materials and Methods*). The specific binding determined in the absence of agonists was defined as 100%. Data are presented as means \pm S.D. from two or three separate experiments carried out in duplicate.

TABLE 2

Binding affinities of oxotremorine, CMI-936, and CMI-1145 for the M_2 and M_4 mAChRs

Binding affinities (K_i values) were determined in [^3H]NMS competition binding assays as described under *Materials and Methods*, using membranes prepared from CHO-K1 cell lines stably expressing the human M_2 or M_4 mAChRs (Dörje et al., 1991). [^3H]NMS was used at a concentration of 50 pM . [^3H]NMS saturation binding studies yielded the following [^3H]NMS affinities (K_d values): M_2 , $545 \pm 82 \text{ pM}$; M_4 , $160 \pm 6 \text{ pM}$ (means \pm S.D.; $n = 3$). IC_{50} values were converted to K_i values by using the Cheng-Prusoff equation (Cheng and Prusoff, 1973). Data are presented as means \pm S.D. from two or three separate experiments carried out in duplicate.

	K_i Values		$K_i\text{-}M_2/K_i\text{-}M_4$
	M_2	M_4	
	μM		
Oxotremorine	0.32 ± 0.08	0.51 ± 0.18	0.63
CMI-936	10.1 ± 1.4	$0.65 \pm 0.10^*$	15.5
CMI-1145	5.2 ± 3.1	$0.80 \pm 0.08^*$	6.5

* Significantly different from the corresponding M_2 receptor value ($p < 0.05$, t test).

Fig. 10, use of an M_2 receptor-specific rat monoclonal antibody (Levey et al., 1995) indicated that M_2 receptors were widely expressed in the spinal cord of wild-type and M_4 single-KO mice. M_2 receptor-specific fibers and puncta were located in neuropil throughout the gray matter, but staining was most intense in lamina II of the dorsal horn (substantia gelatinosa). In lamina IX of the ventral horn, large M_2 receptor-specific puncta covered the outer surface of motor neurons. The plasma membrane of these neurons was also immunoreactive. As expected, spinal cord preparations derived from M_2 KO and M_2/M_4 double-KO mice gave only faint background staining, confirming the specificity of the M_2 receptor antibody (Fig. 10). In contrast, use of an M_4 receptor-specific rabbit polyclonal antibody (Levey et al., 1991), which readily detected M_4 receptors in mouse striatum and other forebrain regions (Hohmann et al., 1995), did not reproducibly lead to M_4 receptor-specific staining of mouse spinal cord tissue (data not shown). In agreement with the results of the [3 H]NMS binding assays (see previous paragraph), this observation indicates that M_4 receptors are expressed at very low levels in the mouse spinal cord.

Discussion

To assess the roles of the M_2 and M_4 mAChRs in muscarinic agonist-induced analgesia, we studied the antinociceptive effects of several centrally active muscarinic agonists in M_2 KO, M_4 KO, and M_2/M_4 double-KO mice. Analgesia measurements were carried out using the tail-flick and hot plate assays. Strikingly, the nonselective muscarinic agonist oxotremorine when administered s.c., was completely devoid of analgesic activity in M_2/M_4 double-KO mice in both assays. As reported previously (Gomez et al., 1999a), the analgesic potency of oxotremorine was significantly reduced in M_2 single-KO mice (Fig. 2, A and D). However, maximum analgesia could still be elicited in M_2 KO mice by increased doses of oxotremorine (Fig. 2, A and D), indicating that non- M_2 mAChRs can also mediate profound antinociception. The

complete absence of muscarinic agonist-induced antinociception in M_2/M_4 double-KO mice indicates that both M_2 and M_4 mAChRs are involved in mediating muscarinic analgesia and that non- M_2/M_4 mAChRs do not contribute to this activity. As observed by Gomez et al. (1999b), oxotremorine showed similar analgesic potencies in M_4 single KO and wild-type mice (Fig. 2, B and E; Table 1). This observation is consistent with the concept that oxotremorine-induced analgesia preferentially involves M_2 mAChRs and that the presence of the intact M_2 receptor signaling pathway in the M_4 KO mice is sufficient to trigger potent analgesic effects.

To examine whether the results obtained with oxotremorine are also applicable to other classes of muscarinic agonists, we carried out analogous studies with two novel muscarinic agonists, CMI-936 and CMI-1145 (Ellis et al., 1999), which are oxadiazole derivatives of the nicotinic receptor agonist, epibatidine. When injected s.c. into mice, both compounds displayed robust analgesic effects that could be partially blocked by i.t. administration of the snake toxin, MT-3 (Ellis et al., 1999). Because the MT-3 toxin displays at least 40-fold higher affinity for M_4 than for the other mAChR subtypes (Jolkkonen et al., 1994; Liang et al., 1996), this observation suggested (but did not prove) that CMI-936 and CMI-1145 might be endowed with a receptor selectivity profile different from that of oxotremorine. In the present study, we therefore carried out radioligand binding studies to determine the affinities of CMI-936 and CMI-1145 for the cloned M_2 and M_4 mAChRs (Dörje et al., 1991). In contrast to oxotremorine, which displayed little receptor subtype preference, CMI-936 and CMI-1145 bound to M_4 receptors with markedly higher affinities (15.5 and 6.5-fold, respectively) than to M_2 receptors (Fig. 8, Table 2). Although the degree of M_4 selectivity displayed by these agents is relatively modest, the two CMI compounds nevertheless are important novel pharmacological tools. It should be noted in this context that another muscarinic agonist, McN-A-343, has also been reported to exhibit a certain degree of M_4 receptor selectivity (Lazareno et al., 1993; Richards and van Giersbergen, 1995).

In contrast to oxotremorine, which displayed a significant reduction in analgesic potency only in the M_2 single-KO mice, CMI-936 and CMI-1145, when given s.c., showed reduced analgesic potencies (by about 2- to 6-fold) in both M_2 and M_4




Fig. 9. Number of [3 H]NMS binding sites in spinal cord tissue from wild-type and mAChR KO mice. Membranes prepared from whole spinal cord of M_2 KO, M_4 KO, and M_2/M_4 double-KO mice and their corresponding wild-type (WT) control mice were incubated with a saturating concentration (2 nM) of the nonselective muscarinic antagonist, [3 H]NMS, as described under *Materials and Methods*. Data are presented as means \pm S.D. from three separate experiments carried out in triplicate (*, $P < 0.05$).




Fig. 10. Immunohistochemical localization of the M_2 mAChR in mouse spinal cord. Slices of mouse cervical spinal cord were incubated with an M_2 receptor-specific rat monoclonal antibody (Levey et al., 1995) and processed for immuno-peroxidase staining as described under *Materials and Methods* (Levey et al., 1995). In preparations from wild-type (WT) and M_4 KO mice, M_2 receptor-specific staining was detected throughout the dorsal and ventral gray matter. M_2 receptor immunoreactivity was particularly rich in the substantia gelatinosa of the dorsal horn (SG) and the outer surface of motor neurons (MN) of lamina IX of the ventral horn. As expected, only background staining was observed in samples derived from M_2 KO and M_2/M_4 double-KO mice.

single-KO mice (Figs. 3 and 4, Table 1), probably reflecting their ability to preferentially bind to M_4 receptors. However, like oxotremorine, the two CMI compounds were completely devoid of analgesic activity in the M_2/M_4 double-KO mice (Figs. 3, C and F, and 4, C and F). These data confirm that both M_2 and M_4 receptors participate in muscarinic agonist-mediated analgesic responses and indicate that the receptor selectivity profile of the specific muscarinic agonist under investigation determines to which extent the two receptors contribute to the observed analgesia response.

To study the relative contribution of spinal and supraspinal pathways to muscarinic agonist-dependent antinociception, we carried out analogous analgesia measurements after i.t. or i.c.v. administration of oxotremorine, CMI-936, and CMI-1145. The results of these studies closely mirrored those obtained after systemic (s.c.) administration of these three agonists. Oxotremorine, when given i.t. or i.c.v. (10 $\mu\text{g}/\text{mouse}$), showed a significantly reduced analgesic response (by $\sim 50\text{--}90\%$) in M_2 KO mice but retained full analgesic activity in M_4 KO mice (Fig. 5, A and B). In contrast, the two CMI compounds, administered i.t. or i.c.v. (20 $\mu\text{g}/\text{mouse}$), showed a significant decrease in analgesic activity (by $\sim 40\text{--}80\%$) in both M_2 and M_4 single-KO mice (Figs. 6, A and B, and 7, A and B). As observed after systemic administration, all three agonists lacked significant analgesic activity in M_2/M_4 double-KO mice when administered i.t. or i.c.v. (Figs. 5C, 6C, and 7C). These findings indicate that both M_2 and M_4 receptors participate in mediating spinal as well as supraspinal analgesic effects and that other mAChRs are not involved in this activity. However, our data do not completely rule out the possibility that stimulation of M_1 , M_3 , or M_5 mAChRs can also reduce pain sensitivity (at least theoretically) by activating a cholinergic pathway which ultimately leads to stimulation of M_2 and/or M_4 receptors.

[^3H]NMS radioligand binding studies with spinal cord preparations from wild-type, M_2 KO, M_4 KO, or M_2/M_4 double-KO mice showed that M_2 receptors represent the majority ($\sim 90\%$) of spinal cord mAChRs (Fig. 9). Immunohistochemical studies revealed that M_2 receptors are abundantly expressed in lamina II of the dorsal horn (Fig. 10), a region in which the primary nociceptive afferent fibers (A δ and C fibers) terminate (Yaksh, 1988; Gillberg et al., 1990; Levine, 1998; Snider and McMahon, 1998, 1999). This observation is consistent with immunohistochemical and radioligand binding studies carried out with rat (Höglund and Baghdoyan, 1997; Yung and Lo, 1997) and human (Potter et al., 1996) tissues demonstrating high levels of M_2 receptor expression in the superficial layers of the dorsal horn. In both [^3H]NMS radioligand binding and immunohistochemical studies, we were unable to reproducibly detect M_4 receptors in mouse spinal cord tissue, suggesting that spinal M_4 receptors are expressed at very low abundance. In agreement with this observation, pharmacological studies indicate that M_4 receptors represent only a minor fraction of the total mAChR population in rat (Höglund and Baghdoyan, 1997) and human (Potter et al., 1996) spinal cord. The low abundance of M_4 receptors in mouse spinal cord may explain why the nonselective muscarinic agonist, oxotremorine, displayed virtually unchanged analgesic activity in M_4 single-KO mice (see discussion above). Analogously, the high density of spinal M_2 receptors is likely to be responsible for the very pronounced reduction in analgesic potency of systemically

administered oxotremorine observed with M_2 single-KO mice in the tail-flick experiments (Fig. 2, A and Table 1).

Lesion studies suggest that mAChRs in the spinal dorsal horn are localized to the nerve terminals of primary afferent pain fibers (Gillberg and Wiksten, 1986). In agreement with this observation, light and electron microscopic studies have detected M_2 and M_4 receptor immunostaining in small type neurons of rat dorsal ganglia believed to be involved in the transmission of pain stimuli to the spinal cord (Bernardini et al., 1999; Haberberger et al., 1999). In addition, electrophysiological experiments (Bleazard and Morris, 1993), combined with electron microscopic studies (Ribeiro da Silva and Cuello, 1990), strongly suggest that presynaptic mAChRs present on pain fibers terminating in lamina II of the dorsal horn function to inhibit the release of excitatory neurotransmitters required for the efficient propagation of pain signals. However, electrophysiological studies also indicate that inhibitory postsynaptic mAChRs located on spinal dorsal horn projection neurons (Urban et al., 1989) or excitatory mAChRs present on spinal GABA-ergic interneurons (Baba et al., 1998) may also contribute to the inhibition of spinal nociceptive pathways. Taken together, these studies suggest that both presynaptic and postsynaptic mechanisms contribute to the antinociceptive effects mediated by spinal M_2 and/or M_4 mAChRs. It has been proposed that spinal μ and δ opioid and α_2 adrenergic receptors mediate their analgesic effects through similar mechanisms (Siddall and Cousins, 1995; Yaksh, 1999).

The distribution of supraspinal (brain) M_2 and M_4 mAChRs has been mapped in great detail in previous studies (see, for example, Levey et al., 1991; Levey, 1993; Hohmann et al., 1995; Gomeza et al., 1999a,b). These studies have shown that M_2 and M_4 receptors are widely expressed throughout the brain, including all major regions predicted to be involved in pain transmission, modulation, and perception (pons/medulla, midbrain, thalamus, cerebral cortex, etc.).

Although spinal M_4 receptors are predicted to be expressed at much lower levels than spinal M_2 receptors, our results suggest that this rather small population of M_4 receptors nevertheless can mediate pronounced analgesic effects. M_4 receptors, unlike M_2 receptors, which are widely expressed both in the body periphery and in the CNS, are primarily expressed in the brain (Levey et al., 1991; Levey, 1993; Vilario et al., 1993; Yasuda et al., 1993; Wolfe and Yasuda, 1995; Gomeza et al., 1999a,b). Thus, from a clinical point of view, the development of selective M_4 receptor agonists as novel analgesic drugs seems particularly attractive. Such agents are predicted to lack significant muscarinic side effects, because the clinically most bothersome unwanted effects of muscarinic agonists, such as changes in heart rate, dry mouth, and impaired smooth muscle activity, are predicted to be mediated primarily by M_2 and M_3 mAChRs (Caulfield, 1993; Wess, 1996; Eglén et al., 1999). In addition, several lines of evidence (Widman et al., 1985; Swedberg et al., 1997; Petry et al., 1998) suggest that the use of muscarinic agonists for the management of chronic pain may be less likely to lead to tolerance, addiction, and physical dependence, severe side effects associated with the use of classic opioid analgesics. As discussed above, the two epibatidine derivatives, CMI-936 and CMI-1145, showed a moderate degree of M_4 receptor selectivity (Table 2). Systematic structural modification of

these or other lead compounds should eventually lead to more selective M₄ receptor agonists suitable for clinical use.

Acknowledgments

We thank H. Rees for expert technical assistance.

References

- Adams JU, Paronis CA, and Holtzman SG (1990) Assessment of relative intrinsic activity of μ -opioid analgesics in vivo by using β -funtaltrexamine. *J Pharmacol Exp Ther* **255**:1027–1032.
- Baba H, Kohno T, Okamoto M, Goldstein PA, Shimoji K, and Yoshimura M (1998) Muscarinic facilitation of GABA release in substantia gelatinosa of the rat spinal dorsal horn. *J Physiol* **508**:83–93.
- Bartolini A, Ghelardini C, Fantetti L, Malcangio M, Malmberg-Aiello P, and Giotti A (1992) Role of muscarinic receptor subtypes in central antinociception. *Br J Pharmacol* **105**:77–82.
- Bernardini N, Levey AI, and Augusti-Tocco G (1999) Rat dorsal root ganglia express m1–m4 muscarinic receptor proteins. *J Peripher Nerv Syst* **4**:222–232.
- Bleazard L and Morris R (1993) The effects of cholinergic agonists and antagonists on C-fibre evoked responses in the substantia gelatinosa of neonatal rat spinal cord slices. *Br J Pharmacol* **110**:1061–1066.
- Caulfield MP (1993) Muscarinic receptors - characterization, coupling and function. *Pharmacol Ther* **58**:319–379.
- Cheng Y and Prusoff WH (1973) Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 per cent inhibition (I_{50}) of an enzymatic reaction. *Biochem Pharmacol* **22**:3099–3108.
- D'Amour FE and Smith DL (1941) A method for determining loss of pain sensation. *J Pharmacol Exp Ther* **72**:74–79.
- Dörje F, Wess J, Lambrecht G, Tacke R, Mutschler E, and Brann MR (1991) Antagonist binding profiles of five cloned human muscarinic receptor subtypes. *J Pharmacol Exp Ther* **256**:727–733.
- Duttaroy A, Kirtman R, Farrell F, Phillips M, Philippe J, Monderson T, and Yoburn BC (1997) The effect of cumulative dosing on the analgesic potency of morphine in mice. *Pharmacol Biochem Behav* **58**:67–71.
- Eglen RM, Choppin A, Dillon MP, and Hegde S (1999) Muscarinic receptor ligands and their therapeutic potential. *Curr Opin Chem Biol* **3**:426–432.
- Eisenach JC (1999) Muscarinic-mediated analgesia. *Life Sci* **64**:549–554.
- Ellis JL, Harman D, Gonzalez J, Spera ML, Liu R, Shen TY, Wypij DM, and Zuo F (1999) Development of muscarinic analgesics derived from epibatidine: role of the M4 receptor subtype. *J Pharmacol Exp Ther* **288**:1143–1150.
- Ghelardini C, Galeotti N, and Bartolini A (2000) Loss of muscarinic antinociception by antisense inhibition of M₁ receptors. *Br J Pharmacol* **129**:1633–1640.
- Gillberg PG, Askmark H, and Aquilonius S-M (1990) Spinal cholinergic mechanisms. *Prog Brain Res* **84**:361–370.
- Gillberg PG, Gordh T Jr, Hartvig P, Jansson I, Pettersson J, and Post C (1989) Characterization of the antinociception induced by intrathecally administered carbachol. *Pharmacol Toxicol* **64**:340–343.
- Gillberg PG and Wiksten B (1986) Effects of spinal cord lesions and rhizotomies on cholinergic and opiate receptor binding sites in rat spinal cord. *Acta Physiol Scand* **126**:575–582.
- Gomez J, Shannon H, Kostenis E, Felder C, Zhang L, Brodtkin J, Grinberg A, Sheng H, and Wess J (1999a) Pronounced pharmacologic deficits in M2 muscarinic acetylcholine receptor knockout mice. *Proc Natl Acad Sci USA* **96**:1692–1697.
- Gomez J, Zhang L, Kostenis E, Felder C, Bymaster F, Brodtkin J, Shannon H, Xia B, Deng C, and Wess J (1999b) Enhancement of D1 dopamine receptor-mediated locomotor stimulation in M4 muscarinic acetylcholine receptor knockout mice. *Proc Natl Acad Sci USA* **96**:10483–10488.
- Green PG and Kitchen I (1986) Antinociception opioids and the cholinergic system. *Prog Neurobiol* **26**:119–146.
- Haberberger R, Henrich M, Couraud JY, and Kummer W (1999) Muscarinic M2-receptors in rat thoracic dorsal root ganglia. *Neurosci Lett* **266**:177–180.
- Haley TJ and McCormick WG (1957) Pharmacological effects produced by intracerebral injection of drugs in the conscious mouse. *Br J Pharmacol* **12**:12–15.
- Hamilton SE, Loose MD, Qi M, Levey AI, Hille B, McKnight GS, Idzerda RL, and Nathanson NM (1997) Disruption of the m1 receptor gene ablates muscarinic receptor-dependent M current regulation and seizure activity in mice. *Proc Natl Acad Sci USA* **94**:13311–13316.
- Harris LS, Dewey WL, Howes JF, Kennedy JS, and Pars H (1969) Narcotic-antagonist analgesics: interactions with cholinergic systems. *J Pharmacol Exp Ther* **169**:17–22.
- Hartvig P, Gillberg PG, Gordh T Jr, and Post C (1989) Cholinergic mechanisms in pain and analgesia. *Trends Pharmacol Sci* **10**(Suppl):75–79.
- Höglund AU and Baghdoyan HA (1997) M2, M3 and M4, but not M1, muscarinic receptor subtypes are present in rat spinal cord. *J Pharmacol Exp Ther* **281**:470–477.
- Hohmann CF, Potter ED, and Levey AI (1995) Development of muscarinic receptor subtypes in the forebrain of the mouse. *J Comp Neurol* **358**:88–101.
- Honda K, Harada A, Takano Y, and Kamiya H (2000) Involvement of M₃ muscarinic receptors of the spinal cord in formalin-induced nociception in mice. *Brain Res* **859**:38–44.
- Howes JF, Harris LS, Dewey WL, and Voyda CA (1969) Brain acetylcholine levels and inhibition of the tail-flick reflex in mice. *J Pharmacol Exp Ther* **169**:23–28.
- Hylden JKL and Wilcox GL (1980) Intrathecal morphine in mice: a new technique. *Eur J Pharmacol* **67**:313–316.
- Iwamoto ET and Marion L (1993) Characterization of the antinociception produced by intrathecally administered muscarinic agonists in rats. *J Pharmacol Exp Ther* **266**:329–338.
- Jolkkonen M, van Giersbergen PL, Hellman U, Wernstedt C, and Karlsson E (1994) A toxin from the green mamba *Dendroaspis angusticeps*: amino acid sequence and selectivity for muscarinic m4 receptors. *FEBS Lett* **352**:91–94.
- Lazareno S, Farries T, and Birdsall NJM (1993) Pharmacological characterization of guanine nucleotide exchange reactions in membranes from CHO cells stably transfected with human muscarinic receptors m1–m4. *Life Sci* **52**:449–456.
- Levey AI (1993) Immunological localization of m1–m5 muscarinic acetylcholine receptors in peripheral tissues and brain. *Life Sci* **52**:441–448.
- Levey AI, Edmunds SM, Hersch SM, Wiley RG, and Heilman CJ (1995) Light and electron microscopic study of m2 muscarinic acetylcholine receptor in the basal forebrain of the rat. *J Comp Neurol* **351**:339–356.
- Levey AI, Kitt CA, Simonds WF, Price DL, and Brann MR (1991) Identification and localization of muscarinic acetylcholine receptor proteins in brain with subtype-specific antibodies. *J Neurosci* **11**:3218–3226.
- Levine JD (1998) New directions in pain research: molecules to maladies. *Neuron* **20**:649–654.
- Liang JS, Carsi-Gabrenas J, Krajewski JL, McCafferty JM, Purkerson SL, Santiago MP, Strauss WL, Valentine HH, and Potter LT (1996) Anti-muscarinic toxins from *Dendroaspis angusticeps*. *Toxicon* **34**:1257–1267.
- Ma HC, Dohi S, Wang YF, Ishizawa Y, and Yanagitate F (2001) The antinociceptive and sedative effects of carbachol and oxydodone administered into brainstem pontine reticular formation and spinal subarachnoid space in rats. *Anesth Analg* **92**:1307–1315.
- Matsui M, Motomura D, Karasawa H, Fujikawa T, Jiang J, Komiya Y, Takahashi S, and Taketo MM (2000) Multiple functional defects in peripheral autonomic organs in mice lacking muscarinic acetylcholine receptor gene for the M₃ subtype. *Proc Natl Acad Sci USA* **97**:9579–9584.
- Naguib M and Yaksh TL (1997) Characterization of muscarinic receptor subtypes that mediate antinociception in the rat spinal cord. *Anesth Analg* **85**:847–853.
- Pert A (1987) Cholinergic and catecholaminergic modulation of nociceptive reactions. Interactions with opiates. *Pain Headache* **9**:1–63.
- Petry NM, Bickel WK, Huddleston J, Tzanis E, and Badger GJ (1998) A comparison of subjective, psychomotor and physiological effects of a novel muscarinic analgesic, LY297802 tartrate and oral morphine in occasional drug users. *Drug Alcohol Depend* **50**:129–136.
- Potter LT, Borenstein DW, and Levey AI (1996) Muscarinic receptors in the superficial dorsal horn of the human, rat and mouse spinal cord (Abstract). *Life Sci* **64**:585.
- Ribeiro da Silva A and Cuello AC (1990) Choline acetyltransferase-immunoreactive profiles are presynaptic to primary sensory fibers in the rat superficial dorsal horn. *J Comp Neurol* **295**:370–384.
- Richards MH and van Giersbergen PLM (1995) Human muscarinic receptors in A9L and CHO cells: activation by full and partial agonists. *Br J Pharmacol* **114**:1241–1249.
- Siddall PJ and Cousins MJ (1995) Pain mechanisms and management: an update. *Clin Exp Pharm Physiol* **22**:679–688.
- Shannon HE, Womer DE, Bymaster FP, Calligaro DO, DeLapp NC, Mitch CH, Ward JS, Whitesitt CA, Swedberg MD, Sheardown MJ, et al. (1997) *In vivo* pharmacology of butylthio[2.2.2] (LY297802)/NMC11–1053, an orally acting antinociceptive muscarinic agonist. *Life Sci* **60**:969–976.
- Snider WD and McMahon SB (1998) Tackling pain at the source: new ideas about nociceptors. *Neuron* **20**:629–632.
- Swedberg MD, Sheardown MJ, Sauerberg P, Olesen PH, Suzdak PD, Hansen KT, Bymaster FP, Ward JS, Mitch CH, Calligaro DO, et al. (1997) Butylthio[2.2.2] (NMC 11–1053/LY297802): an orally active muscarinic agonist analgesic. *J Pharmacol Exp Ther* **281**:876–883.
- Urban L, Willetts J, Murase K, and Randic M (1989) Cholinergic effects on spinal dorsal horn neurons in vitro: an intracellular study. *Brain Res* **500**:12–20.
- Vilaro MT, Mengod G, and Palacios JM (1993) Adv and limitations of the molecular neuroanatomy of cholinergic receptors: the example of multiple muscarinic receptors. *Prog Brain Res* **98**:95–101.
- Wei J, Walton EA, Milici A, and Buccafusco JJ (1994) m1–m5 muscarinic receptor distribution in rat CNS by RT-PCR and HPLC. *J Neurochem* **63**:815–821.
- Wenger GR (1980) Cumulative dose-response curves in behavioral pharmacology. *Pharmacol Biochem Behav* **49**:647–651.
- Wess J (1996) Molecular biology of muscarinic acetylcholine receptors. *Crit Rev Neurobiol* **10**:69–99.
- Widman M, Tucker S, Brase DA, and Dewey WL (1985) Cholinergic agents: antinociception without morphine type dependence in rats. *Life Sci* **36**:2007–2015.
- Wolfe BB and Yasuda RP (1995) Development of selective antisera for muscarinic cholinergic receptor subtypes. *Ann NY Acad Sci* **757**:186–193.
- Woolfe G and Macdonald AD (1944) The evaluation of analgesic action of pethidine hydrochloride (demerol). *J Pharmacol Exp Ther* **80**:300–307.
- Yaksh TL (1988) CNS mechanisms of pain and analgesia. *Cancer Surv* **7**:5–28.
- Yaksh TL (1999) Spinal systems and pain processing: development of novel analgesic drugs with mechanistically defined models. *Trends Pharmacol Sci* **20**:329–337.
- Yamada M, Miyakawa T, Duttaroy A, Yamanaka A, Moriguchi T, Makita R, Ogawa M, Chou CJ, Xia B, Crawley JN, et al. (2001) Mice lacking the M3 muscarinic acetylcholine receptor are hypophagic and lean. *Nature (Lond)* **410**:207–212.
- Yasuda RP, Ciesla W, Flores LR, Wall SJ, Li M, Satkus SA, Weisstein JS, Spagnola BV, and Wolfe BB (1993) Development of antisera selective for m4 and m5 muscarinic cholinergic receptors: distribution of m4 and m5 receptors in rat brain. *Mol Pharmacol* **43**:149–157.
- Yung KKL and Lo YL (1997) Immunocytochemical localization of muscarinic m2 receptor in the rat spinal cord. *Neurosci Lett* **229**:81–84.

Address correspondence to: Dr. Jürgen Wess, Chief, Molecular Signaling Section, Lab. of Bioorganic Chemistry, NIH-NIDDK, Bldg. 8A, Room B1A-05, 8 Center Drive, MSC 0810, Bethesda, MD 20892-0810. E-mail: jwess@helix.nih.gov